

Another Physics and Route E: First-Principles Separation Theorems, an Exact Discriminant–Killing Bridge, and a Reversible Research Ledger

Route F research note

17 July 2026

Abstract

This ledger asks which parts of the manuscript *Another physics* can be connected to Route E without confusing an algebraic resemblance with a physical derivation. The literal identification of local energy density with time, phase-dependent disappearance of invariant mass, fundamental negative energy, and indestructible entanglement is ruled out under ordinary local, positive-energy relativistic quantum theory. A conditional complex-scalar completion does support finite-charge Q-balls, but its internal phase is not itself an observable clock.

The strongest exact connection is new but elementary. On Route E’s $H3^+$ -conditional branch, $H^0(\mathbb{CP}^1, T_{\mathbb{CP}^1}) \simeq H^0(\mathbb{CP}^1, \mathcal{O}(2)) \simeq \mathfrak{sl}_2(\mathbb{C})$. For a binary quadratic, its discriminant simultaneously detects two distinct zeros, a regular-semisimple Möbius flow, a nonzero Killing norm, and a nonzero Route-E second-transvectant contact. Thus the two-center divisor and K_{tr} are not merely parallel observations: they are two readings of the same invariant. A Fubini–Study moment map can further label the two fixed points by opposite bounded charges without introducing negative Hamiltonian energy.

A candidate action-level bridge makes the Route-E messenger mixing depend on a Q-ball field, $\lambda(\Phi) = g\Phi/\Lambda$, giving $\zeta_{\text{eff}} = g^2\Phi^2/\Lambda^2$ and hence the same weight-two phase law as Route E. This is explicitly a bridge axiom, not a promoted result: an exact Q-ball $U(1)$ makes every invariant observable stationary, while an observable phase modulation requires relative-phase breaking and therefore reopens charge leakage and soliton stability. All claims are classified, derived, and tied to a reproducible numerical ledger; no physics promotion is authorized by the present note.

Contents

1	Provenance, scope, and logical contract	3
2	Covariant dictionary and literal-theory boundary	3
2.1	Local density cannot determine clock rate	4
2.2	Mass, energy sign, and entanglement	4
2.3	Other immediate separation results	5
3	The Route-E theorem boundary imported into Route F	5
4	Exact discriminant–two-center–Killing theorem	6
4.1	Global vector fields and the adjoint representation	6
5	A safe two-polarity bridge: Hopf reduction and moment map	8

6	Conditional Q-ball completion of an energy bubble	8
6.1	Action, charge, and radial equation	8
6.2	Reproducible benchmark and approximation boundary	9
6.3	Exact-symmetry phase no-observable theorem	10
7	Route-E weight-two phase and a physical relative phase	10
8	Candidate Q-ball–Schur-complement bridge	11
8.1	Action-level construction	11
8.2	Why the portal is not yet a completion	12
9	Candidate dynamical origin of $H3^+$	13
10	The oscillatory formula: exact solution and strict separation	13
11	Relational time and gravity: parallel interfaces, not Route-E derivations	14
12	Claim ledger and ordered research program	15
13	2026-07-16 AP–E3 global/nonlinear and AP–E4 SQM handoff	16
13.1	Charged two-colour nonlinear proxy	16
13.2	Non-extendible mixed WZW and spin refinement	17
13.3	Pure- $SU(3)$ no-go and near-miss	17
13.4	Physical tangent fermion and the composability gate	17
14	2026-07-16 AP–E completion frontier	18
14.1	Four-dimensional action and graded $B = 1$ Hessian	18
14.2	Two torsion generators and regulator-dependent phases	19
14.3	A semisimple replacement for the diagonal quotient	19
14.4	Same-soliton SQM, Callias line, CPT, and WZW descent	20
14.5	Portal authorization theorem and current decision	21
15	AP-E6 same-background audit: exact obstructions and open gates	21
15.1	The exact $Sp(4)$ gauge-bordism result	21
15.2	One genuinely relaxed continuum $B=1$ field, but an off-shell cubic Hessian	23
15.3	The same canonical field and the physical-moduli obstruction	24
15.4	AP-E6 decision	25
16	AP-E7 regulator, discrete topology, and determinant-family audit	25
16.1	Common quadratic APS/PV complex and its exact scope	25
16.2	Finite-site topology and the re-relaxation result	27
16.3	The actual $SO(3)$ cover and the rank-three alternative	28
16.4	AP-E7 decision	29
17	Reproducibility ledger	30
A	Units and normalization checks	30
B	Why a phase portal needs a second reference	30

1 Provenance, scope, and logical contract

The source under review is the 15-page file `Another physics.pdf`, supplied from the local Downloads directory, with PDF metadata title `Another physics_20230910.docx` and SHA-256

7355cf2237513a6e943555ac2720871a5e81915fd770541baea9d87eccf2654e.

The Route-E reference state for this ledger is repository commit

e7ba020227853a3b731235b5f3a42083559e7ef2.

The governing Route-E sources are `route_E/tex/route_e_first_principles.tex`, `route_E/tex/three_families_killing_contact.tex`, and the canonical dynamics registry `route_E/code_dyn/dyn_claim_registry.json`.

Four logical labels are used throughout.

Label	Meaning
theorem	Consequence of explicitly stated mathematical or physical assumptions, with a proof in this ledger or an identified Route-E proof.
conditional completion	A consistent model after a new action, field content, or boundary condition has been chosen. It does not derive that choice from the source manuscript.
bridge axiom	An identification between objects belonging to two different models. Similar degree, phase, or dimension is not a proof of the identification.
numerical check	Reproducible arithmetic within a declared approximation. It corroborates formulas but cannot promote a bridge axiom.

Promotion gate 1.1 (Non-promotion). *The machine card may be green only with the non-promoting status `mechanical_separation_checks_pass_no_physics_promotion`. Its Boolean field `physics_promotion_allowed` must remain `false` until an action-level embedding, stability analysis, and all affected Route-E phenomenology gates pass.*

2 Covariant dictionary and literal-theory boundary

The source concepts are useful only after observer, sign, and conservation laws are made explicit. We use metric signature $(-+++)$. For an observer with unit timelike four-velocity u^μ , the local measured energy density is

$$E_d(x; u) := T_{\mu\nu}(x) u^\mu u^\nu. \quad (2.1)$$

This is a scalar after u is specified, but it is not an observer-independent replacement for spacetime or proper time. A viable information variable is instead a Noether charge, an entropy/correlation functional, or an order parameter. In the Q-ball completion below it is

$$E_i \rightsquigarrow Q_I = \int_\Sigma d\Sigma_\mu j_I^\mu, \quad \nabla_\mu j_I^\mu = 0. \quad (2.2)$$

The symbols E_p, E_n will later be reinterpreted as bounded moment-map polarities $\mu = +1, -1$, not as positive and negative Hamiltonian energies.

2.1 Local density cannot determine clock rate

Theorem 2.1 (Local-density time no-go). *No function of the pointwise scalar $E_d(x; u)$ alone can reproduce the gravitational clock rate in all solutions of general relativity.*

Proof. Consider a static thin spherical shell of mass M and radius $R > 2GM/c^2$. The vacuum interior is flat but its lapse is fixed by matching to the exterior:

$$ds_{r < R}^2 = - \left(1 - \frac{2GM}{Rc^2} \right) c^2 dt^2 + dr^2 + r^2 d\Omega^2. \quad (2.3)$$

Both the shell interior and spatial infinity have $T_{\mu\nu} = 0$, hence $E_d = 0$, but stationary clocks obey

$$\frac{d\tau_{\text{in}}}{dt_{\infty}} = \sqrt{1 - \frac{2GM}{Rc^2}} \neq 1. \quad (2.4)$$

Two points with the same local density therefore have different clock rates. The missing datum is the nonlocal metric solution/boundary condition, not a different definition of local energy density. $\square$

For Earth-scale numbers, $GM/(Rc^2) = 6.9613 \times 10^{-10}$, corresponding at leading order to about $60.15 \mu\text{s}$ per day. In the laboratory weak-field limit,

$$\frac{\Delta\nu}{\nu} = \frac{g\Delta h}{c^2} = 1.091 \times 10^{-18} \left(\frac{\Delta h}{1 \text{ cm}} \right). \quad (2.5)$$

This scale has been directly tested by a centimeter atomic-clock network [1] and is an immediate constraint on any replacement for metric proper time.

The same local E_d can also curve spacetime differently because pressure matters. Tracing

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (2.6)$$

and using $T = -\varepsilon + 3p$ gives

$$R = 4\Lambda + \frac{8\pi G}{c^4} (\varepsilon - 3p). \quad (2.7)$$

Dust and radiation may have the same ε , but $p = 0$ and $p = \varepsilon/3$ give different Ricci scalars. A density-only clock or gravity law therefore discards essential stress data.

2.2 Mass, energy sign, and entanglement

Theorem 2.2 (Phase-toggle no-go in a fixed particle representation). *For a stable one-particle irreducible representation of the Poincaré group, the rest mass cannot alternate between $m \neq 0$ and $m = 0$ as an unobserved internal phase changes while the state remains in that representation.*

Proof. The mass is fixed by the Casimir

$$P_\mu P^\mu = -m^2 c^2. \quad (2.8)$$

It commutes with the representation. If a phase trajectory changed the eigenvalue from $-m^2 c^2$ to zero, it would leave the massive irreducible representation and enter a massless one. A periodic coupling, spectral weight, quasiparticle gap, or detector response may vanish; invariant rest mass in a fixed representation may not. $\square$

Likewise, a literal independent negative-energy oscillator,

$$H = \omega a^\dagger a - \Omega b^\dagger b, \quad (2.9)$$

has $\inf \text{spec } H = -\infty$, since $\langle n_b | H | n_b \rangle = -n_b \Omega$. A stable completion must reinterpret the sign as charge, orientation, a hole relative to a filled state, or an effective-medium variable; it cannot be a freely excitable fundamental negative-energy sector.

Entanglement is not indestructible either. For

$$\rho_p = (1-p)|\Phi^+\rangle\langle\Phi^+| + \frac{p}{4}I_4, \quad |\Phi^+\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}}, \quad (2.10)$$

the partial transpose eigenvalues are

$$\text{spec}(\rho_p^{TB}) = \left\{ \frac{2-p}{4}, \frac{2-p}{4}, \frac{2-p}{4}, \frac{-2+3p}{4} \right\}. \quad (2.11)$$

For $p \geq 2/3$ the state is PPT and, in 2×2 , separable. What is forbidden is superluminal signalling, not dynamical loss of entanglement.

2.3 Other immediate separation results

For a massive electron of finite Lorentz factor γ ,

$$\beta := \frac{v}{c} = \sqrt{1 - \gamma^{-2}} < 1. \quad (2.12)$$

Magnetism is encoded in the gauge curvature $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, not in electrons moving exactly at c . Event-horizon escape is similarly excluded by causal structure; jets must be launched outside the horizon.

If a proposed galactic potential is parabolic, $\Phi_N(r) = ar^2$, then

$$\frac{v_c^2}{r} = \frac{d\Phi_N}{dr} = 2ar \implies v_c = \sqrt{2a} r, \quad (2.13)$$

which is solid-body rotation, not an asymptotically flat curve. A concrete modified-gravity alternative must fit both dynamics and lensing; recent weak lensing reaches far beyond classical rotation-curve radii [2]. Finally a decay half-life alone supplies no force. Only momentum flux does; for perfectly absorbed radiation

$$F = \frac{P}{c} = 3.3356 \text{ nN} \left(\frac{P}{1 \text{ W}} \right). \quad (2.14)$$

3 The Route-E theorem boundary imported into Route F

Route E assumes a self-carried family space

$$V = H^0(\Sigma, L), \quad L \simeq T_\Sigma, \quad V \simeq H^0(\Sigma, T_\Sigma) = \text{aut}(\Sigma). \quad (3.1)$$

The original invariant-contact hypothesis H3 requires a nondegenerate symmetric invariant bilinear form. It gives only $N_{\text{fam}} \leq 3$, since the genus-one one-dimensional Abelian algebra admits such a

form. The exact selection $N_{\text{fam}} = 3$ requires the explicit stronger hypothesis $H3^+$: the contact is the nondegenerate adjoint-trace/Killing form. On that selected branch,

$$\Sigma \simeq \mathbb{CP}^1, \quad T_{\mathbb{CP}^1} \simeq \mathcal{O}(2), \quad V \simeq \mathfrak{sl}_2(\mathbb{C}), \quad \dim_{\mathbb{C}} V = 3. \quad (3.2)$$

Thus every result below that invokes this branch is conditional on $H3^+$; the subsequent algebra needs no additional physical axiom once the branch is chosen.

Riemann–Roch and Serre duality give the dimension transparently:

$$h^0(\mathcal{O}(2)) - h^1(\mathcal{O}(2)) = \deg \mathcal{O}(2) + 1 = 3, \quad (3.3)$$

$$h^1(\mathcal{O}(2)) = h^0(K \otimes \mathcal{O}(-2)) = h^0(\mathcal{O}(-4)) = 0, \quad (3.4)$$

so $h^0 = 3$. Here $\mathcal{O}(2)$ is a degree-two holomorphic line bundle, not the orthogonal group $O(2)$.

In Route E's spherical basis (T_{+1}, T_0, T_{-1}) ,

$$K_{\text{tr}} = \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad B = 2\sqrt{3} K_{\text{tr}}, \quad K_{\text{tr}}^2 = \frac{1}{3} I. \quad (3.5)$$

This complex symmetric invariant pairing is not a Hermitian energy operator. Its indefinite real-slice signs cannot be read as positive and negative physical energies.

4 Exact discriminant–two-center–Killing theorem

4.1 Global vector fields and the adjoint representation

Let $\xi = v/u$ on an affine chart of $\mathbb{CP}^1$. A holomorphic vector field has the local form $X = q(\xi)\partial_\xi$. On the other chart $\eta = 1/\xi$,

$$q(\xi)\partial_\xi = -\eta^2 q(1/\eta)\partial_\eta. \quad (4.1)$$

Regularity at $\eta = 0$ therefore forces $\deg q \leq 2$:

$$X_q = (a + b\xi + c\xi^2)\partial_\xi. \quad (4.2)$$

The basis

$$E = \partial_\xi, \quad H = -2\xi\partial_\xi, \quad F = -\xi^2\partial_\xi \quad (4.3)$$

obeys

$$[H, E] = 2E, \quad [H, F] = -2F, \quad [E, F] = H, \quad (4.4)$$

which proves $H^0(\mathbb{CP}^1, T) \simeq \mathfrak{sl}_2(\mathbb{C})$.

Homogenizing (4.2) gives the binary quadratic

$$p(u, v) = au^2 + buv + cv^2, \quad \Delta(p) = b^2 - 4ac. \quad (4.5)$$

The representation statement is precisely

$$H^0(\mathbb{CP}^1, \mathcal{O}(2)) \simeq \text{Sym}^2(\mathbb{C}^2)^*, \quad (4.6)$$

and the invariant epsilon tensor identifies the dual with $\text{Sym}^2 \mathbb{C}^2$ for $SL(2)$.

Theorem 4.1 (Discriminant–Killing equivalence). *For a nonzero $X_q \in H^0(\mathbb{CP}^1, T)$, the following are equivalent:*

1. its zero divisor is a sum of two distinct points;
2. $\Delta(p) \neq 0$;
3. the corresponding $\mathfrak{sl}_2(\mathbb{C})$ element is regular semisimple;
4. its Killing norm is nonzero;
5. its Route-E contact norm $K_{\text{tr}}(X_q, X_q)$ is nonzero.

The complement $\Delta = 0$ is the nonzero nilpotent/contact-null cone and has one double zero.

Proof. Take the standard projective infinitesimal matrix

$$A_q = \begin{pmatrix} -b/2 & -c \\ a & b/2 \end{pmatrix}. \quad (4.7)$$

Direct multiplication gives

$$A_q^2 = \frac{\Delta(p)}{4} I. \quad (4.8)$$

Hence $\Delta \neq 0$ gives two nonzero opposite eigenvalues and a regular semisimple element. If $\Delta = 0$ and $A_q \neq 0$, then $A_q^2 = 0$, so the element is nilpotent. The roots of p are distinct exactly when its discriminant is nonzero; when it vanishes, the homogeneous quadratic is a square and has a double projective root. Counting multiplicity, the total degree is always

$$\deg \text{div}(X_q) = c_1(T_{\mathbb{CP}^1})[\mathbb{CP}^1] = \chi(S^2) = 2. \quad (4.9)$$

For $\mathfrak{sl}_2(\mathbb{C})$, the canonical adjoint-trace Killing form obeys $B(A, A) = 4 \text{Tr}_{2 \times 2}(A^2)$, so the displayed matrix gives

$$B(A_q, A_q) = 2\Delta(p). \quad (4.10)$$

No rescaling of Lie-algebra generators is allowed here. Instead, write the same matrix in Route E's spherical basis as

$$A_q = x_+ T_{+1} + x_0 T_0 + x_- T_{-1} = \begin{pmatrix} x_0/2 & -x_+/\sqrt{2} \\ x_-/\sqrt{2} & -x_0/2 \end{pmatrix}, \quad a = \frac{x_-}{\sqrt{2}}, \quad b = -x_0, \quad c = \frac{x_+}{\sqrt{2}}. \quad (4.11)$$

Then $\Delta = x_0^2 - 2x_+x_-$, while Eq. (3.5) gives $B(\mathbf{x}, \mathbf{x}) = 2\sqrt{3} \mathbf{x}^T K_{\text{tr}} \mathbf{x} = 2(x_0^2 - 2x_+x_-) = 2\Delta$. This is an invertible change of coordinates, not a change of bracket. It proves the remaining zero-versus-nonzero equivalence and fixes the normalization. $\square$

The normalization can be checked without words. Write

$$p = \frac{x_-}{\sqrt{2}} u^2 - x_0 uv + \frac{x_+}{\sqrt{2}} v^2, \quad \mathbf{x} = (x_+, x_0, x_-)^T. \quad (4.12)$$

Then

$$\Delta(p) = x_0^2 - 2x_+x_-, \quad (4.13)$$

$$\mathbf{x}^T K_{\text{tr}} \mathbf{x} = \frac{1}{\sqrt{3}} (x_0^2 - 2x_+x_-), \quad (4.14)$$

$$\boxed{B(\mathbf{x}, \mathbf{x}) = 2\Delta(p) = 2\sqrt{3} \mathbf{x}^T K_{\text{tr}} \mathbf{x}.} \quad (4.15)$$

The unnormalized second transvectant is

$$(p, p)_2 = p_{uu}p_{vv} - 2p_{uv}^2 + p_{vv}p_{uu} = 8ac - 2b^2 = -2\Delta(p) = -B(\mathbf{x}, \mathbf{x}), \quad (4.16)$$

so the same invariant is the transvectant singlet, again up to its declared normalization.

Remark 4.2 (What the theorem does not say). *The theorem welds two Route-E structures exactly. It does not identify the two zeros with positive/negative energy, two particles, or two families. It also shows why the regular-semisimple qualifier matters: a generic nonzero section has two distinct centers, but a nilpotent section has one center with multiplicity two.*

5 A safe two-polarity bridge: Hopf reduction and moment map

A creative way to retain the source manuscript's two polarities without a negative-energy instability is to begin with a complex doublet $z = (z_0, z_1) \in \mathbb{C}^2$. At fixed norm, $z^\dagger z = 1$, the state space is S^3 . Quotienting the common phase gives

$$S^3/U(1) \simeq \mathbb{CP}^1. \quad (5.1)$$

The Hopf map is

$$\mathbf{n} = (2\Re z_0^* z_1, 2\Im z_0^* z_1, |z_0|^2 - |z_1|^2), \quad \mathbf{n}^2 = 1. \quad (5.2)$$

For the relative circle action, a normalized Fubini–Study moment map is

$$\mu([z_0 : z_1]) = \frac{|z_0|^2 - |z_1|^2}{|z_0|^2 + |z_1|^2}, \quad -1 \leq \mu \leq 1. \quad (5.3)$$

Its two fixed points satisfy

$$p_+ = [1 : 0], \quad \mu(p_+) = +1; \quad p_- = [0 : 1], \quad \mu(p_-) = -1. \quad (5.4)$$

The numerical endpoint values ± 1 use the displayed generator and the symmetric additive normalization. Rescaling the generator by c rescales them to $\pm c$. What is normalization-independent is that the compact moment image is bounded and that the two fixed points are opposite extrema (up to reversing orientation). This suggests the normalized safe dictionary

$$E_p, E_n \rightsquigarrow \mu = +1, -1 \quad (\text{charge/orientation polarity}), \quad (5.5)$$

while the physical Hamiltonian remains bounded below.

With

$$\omega_{\text{FS}} = \frac{i d\xi \wedge d\bar{\xi}}{(1 + |\xi|^2)^2}, \quad \frac{1}{2\pi} \int_{\mathbb{CP}^1} \omega_{\text{FS}} = 1, \quad (5.6)$$

geometric quantization of Chern number $k \geq 0$ gives

$$L \simeq \mathcal{O}(k), \quad \mathcal{H}_k = H^0(\mathbb{CP}^1, \mathcal{O}(k)) \simeq \text{Sym}^k \mathbb{C}^2, \quad \dim \mathcal{H}_k = k + 1. \quad (5.7)$$

If, and only if, the physical bubble reduction also satisfies Route E's self-carried identification $L = T_{\mathbb{CP}^1} \simeq \mathcal{O}(2)$, then $k = 2$ and $\dim \mathcal{H} = 3$.

Bridge axiom 5.1 (Projective-doublet bridge, BA-P). *The energy-information order parameter is a physical complex doublet; its fixed-charge reduction is $\mathbb{CP}^1$; its Berry/prequantum bundle equals $T_{\mathbb{CP}^1}$; and its chiral zero-mode operator has $\ker D^+ \simeq H^0(T)$, $\ker D^- = 0$, with no extra charged zero modes.*

None of the four clauses follows merely from the Hopf identity (5.1). This bridge is therefore a concrete research target, not a derivation of three families.

6 Conditional Q-ball completion of an energy bubble

6.1 Action, charge, and radial equation

Replace a literal positive/negative-energy bubble by one complex scalar with a healthy kinetic term:

$$S_\Phi = \int d^4x \sqrt{-g} [-g^{\mu\nu} \partial_\mu \Phi^* \partial_\nu \Phi - U(|\Phi|)], \quad (6.1)$$

with

$$U(f) = m^2 f^2 - \lambda f^4 + \frac{\eta}{M^2} f^6, \quad m^2, \lambda, \eta, M^2 > 0. \quad (6.2)$$

The global $U(1)$ current can be normalized so that for

$$\Phi(t, r) = f(r) e^{i\omega t} \quad (6.3)$$

the charge and energy in flat space are

$$Q = 2\omega \int d^3x f^2, \quad (6.4)$$

$$E = \int d^3x [\omega^2 f^2 + (\nabla f)^2 + U(f)]. \quad (6.5)$$

Variation gives the boundary-value problem

$$f'' + \frac{2}{r} f' = \left(m^2 - \omega^2 - 2\lambda f^2 + \frac{3\eta}{M^2} f^4 \right) f, \quad f'(0) = 0, \quad f(\infty) = 0. \quad (6.6)$$

Proposition 6.1 (Existence window for the sextic model). *The thin-wall Q-ball window is nonempty when*

$$0 < \omega_0^2 := \min_{f>0} \frac{U(f)}{f^2} < m^2, \quad (6.7)$$

and for (6.2)

$$f_0^2 = \frac{\lambda M^2}{2\eta}, \quad \omega_0^2 = m^2 - \frac{\lambda^2 M^2}{4\eta}, \quad \omega_0^2 < \omega^2 < m^2. \quad (6.8)$$

Proof. The effective ratio is

$$\frac{U(f)}{f^2} = m^2 - \lambda f^2 + \frac{\eta}{M^2} f^4. \quad (6.9)$$

Differentiation with respect to f^2 gives $-\lambda + 2\eta f^2/M^2 = 0$, hence the displayed f_0 and ω_0 . The upper inequality ensures exponential decay at infinity because the linearized equation has inverse length $\sqrt{m^2 - \omega^2}$; the lower inequality makes the effective potential $U_\omega = U - \omega^2 f^2$ negative somewhere, allowing a nontrivial bounce-like radial trajectory. This is the standard Q-ball criterion [3]; interacting extensions can alter the stability range and must be retested [4]. $\square$

6.2 Reproducible benchmark and approximation boundary

For

$$m = M = 1 \text{ GeV}, \quad \lambda = \eta = 1, \quad Q = 10^6, \quad (6.10)$$

one finds

$$f_0 = \frac{1}{\sqrt{2}} \text{ GeV}, \quad \omega_0 = \frac{\sqrt{3}}{2} \text{ GeV} = 0.8660254 \text{ GeV}. \quad (6.11)$$

At $\omega = \omega_0$,

$$U_\omega(f) = f^2(f^2 - f_0^2)^2, \quad (6.12)$$

and the wall tension in the present kinetic convention is

$$\sigma = 2 \int_0^{f_0} \sqrt{U_\omega(f)} df = \frac{1}{8} \text{ GeV}^3. \quad (6.13)$$

Using $Q = 2\omega_0 f_0^2 (4\pi R^3/3)$ gives

$$R_0 = \left(\frac{3Q}{8\pi\omega_0 f_0^2} \right)^{1/3} = 65.0819045 \text{ GeV}^{-1} = 12.8424157 \text{ fm}. \quad (6.14)$$

Adding the leading surface energy yields

$$\frac{E}{Q} \simeq \omega_0 + \frac{4\pi R_0^2 \sigma}{Q} = 0.872678754 \text{ GeV} < m. \quad (6.15)$$

An independent fixed-step-profile variation minimizes

$$E(R) = \frac{Q^2}{4f_0^2 V(R)} + U(f_0)V(R) + 4\pi R^2 \sigma, \quad V(R) = \frac{4\pi R^3}{3}, \quad (6.16)$$

and gives

$$R = 64.9712654 \text{ GeV}^{-1}, \quad \omega = 0.870457184 \text{ GeV}, \quad E/Q = 0.872667434 \text{ GeV}. \quad (6.17)$$

These are thin-wall/step-profile checks, not an exact numerical solution of (6.6). The exact profile and fluctuation operator remain a promotion gate.

6.3 Exact-symmetry phase no-observable theorem

Theorem 6.2 (No phase clock from an exact Q-ball symmetry). *Let the action and a local observable $\mathcal{O}$ be invariant under the global transformation $\Phi \mapsto e^{i\alpha}\Phi$. On the stationary Q-ball orbit $\Phi = f(r)e^{i\omega t}$, $\mathcal{O}$ is independent of time. Therefore the exact Q-ball phase alone cannot make an invariant mass or visibility periodically appear and disappear.*

Proof. For every t , the field is obtained from $f(r)$ by the symmetry transformation with $\alpha = \omega t$. Thus

$$\mathcal{O}[f(r)e^{i\omega t}] = \mathcal{O}[e^{i\omega t}f(r)] = \mathcal{O}[f(r)]. \quad (6.18)$$

In particular, $|\Phi|^2$, $T_{\mu\nu}$, and any invariant pole mass are stationary. An observable phase requires a second phase reference or an explicitly symmetry-breaking spurion. $\square$

This creates a useful no-free-lunch constraint. Exact $U(1)$ protects Q but hides the absolute phase; a fixed relative-phase term can expose the phase but then generally gives $\nabla_\mu j^\mu \neq 0$, so absolute Q-ball stability is replaced by a calculable lifetime.

7 Route-E weight-two phase and a physical relative phase

Route E reconstructs

$$M_R = -m_D^T m_\nu^{-1} m_D, \quad m_D = v Y_\nu. \quad (7.1)$$

Under a uniform nonzero complex rescaling $Y_\nu \mapsto y Y_\nu$,

$$M_R \mapsto y^2 M_R. \quad (7.2)$$

If the positive scale is defined by a Takagi singular value, $M_* = \sigma_{\max}(M_R) > 0$, then

$$M_* \mapsto |y|^2 M_*. \quad (7.3)$$

For the normalized contact projection this gives the exact covariance

$$\boxed{\zeta \mapsto \frac{y^2}{|y|^2} \zeta = e^{2i \arg y} \zeta.} \quad (7.4)$$

Consequently ζ is invariant under any nonzero real uniform scaling, not only positive scaling; for a complex scaling its phase has weight two. The normalized construction is undefined at $y = 0$, and family-dependent rescalings do not obey (7.4).

The benchmark is

$$\zeta = 0.1076472949 + 0.0736514853i, \quad |\zeta| = 0.1304319033, \quad \arg \zeta = 0.6000380203. \quad (7.5)$$

Uniform phase covariance is not itself observable. If Φ carries weight one and ζ weight two, a rephasing-invariant relative phase is

$$e^{i\delta} := \frac{\zeta^* \Phi^2}{|\zeta| |\Phi|^2}, \quad \delta = 2 \arg \Phi - \arg \zeta. \quad (7.6)$$

A bounded visibility ansatz with the source manuscript's four special phases is then

$$w(\delta) = \frac{1 + \cos \delta}{2} = \cos^2 \left(\arg \Phi - \frac{1}{2} \arg \zeta \right), \quad 0 \leq w \leq 1. \quad (7.7)$$

This changes a coupling or spectral residue, not the Poincaré Casimir. The two naive identifications are numerically inequivalent:

$$\cos^2(\arg \zeta) = 0.68114344, \quad \cos^2(\arg \zeta / 2) = 0.91265707. \quad (7.8)$$

Route E selects neither convention.

For two complex symmetric Majorana structures $M_R = M_V + M_C$, $M_C = \zeta K_{\text{tr}}$, a family-basis congruence sends both as $M \mapsto U^T M U$. If M_V is invertible,

$$\mathcal{R} := M_V^{-1} M_C \mapsto U^{-1} \mathcal{R} U. \quad (7.9)$$

Therefore quantities such as

$$\Im \text{Tr}(\mathcal{R}^n), \quad \arg \det(I + \mathcal{R}) \quad (7.10)$$

are basis invariants that can test a *relative* contact phase. Any future phase claim must be written in such invariant form rather than as an absolute $\arg \zeta$.

8 Candidate Q-ball–Schur-complement bridge

8.1 Action-level construction

Let $N \in V$ and a sterile heavy messenger $X \in V^*$ be left-handed Weyl family vectors. A non-supersymmetric quadratic mass sector with a dynamical scalar-dependent mixing is

$$\begin{aligned} \mathcal{L}_{NX} = & iN^\dagger \bar{\sigma}^\mu D_\mu N + iX^\dagger \bar{\sigma}^\mu \partial_\mu X \\ & - \frac{M_*}{2} [N^T M_V N + 2\lambda(\Phi) X^T N - X^T K_{\text{tr}}^{-1} X] + \text{h.c.} \end{aligned} \quad (8.1)$$

At momenta well below the messenger gap, the algebraic equation is

$$X = \lambda(\Phi) K_{\text{tr}} N. \quad (8.2)$$

Substitution gives

$$\mathcal{L}_{\text{eff}} \supset -\frac{M_*}{2} N^T [M_V + \lambda(\Phi)^2 K_{\text{tr}}] N + \text{h.c.} \quad (8.3)$$

More generally, if the messenger kernel is $-\mu K_{\text{tr}}^{-1}$, then

$$\zeta_{\text{eff}} = \frac{\lambda(\Phi)^2}{\mu}. \quad (8.4)$$

Route E sets $\mu = 1$ by a field rescaling; keeping it visible is useful for charge and normalization audits.

The direct phase-doubling bridge is

$$\lambda(\Phi) = g \frac{\Phi}{\Lambda} \quad \Longrightarrow \quad \boxed{\zeta_{\text{eff}}(x) = g^2 \frac{\Phi(x)^2}{\Lambda^2}}. \quad (8.5)$$

On a Q-ball background,

$$\zeta_{\text{eff}}(r, t) = g^2 \frac{f(r)^2}{\Lambda^2} e^{2i\omega t}. \quad (8.6)$$

This exactly reproduces a localized weight-two coefficient. It is the most direct candidate relation between the two research programs.

Bridge axiom 8.1 (Q-ball contact portal, BA-QK). *The physical heavy mode has the Route-E family representation and kernel in (8.1); the dynamical mixing is $g\Phi/\Lambda$; all charge, gauge, anomaly, and operator-selection rules are satisfied; and the resulting relative phase cannot be removed by a field redefinition.*

8.2 Why the portal is not yet a completion

The following gates are mandatory.

Charge ledger. With $q_\Phi = +1$, the choices $q_X = 0$, $q_N = -1$ make $\Phi X N$ and XX invariant, but $N^T M_V N$ then needs a charge-+2 spurion. If M_V is instead a fixed explicit term, the Q-ball $U(1)$ is broken. If all spurions transform dynamically, the full background may be stationary in a rotating frame and no visibility modulation follows. This tension must be resolved, not hidden in notation.

Tree-level normalization. For canonical messenger kinetic terms, the leading low-energy substitution also gives

$$Z_N = I + |\lambda|^2 K_{\text{tr}}^\dagger K_{\text{tr}} = \left(1 + \frac{|\lambda|^2}{3}\right) I. \quad (8.7)$$

At the Route-E benchmark $|\lambda|^2 = |\zeta|$,

$$\delta Z_N = \frac{|\zeta|}{3} = 0.0434773011. \quad (8.8)$$

This is a tree-level matching effect. The canonical physical test is the complete Weinberg combination $Y M^{-1} Y^T$, not ζ by itself.

Full propagator and interaction. When external energies are not negligible relative to all heavy Takagi singular values, one must use

$$\mathcal{M}_{NX} = M_* \begin{pmatrix} M_V & \lambda I \\ \lambda I & -K_{\text{tr}}^{-1} \end{pmatrix} \quad (8.9)$$

and its kinetic matrix. The original fixed- λ model is Gaussian and does not by itself generate a $|\lambda|^2/(16\pi^2)$ anomalous dimension. The dynamical vertex $g\Phi XN/\Lambda$ supplies a genuine interaction, but its loop matching, cutoff power counting, and counterterms must be calculated.

Selection and leakage. The existing Route-B rule permits the unwanted post-breaking operator XLH . A valid discrete/gauge rule must forbid it and enumerate operators through a declared dimension. Moreover, if neutrinos or messengers carry the Q-ball charge, the soliton may emit them. Absolute stability requires its energy per unit charge to lie below every kinematically available charge-carrying channel; explicit phase locking instead requires a lifetime calculation.

Static benchmark versus rotating background. Route E fits a constant ζ . Equation (8.6) is periodic relative to a fixed M_V . It therefore requires a Floquet/adiabatic seesaw analysis and cannot be inserted into the static fit without rerunning neutrino, threshold, proton-decay, and cosmology gates.

9 Candidate dynamical origin of $H3^+$

The previous sections use $H3^+$ but do not derive it. There is, however, a specific loop mechanism worth testing. Couple a complete heavy family adjoint multiplet to a background family source A_μ^a . Covariance and the Ward identity constrain its vacuum polarization to

$$\Pi_{ab}^{\mu\nu}(p) = (p^2 \eta^{\mu\nu} - p^\mu p^\nu) \Pi(p^2) \text{Tr}_{\text{ad}}(\text{ad } T_a \text{ad } T_b). \quad (9.1)$$

The family tensor is therefore

$$B_{ab}^{\text{Kill}} := \text{Tr}_{\text{ad}}(\text{ad } T_a \text{ad } T_b). \quad (9.2)$$

For a simple algebra, invariant symmetric bilinears form a one-dimensional space, so a nonzero result is proportional to the Killing form. On the Route-E branch this is $2\sqrt{3}K_{\text{tr}}$.

Bridge axiom 9.1 (Adjoint-loop $H3^+$ mechanism, BA- $H3$). *A healthy UV sector contains a complete, non-canceling adjoint current whose low-energy two-point function induces the required family bilinear, and a specified source/messenger map converts that bilinear into the Majorana contact rather than merely a gauge kinetic term.*

Equation (9.1) explains why a Killing tensor is natural once an adjoint loop exists; it does not prove the existence of that loop, its nonzero coefficient, a positive spectral density in the relevant channel, or the conversion to a Majorana mass. $H3^+$ therefore remains physically open.

10 The oscillatory formula: exact solution and strict separation

The source manuscript includes

$$\sin(x^2) + \cos(x^2) = \sin(y^2) + \cos(y^2). \quad (10.1)$$

Using $\sin t + \cos t = \sqrt{2} \sin(t + \pi/4)$, the identity $\sin A = \sin B$ gives

$$A = B + 2\pi n \quad \text{or} \quad A = \pi - B + 2\pi n, \quad n \in \mathbb{Z}. \quad (10.2)$$

With $A = x^2 + \pi/4$, $B = y^2 + \pi/4$, the complete real solution set is

$$\boxed{x^2 - y^2 = 2\pi n} \quad \text{or} \quad \boxed{x^2 + y^2 = \frac{\pi}{2} + 2\pi n}, \quad (10.3)$$

subject to real-coordinate nonemptiness. The circular radii are

$$r_n^2 = \frac{\pi}{2} + 2\pi n, \quad n \geq 0, \quad (10.4)$$

and their exact radial pitch is

$$r_{n+1} - r_n = \frac{2\pi}{r_{n+1} + r_n} \sim \frac{\pi}{r_n}. \quad (10.5)$$

Thus a shrinking interference pitch is a theorem. These are continuous circles/hyperbolae, not discrete atomic orbitals; standard subshell degeneracies are $2(2\ell + 1) = 2, 6, 10, 14, \dots$

There is no direct Route-E identification. The functions $\sin(\xi^2)$ and $\cos(\xi^2)$ have an essential singularity in the $\eta = 1/\xi$ chart, whereas global sections of $T_{\mathbb{CP}^1}$ are exactly degree-at-most-two polynomials. A shared appearance of squares is insufficient to identify the state spaces.

11 Relational time and gravity: parallel interfaces, not Route-E derivations

The source intuition that time may be relational has a standard conditional form. Let a clock C and system S satisfy the stationary constraint

$$(H_C + H_S)|\Psi\rangle = 0. \quad (11.1)$$

If clock states obey $\langle t|H_C = -i\hbar\partial_t\langle t|$, then

$$|\psi(t)\rangle := \langle t|\Psi\rangle \quad (11.2)$$

satisfies

$$0 = \langle t|(H_C + H_S)|\Psi\rangle \quad (11.3)$$

$$= -i\hbar\partial_t\langle\psi(t)| + H_S|\psi(t)\rangle, \quad (11.4)$$

hence

$$i\hbar\partial_t|\psi(t)\rangle = H_S|\psi(t)\rangle. \quad (11.5)$$

This Page–Wootters construction [5] has active modern extensions, including finite-dimensional time-dilation mechanisms [6]. It does not identify time with local E_d , nor does it derive Route E.

A thermodynamic route to Einstein’s equation is likewise conditional. On every local Rindler horizon, impose

$$\delta Q = T_U \delta S, \quad T_U = \frac{\hbar\kappa}{2\pi}, \quad \delta S = \frac{\delta A}{4G\hbar}. \quad (11.6)$$

The Raychaudhuri equation relates the area variation to $R_{\mu\nu}k^\mu k^\nu$, while the heat flux is proportional to $T_{\mu\nu}k^\mu k^\nu$. Requiring the equality for every null vector k^μ gives

$$R_{\mu\nu} + \Phi g_{\mu\nu} = 8\pi G T_{\mu\nu}. \quad (11.7)$$

Stress conservation and the Bianchi identity fix $\Phi = -R/2 + \Lambda$, yielding Einstein’s equation [7, 8]. The horizon entropy law and local equilibrium are inputs. This route is a gravity interface for later work, not evidence for the family/contact bridge.

12 Claim ledger and ordered research program

Claim	Status	Boundary / next gate
E_d alone determines time/gravity	refuted	Thin-shell and pressure counterexamples, Eqs. (2.3)–(2.7).
Invariant mass disappears at four internal phases	refuted	Poincaré Casimir; only coupling/residue/gap may be modulated.
Route-E CP1/O(2) carrier has three sections	conditional theorem	Requires H3 ⁺ to select the branch; H3 alone gives only the upper bound.
Two centers $\Leftrightarrow \Delta \neq 0 \Leftrightarrow K_{\text{tr-norm}}$ nonzero	proved	Exact on the selected CP1 branch, Theorem 4.1.
Opposite center labels without negative energy	candidate bridge	Moment-map polarity is exact; physical identification requires BA-P.
Energy bubble as a Q-ball	conditional completion	Exact radial solution, Hessian, charge leakage, and gravity remain open.
ζ carries phase weight two	proved in Route E	Uniform complex rescaling only; the absolute phase is not observable.
$\zeta_{\text{eff}} = g^2 \Phi^2 / \Lambda^2$	candidate bridge	Requires BA-QK, charge/selection rules, full matching, and rerun fits.
Adjoint loop physically derives H3 ⁺	open mechanism	Killing tensor follows conditionally; existence and Majorana conversion do not.
Oscillatory equation quantizes atoms or gives CP1 sections	refuted	It gives continuous conics and has an essential singularity at CP1 infinity.
Relational time / thermodynamic gravity	parallel conditional tracks	Useful interfaces; neither derives Route E or the bubble map.

The sequential program is deliberately fail-closed.

- AP–E1. Projective-doublet action.** Write a two-component bubble action, identify its constraint and gauge/global quotient, and prove or refute that the physical reduced moduli space is $\mathbb{CP}^1$.
- AP–E2. Discriminant/contact verifier.** Preserve Theorem 4.1 as an exact symbolic regression test, including the regular/nilpotent boundary and Route-E normalization.
- AP–E3. Self-carried geometric quantization.** Derive the Berry curvature and Chern number from AP–E1 rather than imposing them; test whether the prequantum bundle is actually $T_{\mathbb{CP}^1}$.
- AP–E4. Chiral zero modes.** Construct the fluctuation/Dirac operator, prove $\ker D^+ \simeq H^0(T)$, compute $\ker D^-$, extra modes, and the spectral gap.
- AP–E5. Exact Q-ball solution.** Solve Eq. (6.6), compute $Q(\omega)$, $E(Q)$, fixed-charge Hessian modes, gravitational compactness, and all charge-emission thresholds.
- AP–E6. Messenger/contact realization.** Build a gauge- and anomaly-consistent version of Eq. (8.1); enumerate operators, forbid XLH , and verify the full kinetic plus six-by-six matching.
- AP–E7. Physical-phase quotient.** Remove all field rephasings, list basis-invariant CP quanti-

ties such as Eq. (7.10), and test whether a relative phase survives without destroying the Q-ball lifetime.

AP–E8. H_3^+ origin test. Compute the candidate adjoint current correlator and its spectral sign; show whether it yields a Majorana contact or only a kinetic bilinear.

AP–E9. Integrated phenomenology. Rerun Route-E flavor/seesaw, threshold, proton-decay, cosmology, and time-dependent/Floquet gates with a single action. Only this stage can request physics promotion.

AP–E10. Independent gravity track. Constrain any E_d carrier with clock redshift, fifth-force, rotation-curve, and lensing data; do not merge it with AP–E1–9 without an explicit operator connecting the sectors.

13 2026-07-16 AP–E3 global/nonlinear and AP–E4 SQM handoff

This section is a reversible index into three standalone derivation notes. It records the equations needed to reconstruct the decision boundary without replacing their complete proofs and numerical ledgers.

13.1 Charged two-colour nonlinear proxy

The declared six-real-field diagnostic potential is

$$V = \frac{\lambda}{4} (\sigma^2 + \pi^2 + 2|\Delta|^2 - v^2)^2 - h\sigma + \mu_g^2 |\Delta|^2, \quad q_\Delta = 2. \quad (13.1)$$

At the mesonic stationary point, $h/f = m_\pi^2$, and its exact Hessian contains

$$m_\sigma^2 = m_\pi^2 + 2\lambda f^2, \quad m_\Delta^2 = m_\pi^2 + \mu_g^2. \quad (13.2)$$

The requested $(e_g, v_g/\Lambda_c)$ scan is explicitly a matching ansatz, not a UV loop result:

$$\frac{\mu_g^2}{\Lambda_c^2} = \frac{\mu_{\text{PG},0}^2}{\Lambda_c^2} + c_g \left(e_g \frac{v_g}{\Lambda_c} \right)^2. \quad (13.3)$$

The mesonic nonlinear sector solves the $B = 1$ hedgehog with

$$B = -\frac{2}{\pi} \int_0^\infty \sin^2 F F' dx = 1, \quad E_2 - E_4 + 3E_0 = 0, \quad (13.4)$$

and diagonalizes both the generalized radial operator and the charged block

$$\mathcal{H}_{\Delta,\ell} = -\frac{d^2}{dx^2} + \alpha^2 + \chi(1 - \cos F) + \frac{\ell(\ell+1)}{x^2}. \quad (13.5)$$

The deterministic 18/18 card passes the classical radial necessary gate. It deliberately leaves lattice QC2D, the full coupled three-dimensional Hessian, quantum determinants, and global target-space unwinding open. The complete formulas are in `route_f/tex/ap_e3_charged_two_colour_soliton_proxy.tex`.

13.2 Non-extendible mixed WZW and spin refinement

Put $X = S^3 \times S^2$, choose integral generators u_3, u_2 , and let $\widehat{\Omega}_n \in \widehat{H}^5(X; \mathbb{Z})$ have curvature $n\omega_3 \wedge \omega_2$ and characteristic class nu_3u_2 . The bosonic action on every closed spin four-manifold, without an extension of its map, is

$$Z_{\text{bos}}(M, \phi) = \exp\left(2\pi i \widehat{\Omega}_n(\phi_*[M])\right). \quad (13.6)$$

Here $H^4(X; \mathbb{R}/\mathbb{Z}) = 0$, so fixed integral curvature has no ordinary flat ambiguity. Stable splitting instead gives

$$\Omega_4^{\text{Spin}}(S^3 \times S^2) \simeq \mathbb{Z} \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2. \quad (13.7)$$

If ν_3 is the spin-bordism class of the inverse image of a regular point under the S^3 projection and ν_2 is the Arf class of the inverse-image spin surface under the S^2 projection, the general reduced spin refinement is

$$Z_{\epsilon_3, \epsilon_2}(M, \phi) = Z_{\text{bos}}(M, \phi)(-1)^{\epsilon_3\nu_3 + \epsilon_2\nu_2}. \quad (13.8)$$

The bosonic differential-character definition is complete; a microscopic APS/Dai–Freed determinant must still choose (ϵ_3, ϵ_2) before the spin-refined action is fixed.

13.3 Pure- $SU(3)$ no-go and near-miss

Adjoint breaking uses

$$\iota(A, e^{i\alpha}) = \text{diag}(e^{i\alpha}A, e^{-2i\alpha}), \quad \ker \iota = \{(1, 1), (-1, -1)\}, \quad (13.9)$$

so every $[SU(2) \times U(1)]/\mathbb{Z}_2$ term descending from $SU(3)$ obeys

$$2j + q = 0 \pmod{2}. \quad (13.10)$$

A colour singlet therefore has even charge, proving that the original $\mathbf{1}_{+1}$ scalar cannot occur. Since $qq(\phi^\dagger)^m$ requires $2X_q = mX_\phi$, the desired $m = 2$ would require an odd doublet charge to equal an even singlet charge. This is impossible. The closest explicit branch instead has

$$\mathbf{3} \rightarrow \mathbf{2}_{+1} \oplus \mathbf{1}_{-2}, \quad \bar{\mathbf{3}} \rightarrow \mathbf{2}_{-1} \oplus \mathbf{1}_{+2}, \quad M_q = m + Vy, \quad M_\ell = m - 2Vy. \quad (13.11)$$

Choosing $m = -Vy$ makes $M_q = 0$ and $M_\ell = -3Vy$, while an adjoint scalar mass split can keep $\mathbf{1}_{+2}$ light. This is an actual tree-level decoupling but its minimal baryon $qq\phi^\dagger$ is an $\mathcal{O}(1)$ $SU(2)_\phi$ flavour doublet, not the Route-E $\mathcal{O}(2)$ triplet. The full proof is in `route_f/tex/ap_e3_nonextendible_wzw_su3_uv_audit.tex`.

13.4 Physical tangent fermion and the composability gate

A half-BPS non-Abelian vortex independently yields a normalizable $\mathbb{CP}^1$ orientational modulus and a worldline fermion. On $v = 1/w$,

$$\psi^v = -w^{-2}\psi^w, \quad D_t\psi^v = -w^{-2}D_t\psi^w, \quad (13.12)$$

so the Grassmann field is physically tangent-valued. Canonical Spin^c quantization gives

$$Q_E = \sqrt{2}\bar{\partial}_E, \quad \mathcal{H}_{\text{ground}} = H^{0,*}(\mathbb{CP}^1, E). \quad (13.13)$$

In the declared canonical Spin^c re-quantization, the tangent Clifford generator alone corresponds to $E = \mathcal{O}$ and gives one ground state. The source-selected half-form ordering has no normalizable zero mode. A separately derived $E = \mathcal{O}(2)$ gives

$$(h^0, h^1) = (3, 0), \quad \lambda_{n,\pm} = \pm \frac{2}{\sqrt{C}} \sqrt{n(n+3)}, \quad \text{mult} = 2n + 3, \quad n \geq 1. \quad (13.14)$$

The intrinsic 27/27 SQM audit therefore derives a physical tangent fermion, but does not derive the canonical vacuum line and is not yet in the charged-two-colour mother model. AP–E3’s signed $k = +2$ WZW character composes with it only after proving the same $B = 1$, the same $\mathbb{CP}^1$ collective-coordinate map, transgressing the degree-five character along the spatial three-cycle, and pulling the resulting differential line back with degree $+2$. The degree-one Route-E portal is postponed until that theorem or an anomaly-free product compactification is available. See `route_f/tex/ap_e4_moduli_space_sqm.tex`.

14 2026-07-16 AP–E completion frontier

This section integrates the next three requested calculations. It records the common normalization and promotion logic; the standalone notes retain the full proofs, source manifests, and negative controls.

14.1 Four-dimensional action and graded $B = 1$ Hessian

The declared Euclidean lattice theory has the schematic action

$$S_E = S_{g,W/\text{imp}} + S_{\text{gf}} + \bar{c}M_{\text{FPC}} + S_{\chi\Delta} + \bar{\psi}D_W[U, V, z]\psi, \quad (14.1)$$

where $V \in SU(2)_c$, $z \in U(1)_g$, the meson $U \in SU(2)$ is neutral, and the complex diquark has charge two. At $(\bar{V}, \bar{z}, \bar{U}, \bar{\Delta}, \bar{\psi}) = (1, 1, U_{B=1}, 0, 0)$, write the even variables as $\Phi = (A, b, \eta, d_1, d_2)$. The complete formal quadratic structure is

$$\mathbb{H}_{\text{cl}} = \begin{pmatrix} K_{\Phi\Phi} & 0 & 0 \\ 0 & 0 & D_W \\ 0 & -D_W^T & 0 \end{pmatrix}, \quad K_{\Phi\Phi} = \begin{pmatrix} K_{AA} & K_{Ab} & K_{A\eta} & K_{Ad_1} & K_{Ad_2} \\ K_{bA} & K_{bb} & K_{b\eta} & K_{bd_1} & K_{bd_2} \\ K_{\eta A} & K_{\eta b} & K_{\eta\eta} & K_{\eta d_1} & K_{\eta d_2} \\ K_{d_1 A} & K_{d_1 b} & K_{d_1 \eta} & K_{\Delta} & 0 \\ K_{d_2 A} & K_{d_2 b} & K_{d_2 \eta} & 0 & K_{\Delta} \end{pmatrix}, \quad (14.2)$$

together with the ghost block. The one-loop functional and its fermionic collective-coordinate curvature are

$$\Gamma^{(1)} = \frac{1}{2} \log \det' K_{\Phi\Phi} - \log \det' M_{\text{FP}} - \Re \log \det D_W, \quad (14.3)$$

$$\partial_i \partial_j \Gamma_f^{(1)} = \Re \text{Tr} (D^{-1} D_i D^{-1} D_j - D^{-1} D_{ij}). \quad (14.4)$$

This is the requested full *formal* gauge–meson–ghost–fermion Hessian. The numerical certificate is deliberately smaller: it uses a 2^4 trivial-link background, substitutes a frozen positive Gauss–Newton meson block for the full Skyrme Jacobi operator, and evaluates one fermion curvature direction. It therefore cannot be called dynamical lattice QC2D or the complete nonlinear Hessian.

The 24/24 card gives

$$\begin{aligned} B(0) &= 0.999981468235, \\ \lambda_{\min}(K_{\eta,\text{test}}) &= 0.621297848377, & \lambda_{\min}(K_{\Delta}) &= 0.694069172619, \\ \lambda_{\min}(K_{\Delta,\text{control}}) &= -0.349308273811, & \sigma_{\min}(D_W) &= 0.702570765924, \end{aligned} \quad (14.5)$$

with gauge/ghost ξ -corrected residual 1.42×10^{-14} , Schur solve residual 6.30×10^{-16} , log-determinant curvature relative error 1.32×10^{-8} , and free Wilson/Symanzik observed orders 1.99861/3.99628. Monte Carlo, measure positivity, the exact Skyrme Jacobi zero-mode projection, continuum renormalization, FR quantization, and finite-amplitude stability all remain false gates. Full details are in `route_f/tex/ap_e5_4d_qc2d_lattice_hessian.tex`.

14.2 Two torsion generators and regulator-dependent phases

Explicit representatives of the reduced bordism generators are

$$\begin{aligned} G_3 &= (S_R^1 \times S^3, \text{pr}_{S^3}, \text{pt}), \\ G_2 &= (T_{RR}^2 \times S^2, \text{pt}, \text{pr}_{S^2}). \end{aligned} \quad (14.6)$$

Their transverse inverse images are the nonbounding periodic circle and the odd-spin torus. Hence

$$\text{ind}_2 D_{S_R^1} = 1, \quad \text{ind}_2 D_{T_{RR}^2}^+ = \text{Arf}(T_{RR}^2) = 1. \quad (14.7)$$

The chirality-paired ambient four-dimensional product spectra define only a factorized reference phase $(+1, +1)$. Stacking the two explicitly localized defect regulators instead gives the complete character table

$$Z_{\epsilon_3, \epsilon_2}(G_3) = (-1)^{\epsilon_3}, \quad Z_{\epsilon_3, \epsilon_2}(G_2) = (-1)^{\epsilon_2}. \quad (14.8)$$

Thus all four torsion assignments are realizable with the same de Rham curvature. This proves regulator non-uniqueness; it does not select the charged-QC2D phase. That selection still requires the complete Dirac–Yukawa operator, heavy mass signs, and its odd-dimensional mapping-torus APS/Dai–Freed problem.

14.3 A semisimple replacement for the diagonal quotient

For $[SU(2)_c \times U(1)]/\mathbb{Z}_2$, every representation obeys

$$2j + q = 0 \pmod{2}. \quad (14.9)$$

It admits $\mathbf{2}_{+1}$ but forbids $\mathbf{1}_{+1}$; hence any successful candidate must retain the unquotiented global form. Besides the minimal backup $SU(2)_c \times SU(2)_H$, the scan finds the simply connected simple group

$$Sp(4) \equiv USp(4) \simeq Spin(5) \quad (14.10)$$

with branching

$$\begin{aligned} \mathbf{4} &\longrightarrow \mathbf{2}_0 \oplus \mathbf{1}_{+1} \oplus \mathbf{1}_{-1}, \\ \mathbf{5} &\longrightarrow \mathbf{2}_{+1} \oplus \mathbf{2}_{-1} \oplus \mathbf{1}_0, \\ \mathbf{10} &\longrightarrow \mathbf{1}_{+2} \oplus \mathbf{2}_{+1} \oplus \mathbf{3}_0 \oplus \mathbf{1}_0 \oplus \mathbf{2}_{-1} \oplus \mathbf{1}_{-2}. \end{aligned} \quad (14.11)$$

Two Dirac **5**'s provide the two light quark flavours. Two, not one, complex scalar **4**'s are required so their copy index retains the global $SU(2)_\phi$ doublet. Then

$$qq(\phi_I^\dagger \phi_J^\dagger)_{\text{sym}} \in \mathbf{3}_\phi \quad (14.12)$$

is the charge-neutral exactly-two $\mathcal{O}(2)$ operator at the representation-theory level. This does not rederive the earlier unit-cell Gauss rule.

One real adjoint **10** breaks the second $SU(2)$ to $U(1)$. For $M_q = m + qyV$, choosing $m = -yV$ keeps $q = +1$ light and makes the $q = -1, 0$ partners heavy at tree level. The displayed field content has

$$b_0^{Sp(4)} = 11 - \frac{4}{3}(2) - \frac{1}{6}(3) - \frac{1}{3}(1) = \frac{15}{2} > 0. \quad (14.13)$$

Vectorlike and Witten-parity checks pass. The tuning is not radiatively protected, however, and neither the full $\Omega_5^{\text{Spin}}(BSp(4))$ character nor the heavy determinant threshold has been computed. The algebraic light coefficient $N_c X_q = +2$ is therefore not yet an all-scale orientation theorem. See `route_f/tex/ap_e3_aps_global_form_search.tex`; its card passes 58/58.

14.4 Same-soliton SQM, Callias line, CPT, and WZW descent

For a hypothetical same-soliton Fredholm family

$$\mathcal{D}_m^+ = D_{A(m)}^+ + \mathcal{Y}_m, \quad m \in \mathcal{M}_1 \simeq \mathbb{CP}^1, \quad (14.14)$$

let the positive-mass eigenline on $S_\infty^2 \times \mathbb{CP}^1$ satisfy

$$c_1(\mathcal{E}_+) = px + qy. \quad (14.15)$$

The families-Callias push-forward gives, in the fixed orientation convention,

$$\text{rank Ind } \mathcal{D}^+ = \epsilon p, \quad c_1(\det \text{Ind } \mathcal{D}^+) = \epsilon p q y. \quad (14.16)$$

Thus a rank-one $\mathcal{O}(+2)$ determinant template requires $\epsilon p = 1, q = 2$. The finite Berry/kernel-family regression obtains $c_1 = 1.99999999999981$ with unit nonzero gap. It is a topology template, not the missing four-dimensional Yukawa operator or its Callias index.

In one fixed Dolbeault polarization, the conjugate triplet is fixed by Serre duality:

$$E_- = K_{\mathbb{CP}^1} \otimes E_+^\vee = \mathcal{O}(-2) \otimes \mathcal{O}(-2) = \mathcal{O}(-4), \quad (h^0, h^1) = (0, 3). \quad (14.17)$$

The naive inverse $\mathcal{O}(-2)$ instead has $(h^0, h^1) = (0, 1)$. Equation (14.17) closes the mathematical fixed-polarization map, not its microscopic CPT regulator.

On framed configuration space, spatial fiber integration is intrinsically defined:

$$\widehat{\mathcal{L}}_B = \int_{\Sigma_3} \text{ev}^* \widehat{\Omega}_5 \in \widehat{H}^2(\widetilde{\mathcal{C}}_B; \mathbb{Z}). \quad (14.18)$$

It descends through $p: \widetilde{\mathcal{C}}_B \rightarrow \mathcal{C}_B$ only when

$$\widehat{\mathcal{L}}_B \in \text{im } p^*, \quad (14.19)$$

which requires an equivariant differential refinement, horizontal invariant curvature, trivial vertical and large-gauge holonomy, and trivial stabilizer characters. After that descent, its pullback to a same-soliton $\mathbb{CP}^1$ is $\mathcal{O}(+2)$ precisely when

$$\int_{\Sigma_3 \times \mathbb{CP}^1} \text{ev}^* c(\widehat{\Omega}_5) = +2. \quad (14.20)$$

The factorized conditional map yields $nB \deg(s) = 2$, but the present theory has not identified its scalar S^2 with a normalizable soliton modulus and has not proved Eq. (14.19). The same-model card therefore passes 27/27 mathematical controls while keeping every composition flag false. The full derivation is `route_f/tex/ap_e4_same_soliton_callias_descent.tex`.

14.5 Portal authorization theorem and current decision

Let

$$\begin{aligned} C_{\text{lat}} &= \text{dynamical 4d phase plus complete projected quantum Hessian,} \\ C_{\text{APS}} &= \text{selected torsion phases plus protected semisimple all-scale completion,} \\ C_{\text{SQM}} &= \text{same-soliton SQM, Callias/CPT line, and gauge-basic WZW descent.} \end{aligned} \quad (14.21)$$

The user-specified ordering is encoded exactly as

$$\text{portal authorized} \iff C_{\text{lat}} \vee C_{\text{APS}} \vee C_{\text{SQM}}. \quad (14.22)$$

All three booleans are currently false. Consequently no portal operator is written.

When Eq. (14.22) first becomes true, the portal theorem must still exhibit a map

$$F : \mathcal{M}_1 \longrightarrow \Sigma_E, \quad \deg F = +1, \quad F^* \mathcal{O}_{\Sigma_E}(2) \simeq \mathcal{O}_{\mathcal{M}_1}(2), \quad (14.23)$$

fix its orientation, show that its operator is gauge and anomaly consistent, and keep every coupling below the retained bosonic, fermionic, and Landau gaps. It must also preserve the physical CPT pairing and introduce no extra zero modes. These are recorded as future prerequisites, not assumed facts.

The integrated promotion card `route_f/code/verify_ap_e5_completion_frontier.py` passes 20/20 and sets

$$\begin{aligned} \text{any_preportal_route_closed} &= \text{false,} \\ \text{degree_one_portal.constructed} &= \text{false,} \\ \text{physics_promotion_allowed} &= \text{false} \end{aligned}$$

(14.24)

15 AP-E6 same-background audit: exact obstructions and open gates

This section supersedes the completion-frontier status immediately above. It records the next execution round without weakening the portal ordering. The compact group denoted $Sp(4)$ in the physics convention is $USp(4)$, or $Sp(2)$ in the mathematical convention used in classifying spaces.

15.1 The exact $Sp(4)$ gauge-bordism result

The low-degree integral cohomology and spin-bordism coefficients are

$$\begin{aligned} H^*(BSp(2); \mathbb{Z}) &= \mathbb{Z}[q_1, q_2], & |q_1| &= 4, & |q_2| &= 8, \\ (\Omega_q^{\text{Spin}})_{q=0}^5 &= (\mathbb{Z}, \mathbb{Z}_2, \mathbb{Z}_2, 0, \mathbb{Z}, 0). \end{aligned} \quad (15.1)$$

In the homological Atiyah–Hirzebruch spectral sequence

$$E_{p,q}^2 = H_p(BSp(2); \Omega_q^{\text{Spin}}) \implies \Omega_{p+q}^{\text{Spin}}(BSp(2)), \quad (15.2)$$

the only nonzero term with $p + q = 5$ is

$$E_{4,1}^2 = H_4(BSp(2); \mathbb{Z}_2) = \mathbb{Z}_2. \quad (15.3)$$

The possible d_2 source and target use H_6 and H_2 , both zero; the d_3 target uses H_1 , and the only later possible map from this torsion group to $E_{0,4}^r \simeq \mathbb{Z}$ is zero. There is no extension ambiguity on this diagonal. Hence

$$\boxed{\Omega_5^{\text{Spin}}(BSp(2)) \simeq \mathbb{Z}_2.} \quad (15.4)$$

A geometric representative is a unit $Sp(2)$ instanton on S^4 times the nonbounding spin circle. Restriction to the instanton $SU(2)_c$ gives

$$\begin{aligned} \mathbf{4} &\downarrow \mathbf{2} \oplus \mathbf{1} \oplus \mathbf{1}, & \mathbf{5} &\downarrow \mathbf{2} \oplus \mathbf{2} \oplus \mathbf{1}, \\ \mathbf{10} &\downarrow \mathbf{3} \oplus \mathbf{2} \oplus \mathbf{2} \oplus \mathbf{1} \oplus \mathbf{1} \oplus \mathbf{1}. \end{aligned} \quad (15.5)$$

With the doublet index normalized to one, the instanton indices are $(1, 2, 6)$. The corresponding mod-two characters therefore are

$$(\chi_4, \chi_5, \chi_{10}) = (-1, +1, +1). \quad (15.6)$$

Thus the displayed two Dirac $\mathbf{5}$'s are trivial on the unique gauge bordism generator. This closes the gauge-bordism subgate only.

It does not select the two target-space signs. Independently,

$$\tilde{\Omega}_4^{\text{Spin}}(S^3 \times S^2) \simeq \mathbb{Z}_2\{S_R^1 \times S^3\} \oplus \mathbb{Z}_2\{T_{\text{Arf}=1}^2 \times S^2\}. \quad (15.7)$$

Without a common microscopic mass-family lift and APS boundary problem, invertible torsion sectors can flip either character. The gauge character in Eq. (15.6) and the ordered target pair are different questions.

The Euclidean note verifies gamma-five reality, two-flavour determinant positivity away from zeroes, and the Pauli–Villars moment identities

$$c_a = (1, -3, 3, -1), \quad \sum_a c_a a^j = 0 \quad (j = 0, 1, 2). \quad (15.8)$$

It does not specify the full interpolation $K(t)$, regulator statistics, APS domain, finite local counterterms, or regulated gauge–scalar–ghost complex. Accordingly these are partial controls, not a complete Euclidean regulator.

At the chosen light branch the infrared orientation remains

$$k_{\text{IR}} = N_c X_q B = 2 \cdot 1 \cdot 1 = +2. \quad (15.9)$$

For a uniform flavour action on a complete $\mathbf{5}$, however, the algebraic charge trace is only the conditional obstruction

$$k_{\text{full}} = 2(+1) + 2(-1) + 0 = +2 - 2 = 0. \quad (15.10)$$

The local pure-gauge shifts are periodic, but no unbroken-group or gravitational APS determinant ratio has been evaluated. Therefore Eq. (15.10) is not an actual heavy-determinant matching computation and neither target eta sign is selected.

15.2 One genuinely relaxed continuum B=1 field, but an off-shell cubic Hessian

The declared classical sector is enlarged to a unit $O(4)$ meson field n , a neutral breathing field s , and a quadratic charge-two field. Its continuum energy is

$$E[n, s] = \int d^3x \left\{ \frac{1}{2} |\partial_i n|^2 + \frac{\kappa}{2} \sum_{i < j} (|\partial_i n|^2 |\partial_j n|^2 - (\partial_i n \cdot \partial_j n)^2) + \frac{1}{2} |\nabla s|^2 + \frac{1}{2} m_s^2 (s-1)^2 + \beta^2 s^2 (1-n_0) \right\}. \quad (15.11)$$

For $n = (\cos F, \hat{x} \sin F)$, direct variation gives

$$(r^2 + 2\kappa \sin^2 F) F'' = -2rF' - 2 \sin F \cos F (\kappa F'^2 - 1) + \frac{2\kappa \sin^3 F \cos F}{r^2} + \beta^2 r^2 s^2 \sin F, \\ s'' + \frac{2}{r} s' = m_s^2 (s-1) + 2\beta^2 s (1 - \cos F). \quad (15.12)$$

With $(\beta, m_s, \kappa) = (1/2, 5/2, 1)$, a nonlinear BVP solve on $R = 8, 10, 12, 16$ converges to

$$E/(4\pi) = 6.1282155, \quad s(0) = 0.91712896, \quad B = +1, \quad (15.13)$$

and obeys the generalized Derrick identity. The canonical 4097-point little-endian float64 array (r, F, s) has SHA-256

81104f59a5f3f4337739fc2a217cafc8da91581c9f1fb40b4fdb6ce0f948d59.

This proves stationarity within the coupled hedgehog sector. It is not a Palais symmetric-criticality proof for every three-dimensional variation.

For the separately declared cubic action, the analytic constrained Hessian is

$$H_{\text{tan}} = T^T \left[H_{\text{emb}} - \bigoplus_x (n_x \cdot g_x) \mathbf{1}_4 \right] T, \quad H_{\text{proj}} = P_{\text{coll}} H_{\text{tan}} P_{\text{coll}}. \quad (15.14)$$

Every $n + s$ entry is differentiated from one finite-difference energy; sparse symmetry, geodesic second differences, and the translation/isorotation projector are checked independently. The quadratic diquark and free toron/ghost operators use separately declared grids and boundaries. They are audited controls, not blocks of one aggregate matrix. Thus the $n + s$ second variation is exact while the complete declared-sector sparse-Hessian gate remains false.

It is not yet a physical stability operator. At fixed $L = 8$, the sampled continuum solution gives

N	a	B_a	$\ \text{grad } E_a\ _{2,\text{dens}}$
9	1.000000	0.168018	0.34766
13	0.666667	0.440012	0.43018
17	0.500000	0.625354	0.42171

The background is therefore off shell for the cubic action and its topology estimator has not converged to one. The observed negative projected curvatures are useful falsifiers of premature stability claims, but they cannot be promoted to soliton instability eigenvalues. No interacting colour links, fermion determinant, four-dimensional importance sampling, or continuum quantum limit is present. Moreover, the cubic boundary samples the finite $R = 16$ profile rather than imposing the exact vacuum; at $L = 8$ its maximum boundary deviation is about 4.93×10^{-2} . This boundary mismatch is recorded as part of the off-shell diagnostic.

15.3 The same canonical field and the physical-moduli obstruction

The Yukawa/Callias audit consumes the checksum above rather than re-solving an older profile. The declared constituent Yukawa term couples to F but not to s ; this decoupling is part of the action, not a silent profile replacement:

$$\mathcal{D}_E = \gamma^\mu D_\mu + M(P_R U_F + P_L U_F^\dagger), \quad H_F = -i\alpha^i D_i + \beta M(\cos F + i\gamma_5 \boldsymbol{\tau} \cdot \hat{\mathbf{x}} \sin F). \quad (15.15)$$

The true localized orbit is found before any family index is assigned. If $A \in SU(2)_V$ fixes U_F , then at a radius with $\sin F \neq 0$, $A(\boldsymbol{\tau} \cdot \hat{\mathbf{x}})A^{-1} = \boldsymbol{\tau} \cdot \hat{\mathbf{x}}$ for every direction. Hence A commutes with all three Pauli matrices and

$$\text{Stab}_{SU(2)_V}(U_F) = \mathbb{Z}_2, \quad \mathcal{M}_{\text{rot}} = SU(2)/\mathbb{Z}_2 \simeq SO(3), \quad (15.16)$$

not $\mathbb{CP}^1$. Quotienting combined space–isospin rotations by the diagonal hedgehog stabilizer gives the same $SO(3)$, not a second orbit. The separate scalar-vacuum $\mathbb{CP}^1$ is spatially uniform, with

$$\|\delta_m \phi\|_{[-L, L]^3}^2 = 8L^3 |\delta_m \phi|^2, \quad (15.17)$$

so it is a bulk Goldstone direction rather than a normalizable soliton modulus.

At infinity Eq. (15.15) has a constant invertible mass. Its positive boundary eigenbundle on S_∞^2 is therefore trivial,

$$c_1(E_+) = 0, \quad \text{ind}_{\text{Callias}} = 0. \quad (15.18)$$

This static statement does not contradict spectral flow along a four-dimensional interpolation; spectral flow equal to B does not force an endpoint zero mode.

There is also a representation-level obstruction to the former two-band template. A globally nondegenerate 2×2 Hermitian mass with one positive band on $X = S_\infty^2 \times \mathbb{CP}^1$ defines $f : X \rightarrow \mathbb{CP}^1$, so its eigenline satisfies

$$c_1(E_+)^2 = f^*(u^2) = 0. \quad (15.19)$$

The desired class instead obeys

$$c_1(E_+) = x + 2y, \quad c_1(E_+)^2 = 4xy \neq 0. \quad (15.20)$$

Thus it requires at least three trivial bands, a nontrivial ambient bundle, or a genuinely infinite-dimensional family, followed by a new uniform-gap proof. On the actual $SO(3)$, $H^2(SO(3); \mathbb{Z}) = \mathbb{Z}_2$, so only a torsion line, not a free $\mathcal{O}(2)$ class, is available.

Finally, the correctly typed universal target character is

$$\widehat{\Xi}_5 = 2 \text{pr}_U^* \widehat{\omega}_3 \smile \text{pr}_s^* \widehat{c}_1(H) \in \widehat{H}^5(SU(2) \times \mathbb{CP}^1; \mathbb{Z}). \quad (15.21)$$

Only after pulling it back by the evaluation map may one form

$$\int_{\Sigma_3} \text{ev}^* \widehat{\Xi}_5 \in \widehat{H}^2(\mathcal{C}_{\text{restricted}}; \mathbb{Z}). \quad (15.22)$$

This establishes a typed local ansatz. It does not yet construct the full equivariant differential cocycle, prove all vertical and large-gauge holonomies, extend through Higgs zeroes and defects, or pull an $\mathcal{O}(2)$ line back to a physical same-soliton $\mathbb{CP}^1$. The finite CPT determinant identity is likewise an algebraic conjugation control, not a microscopic global regulator.

15.4 AP-E6 decision

With the stronger definitions forced by this round, let

$$\begin{aligned}
C_{Sp} &= (\text{complete regulator}) \wedge (\text{both target eta signs}) \\
&\quad \wedge (\text{full heavy APS match}) \wedge (\text{strong phase}), \\
C_{B1} &= (\text{same stationary lattice solution}) \wedge (B_a \rightarrow 1) \\
&\quad \wedge (\text{full projected super-Hessian}) \wedge (\text{4d dynamics}), \\
C_{\text{Cal}} &= (\text{physical moduli}) \wedge (\text{actual gapped family}) \\
&\quad \wedge (\text{CPT determinant line}) \wedge (\text{gauge-basic descent}).
\end{aligned}$$

The exact gauge-bordism subgate, continuum BVP, sparse differentiation, and moduli/two-band no-go are genuine advances, but

$$C_{Sp} = C_{B1} = C_{\text{Cal}} = \text{false}, \quad \boxed{\text{portal not started.}} \quad (15.23)$$

The canonical standalone records are `route_f/tex/ap_e6_sp4_eta_bordism_threshold.tex`, `route_f/tex/ap_e6_relaxed_b1_multigrid_hessian.tex`, and `route_f/tex/ap_e6_same_soliton_yukawa_callias.tex`, together with their machine-readable cards and the independent AP-E6 frontier verifier.

16 AP-E7 regulator, discrete topology, and determinant-family audit

This section supersedes the AP-E6 execution status while retaining every negative boundary proved there. It executes the three requested alternatives in order. The result is deliberately asymmetric: the common quadratic regulator, a pure heavy threshold, the finite-site topology theorem, the classification of the physical $SO(3)$ collective-coordinate lines, and a minimal rank-three mass family are exact. None of them by itself closes a same-model pre-portal physics lane.

16.1 Common quadratic APS/PV complex and its exact scope

For a smooth background $\bar{X} = (\bar{A}, \bar{\Sigma}, \bar{\Phi})$, let the infinitesimal gauge action be

$$\mathcal{R}_{\bar{X}} \epsilon = (D_{\bar{A}} \epsilon, [\bar{\Sigma}, \epsilon], -\epsilon \bar{\Phi}). \quad (16.1)$$

Using background Feynman gauge, the complete bare quadratic boson/ghost operators are defined without deleting mixing terms:

$$\Delta_B = S_{\text{bos}}^{(2)}[\bar{X}] + \mathcal{R}_{\bar{X}} \mathcal{R}_{\bar{X}}^\dagger, \quad \Delta_{\text{gh}} = \mathcal{R}_{\bar{X}}^\dagger \mathcal{R}_{\bar{X}}. \quad (16.2)$$

They act on the stated H^2 Sobolev domains after background stabilizers are projected. The fermion amplitude is built from $H = \gamma_5 \mathcal{K}$. Spinor charge conjugation is retained only as a reference/scalar-mass algebraic control; no fixed-Higgs Kramers symmetry is assumed for a general adjoint mass.

The common determinant prescription uses

$$c_a = (1, -3, 3, -1), \quad \Lambda_a^2 = a \Lambda^2, \quad a = 0, 1, 2, 3, \quad (16.3)$$

for which

$$\sum_a c_a a^r = 0 \quad (r = 0, 1, 2), \quad \sum_a c_a a^3 = -6, \quad \sum_a c_a \log(\lambda + a\Lambda^2) = -\frac{2\Lambda^6}{\lambda^3} + O(\lambda^{-4}). \quad (16.4)$$

At the level of zeta determinants the regulated quadratic amplitude is

$$\begin{aligned} Z_{\text{quad}}^{\text{PV}}(\Lambda) &= \tau_F \prod_{a=0}^3 \det'_{\theta_B}(\Delta_B + a\Lambda^2)^{-c_a/2} \det'(\Delta_{\text{gh}} + a\Lambda^2)^{c_a} \\ &\quad \times \det'(H^2 + a\Lambda^2)^{N_f c_a/2}, \quad N_f = 2. \end{aligned} \quad (16.5)$$

Here $\theta_B = \pi - \varepsilon$ is the upper lateral cut at a saddle. Equation (16.5) is a determinant prescription; it is not a claim that every half-integral power has already been realized by a local second-order Grassmann PV field. The finite part subtracts the integrated supercomplex A_4 coefficient at a fixed scale and declares positive reference masses to have zero local gauge, Hopf, and gravitational theta angles.

On $Y_5 = M_4 \times [0, 1]$, the interpolation operator and its literal APS domain are

$$\begin{aligned} \mathcal{D}_5^+ &= \partial_t + H(t), \\ \text{Dom}_{\text{APS},0}(\mathcal{D}_5^+) &= \{u \in H^1(Y_5) : P_{[0,\infty)}(H_0)u|_0 = 0, \quad P_{(-\infty,0)}(H_1)u|_1 = 0\}. \end{aligned} \quad (16.6)$$

The physical determinant-sign formula uses the fixed cut 0 and gapped endpoints:

$$\tau_F[X(t)] = \exp(i\pi N_f \text{SF}_0\{H(t)\}). \quad (16.7)$$

An arbitrary common resolvent cut still defines a Fredholm APS problem, but its spectral flow cannot replace Eq. (16.7) without the corresponding endpoint spectral-section orientation term. This distinction is part of the regulator, not a convention that may be discarded.

The constant negative-mass heavy bundle is

$$E_h = (V_c \otimes L^{-1}) \oplus \mathbf{1}, \quad x = c_1(L), \quad (16.8)$$

so the one-flavour index is

$$I_h = \int_{M_4} \widehat{A}(TM_4) \text{ch}(E_h)|_4 = \int_{M_4} \left[-c_2(V_c) + x^2 - \frac{1}{8}p_1(TM_4) \right] \in \mathbb{Z}. \quad (16.9)$$

The rank-two charged branch contributes $-p_1/12$, and the neutral singlet contributes $-p_1/24$; hence gravity has not been omitted. For two identical Dirac flavours,

$$\boxed{\tau_h^{\text{pure}} = \exp(i\pi 2I_h) = 1} \quad (16.10)$$

on every closed spin four-manifold and every unbroken $SU(2)_c \times U(1)_H$ bundle. This closes the pure unbroken-gauge/gravity threshold, not a target-dependent mixed WZW determinant.

On the separated Higgs orbit, one can consistently declare

$$h = \widehat{\Sigma}_5/V, \quad P_{\pm} = \frac{h^2 \pm h}{2}, \quad P_0 = 1 - h^2, \quad (\text{rank } P_+, \text{rank } P_-, \text{rank } P_0) = (2, 2, 1). \quad (16.11)$$

A GW-compatible target source built from these projectors has even copies on both target generators and therefore conditionally yields $(+1, +1)$ in the no-stacked-sector convention. It is not

promoted to a finite overlap Dirac–Yukawa theorem until the ordinary/GW chiral projectors, five-dimensional kernel, and determinant orientation are specified in one operator.

There is a sharp reason this source does not restore the original deep-UV claim. Suppose an exact separated rank-two projector $C(\Sigma)$ were continuous and $Sp(4)$ -equivariant through $\Sigma = 0$. Since the complex $\mathbf{5}$ is irreducible, Schur’s lemma gives

$$C(0) = c\mathbf{1}_5, \quad \text{Tr}(HC(0)) = 0, \quad \text{Tr}(HP_+) = 2. \quad (16.12)$$

If C remains a projector, $c = 0$ or 1 , so its rank is zero or five, never two. Thus the exact branch projector must lose its gap, continuity, or equivariance, or acquire new UV data. This theorem does not prohibit every non-idempotent equivariant coupling; it prohibits the exact separated rank-two selector needed by the proposed light-only family. A uniform exact flavour action on the complete $\mathbf{5}$ instead has

$$\kappa_{\text{full}} = 2(+1) + 2(-1) + 0 = 0, \quad \kappa_{\text{light}} = +2, \quad \kappa_{\text{heavy}} = -2. \quad (16.13)$$

Therefore neither construction is yet an all-scale derivation of $k = +2$.

16.2 Finite-site topology and the re-relaxation result

The fixed-boundary finite-site configuration space of the declared naive $n + s$ action is

$$\mathcal{C}_N = (S^3)^{(N-2)^3} \times \mathbb{R}^{(N-2)^3}. \quad (16.14)$$

Both factors are path-connected and a finite product of path-connected spaces is path-connected. Hence no continuous integer-valued function on all of $\mathcal{C}_N$ can distinguish an exact $B = 1$ component from the vacuum. A finite- a topological sector requires an admissibility restriction, topology-preserving interpolation, or infinite dislocation barrier; it is not supplied by the naive sampled action.

For diagnostics, each cube is split into the six Freudenthal tetrahedra. For a generic target $q \in S^3$, let

$$V_T = [n_{x_0} \ n_{x_1} \ n_{x_2} \ n_{x_3}], \quad c_T = V_T^{-1}q \quad \text{for } \det V_T \neq 0. \quad (16.15)$$

For a singular tetrahedron, $\text{span } V_T$ is a proper subspace of $\mathbb{R}^4$. The generic regular target is chosen outside the finite union of these spans, so a singular cell has no preimage and contributes zero; an exceptional target is equivalently handled by an arbitrarily small regular perturbation. The numerical implementation treats $|\det V_T| < 10^{-13}$ as exceptional and separately verifies three-target agreement and a nonzero-determinant margin for every counted hit. The normalized affine tetrahedron hits q exactly when all entries of c_T are positive, and its signed contribution is

$$d_T(q) = \text{sgn}(p) \text{sgn } \det V_T, \quad B_{\text{geom}} = - \sum_T d_T(q). \quad (16.16)$$

Three regular targets must agree. The admissibility margin is

$$\delta_a = \min_T \min_{\lambda_i \geq 0, \sum_i \lambda_i = 1} \left\| \sum_{i=0}^3 \lambda_i n_{x_i} \right\|. \quad (16.17)$$

Only on $\delta_a > 0$ is the normalized interpolation nonsingular and the degree locally constant.

Five independent full-grid relaxations give:

N	L	a	E_{initial}	$E_\star$	B_{geom}
7	6	1.00	43.95516	9.07×10^{-16}	0
9	6	0.75	54.15076	5.33×10^{-17}	0
11	6	0.60	61.09804	8.07×10^{-16}	0
9	8	1.00	43.16126	1.77×10^{-14}	0
11	10	1.00	42.97910	2.04×10^{-15}	0

The maximum final free projected-gradient density is 1.84×10^{-8} : the optimizer has converged, but to the vacuum. Pinning only $n(0) = -1$ also gives $B_{\text{geom}} = 0$ through a degenerate tetrahedron. Admissibility-guarded searches at $\delta_{\text{floor}} = 0.10, 0.05, 0.02$ retain $B_{\text{geom}} = 1$, but stop on the respective floor with gradient densities 0.473, 0.438, 0.418. They do not find an interior stationary point.

An explicitly artificial control fixes the central $3 \times 3 \times 3$ hedgehog n -values while re-relaxing all remaining variables independently on each grid. Its free constrained residual is below 6.2×10^{-8} , its three target degrees equal one, and its exact restricted $n + s$ Hessian has positive lowest eigenvalues 2.2643, 1.8043, 1.5002, 1.1611, 0.7563 on the five grids. The corresponding full unanchored residuals are 0.24–0.66; therefore these are restricted controls, not soliton stability eigenvalues.

On the same sites the audit also assembles a declared quadratic diquark block, free Dirichlet gauge and ghost Laplacians, and

$$K_F = D_W^\dagger D_W \succeq 0. \quad (16.18)$$

Their positive spectra are useful implementation checks. However K_F is not the second derivative of $-\log \det D_W$, the interacting gauge–meson cross blocks are absent, and a ghost operator is not an ordinary positive bosonic Hessian block. Consequently the direct sum is not a physical BRST super-Hessian. The exact finite-site conclusion is stronger and more negative: the naive action has no exact topological component, and its tested unconstrained minimizers unwind.

16.3 The actual $SO(3)$ cover and the rank-three alternative

For the true localized orbit $SO(3) \simeq \mathbb{RP}^3$, the cellular boundary maps are 0, 2, 0. Dualizing gives

$$H^2(SO(3); \mathbb{Z}) = \mathbb{Z}_2 = \{0, t\}, \quad t = \beta(a), \quad 2t = 0, \quad (16.19)$$

where $a \in H^1(SO(3); \mathbb{Z}_2)$ is the cover class. The unique nontrivial complex line is the FR line

$$\mathcal{L}_{\text{FR}} = SU(2) \times_{\mathbb{Z}_2, \text{sign}} \mathbb{C}, \quad (A, z) \sim (-A, -z). \quad (16.20)$$

A generator of $\pi_1(SO(3))$ lifts from $A = \mathbf{1}$ to $A = -\mathbf{1}$, so

$$\text{Hol}_\gamma(\mathcal{L}_{\text{FR}}) = -1, \quad \text{Hol}_\gamma(\mathcal{L}_{\text{FR}}^{\otimes 2}) = +1. \quad (16.21)$$

For a real skew-adjoint family,

$$\text{Hol}_\gamma(\text{Pf}) = (-1)^{\text{ind}_2(\mathcal{D}_{Y_\gamma})}, \quad \text{Det} \simeq \text{Pf}^{\otimes 2}. \quad (16.22)$$

Topology therefore classifies the two candidates but does not compute the actual same-Yukawa mapping-torus mod-two index. The standard conditional Skyrmin rule gives

$$(-1)^{N_c B} = (-1)^{2 \cdot 1} = +1, \quad (16.23)$$

and thus favours the trivial/bosonic sector, but it is not substituted for the missing operator computation. Inversion preserves the torsion class and admits the topological Real lift

$$J[A, z] = [A^{-1}, \bar{z}], \quad J^2 = 1, \quad (16.24)$$

while the microscopic CPT action on spin, gauge, regulator, domain, and Pfaffian orientation remains open.

The torsion carrier cannot reproduce the old $\mathcal{O}(2)$ mechanism: $t \otimes \mathbb{R} = 0$ and a flat representative has zero curvature, whereas $c_1(\mathcal{O}(2))$ is a nonzero free class. Moreover the lowest odd FR rotor block has $j = 1/2$ and full Peter–Weyl multiplicity four, not the three holomorphic sections of $\mathcal{O}(2)$.

The independent rank-three branch does remove the AP-E6 two-band obstruction at the bundle level. On $X = \mathbb{CP}_z^1 \times \mathbb{CP}_w^1$, set

$$s = (z_0 w_0^2, z_1 w_1^2, z_0 w_1^2 + z_1 w_0^2). \quad (16.25)$$

These three sections of $\mathcal{O}(1, 2)$ have no common zero. With

$$e_+ = \frac{\bar{s}}{\|s\|}, \quad P_+ = e_+ e_+^\dagger, \quad \mathcal{M} = m(2P_+ - \mathbf{1}_3), \quad m > 0, \quad (16.26)$$

one has the exact global identities

$$\begin{aligned} \text{spec } \mathcal{M} &= \{+m, -m, -m\}, & \mathcal{M}^2 &= m^2 \mathbf{1}_3, \\ c_1(E_+) &= x + 2y, & c_1(E_-) &= -(x + 2y), & c_2(E_-) &= 4xy. \end{aligned} \quad (16.27)$$

Thus $E_+ \oplus E_- = X \times \mathbb{C}^3$, and rank three is minimal because the $\mathbb{CP}^2$ hyperplane class can have nonzero square while the $\mathbb{CP}^1$ two-band class cannot. A scan of 11,664 points gives $\min \|s\|^2 = 0.25$, projector residual 5.64×10^{-16} , spectral residual 2.89×10^{-15} , exact gap $m = 1.7$, and discrete Berry Chern numbers (1.0000000000000003, 1.9999999999999998).

This is a genuine topological mass-family theorem, but its second $\mathbb{CP}_w^1$ factor is still a spectator rather than a localized normalizable modulus. No charged two-colour representation has yet produced its three bands or the bidegree-(1, 2) Yukawa projector, and the finite matrix gap is not a uniform Fredholm gap of the same soliton operator.

16.4 AP-E7 decision

Let C_{APS} require the common regulator, the original target-family lift, both actual target eta signs, the mixed heavy determinant, and all-scale $k = +2$. Let C_{B1} require an unanchored admissible stationary sequence, the interacting physical super-Hessian, and a dynamical continuum limit. Let C_{fam} require either the actual $SO(3)$ Pfaffian route or a physical rank-three same-soliton family, together with microscopic CPT and gauge-basic WZW descent. The current result is

$$C_{\text{APS}} = C_{B1} = C_{\text{fam}} = \text{false}, \quad \boxed{\text{degree-one Route-E portal not started.}} \quad (16.28)$$

The mathematical advances are retained as reusable lemmas; they are not promotion substitutes. The canonical records are the three `route_f/tex/ap_e7_*` notes, their machine-readable cards, and the independent AP-E7 execution-frontier verifier.

17 Reproducibility ledger

The independent arithmetic/symbolic card is

```
route_f/code/verify_another_physics_route_e_bridge.py.
```

Run from the repository root:

```
python3 route_f/code/verify_another_physics_route_e_bridge.py
```

It writes

```
route_f/output/another_physics_route_e_bridge.json,
route_f/output/another_physics_route_e_bridge.md.
```

The current result is 27/27 mechanical checks, including

$$\max |K_{\text{tr}}^2 - I/3| = 1.11 \times 10^{-16}, \quad |2\Delta - B| = 0, \quad (17.1)$$

zero generator-invariance and canonical factor-two residuals, exact agreement between the loaded Route-E ζ card and the declared benchmark, complex weight-two covariance error 2.80×10^{-17} , the two independent Q-ball energy estimates in Sec. 6, and oscillatory-level-set residuals below 2.7×10^{-14} . The three earlier successor cards pass 18/18, 38/38, and 27/27, respectively. The new completion-frontier cards pass 24/24, 58/58, and 27/27, while their integrated authorization gate passes 20/20. The AP-E6 follow-up cards pass 39/39, 20/20, and 25/25; their fresh-manifest, same-checksum frontier conjunction passes 15/15 while keeping every complete-lane boolean false. The AP-E7 regulator, discrete-topology, and $SO(3)$ /rank-three cards pass 42/42, 16/16, and 18/18, respectively. Their independent same-checksum execution-frontier conjunction passes 16/16, with the two family alternatives represented by a disjunction but every condition within each alternative retained conjunctively. All complete-lane, portal, and promotion booleans remain false. The JSON explicitly retains all cross-theory identifications as non-derived bridge axioms and sets `physics_promotion_allowed=false`.

A Units and normalization checks

In $\hbar = c = 1$ units, $[\Phi] = [f] = [m] = [M] = 1$, $[U] = 4$, $[\sigma] = 3$, $[R] = -1$, while Q, λ, η are dimensionless. Therefore every term in (6.2) has mass dimension four and every term in (6.16) has dimension one. For (8.5), g and ζ_{eff} are dimensionless and Λ has dimension one.

The Q-ball charge convention differs by factors of two across the literature depending on whether $\Phi = \phi e^{i\omega t}$ or $\Phi = \phi e^{i\omega t}/\sqrt{2}$. Equations (6.4)–(6.16) use one convention consistently; mixing conventions would change Q , f_0 , and R factors and is forbidden in the verifier.

B Why a phase portal needs a second reference

Suppose $\zeta_{\text{eff}} = c\Phi^2$ and define the putative invariant $I = \zeta_{\text{eff}}^* \Phi^2 / (|\zeta_{\text{eff}}| |\Phi|^2)$. Then

$$I = \frac{c^* \Phi^* \Phi^2}{|c| |\Phi|^4} = \frac{c^*}{|c|}, \quad (\text{B.1})$$

which is constant. The field cannot serve as its own phase reference. A nontrivial observable requires a second spurion or matrix structure, such as M_V in $M_R = M_V + \zeta_{\text{eff}} K_{\text{tr}}$. If that reference does not transform, it explicitly breaks the Q-ball $U(1)$; if it does transform, one must solve the coupled background and identify an invariant relative phase. This algebra is the sharp reason that phase doubling alone is not a visibility mechanism.

C Minimal falsification criteria

The bridge program should be abandoned or revised if any of the following occurs:

1. the reduced bubble moduli space is not $\mathbb{CP}^1$, or its physical bundle has Chern number other than two;
2. the zero-mode operator has extra unpaired charged modes or does not realize $H^0(T)$;
3. the exact fixed-charge soliton has a negative Hessian mode or E/Q above an allowed charge carrier;
4. every phase in $M_V + \zeta K_{\text{tr}}$ is removable, leaving no invariant quantity such as (7.10);
5. the required explicit breaking makes the soliton lifetime shorter than the intended physical process;
6. a valid selection rule cannot forbid XLH and dangerous operators;
7. the embedded model fails Route-E flavor, threshold, proton-decay, or cosmology gates;
8. a gravity carrier violates the redshift relation (2.5), lensing, or fifth-force bounds.

References

- [1] Xin Zheng, Jonathan Dolde, Matthew C. Cambria, Hong Ming Lim, and Shimon Kolkowitz. A lab-based test of the gravitational redshift with a miniature clock network. *Nature Communications*, 14:4886, 2023.
- [2] Tobias Mistele, Stacy McGaugh, Federico Lelli, James Schombert, and Pengfei Li. Indefinitely flat circular velocities and the baryonic tully–fisher relation from weak lensing. *The Astrophysical Journal Letters*, 969:L3, 2024.
- [3] Sidney Coleman. Q-balls. *Nuclear Physics B*, 262(2):263–283, 1985.
- [4] Yu Hamada, Kiyoharu Kawana, TaeHun Kim, and Philip Lu. Q-balls in the presence of attractive force. *Journal of High Energy Physics*, 2024(8):242, 2024.
- [5] Don N. Page and William K. Wootters. Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, 27:2885–2892, 1983.
- [6] Dario Cafasso, Nicola Pranzini, Jorge Yago Malo, Vittorio Giovannetti, and Marilù Chiofalo. Quantum time and the time-dilation induced interaction transfer mechanism. *Physical Review D*, 110(10):106014, 2024.
- [7] Ted Jacobson. Thermodynamics of spacetime: The einstein equation of state. *Physical Review Letters*, 75:1260–1263, 1995.
- [8] Ted Jacobson. Entanglement equilibrium and the einstein equation. *Physical Review Letters*, 116:201101, 2016.