

A Spin^c Projection-Dolbeault Derivation of Three Standard Model Generations in PSLT

Bo-Yu Chen¹

¹*Independent Researcher*

(Dated: May 1, 2026)

We formulate the Projection Spectral Layer Theory (PSLT) as a first-principles projection-spectral derivation of the three-generation multiplicity of the Standard Model chiral family representation, conditional on the PSLT two-center projection data. The compact projection surface is $C = \mathbb{CP}^1$, the two projection branch points define the divisor $R = p_+ + p_-$, and the generation line bundle is $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$. The internal fermion operator is specified to be the Spin^c Dolbeault–Dirac operator

$$\mathcal{D}_R = \sqrt{2} \left(\bar{\partial}_{L_R} + \bar{\partial}_{L_R}^\dagger \right)$$

acting on $\Lambda^{0,*} T^* C \otimes L_R$. Hodge theory gives

$$\text{Ker } \mathcal{D}_R^+ \cong H^0(C, L_R), \quad \text{Ker } \mathcal{D}_R^- \cong H^1(C, L_R).$$

For $C = \mathbb{CP}^1$ and $L_R = \mathcal{O}(2)$, Riemann–Roch and Serre duality give

$$h^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3, \quad h^1(\mathbb{CP}^1, \mathcal{O}(2)) = 0.$$

Provided the SM-charged chiral carrier is exactly this Spin^c generation sector and all other SM-charged internal sectors are absent from the kernel or fully paired and gapped, the low-energy four-dimensional chiral spectrum contains exactly three Standard Model families and no unpaired light SM-charged exotics. The same two-center geometry supplies the conformal factor Ω from a variational Poisson constraint, the scalar spectral operator $[-\nabla^2 + V_{\text{eff}}]\psi_N = \omega_N^2 \psi_N$, and the scalar spectral data used in an effective formation/visibility probability law

$$P_N(t) = \frac{B_N g_N (1 - e^{-\Gamma_N t})}{\sum_{K=1}^3 B_K g_K (1 - e^{-\Gamma_K t})}.$$

Thus the generation count is a Spin^c index-theoretic consequence of the two-center projection divisor, while (g_N, Γ_N, B_N) describe degeneracy, kinetics, and visible-sector overlap of the three protected modes.

CONTENTS

I. Core Structure	3
II. Parent Projection Data	3

	2
A. SM-charged fermion sector	4
III. One Standard Model Family	5
IV. Spin^c Dolbeault–Dirac Zero Modes	6
V. Three-Generation Theorem	7
VI. Canonical Generation Basis	8
VII. Two-Center Conformal Geometry	9
VIII. Geometry-to-Spectrum Map	10
IX. Degeneracy, Kinetics, Visibility, and Formation	11
X. Conclusion	12
Acknowledgments	13
References	13

I. CORE STRUCTURE

The construction has three logically distinct layers:

1. the *SM charge layer*, where one minimal Standard Model family is fixed by gauge invariance, Yukawa invariance, and anomaly cancellation;
2. the *internal fermion layer*, where the Spin^c Dolbeault–Dirac operator on the projection surface supplies the chiral zero-mode multiplicity;
3. the *formation and visibility layer*, where the two-center conformal geometry determines spectral rates and visible-sector overlaps.

The core Hilbert-space statement is

$$\boxed{\mathcal{H}_{\text{chiral}}^{4\text{D}} = \mathcal{R}_{\text{SM}} \otimes \text{Ker } \mathcal{D}_R^+ = \mathcal{R}_{\text{SM}} \otimes H^0(\mathbb{CP}^1, \mathcal{O}(p_+ + p_-))}. \quad (1)$$

Since $p_+ + p_-$ has degree two,

$$\mathcal{O}(p_+ + p_-) \simeq \mathcal{O}(2), \quad \dim H^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3. \quad (2)$$

The low-energy chiral spectrum is therefore

$$\boxed{\mathcal{H}_{\text{chiral}}^{4\text{D}} = \mathcal{R}_{\text{SM}}^{(1)} \oplus \mathcal{R}_{\text{SM}}^{(2)} \oplus \mathcal{R}_{\text{SM}}^{(3)}}. \quad (3)$$

The family count is the dimension of the positive internal Spin^c fermion zero-mode space.

II. PARENT PROJECTION DATA

Let M_4 be four-dimensional spacetime and let the compact projection surface be

$$C \simeq \mathbb{CP}^1. \quad (4)$$

The two projection branch points define the divisor

$$R = p_+ + p_-, \quad L_R = \mathcal{O}(R) \simeq \mathcal{O}(2). \quad (5)$$

The parent action is organized as

$$S_{\text{FP}} = S_{\text{grav}} + S_{\text{gauge}} + S_{\text{Higgs}} + S_{\text{fermion}} + S_{\Omega} + S_{\text{proj}} + S_{\text{vis}}. \quad (6)$$

The projection sector fixes the topological class of L_R :

$$S_{\text{proj}} = \int_C \lambda_R \left[\frac{i}{2\pi} F_{A_R} - \delta_{p_+}^{(2)} - \delta_{p_-}^{(2)} \right], \quad (7)$$

so that

$$c_1(L_R) = [p_+] + [p_-], \quad \deg L_R = 2. \quad (8)$$

A. SM-charged fermion sector

The SM-charged internal fermion carrier is assumed to be exactly

$$\mathcal{E}_{\text{gen}} = \mathcal{R}_{\text{SM}} \otimes \mathcal{S}_C, \quad \mathcal{S}_C = \Lambda^{0,*} T^* C \otimes L_R. \quad (9)$$

Let

$$X = M_4 \times C. \quad (10)$$

The generation-sector product Dirac operator is

$$\boxed{\mathcal{D}_X = \mathcal{D}_{M_4}^{\text{SM}} \otimes 1 + \gamma_5 \otimes \mathcal{D}_R, \quad \Gamma_X = \gamma_5 \otimes \Gamma_C.} \quad (11)$$

Here $\mathcal{D}_{M_4}^{\text{SM}}$ includes the four-dimensional SM gauge connection on $\mathcal{R}_{\text{SM}}$, Γ_C is the internal Spin^c chirality, and Γ_X is the product chirality. The generation-sector fermion action is

$$\boxed{S_{\text{fermion}}^{\text{gen}} = \int_{M_4 \times C} \text{dvol}_{M_4} \text{dvol}_C \bar{\Psi} i \mathcal{D}_X \Psi.} \quad (12)$$

The internal mode expansion is

$$\Psi(x, y) = \sum_{\alpha} \psi_{\alpha}^{+}(x) u_{\alpha}^{+}(y) + \sum_{\beta} \psi_{\beta}^{-}(x) u_{\beta}^{-}(y) + \sum_{\lambda > 0} \Psi_{\lambda}(x) \chi_{\lambda}(y), \quad (13)$$

where

$$\mathcal{D}_R u_{\alpha}^{+} = 0, \quad u_{\alpha}^{+} \in \text{Ker } \mathcal{D}_R^{+}, \quad \mathcal{D}_R u_{\beta}^{-} = 0, \quad u_{\beta}^{-} \in \text{Ker } \mathcal{D}_R^{-}. \quad (14)$$

After a fixed product-chirality projection, internal positive and negative zero modes produce opposite four-dimensional Weyl chiralities. We choose the product chirality so that $\text{Ker } \mathcal{D}_R^{+}$ carries the left-handed SM family representation $\mathcal{R}_{\text{SM}}$; $\text{Ker } \mathcal{D}_R^{-}$ would supply the opposite four-dimensional chirality in the same generation sector. Thus the chain is explicit:

$$\text{internal } \text{Spin}^c \text{ zero mode} \implies \text{massless 4D Weyl fermion} \implies \text{SM chiral generation.} \quad (15)$$

All other SM-charged internal sectors are required to be vectorlike and genuinely gapped:

$$\mathcal{E}_{\text{fermion}}^{\text{SM}} = \mathcal{E}_{\text{gen}} \oplus \mathcal{E}_{\text{vec}}. \quad (16)$$

The condition is stronger than $\text{ind } \mathcal{D}_{\text{vec}} = 0$. It requires either

$$\text{Ker } \mathcal{D}_{\text{vec}} = 0, \quad (17)$$

or

$$\begin{aligned} \dim \text{Ker } \mathcal{D}_{\text{vec}}^+ &= \dim \text{Ker } \mathcal{D}_{\text{vec}}^-, \\ M_{\text{vec}} : \text{Ker } \mathcal{D}_{\text{vec}}^+ &\longrightarrow \text{Ker } \mathcal{D}_{\text{vec}}^-, \quad M_{\text{vec}} \text{ is full rank and gauge invariant.} \end{aligned} \quad (18)$$

Index zero only excludes extra net chirality. Equation (17) or Eq. (18) is the additional low-energy condition that excludes unpaired massless SM-charged vectorlike exotics. SM-neutral sectors may be present, but they do not carry generation quantum numbers:

$$\mathcal{E}_{\text{fermion}} = \mathcal{E}_{\text{gen}} \oplus \mathcal{E}_{\text{vec}} \oplus \mathcal{E}_{\text{neutral}}. \quad (19)$$

This is the exclusivity condition needed for an exact no-fourth-generation theorem. Without it, an additional SM-charged internal bundle with nonzero index could produce extra chiral matter.

III. ONE STANDARD MODEL FAMILY

The visible gauge group is

$$G_{\text{SM}} = SU(3)_c \times SU(2)_L \times U(1)_Y. \quad (20)$$

For one left-handed Weyl family, take

$$Q : (3, 2)_{Y_Q}, \quad u^c : (\bar{3}, 1)_{Y_u}, \quad d^c : (\bar{3}, 1)_{Y_d}, \quad L : (1, 2)_{Y_L}, \quad e^c : (1, 1)_{Y_e}, \quad (21)$$

with Higgs

$$H : (1, 2)_h. \quad (22)$$

Set $Y_Q = q$. Gauge invariance of the Yukawa couplings gives

$$QH u^c : \quad q + h + Y_u = 0, \quad QH^\dagger d^c : \quad q - h + Y_d = 0, \quad (23)$$

$$LH^\dagger e^c : \quad Y_L - h + Y_e = 0. \quad (24)$$

Hence

$$Y_u = -q - h, \quad Y_d = -q + h, \quad Y_e = h - Y_L. \quad (25)$$

The $SU(3)^2 U(1)$ anomaly gives

$$2Y_Q + Y_u + Y_d = 0, \quad (26)$$

which is automatic after Eq. (25). The $SU(2)^2 U(1)$ anomaly gives

$$3Y_Q + Y_L = 0, \quad Y_L = -3q. \quad (27)$$

The mixed gravitational anomaly requires

$$6Y_Q + 3Y_u + 3Y_d + 2Y_L + Y_e = 0. \quad (28)$$

Substitution gives

$$6q + 3(-q - h) + 3(-q + h) + 2(-3q) + (h + 3q) = 0, \quad (29)$$

so

$$h = 3q. \quad (30)$$

Therefore

$$Y_u = -4q, \quad Y_d = 2q, \quad Y_L = -3q, \quad Y_e = 6q, \quad Y_H = 3q. \quad (31)$$

The $U(1)^3$ anomaly cancels:

$$6q^3 + 3(-4q)^3 + 3(2q)^3 + 2(-3q)^3 + (6q)^3 = 0. \quad (32)$$

With $q = 1/6$,

$$\boxed{(Y_Q, Y_u, Y_d, Y_L, Y_e, Y_H) = \left(\frac{1}{6}, -\frac{2}{3}, \frac{1}{3}, -\frac{1}{2}, 1, \frac{1}{2}\right)}. \quad (33)$$

A sterile $\nu^c : (1, 1)_0$ may be added without changing these anomaly equations.

Proposition 1. *Within the minimal one-family chiral field content above, and assuming the Yukawa couplings $QHuc$, $QH^\dagger d^c$, and $LH^\dagger e^c$, the hypercharge pattern is fixed up to the overall normalization of $U(1)_Y$.*

IV. SPIN^c DOLBEAULT–DIRAC ZERO MODES

The internal fermion operator is not the ordinary spin Dirac operator. It is the Spin^c Dolbeault–Dirac operator

$$\boxed{\mathcal{D}_R = \sqrt{2} \left(\bar{\partial}_{L_R} + \bar{\partial}_{L_R}^\dagger \right)} \quad (34)$$

acting on

$$\mathcal{S}_C = \Lambda^{0,*} T^* C \otimes L_R. \quad (35)$$

We use the canonical Spin^c structure on the complex curve C , whose untwisted spinor bundle is $\Lambda^{0,*} T^* C$. The line bundle L_R is an auxiliary twisting bundle for the Dolbeault–Dirac operator, not the Spin^c determinant line itself. The relevant index is therefore the Dolbeault index

$$\text{ind } \mathcal{D}_R = \chi(C, L_R) = h^0(C, L_R) - h^1(C, L_R). \quad (36)$$

Its internal chirality decomposition is

$$\mathcal{S}_C^+ = \Lambda^{0,0} T^* C \otimes L_R, \quad \mathcal{S}_C^- = \Lambda^{0,1} T^* C \otimes L_R. \quad (37)$$

By Hodge theory,

$$\boxed{\text{Ker } \mathcal{D}_R^+ \cong H^0(C, L_R), \quad \text{Ker } \mathcal{D}_R^- \cong H^1(C, L_R).} \quad (38)$$

This is the step that converts a holomorphic-section count into an internal fermion zero-mode theorem.

Remark 1. The Spin^c choice is essential for the stated bundle $L_R = \mathcal{O}(2)$. For an ordinary spin Dirac operator on $\mathbb{CP}^1$, the spin bundle contributes $K_C^{1/2} = \mathcal{O}(-1)$. Twisting an ordinary spin Dirac operator only by $\mathcal{O}(2)$ would count

$$H^0(\mathbb{CP}^1, K_C^{1/2} \otimes \mathcal{O}(2)) = H^0(\mathbb{CP}^1, \mathcal{O}(1)),$$

which is two-dimensional. Equivalently, an ordinary spin-Dirac presentation would need the twist

$$E = K_C^{-1/2} \otimes L_R = \mathcal{O}(1) \otimes \mathcal{O}(2) = \mathcal{O}(3)$$

so that $K_C^{1/2} \otimes E = L_R$.

V. THREE-GENERATION THEOREM

Theorem 1 (PSLT Spin^c three-generation theorem). *Let $C = \mathbb{CP}^1$, $R = p_+ + p_-$, and $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$. Let the SM-charged generation carrier be*

$$\mathcal{R}_{\text{SM}} \otimes (\Lambda^{0,*} T^* C \otimes L_R),$$

with internal operator $\mathcal{D}_R$ as in Eq. (34). Assume the product Dirac action is Eq. (12), the product chirality is Eq. (11), every other SM-charged internal sector satisfies the full vectorlike gap condition of Eq. (17) or Eq. (18), and all remaining chiral sectors are SM-neutral. Then the low-energy four-dimensional chiral SM spectrum contains exactly three generations:

$$N_{\text{gen}} = \dim \text{Ker } \mathcal{D}_R^+ - \dim \text{Ker } \mathcal{D}_R^- = 3. \quad (39)$$

Moreover, since $\text{Ker } \mathcal{D}_R^- = 0$, the positive internal zero-mode space itself is three-dimensional.

Proof. Riemann–Roch for a line bundle L on a compact Riemann surface gives

$$h^0(C, L) - h^1(C, L) = \deg L + 1 - g(C). \quad (40)$$

For $C = \mathbb{CP}^1$, $g(C) = 0$. Because $R = p_+ + p_-$, $\deg L_R = 2$. Therefore

$$h^0(C, L_R) - h^1(C, L_R) = 3. \quad (41)$$

Serre duality gives

$$h^1(C, L_R) = h^0(C, K_C \otimes L_R^{-1}). \quad (42)$$

On $\mathbb{CP}^1$,

$$K_C = \mathcal{O}(-2), \quad L_R^{-1} = \mathcal{O}(-2), \quad (43)$$

so

$$K_C \otimes L_R^{-1} = \mathcal{O}(-4). \quad (44)$$

A negative-degree line bundle on $\mathbb{CP}^1$ has no nonzero holomorphic sections. Thus

$$h^1(C, L_R) = 0, \quad h^0(C, L_R) = 3. \quad (45)$$

Using Eq. (38),

$$\dim \text{Ker } \mathcal{D}_R^+ = 3, \quad \dim \text{Ker } \mathcal{D}_R^- = 0. \quad (46)$$

The product Dirac reduction in Eqs. (11)–(14) turns these internal zero modes into massless four-dimensional Weyl fermions. The vectorlike gap condition prevents additional SM-charged chiral generations and also removes unpaired massless SM-charged vectorlike exotics. $\square$

In an affine coordinate ζ on $\mathbb{CP}^1$,

$$H^0(\mathbb{CP}^1, \mathcal{O}(2)) = \text{span}\{1, \zeta, \zeta^2\}. \quad (47)$$

This span is only a vector space until a physical basis is chosen. The family count is basis-independent; the labels $N = 1, 2, 3$ require a canonical Hermitian operator.

VI. CANONICAL GENERATION BASIS

Let P_R be the Bergman projector from $L^2(C, L_R)$ onto $H^0(C, L_R)$, using the Fubini–Study metric on $C = \mathbb{CP}^1$ and the induced Hermitian metric on L_R . Choose a Möbius coordinate ζ_R adapted to the two-center divisor, so that the two branch points are placed at 0 and ∞ . The associated axial moment-map function is

$$\mu_R(\zeta_R, \bar{\zeta}_R) = \frac{|\zeta_R|^2 - 1}{|\zeta_R|^2 + 1}. \quad (48)$$

Define the finite-dimensional Toeplitz operator

$$\boxed{\mathcal{T}_R = P_R M_{\mu_R} P_R} \quad (49)$$

on $H^0(C, L_R)$, where M_{μ_R} denotes multiplication by μ_R . The generation basis is defined by

$$\mathcal{T}_R u_N = \tau_N u_N, \quad u_N \in H^0(C, L_R), \quad N = 1, 2, 3. \quad (50)$$

For the symmetric two-center placement and the standard Fubini–Study/Bergman inner product, this basis is equivalent, up to normalization and ordering, to the spin-one monomial basis

$$1, \quad \zeta, \quad \zeta^2. \quad (51)$$

The Toeplitz definition is the physical statement; the monomial basis is only a convenient coordinate representation.

VII. TWO-CENTER CONFORMAL GEOMETRY

The two-center conformal factor is obtained from

$$S_\Omega = \frac{\kappa}{2} \int_{\mathbb{R}^3} d^3x |\nabla \Omega|^2 - 4\pi\kappa \int_{\mathbb{R}^3} d^3x \sigma(x)(\Omega - 1), \quad (52)$$

with

$$\sigma(x) = a [\rho_\varepsilon(x - x_+) + \rho_\varepsilon(x - x_-)]. \quad (53)$$

Variation gives

$$\delta S_\Omega = -\kappa \int_{\mathbb{R}^3} d^3x (\nabla^2 \Omega + 4\pi\sigma) \delta\Omega, \quad (54)$$

hence

$$\boxed{\nabla^2 \Omega = -4\pi\sigma.} \quad (55)$$

Let $(\rho, z_{\text{ax}}, \varphi)$ be cylindrical coordinates on $\mathbb{R}^3$, with centers at

$$x_\pm : \quad \rho = 0, \quad z_{\text{ax}} = \pm D/2. \quad (56)$$

For

$$\rho_\varepsilon(r) = \frac{3\varepsilon^2}{4\pi(r^2 + \varepsilon^2)^{5/2}}, \quad \int_{\mathbb{R}^3} d^3x \rho_\varepsilon = 1, \quad (57)$$

the asymptotically flat solution is

$$\boxed{\Omega(\rho, z_{\text{ax}}; D) = 1 + a \left(\frac{1}{r_+} + \frac{1}{r_-} \right), \quad r_\pm = \sqrt{\rho^2 + (z_{\text{ax}} \mp D/2)^2 + \varepsilon^2}.} \quad (58)$$

VIII. GEOMETRY-TO-SPECTRUM MAP

We use the mostly-plus convention

$$\eta_{\mu\nu} = \text{diag}(-, +, +, +), \quad \square_\eta = -\partial_t^2 + \nabla^2, \quad (59)$$

and the curvature convention

$$R^\rho{}_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}. \quad (60)$$

For the static conformally flat metric

$$g_{\mu\nu} = \Omega^2(x) \eta_{\mu\nu}, \quad (61)$$

this gives

$$R = -6\Omega^{-3}\nabla^2\Omega. \quad (62)$$

Consider

$$(\square_g - m_0^2 - \xi R)\Phi = 0, \quad \Phi(t, x) = e^{-i\omega t} \Omega^{-1}(x) \psi(x). \quad (63)$$

Using

$$\left(\square_g - \frac{1}{6}R \right) (\Omega^{-1}\psi) = \Omega^{-3} \square_\eta \psi, \quad (64)$$

one obtains

$$[-\nabla^2 + m_0^2 \Omega^2 + (1 - 6\xi) \Omega^{-1} \nabla^2 \Omega] \psi = \omega^2 \psi. \quad (65)$$

Define

$$V_{\text{eff}} = m_0^2 \Omega^2 + (1 - 6\xi) \Omega^{-1} \nabla^2 \Omega. \quad (66)$$

Then

$$\boxed{[-\nabla^2 + V_{\text{eff}}] \psi_n = \omega_n^2 \psi_n.} \quad (67)$$

This scalar spectral problem governs formation barriers and visible overlap profiles. It does not determine the number of chiral SM generations; that number is fixed by $\mathcal{D}_R$.

IX. DEGENERACY, KINETICS, VISIBILITY, AND FORMATION

For the protected chiral sector, let $\Pi_N = |u_N\rangle\langle u_N|$ be the projector onto the Toeplitz eigenmode u_N . The chiral degeneracy is

$$g_N^{\text{chiral}} = \text{Tr}_{\mathcal{H}_{\text{gen}}} \Pi_N, \quad \mathcal{H}_{\text{gen}} = H^0(\mathbb{CP}^1, \mathcal{O}(2)). \quad (68)$$

Thus

$$\boxed{g_1 = g_2 = g_3 = 1, \quad g_{N \geq 4}^{\text{chiral}} = 0.} \quad (69)$$

For kinetics, the two centers define a collective-coordinate subspace

$$\mathcal{C}_N = \text{span}\{|N, +\rangle, |N, -\rangle\}. \quad (70)$$

Let P_N^{cc} project onto $\mathcal{C}_N$, $Q_N^{\text{cc}} = 1 - P_N^{\text{cc}}$, and let $\mathcal{K}_N$ be the Euclidean fluctuation operator in this sector. The Feshbach–Schur effective matrix is

$$\mathcal{M}_N = P_N^{\text{cc}} \mathcal{K}_N P_N^{\text{cc}} - P_N^{\text{cc}} \mathcal{K}_N Q_N^{\text{cc}} (Q_N^{\text{cc}} \mathcal{K}_N Q_N^{\text{cc}})^{-1} Q_N^{\text{cc}} \mathcal{K}_N P_N^{\text{cc}}. \quad (71)$$

In the two-center basis,

$$\mathcal{M}_N = \begin{pmatrix} \gamma_{N,+} & \chi_N \sqrt{\gamma_{N,+} \gamma_{N,-}} \\ \chi_N \sqrt{\gamma_{N,+} \gamma_{N,-}} & \gamma_{N,-} \end{pmatrix}, \quad (72)$$

with WKB channel factors

$$\gamma_{N,\pm} = A_{N,\pm} \exp[-2S_{N,\pm}^{\text{WKB}}], \quad S_{N,\pm}^{\text{WKB}} = \int_{\mathcal{B}_{\pm}} ds \sqrt{V_{\text{eff}}(s) - \omega_N^2}. \quad (73)$$

The formation rate is

$$\boxed{\Gamma_N(D, \eta) = \max\{0, \lambda_+(\mathcal{M}_N(D, \eta))\}}, \quad (74)$$

where

$$\lambda_+(\mathcal{M}_N) = \frac{\gamma_{N,+} + \gamma_{N,-}}{2} + \frac{1}{2} \sqrt{(\gamma_{N,+} - \gamma_{N,-})^2 + 4\chi_N^2 \gamma_{N,+} \gamma_{N,-}}. \quad (75)$$

For visibility, let $\pi_D : \mathbb{R}^3 \rightarrow C$ denote the projection map from the spatial two-center chart to the compact projection surface, and let

$$\hat{u}_N(\rho, z_{\text{ax}}; D) = u_N(\pi_D(\rho, z_{\text{ax}})) \quad (76)$$

with the frame/Jacobian factors included in W_{frame} . The conformal Dirac and Higgs fields scale as

$$\Psi = \Omega^{-3/2} \psi, \quad H = \Omega^{-1} \phi. \quad (77)$$

Hence

$$\sqrt{-g} \bar{\Psi} H \Psi \propto \Omega^4 \cdot \Omega^{-3/2} \cdot \Omega^{-1} \cdot \Omega^{-3/2} = \Omega^0. \quad (78)$$

A mode-resolved Yukawa overlap may therefore be written

$$y_N^{\text{eff}}(D) = y_0 \int 2\pi\rho \, d\rho \, dz_{\text{ax}} \hat{u}_N(\rho, z_{\text{ax}}; D) f_L(\rho, z_{\text{ax}}) f_R(\rho, z_{\text{ax}}) W_{\text{frame}}. \quad (79)$$

The normalized visible weight is

$$B_N = \frac{|y_N^{\text{eff}}|^2}{\sum_{I=1}^3 |y_I^{\text{eff}}|^2}. \quad (80)$$

Equivalently,

$$B_N = \frac{\langle u_N | \mathcal{O}_{\text{vis}}^\dagger \mathcal{O}_{\text{vis}} | u_N \rangle}{\sum_{I=1}^3 \langle u_I | \mathcal{O}_{\text{vis}}^\dagger \mathcal{O}_{\text{vis}} | u_I \rangle}. \quad (81)$$

Finally, the formation law

$$\frac{dp_N}{dt} = \Gamma_N(1 - p_N), \quad p_N(0) = 0 \quad (82)$$

has solution

$$p_N(t) = 1 - e^{-\Gamma_N t}. \quad (83)$$

Conditioning on the observation of one of the protected chiral layers gives

$$P_N(t; D, \eta) = \frac{B_N g_N (1 - e^{-\Gamma_N(D, \eta)t})}{\sum_{K=1}^3 B_K g_K (1 - e^{-\Gamma_K(D, \eta)t})}. \quad (84)$$

X. CONCLUSION

The two-center PSLT projection data determine the divisor $R = p_+ + p_-$ and the line bundle $L_R = \mathcal{O}(R) = \mathcal{O}(2)$ on $C = \mathbb{CP}^1$. The SM-charged internal fermions are governed by the product Dirac action on $M_4 \times C$, with the internal Spin^c Dolbeault–Dirac operator acting on $\Lambda^{0,*} T^* C \otimes L_R$. The bundle L_R is the auxiliary Dolbeault twist, and the positive and negative internal zero modes are $H^0(C, L_R)$ and $H^1(C, L_R)$. Riemann–Roch gives index three, and Serre duality gives $H^1(C, L_R) = 0$. With all additional SM-charged sectors either absent from the kernel or fully paired by a gauge-invariant mass map, the theory therefore has exactly three low-energy chiral SM generations and no unpaired light SM-charged exotics.

The scalar two-center spectral operator and the quantities $(g_N, \Gamma_N, B_N, P_N)$ then describe how these three protected modes form and couple to the visible sector. They are phenomenological structures on top of the Spin^c generation theorem, not substitutes for it.

ACKNOWLEDGMENTS

The author acknowledges the mathematical foundations of Riemann surface theory, Spin^c Dirac operators, index theory, and Standard Model anomaly cancellation used in this work.

-
- [1] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics Vol. 52, Springer (1977).
 - [2] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley (1978).
 - [3] O. Forster, *Lectures on Riemann Surfaces*, Graduate Texts in Mathematics Vol. 81, Springer (1981).
 - [4] H. B. Lawson and M.-L. Michelsohn, *Spin Geometry*, Princeton University Press (1989).
 - [5] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., Institute of Physics Publishing (2003).
 - [6] R. A. Bertlmann, *Anomalies in Quantum Field Theory*, Oxford University Press (1996).
 - [7] A. Bilal, “Lectures on Anomalies,” arXiv:0802.0634.
 - [8] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press (1982).
 - [9] H. Feshbach, “Unified theory of nuclear reactions,” *Annals Phys.* **5**, 357 (1958).
 - [10] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press (1978).