

A Conditional Spin^c Dolbeault Index Theorem for Three Standard Model Family Copies in a Two-Center Projection Model

Bo-Yu Chen¹

¹*Independent Researcher*

(Dated: May 2, 2026)

Abstract

We formulate a conditional Spin^c index-theoretic derivation of the three-family multiplicity of the Standard Model chiral representation in Projection Spectral Layer Theory (PSLT). The input is the PSLT two-center projection datum: the compact projection surface $C = \mathbb{CP}^1$, the branch divisor $R = p_+ + p_-$, the generation line bundle $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$, and the protected SM-charged carrier $\mathcal{R}_{\text{SM}} \otimes (\Lambda^{0,*} T^* C \otimes L_R)$. On this carrier the internal fermion operator is the Spin^c Dolbeault–Dirac operator

$$\mathcal{D}_R = \sqrt{2} \left(\bar{\partial}_{L_R} + \bar{\partial}_{L_R}^\dagger \right)$$

and Hodge theory gives

$$\text{Ker } \mathcal{D}_R^+ \cong H^0(C, L_R), \quad \text{Ker } \mathcal{D}_R^- \cong H^1(C, L_R).$$

For $C = \mathbb{CP}^1$ and $L_R = \mathcal{O}(2)$, Riemann–Roch and Serre duality give

$$h^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3, \quad h^1(\mathbb{CP}^1, \mathcal{O}(2)) = 0.$$

Provided all other SM-charged internal sectors are absent from the kernel or fully paired and gapped, the protected SM-charged generation sector contributes exactly three unpaired SM family copies to the four-dimensional low-energy chiral spectrum. The novelty is not the Riemann–Roch computation itself, but the identification of the two-center projection divisor as the generation line bundle for the protected SM-charged Spin^c sector. Scalar spectral data and the quantities $(g_N^{\text{chiral}}, \Gamma_N^{\text{eff}}, B_N)$ are kept as a post-index formation/visibility interface: they assign profiles, rates, and visible overlaps to the three protected modes, but they do not determine the family count.

I. INTRODUCTION AND CORE STRUCTURE

The Standard Model contains three chiral copies of the same anomaly-free family representation. This paper addresses only the multiplicity problem inside a conditional PSLT projection model: given a two-center projection surface, a generation line bundle, and a protected Spin^c carrier for SM-charged fermions, it asks whether the number of protected chiral family copies is fixed. It does not derive the two-center projection data from a UV-complete quantum-gravity vacuum, and it does not attempt to explain Yukawa hierarchies, CKM or PMNS mixing, or the detailed mass spectrum.

Index-theoretic generation counting is a standard tool in field theory and compactification problems [1, 5]. Related cohomological or topological mechanisms for chiral matter appear

in heterotic, brane, and F-theory constructions [10–15]. The specific claim here is narrower: within the PSLT protected generation sector, the two-center branch divisor $R = p_+ + p_-$ is identified with the generation line bundle $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$, so the SM family multiplicity is the Spin^c Dolbeault index of $(\mathbb{CP}^1, \mathcal{O}(2))$. The novelty is therefore the physical identification of the PSLT two-center projection divisor as the generation line bundle, not the elementary Riemann–Roch calculation $h^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3$ by itself.

The construction has three logically distinct layers:

1. the *SM charge layer*, where, under the minimal one-family field-content and Yukawa assumptions, the Standard Model hypercharge pattern is fixed by gauge invariance and anomaly cancellation;
2. the *internal fermion layer*, where the Spin^c Dolbeault–Dirac operator on the projection surface supplies the chiral zero-mode multiplicity;
3. the *formation and visibility layer*, where the two-center conformal geometry supplies scalar spectral data for effective rates and visible-sector overlaps.

The data needed for the Spin^c generation-multiplicity theorem are only

$$\boxed{\mathfrak{D}_{\text{gen}} = (C, R, L_R, \mathscr{D}_R).} \quad (1)$$

They fix $\text{Ker } \mathscr{D}_R^+$, hence the number of protected chiral generations. The effective formation/visibility layer uses additional interface data

$$\boxed{\mathfrak{D}_{\text{int}}(D) = (\pi_D, \mathfrak{j}_D, w_D, \mathcal{J}_{\text{vis}}).} \quad (2)$$

These interface data assign scalar profiles, rates, and visible overlaps to the protected modes; they are not inputs to the index-theoretic statement $N_{\text{gen}} = 3$. The core Hilbert-space statement is

$$\boxed{\mathcal{H}_{\text{chiral}}^{4\text{D}} = \mathcal{R}_{\text{SM}} \otimes \text{Ker } \mathscr{D}_R^+ = \mathcal{R}_{\text{SM}} \otimes H^0(\mathbb{CP}^1, \mathcal{O}(p_+ + p_-)).} \quad (3)$$

Since $p_+ + p_-$ has degree two,

$$\mathcal{O}(p_+ + p_-) \simeq \mathcal{O}(2), \quad \dim H^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3. \quad (4)$$

The low-energy chiral spectrum is therefore

$$\boxed{\mathcal{H}_{\text{chiral}}^{4\text{D}} = \mathcal{R}_{\text{SM}}^{(1)} \oplus \mathcal{R}_{\text{SM}}^{(2)} \oplus \mathcal{R}_{\text{SM}}^{(3)}} \quad (5)$$

The family count is the dimension of the positive internal Spin^c fermion zero-mode space.




FIG. 1. Logical architecture of the PSLT Spin^c generation theorem. The Standard Model charge layer fixes one anomaly-free chiral family representation $\mathcal{R}_{\text{SM}}$. The protected internal fermion layer uses the two-center projection data $C = \mathbb{CP}^1$, $R = p_+ + p_-$, $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$, and the Spin^c Dolbeault–Dirac operator $\mathscr{D}_R$ to prove the family multiplicity through $\text{Ker } \mathscr{D}_R^+ \simeq H^0(C, L_R)$. The scalar spectral, kinetic, and visibility data form a post-index effective interface and are not inputs to the theorem $N_{\text{gen}} = 3$.

A. Notation conventions

Throughout the paper, D denotes the two-center separation parameter. Dirac-type operators are written with script letters, such as $\mathscr{D}_R$, $\mathscr{D}_X$, $\mathscr{D}_{\text{vec}}$, and $\mathscr{D}_{M_4}^{\text{SM}}$. The Bergman projector onto $H^0(C, L_R)$ is P_R , the collective-coordinate projector is P_N^{cc} , the one-dimensional Toeplitz-mode projector is $\Pi_N = |u_N\rangle\langle u_N|$, and the observed layer probability is $P_N(t; D, \eta)$. The protected zero-mode degeneracy is denoted g_N^{chiral} ; any unadorned degeneracy symbol is avoided in the chiral family count.

II. TWO-CENTER PROJECTION DATA AND TOPOLOGICAL CONSTRAINT

Let M_4 be four-dimensional spacetime and let the compact projection surface be

$$C \simeq \mathbb{CP}^1. \quad (6)$$

The two projection branch points define the divisor

$$R = p_+ + p_-, \quad L_R = \mathcal{O}(R) \simeq \mathcal{O}(2). \quad (7)$$

For bookkeeping we denote a possible effective decomposition by

$$S_{\text{eff}} = S_{\text{grav}} + S_{\text{gauge}} + S_{\text{Higgs}} + S_{\text{fermion}} + S_{\Omega}^{\text{cov}} + S_{\text{proj}} + S_{\text{vis}}. \quad (8)$$

TABLE I. Assumption boundary and theorem scope. The theorem proves the three-family multiplicity inside the protected PSLT Spin^c generation sector. Effective scalar spectral quantities and formation/visibility weights are post-index interface data and are not used to prove $N_{\text{gen}} = 3$.

Object or claim	Status in this paper	Role
$C = \mathbb{CP}^1$	PSLT two-center projection	Input
	datum	
$R = p_+ + p_-$	PSLT branch divisor	Input
$L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$	Determined by R on $\mathbb{CP}^1$	Generation line bundle
$\Lambda^{0,*}T^*C \otimes L_R$	Assumed protected carrier	Spin^c internal fermion sector
$\mathcal{D}_R$	Specified Spin^c	Counts protected zero modes
	Dolbeault–Dirac operator	
$h^0 = 3, h^1 = 0$	Theorem consequence	Three protected chiral copies
Vectorlike gap for other	Assumption	Excludes extra light
SM-charged sectors		SM-charged exotics
$\Omega, V_{\text{eff}}, \Gamma_N^{\text{eff}}, B_N, P_N$	Effective post-index	Formation/visibility, not
	interface	family-count proof
UV derivation of	Not proven here	Future strengthening
(C, R, L_R)		

No UV-complete parent action is derived in this paper. The theorem uses only the projection data and the cohomological constraint below. The projection sector fixes the topological class of L_R , not a pointwise equality between a smooth curvature form and a delta-current. The robust condition is the cohomological constraint

$$\left[\frac{i}{2\pi} F_{A_R} \right] = \text{PD}(R) = [p_+] + [p_-] \in H^2(C, \mathbb{Z}), \quad R = p_+ + p_-. \quad (9)$$

Equivalently,

$$c_1(L_R) = \text{PD}(R), \quad \deg L_R = 2. \quad (10)$$

When an action-like expression is useful, it should be read as a topological constraint or as a regulated current limit:

$$S_{\text{proj}}^{\text{top}} = \int_C \lambda_R \left[\frac{i}{2\pi} F_{A_R} - J_R^{(\varepsilon)} \right], \quad (11)$$




FIG. 2. Two-center projection geometry and the generation divisor. The spatial chart contains two ordered centers $x_{\pm}$ at $z_{\text{ax}} = \pm D/2$, with regulated distances $r_{\pm}$. The projection map π_D sends the regulated two-center chart to $C \simeq \mathbb{CP}^1$, and in the collapsed-core limit maps the centers to the branch points p_+ and p_- . The divisor $R = p_+ + p_-$ has degree two and defines the generation line bundle $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$. The midplane normalization $z_{\text{ax}} = 0 \Rightarrow |\zeta_R| = 1$ fixes the residual scale of the affine coordinate.

where $J_R^{(\varepsilon)}$ is any smooth representative of $\text{PD}(R)$ that tends, as a current, to $\delta_{p_+}^{(2)} + \delta_{p_-}^{(2)}$. The topological input is Eq. (9); Eq. (11) is only a regulated implementation of the same cohomology class.

A. SM-charged fermion sector

The SM-charged internal fermion carrier is assumed to be exactly

$$\mathcal{E}_{\text{gen}} = \mathcal{R}_{\text{SM}} \otimes \mathcal{S}_C, \quad \mathcal{S}_C = \Lambda^{0,*} T^* C \otimes L_R. \quad (12)$$

Let

$$X = M_4 \times C. \quad (13)$$

The generation-sector product Dirac operator is

$$\mathcal{D}_X = \mathcal{D}_{M_4}^{\text{SM}} \otimes 1 + \gamma_5 \otimes \mathcal{D}_R, \quad \Gamma_X = \gamma_5 \otimes \Gamma_C. \quad (14)$$

Here $\mathcal{D}_{M_4}^{\text{SM}}$ includes the four-dimensional SM gauge connection on $\mathcal{R}_{\text{SM}}$, Γ_C is the internal Spin^c chirality, and Γ_X is the product chirality. The generation-sector fermion action is

$$S_{\text{fermion}}^{\text{gen}} = \int_{M_4 \times C} \text{dvol}_{M_4} \text{dvol}_C \bar{\Psi} i \mathcal{D}_X \Psi. \quad (15)$$

The internal mode expansion is

$$\Psi(x, y) = \sum_{\alpha} \psi_{\alpha}^{+}(x) u_{\alpha}^{+}(y) + \sum_{\beta} \psi_{\beta}^{-}(x) u_{\beta}^{-}(y) + \sum_{\lambda > 0} \Psi_{\lambda}(x) \chi_{\lambda}(y), \quad (16)$$

where

$$\mathcal{D}_R u_{\alpha}^{+} = 0, \quad u_{\alpha}^{+} \in \text{Ker } \mathcal{D}_R^{+}, \quad \mathcal{D}_R u_{\beta}^{-} = 0, \quad u_{\beta}^{-} \in \text{Ker } \mathcal{D}_R^{-}. \quad (17)$$

After a fixed product-chirality projection, internal positive and negative zero modes produce opposite four-dimensional Weyl chiralities. We choose the product chirality so that $\text{Ker } \mathcal{D}_R^{+}$ carries the left-handed SM family representation $\mathcal{R}_{\text{SM}}$; $\text{Ker } \mathcal{D}_R^{-}$ would supply the opposite four-dimensional chirality in the same generation sector. Thus the chain is explicit:

$$\begin{aligned} \text{internal Spin}^c \text{ zero mode} &\implies \text{massless 4D Weyl fermion} \\ &\implies \text{SM chiral generation.} \end{aligned} \quad (18)$$

All other SM-charged internal sectors are required to be vectorlike and genuinely gapped:

$$\mathcal{E}_{\text{fermion}}^{\text{SM}} = \mathcal{E}_{\text{gen}} \oplus \mathcal{E}_{\text{vec}}. \quad (19)$$

The condition is stronger than $\text{ind } \mathcal{D}_{\text{vec}} = 0$. It requires either

$$\text{Ker } \mathcal{D}_{\text{vec}} = 0, \quad (20)$$

or

$$\begin{aligned} \dim \text{Ker } \mathcal{D}_{\text{vec}}^{+} &= \dim \text{Ker } \mathcal{D}_{\text{vec}}^{-}, \\ M_{\text{vec}} : \text{Ker } \mathcal{D}_{\text{vec}}^{+} &\longrightarrow \text{Ker } \mathcal{D}_{\text{vec}}^{-}, \quad M_{\text{vec}} \text{ is full rank and gauge invariant.} \end{aligned} \quad (21)$$

Index zero only excludes extra net chirality. Equation (20) or Eq. (21) is the additional low-energy condition that excludes unpaired massless SM-charged vectorlike exotics. SM-neutral sectors may be present, but they do not carry generation quantum numbers:

$$\mathcal{E}_{\text{fermion}} = \mathcal{E}_{\text{gen}} \oplus \mathcal{E}_{\text{vec}} \oplus \mathcal{E}_{\text{neutral}}. \quad (22)$$

This is the exclusivity condition needed for the conditional no-fourth-family statement within the protected SM-charged sector. Without it, the Spin^c theorem proves only the three-dimensionality of the protected generation sector, not the absence of additional SM-charged chiral sectors.

III. ONE STANDARD MODEL FAMILY

The local anomaly equations use the Standard Model Lie algebra, with standard anomaly conventions [7, 8]. When a global group and charge lattice are needed, we take

$$G_{\text{SM}} = \frac{SU(3)_c \times SU(2)_L \times U(1)_Y}{\mathbb{Z}_6}. \quad (23)$$

This global choice fixes the conventional hypercharge unit used below; without it, the local equations determine hypercharge only up to an overall $U(1)_Y$ normalization. For one left-handed Weyl family, take

$$Q : (3, 2)_{Y_Q}, \quad u^c : (\bar{3}, 1)_{Y_u}, \quad d^c : (\bar{3}, 1)_{Y_d}, \quad L : (1, 2)_{Y_L}, \quad e^c : (1, 1)_{Y_e}, \quad (24)$$

with Higgs

$$H : (1, 2)_h. \quad (25)$$

Set $Y_Q = q$. Gauge invariance of the Yukawa couplings gives

$$QH u^c : \quad q + h + Y_u = 0, \quad QH^\dagger d^c : \quad q - h + Y_d = 0, \quad (26)$$

$$LH^\dagger e^c : \quad Y_L - h + Y_e = 0. \quad (27)$$

Hence

$$Y_u = -q - h, \quad Y_d = -q + h, \quad Y_e = h - Y_L. \quad (28)$$

The $SU(3)^2 U(1)$ anomaly gives

$$2Y_Q + Y_u + Y_d = 0, \quad (29)$$

which is automatic after Eq. (28). The pure color anomaly also cancels:

$$\mathcal{A}_{SU(3)^3} = 2A(3) + A(\bar{3}) + A(\bar{3}) = 2A(3) - A(3) - A(3) = 0, \quad (30)$$

where the factor of two comes from the two $SU(2)_L$ components of Q , and $A(\bar{3}) = -A(3)$.

The $SU(2)^2 U(1)$ anomaly gives

$$3Y_Q + Y_L = 0, \quad Y_L = -3q. \quad (31)$$

The mixed gravitational anomaly requires

$$6Y_Q + 3Y_u + 3Y_d + 2Y_L + Y_e = 0. \quad (32)$$

Substitution gives

$$6q + 3(-q - h) + 3(-q + h) + 2(-3q) + (h + 3q) = 0, \quad (33)$$

so

$$h = 3q. \quad (34)$$

Therefore

$$Y_u = -4q, \quad Y_d = 2q, \quad Y_L = -3q, \quad Y_e = 6q, \quad Y_H = 3q. \quad (35)$$

The $U(1)^3$ anomaly cancels:

$$6q^3 + 3(-4q)^3 + 3(2q)^3 + 2(-3q)^3 + (6q)^3 = 0. \quad (36)$$

There is no Witten global $SU(2)$ anomaly [9]. One family contains

$$3 \text{ color copies of } Q + 1 \text{ lepton doublet} = 4 \quad (37)$$

left-handed $SU(2)_L$ Weyl doublets, which is even. For three families the number is 12, still even. With $q = 1/6$,

$$\boxed{(Y_Q, Y_u, Y_d, Y_L, Y_e, Y_H) = \left(\frac{1}{6}, -\frac{2}{3}, \frac{1}{3}, -\frac{1}{2}, 1, \frac{1}{2}\right)}. \quad (38)$$

A sterile $\nu^c : (1, 1)_0$ may be added without changing these anomaly equations.

Proposition 1. *Within the minimal one-family chiral field content above, and assuming the Yukawa couplings $QHuc$, $QH^\dagger d^c$, and $LH^\dagger e^c$, the local anomaly equations fix the hypercharge pattern up to the overall normalization of $U(1)_Y$. The choice in Eq. (23) fixes the conventional unit $q = 1/6$.*

This proposition fixes the one-family representation that is later tensored with the Spin^c zero-mode space; it does not determine the number of families.

IV. SPIN^c DOLBEAULT–DIRAC ZERO MODES

The internal fermion operator is not the ordinary spin Dirac operator. It is the Spin^c Dolbeault–Dirac operator

$$\boxed{\mathcal{D}_R = \sqrt{2} \left(\bar{\partial}_{L_R} + \bar{\partial}_{L_R}^\dagger \right)} \quad (39)$$

acting on

$$\mathcal{S}_C = \Lambda^{0,*} T^* C \otimes L_R. \quad (40)$$

Remark 1 (BPS-compatible origin of the Dolbeault representative). In a BPS or worldline-supersymmetric realization of this protected generation sector, massless states are represented by Q -cohomology classes, with

$$Q^2 = 0, \quad H = \{Q, Q^\dagger\}. \quad (41)$$

For the holomorphic carrier

$$\mathcal{H}_{\text{int}} = \Omega^{0,*}(C, L_R), \quad (42)$$

locality, first-order gauge covariance, and compatibility with the complex structure select a Cauchy–Riemann type representative

$$Q = \bar{\partial}_{L_R} \quad (43)$$

up to holomorphic bundle isomorphism and gauge choice. The corresponding Hermitian Dirac representative is therefore

$$\mathcal{D}_R = \sqrt{2}(Q + Q^\dagger) = \sqrt{2}(\bar{\partial}_{L_R} + \bar{\partial}_{L_R}^\dagger). \quad (44)$$

This remark is only a BPS-compatible motivation for the protected PSLT sector used below; the theorem remains conditional on the two-center projection data, the holomorphic carrier $\Omega^{0,*}(C, L_R)$, and the absence of additional flux, superpotential, Ext-complex, or torsion deformations that would replace the Dolbeault differential by a different protected complex. It is not an unconditional vacuum-selection theorem for all string or quantum-gravity compactifications.

We use the canonical Spin^c structure on the complex curve C , whose untwisted spinor bundle is $\Lambda^{0,*}T^*C$. The line bundle L_R is an auxiliary twisting bundle for the Dolbeault–Dirac operator, not the Spin^c determinant line itself. The relevant index is therefore the Dolbeault index

$$\text{ind } \mathcal{D}_R = \chi(C, L_R) = h^0(C, L_R) - h^1(C, L_R). \quad (45)$$

Its internal chirality decomposition is

$$\mathcal{S}_C^+ = \Lambda^{0,0}T^*C \otimes L_R, \quad \mathcal{S}_C^- = \Lambda^{0,1}T^*C \otimes L_R. \quad (46)$$

By Hodge theory,

$$\boxed{\text{Ker } \mathcal{D}_R^+ \cong H^0(C, L_R), \quad \text{Ker } \mathcal{D}_R^- \cong H^1(C, L_R).} \quad (47)$$

This is the step that converts a holomorphic-section count into an internal fermion zero-mode theorem.

Remark 2. The Spin^c choice is essential for the stated bundle $L_R = \mathcal{O}(2)$. For an ordinary spin Dirac operator on $\mathbb{CP}^1$, the spin bundle contributes $K_C^{1/2} = \mathcal{O}(-1)$. Twisting an ordinary spin Dirac operator only by $\mathcal{O}(2)$ would count

$$H^0(\mathbb{CP}^1, K_C^{1/2} \otimes \mathcal{O}(2)) = H^0(\mathbb{CP}^1, \mathcal{O}(1)),$$

which is two-dimensional. Equivalently, an ordinary spin-Dirac presentation would need the twist

$$E = K_C^{-1/2} \otimes L_R = \mathcal{O}(1) \otimes \mathcal{O}(2) = \mathcal{O}(3)$$

so that $K_C^{1/2} \otimes E = L_R$.




FIG. 3. Spin^c index-theoretic origin of the three protected families. The two-center divisor fixes $L_R = \mathcal{O}(2)$. The protected internal fermion operator is the Spin^c Dolbeault–Dirac operator, whose positive and negative kernels are identified by Hodge theory with $H^0(C, L_R)$ and $H^1(C, L_R)$. For $C = \mathbb{CP}^1$, Riemann–Roch and Serre duality give $h^0 = 3$ and $h^1 = 0$, so the positive internal zero-mode space produces three four-dimensional SM chiral family copies.




FIG. 4. Why the Spin^c Dolbeault–Dirac representative is essential. The protected sector used here is the canonical Spin^c bundle $\Lambda^{0,*}T^*C$ twisted by $L_R = \mathcal{O}(2)$, and it counts $H^0(\mathbb{CP}^1, \mathcal{O}(2))$, giving three positive zero modes. An ordinary spin Dirac operator on $\mathbb{CP}^1$ includes $K_C^{1/2} = \mathcal{O}(-1)$; twisting it only by $\mathcal{O}(2)$ would count $H^0(\mathbb{CP}^1, \mathcal{O}(1))$, which is two-dimensional. Thus the Spin^c carrier is part of the stated protected sector, not an interchangeable notation.

V. THREE-GENERATION THEOREM

Remark 3 (Assumption boundary). The theorem below proves the multiplicity of unpaired SM-charged chiral zero modes inside the protected PSLT generation sector specified by $(C, R, L_R, \mathcal{D}_R)$. It does not derive the two-center projection data or the protected carrier from a UV-complete compactification. Those are model inputs; the statement proved here is the resulting Spin^c index theorem.

Theorem 1 (PSLT Spin^c three-generation theorem). *Let $C = \mathbb{CP}^1$, $R = p_+ + p_-$, and $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$. Let the SM-charged generation carrier be*

$$\mathcal{R}_{\text{SM}} \otimes (\Lambda^{0,*} T^* C \otimes L_R),$$

with internal operator $\mathcal{D}_R$ as in Eq. (39). Assume the product Dirac action is Eq. (15), the product chirality is Eq. (14), every other SM-charged internal sector satisfies the full vectorlike gap condition of Eq. (20) or Eq. (21), and all remaining chiral sectors are SM-neutral. Then the low-energy four-dimensional chiral SM spectrum contains exactly three unpaired SM-charged chiral generations:

$$N_{\text{gen}} = \dim \text{Ker } \mathcal{D}_R^+ - \dim \text{Ker } \mathcal{D}_R^- = 3. \quad (48)$$

Moreover, since $\text{Ker } \mathcal{D}_R^- = 0$, the positive internal zero-mode space itself is three-dimensional.

Proof. Riemann–Roch for a line bundle L on a compact Riemann surface gives [2–4]

$$h^0(C, L) - h^1(C, L) = \deg L + 1 - g(C). \quad (49)$$

For $C = \mathbb{CP}^1$, $g(C) = 0$. Because $R = p_+ + p_-$, $\deg L_R = 2$. Therefore

$$h^0(C, L_R) - h^1(C, L_R) = 3. \quad (50)$$

Serre duality gives

$$h^1(C, L_R) = h^0(C, K_C \otimes L_R^{-1}). \quad (51)$$

On $\mathbb{CP}^1$,

$$K_C = \mathcal{O}(-2), \quad L_R^{-1} = \mathcal{O}(-2), \quad (52)$$

so

$$K_C \otimes L_R^{-1} = \mathcal{O}(-4). \quad (53)$$

A negative-degree line bundle on $\mathbb{CP}^1$ has no nonzero holomorphic sections. Thus

$$h^1(C, L_R) = 0, \quad h^0(C, L_R) = 3. \quad (54)$$

Using Eq. (47),

$$\dim \text{Ker } \mathcal{D}_R^+ = 3, \quad \dim \text{Ker } \mathcal{D}_R^- = 0. \quad (55)$$

The product Dirac reduction in Eqs. (14)–(17) turns these internal zero modes into massless four-dimensional Weyl fermions. The stated exclusivity assumption prevents additional nonzero-index SM-charged generation sectors, while the vectorlike gap condition removes unpaired massless SM-charged vectorlike exotics. $\square$

In an affine coordinate ζ on $\mathbb{CP}^1$,

$$H^0(\mathbb{CP}^1, \mathcal{O}(2)) = \text{span}\{1, \zeta, \zeta^2\}. \quad (56)$$

This span is only a vector space until a physical basis is chosen. The family count is basis-independent; the labels $N = 1, 2, 3$ require a canonical Hermitian operator.

VI. CANONICAL GENERATION BASIS

The two branch points alone determine the divisor and the family count, but a physical basis also needs the normalization of the affine coordinate on C . Placing p_+ at 0 and p_- at ∞ leaves the residual freedom $\zeta_R \mapsto a\zeta_R$. In PSLT this freedom is fixed by the two-center projection map from the spatial chart to the compact projection surface.

A. Projection map and coordinate normalization

Let $(\rho, z_{\text{ax}}, \varphi)$ be cylindrical coordinates on $\mathbb{R}^3$, with the ordered centers

$$x_{\pm} : \quad \rho = 0, \quad z_{\text{ax}} = \pm D/2. \quad (57)$$

With regulator $\varepsilon > 0$, define

$$r_{\pm} = \sqrt{\rho^2 + (z_{\text{ax}} \mp D/2)^2 + \varepsilon^2}. \quad (58)$$

The physical affine coordinate is

$$\boxed{\zeta_R(\rho, z_{\text{ax}}, \varphi; D, \varepsilon) = \left(\frac{r_+}{r_-}\right)^{1/2} e^{i\varphi}.} \quad (59)$$

Strictly, Eq. (59) is a cylindrical-coordinate representative on $\mathbb{R}^3 \setminus A_{\text{ax}}$, where

$$A_{\text{ax}} := \{\rho = 0\}, \quad (60)$$

because the angular coordinate φ is undefined on the axis. The axial set has zero measure with respect to $d\mu_{\text{cyl}} = \rho d\rho dz_{\text{ax}} d\varphi$, so the pullback construction used in the visibility integrals may be read in an L^2 -almost-everywhere sense. Equivalently, one may smooth the axial phase and take the regularized limit. With this convention, we write the projection map as a map from the spatial chart:

$$\boxed{\pi_D : \mathbb{R}^3 \longrightarrow C \simeq \mathbb{CP}^1, \quad \pi_D(\rho, z_{\text{ax}}, \varphi) = [1 : \zeta_R(\rho, z_{\text{ax}}, \varphi; D, \varepsilon)].} \quad (61)$$

In the collapsed-core limit $\varepsilon \rightarrow 0$,

$$\pi_D(x_+) = p_+ = [1 : 0], \quad \pi_D(x_-) = p_- = [0 : 1]. \quad (62)$$

For finite ε , the source cores are mapped to regulated neighborhoods of the branch points. The normalization is fixed by the midplane condition

$$z_{\text{ax}} = 0 \implies |\zeta_R| = 1, \quad (63)$$

and the residual $U(1)$ phase is fixed by the origin of the axial angle φ . Thus the earlier scale freedom $\zeta_R \mapsto a\zeta_R$ is removed by the physical projection geometry rather than by an arbitrary Gram–Schmidt convention.

B. Toeplitz basis

Let P_R be the Bergman projector from $L^2(C, L_R)$ onto $H^0(C, L_R)$, using the Fubini–Study metric on $C = \mathbb{CP}^1$ and the induced Hermitian metric on L_R . Using the coordinate ζ_R fixed by Eq. (59), the associated axial moment-map function is

$$\mu_R(\zeta_R, \bar{\zeta}_R) = \frac{|\zeta_R|^2 - 1}{|\zeta_R|^2 + 1}. \quad (64)$$

Let M_{μ_R} denote multiplication by μ_R . Define the finite-dimensional Toeplitz operator on $H^0(C, L_R)$ by

$$\boxed{\mathcal{T}_R = P_R M_{\mu_R} P_R} \quad (65)$$

This is a finite-dimensional Toeplitz quantization of the axial moment-map observable [16]. The generation basis is defined by

$$\mathcal{T}_R u_N = \tau_N u_N, \quad u_N \in H^0(C, L_R), \quad N = 1, 2, 3. \quad (66)$$

For the symmetric two-center placement and the standard Fubini–Study/Bergman inner product, this basis is equivalent, up to normalization and ordering, to the spin-one monomial basis

$$1, \quad \zeta, \quad \zeta^2. \quad (67)$$

In this symmetric realization $\mathcal{T}_R$ is diagonal in the monomial basis and has three distinct axial weights; after the orientation of the ordered pair (p_+, p_-) is fixed, it canonically orders the three one-dimensional eigenspaces. The Toeplitz definition is the physical statement for labeling the already three-dimensional protected space; it is not a substitute for the Riemann–Roch and Serre-duality dimension argument. The monomial basis is only a convenient coordinate representation.




FIG. 5. Canonical family labels from the two-center Toeplitz operator. The family count $\dim H^0(\mathbb{CP}^1, \mathcal{O}(2)) = 3$ is basis-independent, but physical labels require a canonical Hermitian operator. The two-center projection fixes the affine coordinate ζ_R , hence the axial moment-map function μ_R . Projecting multiplication by μ_R onto the protected holomorphic section space gives the Toeplitz operator $\mathcal{T}_R = P_R M_{\mu_R} P_R$. Its eigenvectors define the canonical generation basis, equivalent for the symmetric two-center placement to the spin-one monomial basis $1, \zeta, \zeta^2$ up to normalization and ordering.

VII. TWO-CENTER CONFORMAL GEOMETRY

The two-center conformal factor is obtained in the static sector. The covariant effective term S_Ω^{cov} in Eq. (8) is not specified by the spatial expression below; rather, under the static two-center ansatz it reduces to the three-dimensional variational energy functional

$$E_\Omega = \frac{\kappa}{2} \int_{\mathbb{R}^3} d^3x |\nabla \Omega|^2 - 4\pi\kappa \int_{\mathbb{R}^3} d^3x \sigma(x)(\Omega - 1), \quad (68)$$

with

$$\sigma(x) = a [\rho_\varepsilon(x - x_+) + \rho_\varepsilon(x - x_-)]. \quad (69)$$

Variation gives

$$\delta E_\Omega = -\kappa \int_{\mathbb{R}^3} d^3x (\nabla^2 \Omega + 4\pi\sigma) \delta \Omega, \quad (70)$$

hence

$$\boxed{\nabla^2 \Omega = -4\pi\sigma.} \quad (71)$$

Let $(\rho, z_{\text{ax}}, \varphi)$ be cylindrical coordinates on $\mathbb{R}^3$, with centers at

$$x_\pm : \quad \rho = 0, \quad z_{\text{ax}} = \pm D/2. \quad (72)$$

For

$$\rho_\varepsilon(r) = \frac{3\varepsilon^2}{4\pi(r^2 + \varepsilon^2)^{5/2}}, \quad \int_{\mathbb{R}^3} d^3x \rho_\varepsilon = 1, \quad (73)$$

the asymptotically flat solution is

$$\Omega(\rho, z_{\text{ax}}; D) = 1 + a \left(\frac{1}{r_+} + \frac{1}{r_-} \right), \quad r_\pm = \sqrt{\rho^2 + (z_{\text{ax}} \mp D/2)^2 + \varepsilon^2}. \quad (74)$$

VIII. GEOMETRY-TO-SPECTRUM MAP

We use the mostly-plus convention [6, 17]

$$\eta_{\mu\nu} = \text{diag}(-, +, +, +), \quad \square_\eta = -\partial_t^2 + \nabla^2, \quad (75)$$

and the curvature convention

$$R^\rho{}_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}. \quad (76)$$

For the static conformally flat metric

$$g_{\mu\nu} = \Omega^2(x) \eta_{\mu\nu}, \quad (77)$$

this gives

$$R = -6\Omega^{-3}\nabla^2\Omega. \quad (78)$$

Consider

$$(\square_g - m_0^2 - \xi R)\Phi = 0, \quad \Phi(t, x) = e^{-i\omega t} \Omega^{-1}(x) \psi(x). \quad (79)$$

Using

$$\left(\square_g - \frac{1}{6}R \right) (\Omega^{-1}\psi) = \Omega^{-3} \square_\eta \psi, \quad (80)$$

one obtains

$$[-\nabla^2 + m_0^2 \Omega^2 + (1 - 6\xi) \Omega^{-1} \nabla^2 \Omega] \psi = \omega^2 \psi. \quad (81)$$

Define

$$V_{\text{eff}} = m_0^2 \Omega^2 + (1 - 6\xi) \Omega^{-1} \nabla^2 \Omega. \quad (82)$$

Then

$$[-\nabla^2 + V_{\text{eff}}] \psi_n = \omega_n^2 \psi_n. \quad (83)$$

This scalar spectral problem governs formation barriers and visible overlap profiles. It does not determine the number of chiral SM generations; that number is fixed by $\mathcal{D}_R$.

Already fixed by the Spin^c theorem: protected labels $N = 1, 2, 3$ and modes $u_N \in H^0(\mathbb{CP}^1, \mathcal{O}(2))$.




FIG. 6. Post-index scalar spectral interface for formation and visibility. After the Spin^c index theorem has fixed the protected generation space to three modes, the two-center scalar geometry supplies an effective interface for profiles, rates, and visible overlaps. The conformal factor Ω determines V_{eff} , whose scalar eigenmodes ψ_n are connected to the protected labels only through scalarization weights and hard/soft assignment. The probability law $P_N(t)$ describes post-index formation and visibility of $N = 1, 2, 3$; it is not an additional input to the proof of the family multiplicity.

IX. DEGENERACY, KINETICS, VISIBILITY, AND FORMATION

The structures in this section are an effective formation/visibility model built on the Spin^c generation theorem. They use scalar spectral data and two-center collective coordinates to parametrize rates and overlaps; they do not supply an independent proof of the number of chiral SM generations. The interface data used below are precisely $\mathfrak{D}_{\text{int}}(D)$ from Eq. (2); changing them changes the effective formation/visibility model, not the Spin^c multiplicity result.

For the protected chiral sector, let $\Pi_N = |u_N\rangle\langle u_N|$ be the projector onto the Toeplitz eigenmode u_N . The chiral degeneracy is

$$g_N^{\text{chiral}} = \text{Tr}_{\mathcal{H}_{\text{gen}}} \Pi_N, \quad \mathcal{H}_{\text{gen}} = H^0(\mathbb{CP}^1, \mathcal{O}(2)). \quad (84)$$

Thus

$$\boxed{g_1^{\text{chiral}} = g_2^{\text{chiral}} = g_3^{\text{chiral}} = 1, \quad g_{N \geq 4}^{\text{chiral}} = 0.} \quad (85)$$

A. Generation-to-spatial mode assignment

The protected generation basis $u_N \in H^0(C, L_R)$ and the scalar spatial eigenmodes ψ_n of Eq. (83) are not the same objects. The former are Spin^c zero modes on C ; the latter are eigenfunctions of a Schrödinger operator on the spatial two-center chart. We reserve $N = 1, 2, 3$ for protected generation labels and n for scalar spectral labels. Define

$$\mathcal{H}_{\text{gen}} = H^0(C, L_R), \quad \mathcal{H}_\Omega(D) = L^2(\mathbb{R}^3, d\mu_{\text{cyl}}), \quad d\mu_{\text{cyl}} = \rho d\rho dz_{\text{ax}} d\varphi, \quad (86)$$

and

$$L_\Omega(D) := -\nabla^2 + V_{\text{eff}}(\cdot; D), \quad L_\Omega(D)\psi_n(\cdot; D) = \omega_n^2(D)\psi_n(\cdot; D). \quad (87)$$

We use the Friedrichs self-adjoint realization of $L_\Omega(D)$ associated with the semibounded quadratic form of Eq. (87) on $\mathcal{H}_\Omega(D)$, after the regulated two-center potential $V_{\text{eff}}(\cdot; D)$ has been fixed. In finite-volume numerical applications, $L_\Omega(D)$ denotes the corresponding self-adjoint realization with the stated boundary conditions. The notation $dE_\lambda(D)$ below is the projection-valued spectral measure of this self-adjoint operator [19]. The projection map π_D pulls the generation bundle to the spatial chart:

$$\widehat{L}_R := \pi_D^* L_R, \quad \widehat{h}_R := \pi_D^* h_R, \quad \widehat{u} := \pi_D^* u \in \Gamma(\mathbb{R}^3, \widehat{L}_R). \quad (88)$$

To attach scalar barrier data to a protected generation, introduce the linear raw scalarization operator

$$\boxed{\widetilde{\mathcal{C}}_D : \mathcal{H}_{\text{gen}} \longrightarrow \mathcal{H}_\Omega(D), \quad \widetilde{\mathcal{C}}_D u = w_D^{1/2} \langle j_D, \pi_D^* u \rangle_{\widehat{h}_R}.} \quad (89)$$

Here $j_D \in \Gamma(\mathbb{R}^3, \widehat{L}_R)$ is the locking/probe section supplied by the two-center projection data, and $w_D \geq 0$ is a scalar density weight. With our convention that the Hermitian pairing is antilinear in the first entry and linear in the second, $\widetilde{\mathcal{C}}_D$ is linear in u . The raw scalar profile is assumed nonzero on the protected modes u_N . The normalized profile map is the generally nonlinear map

$$\boxed{c_D(u) = \frac{\widetilde{\mathcal{C}}_D u}{\|\widetilde{\mathcal{C}}_D u\|_{\mathcal{H}_\Omega(D)}}.} \quad (90)$$

The spectral weights of a generation mode against scalar eigenmodes are

$$\boxed{\mu_{nN}(D) = \left| \langle \psi_n(\cdot; D), c_D(u_N) \rangle_{\mathcal{H}_\Omega(D)} \right|^2.} \quad (91)$$

For a normalized discrete scalar spectral resolution, $\sum_n \mu_{nN} = 1$. If the scalar operator has a continuous component, Eq. (91) is replaced by the projection-valued spectral measure

$$d\mu_N^D(\lambda) = \|dE_\lambda(D) c_D(u_N)\|_{\mathcal{H}_\Omega(D)}^2. \quad (92)$$

Only when a unique scalar mode dominates do we define a hard mode-locking map. The condition is

$$n(N; D) = \arg \max_n \mu_{nN}(D), \quad \mu_{n(N; D), N}(D) - \sup_{m \neq n(N; D)} \mu_{mN}(D) \geq \Delta_{\text{lock}} > 0. \quad (93)$$

In that regime,

$$\mathcal{I}_D(u_N) := \psi_{n(N; D)}(\cdot; D), \quad \psi_N^{\text{lock}} := \psi_{n(N; D)}, \quad \omega_N(D) := \omega_{n(N; D)}(D). \quad (94)$$

If Eq. (93) fails, the generation must be assigned scalar data softly:

$$\Gamma_N^{\text{soft}}(D, \eta) = \sum_n \mu_{nN}(D) \Gamma_n(D, \eta), \quad (95)$$

or, for a continuous spectral component,

$$\Gamma_N^{\text{soft}}(D, \eta) = \int \Gamma(\lambda; D, \eta) d\mu_N^D(\lambda). \quad (96)$$

Here Γ_n is the scalar-mode formation rate for the n th eigenmode, computed by the same two-center Schur/WKB construction. Thus the Spin^c theorem fixes u_N , while $\tilde{\mathcal{C}}_D$, c_D , μ_{nN} , and, only under Eq. (93), $\mathcal{I}_D$, supply the scalar formation interface. All hard-locking quantities ω_N , Γ_N , and $|N, \pm\rangle$ below are understood only after Eq. (93); the final probability law is written with Γ_N^{eff} to cover both the hard-locking and soft-assignment regimes.

For kinetics in the hard-locking regime, the locked scalar mode ψ_N^{lock} and the two centers define a collective-coordinate subspace

$$\mathcal{C}_N = \text{span}\{|N, +\rangle, |N, -\rangle\}. \quad (97)$$

Let P_N^{cc} project onto $\mathcal{C}_N$, $Q_N^{\text{cc}} = 1 - P_N^{\text{cc}}$, and let $\mathcal{K}_N(D, \eta)$ be the Euclidean fluctuation operator in this sector. The dimensionless parameter η denotes the two-center overlap controlling kinetic mixing. For the regulated source model, we take

$$\boxed{\eta(D, \varepsilon) = \int_{\mathbb{R}^3} d^3x \sqrt{\rho_\varepsilon(x - x_+) \rho_\varepsilon(x - x_-)}, \quad 0 \leq \eta \leq 1.} \quad (98)$$

Since ρ_ε is normalized, Eq. (98) is the Hellinger affinity of the two regulated source densities. The bound $0 \leq \eta \leq 1$ follows from Cauchy–Schwarz, with $\eta = 1$ only for coincident identical profiles. It enters the fluctuation operator, WKB prefactors, and center mixing coefficient:

$$\mathcal{K}_N = \mathcal{K}_N(D, \eta), \quad A_{N,\pm} = A_{N,\pm}(D, \eta), \quad \chi_N = \chi_N(D, \eta). \quad (99)$$

The Feshbach–Schur step assumes a gap on the eliminated complement:

$$0 \notin \text{Spec}(Q_N^{\text{cc}} \mathcal{K}_N(D, \eta) Q_N^{\text{cc}}). \quad (100)$$

If this gap closes, the inverse below must be replaced by the corresponding reduced resolvent or pseudoinverse. Under Eq. (100), the Feshbach–Schur effective matrix is [18]

$$\mathcal{M}_N = P_N^{\text{cc}} \mathcal{K}_N P_N^{\text{cc}} - P_N^{\text{cc}} \mathcal{K}_N Q_N^{\text{cc}} (Q_N^{\text{cc}} \mathcal{K}_N Q_N^{\text{cc}})^{-1} Q_N^{\text{cc}} \mathcal{K}_N P_N^{\text{cc}}. \quad (101)$$

In the two-center basis,

$$\mathcal{M}_N = \begin{pmatrix} \gamma_{N,+} & \chi_N \sqrt{\gamma_{N,+} \gamma_{N,-}} \\ \chi_N \sqrt{\gamma_{N,+} \gamma_{N,-}} & \gamma_{N,-} \end{pmatrix}, \quad (102)$$

with WKB channel factors

$$\gamma_{N,\pm} = A_{N,\pm} \exp[-2S_{N,\pm}^{\text{WKB}}], \quad S_{N,\pm}^{\text{WKB}} = \int_{\mathcal{B}_{N,\pm}} ds \sqrt{(V_{\text{eff}}(s) - \omega_N^2)_+}. \quad (103)$$

Here

$$\mathcal{B}_{N,\pm} = \{s \text{ on the } \pm \text{ channel} : V_{\text{eff}}(s) > \omega_N^2\}, \quad (104)$$

and $(x)_+ = \max\{x, 0\}$. The formation rate is

$$\boxed{\Gamma_N(D, \eta) = \max\{0, \lambda_+(\mathcal{M}_N(D, \eta))\}}, \quad (105)$$

where

$$\lambda_+(\mathcal{M}_N) = \frac{\gamma_{N,+} + \gamma_{N,-}}{2} + \frac{1}{2} \sqrt{(\gamma_{N,+} - \gamma_{N,-})^2 + 4\chi_N^2 \gamma_{N,+} \gamma_{N,-}}. \quad (106)$$

For visibility, use the same pullback bundle from Eq. (88). For the N th generation, the spatial representative is the section

$$\hat{u}_N := \pi_D^* u_N \in \Gamma(\mathbb{R}^3, \hat{L}_R), \quad (107)$$

not a scalar function. The visible kernel is correspondingly a bundle-valued source

$$\mathcal{J}_{\text{vis}}(D, \varepsilon) \in \Gamma(\mathbb{R}^3, \hat{L}_R). \quad (108)$$

The φ -dependence is therefore not discarded; in axially reduced formulas it has already been projected into the selected angular channel and absorbed into the chosen representative of $\mathcal{J}_{\text{vis}}$ and the measure factor. The conformal Dirac and Higgs fields scale as

$$\Psi = \Omega^{-3/2}\psi, \quad H = \Omega^{-1}\phi. \quad (109)$$

Hence

$$\sqrt{-g} \bar{\Psi} H \Psi \propto \Omega^4 \cdot \Omega^{-3/2} \cdot \Omega^{-1} \cdot \Omega^{-3/2} = \Omega^0. \quad (110)$$

A mode-resolved Yukawa overlap is the Hermitian pairing on the pullback bundle:

$$y_N^{\text{eff}}(D, \varepsilon) = y_0 \int_{\mathbb{R}^3} \langle \hat{u}_N(x; D, \varepsilon), \mathcal{J}_{\text{vis}}(x; D, \varepsilon) \rangle_{\hat{h}_R} d\mu_{\text{cyl}}(x), \quad (111)$$

We use the convention that the Hermitian pairing is antilinear in the first entry. With this mathematical convention, y_N^{eff} is an antilinear amplitude in the generation vector u_N . This does not affect B_N , which depends only on $|y_N^{\text{eff}}|^2$. If y_N^{eff} is later used as a phase-sensitive Yukawa or Wilson coefficient and one wants a convention linear in u_N , one may equivalently use the opposite pairing order, $\int \langle \mathcal{J}_{\text{vis}}, \hat{u}_N \rangle_{\hat{h}_R} d\mu_{\text{cyl}}$; the normalized visibility B_N is unchanged. In a local frame $\hat{e}_R = \pi_D^* e_R$, write

$$\hat{u}_N = U_N \hat{e}_R, \quad \mathcal{J}_{\text{vis}} = J_{\text{vis}} \hat{e}_R, \quad \hat{h}_R(\hat{e}_R, \hat{e}_R) = H_R. \quad (112)$$

Then

$$\langle \hat{u}_N, \mathcal{J}_{\text{vis}} \rangle_{\hat{h}_R} = H_R \overline{U_N} J_{\text{vis}}. \quad (113)$$

This is independent of the frame. Indeed, under $\hat{e}'_R = g \hat{e}_R$, with $g : \mathbb{R}^3 \rightarrow \mathbb{C}^*$,

$$U'_N = g^{-1} U_N, \quad J'_{\text{vis}} = g^{-1} J_{\text{vis}}, \quad H'_R = |g|^2 H_R, \quad (114)$$

and hence

$$H'_R \overline{U'_N} J'_{\text{vis}} = H_R \overline{U_N} J_{\text{vis}}. \quad (115)$$

The earlier scalar-looking formula is only this local trivialized representative:

$$y_N^{\text{eff}}(D, \varepsilon) = y_0 \int_0^\infty \rho d\rho \int_{-\infty}^\infty dz_{\text{ax}} \int_0^{2\pi} d\varphi \overline{U_N(\rho, z_{\text{ax}}, \varphi; D, \varepsilon)} \\ \times f_L(\rho, z_{\text{ax}}) f_R(\rho, z_{\text{ax}}) W_{\text{frame}}(\rho, z_{\text{ax}}, \varphi; D, \varepsilon). \quad (116)$$

Here W_{frame} is not an additional invariant object; it denotes the local representative of the bundle frame, Hermitian metric H_R , visible source J_{vis} , angular-channel projection, Jacobian

convention, and any remaining complex-conjugation convention. When the visible kernel has already been reduced to an $m = 0$ axial channel, Eq. (116) reduces to the earlier $2\pi\rho d\rho dz_{\text{ax}}$ form with the angular integral included in W_{frame} . The normalized visible weight is

$$B_N = \frac{|y_N^{\text{eff}}|^2}{\sum_{I=1}^3 |y_I^{\text{eff}}|^2}. \quad (117)$$

We assume the visible kernel is not orthogonal to the entire protected generation space, so that

$$\sum_{I=1}^3 |y_I^{\text{eff}}|^2 > 0. \quad (118)$$

Under this nonzero-overlap condition the normalized weights B_N are well defined. For the operator notation in the equivalent form, define the linear visible-overlap functional

$$\mathcal{O}_{\text{vis}} : \mathcal{H}_{\text{gen}} \longrightarrow \mathbb{C}, \quad (\mathcal{O}_{\text{vis}} u) = y_0 \int_{\mathbb{R}^3} \langle \mathcal{J}_{\text{vis}}(x; D, \varepsilon), \pi_D^* u(x) \rangle_{\hat{h}_R} d\mu_{\text{cyl}}(x). \quad (119)$$

This uses the opposite pairing order from Eq. (111), so it is linear in u under the stated Hermitian convention. The two overlap conventions differ only by pairing order and conjugation convention and give

$$|\mathcal{O}_{\text{vis}} u_N|^2 = |y_N^{\text{eff}}|^2. \quad (120)$$

Thus $\mathcal{O}_{\text{vis}}^\dagger \mathcal{O}_{\text{vis}}$ is the induced rank-one positive operator on $\mathcal{H}_{\text{gen}}$, and equivalently

$$B_N = \frac{\langle u_N | \mathcal{O}_{\text{vis}}^\dagger \mathcal{O}_{\text{vis}} | u_N \rangle}{\sum_{I=1}^3 \langle u_I | \mathcal{O}_{\text{vis}}^\dagger \mathcal{O}_{\text{vis}} | u_I \rangle}. \quad (121)$$

Finally, at the coarse-grained effective level, define the rate used by the conditional probability law as

$$\Gamma_N^{\text{eff}}(D, \eta) = \begin{cases} \Gamma_N(D, \eta), & \text{in the hard-locking regime of Eq. (93),} \\ \Gamma_N^{\text{soft}}(D, \eta), & \text{otherwise.} \end{cases} \quad (122)$$

Assume the corresponding single-rate Markovian formation law

$$\frac{dp_N}{dt} = \Gamma_N^{\text{eff}}(1 - p_N), \quad p_N(0) = 0 \quad (123)$$

has solution

$$p_N(t) = 1 - e^{-\Gamma_N^{\text{eff}} t}. \quad (124)$$

Conditioning on the observation of one of the protected chiral layers gives

$$P_N(t; D, \eta) = \frac{B_N g_N^{\text{chiral}} \left(1 - e^{-\Gamma_N^{\text{eff}}(D, \eta)t}\right)}{\sum_{K=1}^3 B_K g_K^{\text{chiral}} \left(1 - e^{-\Gamma_K^{\text{eff}}(D, \eta)t}\right)}. \quad (125)$$

Equation (125) is understood for $t > 0$ and for parameter values for which the denominator is nonzero. At $t = 0$, or if all effective formation rates vanish, the conditional observation probability is not defined. When the rate-weighted denominator is nonzero, the right limit is

$$P_N(0^+; D, \eta) = \frac{B_N g_N^{\text{chiral}} \Gamma_N^{\text{eff}}(D, \eta)}{\sum_{K=1}^3 B_K g_K^{\text{chiral}} \Gamma_K^{\text{eff}}(D, \eta)}. \quad (126)$$

X. LIMITATIONS AND OUTLOOK

The result should be read as a conditional field-theoretic and index-theoretic model, not as a complete theory of flavor. Several inputs remain outside the theorem. First, the paper assumes the PSLT two-center projection data (C, R) rather than deriving $C = \mathbb{CP}^1$ and $R = p_+ + p_-$ from a UV-complete quantum-gravity vacuum. Second, the protected SM-charged carrier $\mathcal{R}_{\text{SM}} \otimes (\Lambda^{0,*} T^* C \otimes L_R)$ and the Spin^c Dolbeault representative are taken as the protected generation sector; the BPS remark provides a compatible motivation, not a universal vacuum-selection theorem. Third, the exact three-family conclusion requires the stated exclusivity and gap assumptions for all additional SM-charged internal sectors. Fourth, the theorem fixes the multiplicity of protected family copies but does not determine Yukawa hierarchies, CKM or PMNS mixing, neutrino masses, or the detailed spectrum of massive vectorlike states.

The scalar two-center spectral operator and the effective quantities $(g_N^{\text{chiral}}, \Gamma_N^{\text{eff}}, B_N, P_N)$ should therefore be viewed as a post-index phenomenological interface. They can be used to assign formation rates and visible-sector overlaps to the three protected modes, but they are not inputs to the Spin^c proof of $N_{\text{gen}} = 3$. Future work would need to derive the projection data and protected carrier from a more microscopic PSLT or compactification construction, and then connect the overlap functional to realistic flavor textures.

XI. CONCLUSION

The two-center PSLT projection data determine the divisor $R = p_+ + p_-$ and the line bundle $L_R = \mathcal{O}(R) \simeq \mathcal{O}(2)$ on $C = \mathbb{CP}^1$. The SM-charged internal fermions are governed by the product Dirac action on $M_4 \times C$, with the internal Spin^c Dolbeault–Dirac operator acting on $\Lambda^{0,*}T^*C \otimes L_R$. The bundle L_R is the auxiliary Dolbeault twist, and the positive and negative internal zero modes are $H^0(C, L_R)$ and $H^1(C, L_R)$. Riemann–Roch gives index three, and Serre duality gives $H^1(C, L_R) = 0$. Thus, conditional on the PSLT two-center projection data and the stated gap/exclusivity assumptions, the protected SM-charged generation sector contributes exactly three low-energy chiral SM family copies, with no additional unpaired light SM-charged exotics arising from the allowed extra sectors.

After the raw scalarization operator $\tilde{\mathcal{C}}_D$, the normalized profile map c_D , the spectral weights μ_{nN} , and, when the locking gap exists, the hard map $\mathcal{I}_D : u_N \mapsto \psi_N^{\text{lock}}$, the scalar two-center spectral operator and the quantities $(g_N^{\text{chiral}}, \Gamma_N^{\text{eff}}, B_N, P_N)$ define an effective model for how these three protected modes form and couple to the visible sector. They are coarse-grained phenomenological structures on top of the Spin^c generation theorem, not substitutes for it.

ACKNOWLEDGMENTS

The author reports no external funding for this work.

-
- [1] M. F. Atiyah and I. M. Singer, “The index of elliptic operators: I,” *Annals Math.* **87**, 484 (1968).
 - [2] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics Vol. 52, Springer (1977).
 - [3] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley (1978).
 - [4] O. Forster, *Lectures on Riemann Surfaces*, Graduate Texts in Mathematics Vol. 81, Springer (1981).
 - [5] H. B. Lawson and M.-L. Michelsohn, *Spin Geometry*, Princeton University Press (1989).
 - [6] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., Institute of Physics Publishing (2003).
 - [7] R. A. Bertlmann, *Anomalies in Quantum Field Theory*, Oxford University Press (1996).

- [8] A. Bilal, “Lectures on Anomalies,” arXiv:0802.0634.
- [9] E. Witten, “An $SU(2)$ anomaly,” Phys. Lett. B **117**, 324 (1982).
- [10] P. Candelas, G. T. Horowitz, A. Strominger, and E. Witten, “Vacuum configurations for superstrings,” Nucl. Phys. B **258**, 46 (1985).
- [11] E. Witten, “New issues in manifolds of $SU(3)$ holonomy,” Nucl. Phys. B **268**, 79 (1986).
- [12] V. Braun, Y.-H. He, B. A. Ovrut, and T. Pantev, “A heterotic standard model,” Phys. Lett. B **618**, 252 (2005), arXiv:hep-th/0501070.
- [13] V. Bouchard and R. Donagi, “An $SU(5)$ heterotic standard model,” Phys. Lett. B **633**, 783 (2006), arXiv:hep-th/0512149.
- [14] R. Blumenhagen, B. Kors, D. Lüst, and S. Stieberger, “Four-dimensional string compactifications with D-branes, orientifolds and fluxes,” Phys. Rept. **445**, 1 (2007), arXiv:hep-th/0610327.
- [15] C. Beasley, J. J. Heckman, and C. Vafa, “GUTs and exceptional branes in F-theory I,” JHEP **01**, 058 (2009), arXiv:0802.3391.
- [16] M. Bordemann, E. Meinrenken, and M. Schlichenmaier, “Toeplitz quantization of Kähler manifolds and $\mathfrak{gl}(N)$, $N \rightarrow \infty$ limits,” Commun. Math. Phys. **165**, 281 (1994).
- [17] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press (1982).
- [18] H. Feshbach, “Unified theory of nuclear reactions,” Annals Phys. **5**, 357 (1958).
- [19] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press (1978).