

Breakpoint model

1 Subject specific, single breakpoint, single slope

1.1 Direct optimisation

Consider a response variable Y and an explanatory variable X related by:

$$Y = \beta X - \beta(X - \psi)_+ + \varepsilon$$

where $(\beta, \psi) \in \mathbb{R}^2$, $\varepsilon \sim \mathcal{N}(0, \sigma^2)$, and $(x)_+ = x$ if $x > 0$ and 0 otherwise. Introduce $\Theta = (\beta, \psi, \sigma^2)$, we can express the likelihood relative to n iid observations as:

$$\mathcal{L}(\Theta) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(Y_i - \beta X_i + \beta(X_i - \psi)_+)^2}{2\sigma^2}\right)$$

and the log-likelihood as:

$$\ell(\Theta) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \sum_{i=1}^n \frac{(Y_i - \beta X_i + \beta(X_i - \psi)_+)^2}{2\sigma^2}$$

Maximizing the likelihood with respect to $\Theta_\mu = (\beta, \psi)$ is equivalent to minimizing the residual sum of squared (RSS):

$$RSS(\Theta_\mu) = \sum_{i=1}^n (Y_i - \beta X_i + \beta(X_i - \psi)_+)^2$$

One approach to estimate the model parameters is to directly minimize this log-likelihood using, for instance grid search. This may be numerically inefficient but does not require any smoothness assumption on the likelihood.

- this is what `optim.lmbreak_NLMINB` in the `lmbreak` package does (set argument `control` to `list(optimizer = "nlnminb")`). `nlnminb` seems to employ a quasi Newton method though, some does require some smoothness.

1.2 Gradient descent

One can re-write the residual sum of squares as:

$$RSS(\Theta_\mu) = \sum_{i=1}^n \varepsilon_i(\psi)^2 = Y(I - H(\psi))(I - H(\psi))^T Y^T = Y(I - H(\psi))Y^T$$

where $H(\psi) = Z(\psi) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi)$ is the hat matrix and $Z(\psi) = (X - (X - \psi)_+)$. It is idempotent and so is $I - H$ where I denote the identity matrix. Therefore a possible derivative w.r.t. ψ of the RSS is:

$$\begin{aligned} \frac{\partial RSS(\Theta_\mu)}{\partial \psi} &= -Y \frac{\partial H(\psi)}{\partial \psi} Y^T \\ &= -Y \frac{\partial Z(\psi)}{\partial \psi} (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \\ &\quad + Y Z(\psi) (Z^T(\psi)Z(\psi))^{-1} \left(\frac{\partial Z^T(\psi)}{\partial \psi} Z(\psi) + Z^T(\psi) \frac{\partial Z(\psi)}{\partial \psi} \right) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \\ &\quad - Y Z(\psi) (Z^T(\psi)Z(\psi))^{-1} \frac{\partial Z^T(\psi)}{\partial \psi} Y^T \end{aligned}$$

The first and last term being scalar and transpose of each other they are equal. Similarly the two middle terms are also scalar and transpose of each other so equal. Therefore:

$$\begin{aligned} \frac{\partial RSS(\Theta_\mu)}{\partial \psi} &= -2Y \frac{\partial Z(\psi)}{\partial \psi} (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \\ &\quad + 2Y Z(\psi) (Z^T(\psi)Z(\psi))^{-1} \left(Z^T(\psi) \frac{\partial Z(\psi)}{\partial \psi} \right) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \\ &= 2Y \left(-\frac{\partial Z(\psi)}{\partial \psi} + Z(\psi) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) \frac{\partial Z(\psi)}{\partial \psi} \right) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \\ &= 2Y \left(Z(\psi) (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) - I \right) \frac{\partial Z(\psi)}{\partial \psi} (Z^T(\psi)Z(\psi))^{-1} Z^T(\psi) Y^T \end{aligned}$$

So it only remains to evaluate the (sub-)derivatives of $X(\psi)$:

$$\frac{\partial Z(\psi)}{\partial \psi} = \frac{\partial (X - \psi) \mathbb{1}_{X > \psi}}{\partial \psi} = -\mathbb{1}_{X > \psi} - \mathbb{1}_{X = \psi} \delta = -(\mathbb{1}_{X > \psi} + \mathbb{1}_{X = \psi} \delta)$$

where $\delta \in [0, 1]$.

So a second approach to estimate the model parameters is to perform a gradient descent based on sub-derivatives

- this is what `optim.lmbreak_BFGS` in the `lmbreak` package does (set argument `control` to `list(optimizer = "BFGS")`). δ is then randomly sampled in a uniform distribution whose support is $[0, 1]$.

Note: another approach to 'profile-out' the β , is to first plug-in the OLS estimator:

$$\tilde{\beta}(\psi) = \frac{\sum_{i=1}^n X_i Y_i - (X_i - \psi)_+ Y_i}{\sum_{i=1}^n (X_i + (X_i - \psi)_+)^2} = \frac{\sum_{i=1}^n \mathbf{1}_{X_i \leq \psi} X_i Y_i}{\sum_{i=1}^n \mathbf{1}_{X_i \leq \psi} X_i^2}$$

and then minimize w.r.t. ψ :

$$\ell(\psi) = \sum_{i=1}^n (Y_i - \tilde{\beta}(\psi) X_i + \tilde{\beta}(\psi) (X_i - \psi)_+)^2$$

So the first 'derivative' should solve:

$$\begin{aligned} 0 &= -2 \sum_{i=1}^n \left[\frac{\partial \tilde{\beta}(\psi)}{\partial \psi} (X_i - (X_i - \psi)_+) - \tilde{\beta}(\psi) \frac{\partial (X_i - \psi)_+}{\partial \psi} \right] (Y_i - \tilde{\beta}(\psi) X_i + \tilde{\beta}(\psi) (X_i - \psi)_+) \\ 0 &= \frac{\partial \tilde{\beta}(\psi)}{\partial \psi} \sum_{i=1}^n (X_i - (X_i - \psi)_+) (Y_i - \tilde{\beta}(\psi) X_i + \tilde{\beta}(\psi) (X_i - \psi)_+) \\ &\quad - \tilde{\beta}(\psi) \sum_{i=1}^n \frac{\partial (X_i - \psi)_+}{\partial \psi} (Y_i - \tilde{\beta}(\psi) X_i + \tilde{\beta}(\psi) (X_i - \psi)_+) \end{aligned}$$

Noticing that $\tilde{\beta}(\psi)$ is a step function with jumps at $(X_i)_{i \in \{1, \dots, n\}}$. If the breakpoint does not coincide with any datapoint, it has derivative 0 w.r.t. ψ . Under such assumption, the previous expression simplifies into:

$$\begin{aligned} 0 &= -\tilde{\beta}(\psi) \sum_{i=1}^n \mathbf{1}_{X_i \geq \psi} (Y_i - \tilde{\beta}(\psi) \psi) \\ \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{X_i \geq \psi} Y_i &= \tilde{\beta}(\psi) \psi \end{aligned}$$

So the breakpoint should be such that the average post-breakpoint value equal the fitted plateau.

1.3 Example

Simulate some data:

```
library(lmbreak)

set.seed(10)
df10 <- simBreak(c(1, 100), breakpoint = c(0,2,4),
                 slope = c(1,0), sigma = 0.05)
```

Automatic fit of the breakpoint model:

```
e.lmbreak10 <- lmbreak(Y ~ 0 + bp(X, pattern = "10", start = c(2)),
                      data = df10)
logLik(e.lmbreak10)
```

```
[1] 161.5047
```

Estimates:

```
model.tables(e.lmbreak10)
```

	X	duration	intercept	slope
1	0.05773562	1.940347	0.0579186	1.003169
2	1.99808295	1.847127	2.0044152	0.000000
3	3.84520966	NA	2.0044146	NA

OLS:

```
XX <- df10$X - pmax(df10$X-coef(e.lmbreak10),0)
solve(t(XX) %*% XX) %*% t(XX) %*% df10$Y
```

```
[,1]
[1,] 1.003169
```

Explicit OLS:

```
sum((df10$X < coef(e.lmbreak10))*df10$X*df10$Y)/sum((df10$X < coef(e.
lmbreak10))*df10$X^2)
```

```
[1] 1.003169
```

Full likelihood:

```
calcLogLik <- function(theta){
  beta <- theta["beta"]
  psi <- theta["psi"]
  sigma <- exp(theta["logsigma"])
  out <- NROW(df10)/2 * log(2*pi) + NROW(df10)/2 * log(sigma) + sum((df10$
    Y - beta * df10$X + beta * pmax(df10$X - psi,0))^2)/(2*sigma)
  return(unname(out))
}
## By default optim performs minimization
theta.init <- c(beta = 0, psi = 1, logsigma = 0.5)
res.opt <- optim(par = theta.init, fn = calcLogLik)
c(res.opt$par, value = res.opt$value)
```

beta	psi	logsigma	value
1.003165	1.998090	-6.067996	-161.504705

We retrieve the estimate and log-likelihood of `lmbreak`.

2 Mixed model, single breakpoint, single slope

2.1 Example

We will consider a dataset of 20 subjects with 25 observations each:

```
library(lmbreak)
library(mvtnorm)


## data generating parameters
n.subject <- 50
sigma2T <- 0.005
betaT <- 1
psiT <- 2
MtauT <- tcrossprod(c(0.2,0.1)) * matrix(c(1,0.5,0.5,1),2,2)
seqTime <- seq(0,4,length.out=25)

set.seed(10)
RE <- rmvnorm(n.subject, mean = c(psiT,betaT), sigma = MtauT)

ls.sim <- lapply(1:n.subject, function(iS){ ## iS <- 1
  iSim <- simBreak(c(1, length(seqTime)),
    breakpoint = c(min(seqTime),RE[iS,1],max(seqTime)),
    slope = c(RE[iS,2],0), sigma = sqrt(sigma2T),
    rdist.X = seqTime)

  iSim$id <- iS
  iSim
})
df.sim <- do.call(rbind,ls.sim)
```

`lmbreak` version 0.1.2



2.2 Model

Consider a set of n subjects each undergoing T measurements at equally spaced timepoints $1, \dots, T$ of an outcome variable Y . We consider the following random slope and random breakpoint model:

$$Y_{i,t} = \beta_i t - \beta_i(t - \psi_i)_+ + \varepsilon_{i,t}$$

where $\beta_i = \beta + u_i$ and $\psi_i = \psi + v_i$ and:

$$\begin{bmatrix} u_i \\ v_i \\ \varepsilon_{i,t} \end{bmatrix} = \mathcal{N} \left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \tau_{u,u} & \tau_{u,v} & 0 \\ \tau_{u,v} & \tau_{v,v} & 0 \\ 0 & 0 & \sigma^2 \end{bmatrix} \right)$$

We further denote $\Sigma_\tau = \begin{bmatrix} \tau_{u,u} & \tau_{u,v} \\ \tau_{u,v} & \tau_{v,v} \end{bmatrix}$ whose inverse is $\Sigma_\tau^{-1} = \begin{bmatrix} w_{u,u} & w_{u,v} \\ w_{u,v} & w_{v,v} \end{bmatrix} = \frac{1}{\tau_{u,u}\tau_{v,v} - \tau_{u,v}^2} \begin{bmatrix} \tau_{v,v} & -\tau_{u,v} \\ -\tau_{u,v} & \tau_{u,u} \end{bmatrix}$.

We now can re-write the model as:

$$\begin{aligned} Y_{i,t} &= \beta(t - (t - \psi - v_i)_+) + u_i(t - (t - \psi - v_i)_+) + \varepsilon_{i,t} \\ &= \mu(t, \beta, \psi, u_i, v_i) + \varepsilon_{i,t} \end{aligned}$$

where $\mu(t, \beta, \psi, u_i, v_i) = \beta t + u_i t$ when $t \leq \psi + v_i$
and $\mu(t, \beta, \psi, u_i, v_i) = \beta(\psi + v_i) + u_i(\psi + v_i)$ otherwise.

2.3 Likelihood

2.3.1 Bivariate integral

Introduce $\Theta = (\beta, \psi, \tau_{u,u}, \tau_{u,v}, \tau_{v,v}, \sigma^2)$, we can express the likelihood relative to n iid subjects as:

$$\mathcal{L}(\Theta) = \prod_{i=1}^n \prod_{t=1}^T \int_{u_i=-\infty}^{+\infty} \int_{v_i=-\infty}^{+\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left(-\frac{(Y_i - \mu(t, \beta, \psi, u_i, v_i))^2}{2\sigma^2} \right) f_\tau(u, v) dv_i du_i$$

$$\begin{aligned} \text{where } f_\tau(u, v) &= \frac{1}{2\pi |\Sigma_\tau|^{1/2}} \exp \left(-\frac{1}{2} \begin{bmatrix} u & v \end{bmatrix} \Sigma_\tau^{-1} \begin{bmatrix} u \\ v \end{bmatrix} \right) \\ &= \frac{1}{2\pi \sqrt{\tau_{u,u}\tau_{v,v} - \tau_{u,v}^2}} \exp \left(-\frac{u^2 w_{u,u} + 2uv w_{u,v} + v^2 w_{v,v}}{2} \right) \end{aligned}$$

One could evaluate this log-likelihood using numerical integration but this will typically be very slow:

- e.g. for a single subject at a single timepoint:

```
system.time( ## single subject
  resIntegral2 <- logLik_mixed10(
    Y = df.sim[df.sim$id==1,"Y"], t = df.sim[df.sim$id==1,"X"],
    beta = betaT, psi = psiT, tau_uu = MtauT, sigma2 = sigma2T,
    integration = "bivariate")
)
```

```
bruger    system forløbet
11.64      0.61      12.30
```

```
head(resIntegral2)
```

	integral1	error1	integral2	error2	integral	error	logLik
1	NA	NA	NA	NA	4.220886	4.167256e-05	1.4400450
2	NA	NA	NA	NA	4.810385	4.781720e-05	1.5707771
3	NA	NA	NA	NA	2.947878	2.925916e-05	1.0810858
4	NA	NA	NA	NA	3.071459	3.067560e-05	1.1221526
5	NA	NA	NA	NA	2.468231	2.417586e-05	0.9035015
6	NA	NA	NA	NA	2.181033	2.144690e-05	0.7797984

Since standard optimizers may evaluate the likelihood a large number of times this direct approach is not really feasible (too slow).

2.3.2 Nested univariate integrals

To simplify the numerical integration for a bivariate integral to univariate integral, we start by splitting second integral depending on whether $t \leq \psi + v_i$ i.e. $v_i \geq t - \psi$:

$$\begin{aligned}
\mathcal{L}(\Theta) &= \prod_{i=1}^n \prod_{t=1}^T \int_{u_i=-\infty}^{+\infty} \int_{v_i=t-\psi}^{+\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(Y_i - \beta t - u_i t)^2}{2\sigma^2}\right) f_{\tau}(u_i, v_i) dv_i du_i \\
&\quad + \prod_{i=1}^n \prod_{t=1}^T \int_{u_i=-\infty}^{+\infty} \int_{v_i=-\infty}^{t-\psi} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(Y_i - \beta(\psi + v_i) - u_i(\psi + v_i))^2}{2\sigma^2}\right) f_{\tau}(u_i, v_i) dv_i du_i \\
&= \frac{1}{(2\pi)^{3nT/2} \sigma^{nT} (\tau_{u,u} \tau_{v,v} - \tau_{u,v}^2)^{nT/2}} \prod_{i=1}^n \prod_{t=1}^T (\mathcal{L}_{1,i,t}(\Theta) + \mathcal{L}_{2,i,t}(\Theta))
\end{aligned}$$

where we introduce the notation:

$$\begin{aligned}
\mathcal{L}_{1,i,t}(\Theta) &= \int_{u_i=-\infty}^{+\infty} \int_{v_i=t-\psi}^{+\infty} \exp\left(-\frac{(Y_i - \beta t - u_i t)^2}{2\sigma^2} - \frac{u_i^2 w_{u,u} + 2u_i v_i w_{u,v} + v_i^2 w_{v,v}}{2}\right) dv_i du_i \\
\mathcal{L}_{2,i,t}(\Theta) &= \int_{u_i=-\infty}^{+\infty} \int_{v_i=-\infty}^{t-\psi} \exp\left(-\frac{(Y_i - \beta(\psi + v_i) - u_i(\psi + v_i))^2}{2\sigma^2} - \frac{u_i^2 w_{u,u} + 2u_i v_i w_{u,v} + v_i^2 w_{v,v}}{2}\right) dv_i du_i
\end{aligned}$$

Using that:

$$\begin{aligned}
\int_a^b \exp\left(-\frac{1}{2}(Av_i^2 + 2Bv_i)\right) dv_i &= \int_a^b \exp\left(-\frac{A}{2}(v_i^2 + 2\frac{B}{A}v_i)\right) dv_i \\
&= \int_a^b \exp\left(-\frac{A}{2}\left(v_i + \frac{B}{A}\right)^2 + \frac{B^2}{2A}\right) dv_i \\
&= \sqrt{\frac{2\pi}{A}} \exp\left(\frac{B^2}{2A}\right) \frac{1}{\sqrt{2\pi A^{-1}}} \int_a^b \exp\left(-\frac{(v_i^2 - -\frac{B}{A})^2}{2A^{-1}}\right) dv_i \\
&= \sqrt{\frac{2\pi}{A}} \exp\left(\frac{B^2}{2A}\right) \left(\Phi_{\mu=-\frac{B}{A}, \sigma^2=A^{-1}}(b) - \Phi_{\mu=-\frac{B}{A}, \sigma^2=A^{-1}}(a)\right) \\
&= \sqrt{\frac{2\pi}{A}} \exp\left(\frac{B^2}{2A}\right) \left(\Phi\left(\frac{bA+B}{\sqrt{A}}\right) - \Phi\left(\frac{aA+B}{\sqrt{A}}\right)\right)
\end{aligned}$$

where $\Phi_{\mu, \sigma^2}(\bullet) = \Phi\left(\frac{\bullet - \mu}{\sigma}\right)$ denotes the Cumulative Distribution Function (CDF) of a normal distribution with mean μ and variance σ^2 . For the first term, using $a = t - \psi$, $b = +\infty$, $A = w_{v,v}$, and $B = u_i w_{u,v}$, we obtain:

$$\mathcal{L}_{1,i,t}(\Theta) = \sqrt{\frac{2\pi}{w_{v,v}}} \int_{-\infty}^{+\infty} \exp\left(-\frac{(Y_i - \beta t - u_i t)^2}{2\sigma^2} - \frac{u_i^2(w_{u,u} - \frac{w_{u,v}^2}{w_{v,v}})}{2}\right) \left(1 - \Phi\left(\frac{w_{v,v}(t - \psi) + u_i w_{u,v}}{\sqrt{w_{v,v}}}\right)\right) du_i$$

which can be evaluated using numerical intergration (e.g. `stats::integrate` and `stats::pnorm` - see function next page). A similar approach can be used to evaluate the second term.

$$\begin{aligned}
\mathcal{L}_{2,i,t}(\Theta) &= \int_{u_i=-\infty}^{+\infty} \int_{v_i=-\infty}^{t-\psi} \exp\left(-\frac{(Y_i - \psi(\beta + u_i) - v_i(\beta + u_i))^2}{2\sigma^2} - \frac{u_i^2 w_{u,u} + 2u_i v_i w_{u,v} + v_i^2 w_{v,v}}{2}\right) dv_i du_i \\
&= \int_{u_i=-\infty}^{+\infty} \int_{v_i=-\infty}^{t-\psi} \exp\left(-\frac{(Y_i - \psi(\beta + u_i))^2 + v_i^2(\beta + u_i)^2 - 2v_i(\beta + u_i)(Y_i - \psi(\beta + u_i))}{2\sigma^2} - \frac{u_i^2 w_{u,u} + 2u_i v_i w_{u,v} + v_i^2 w_{v,v}}{2}\right) dv_i du_i
\end{aligned}$$

Using $a = -\infty$, $b = t - \psi$, $A = w_{v,v} + \frac{(\beta + u_i)^2}{\sigma^2}$, and $B = u_i w_{u,v} - h(u_i)$ where $h(u_i) = \frac{(\beta + u_i)(Y_i - \psi(\beta + u_i))}{\sigma^2}$, we obtain for $\mathcal{L}_{2,i,t}(\Theta)$:

$$\sqrt{2\pi} \int_{-\infty}^{+\infty} \exp\left(-\frac{(Y_i - \psi(\beta + u_i))^2}{2\sigma^2} - \frac{u_i^2 w_{u,u} - \frac{(u_i w_{u,v} - h(u_i))^2}{w_{v,v} + \sigma^{-2}(\beta + u_i)^2}}{2}\right) \frac{\Phi\left(\frac{(w_{v,v} + \sigma^{-2}(\beta + u_i)^2)(t - \psi) + u_i w_{u,v} - h(u_i)}{\sqrt{w_{v,v} + \sigma^{-2}(\beta + u_i)^2}}\right)}{\sqrt{w_{v,v} + \sigma^{-2}(\beta + u_i)^2}} du_i$$

This approach is (much) more numerically efficient:

```
system.time( ## whole dataset
  resIntegral <- logLik_mixed10(
    Y = df.sim$Y, t = df.sim$X,
    beta = betaT, psi = psiT, sigma2 = sigma2T,
    tau_uu = MtauT[1,1], tau_uv = MtauT[1,2], tau_vv = MtauT[2,2],
    integration = "univariate")
)
```

```
bruger    system forløbet
  0.22      0.00      0.22
```

```
head(resIntegral)
```

	integral1	error1	integral2	error2	integral	error	logLik
1	4.220895	1.368832e-06	1.086702e-88	1.821438e-88	4.220895	NA	1.4400472
2	4.810386	1.252909e-03	1.005836e-78	1.705596e-78	4.810386	NA	1.5707774
3	2.947875	9.400217e-03	5.921995e-68	9.665419e-68	2.947875	NA	1.0810846
4	3.071454	9.777754e-03	1.220746e-61	2.099327e-61	3.071454	NA	1.1221511
5	2.468229	1.075678e-02	8.763755e-49	1.579133e-48	2.468229	NA	0.9035010
6	2.181034	3.596076e-05	6.274525e-40	1.146041e-39	2.181034	NA	0.7797989

and leads to nearly identical results:

```
quantile(resIntegral$logLik[1:NROW(resIntegral2)]-resIntegral2$logLik)
```

	0%	25%	50%	75%	100%
	-1.216897e-04	-8.942998e-07	5.361709e-05	2.670979e-04	3.104343e-03

Total log-likelihood can be obtained by summing the logLik over rows:

```
sum(resIntegral$logLik)
```

```
[1] 59.66285
```

or, equivalently, setting the argument indiv to FALSE:

```
logLik_mixed10(
  Y = df.sim$Y, t = df.sim$X,
  beta = betaT, psi = psiT, sigma2 = sigma2T,
  tau_uu = MtauT[1,1], tau_uv = MtauT[1,2], tau_vv = MtauT[2,2],
  indiv = FALSE) ## uses univariate-fast by default
```

```
[1] 59.66285
```

2.4 Direct optimisation

We can now search for the model parameters maximizing the likelihood:

- using a transformation to constrain the breakpoint to be within the range of observation times, the variance-covariance matrix of the random effect to be positive, and the variance to be positive

```
res.optim <- optim(
  par = c(beta = 1,
          psi.logit = qlogis(0.5),
          tau_uu.log = log(MtauT[1,1]),
          rho_uv = atanh(cov2cor(MtauT)[1,2]),
          tau_vv.log = log(MtauT[2,2]),
          sigma2.log = log(0.05)),
  fn = function(theta){
    theta["psi"] <- max(df.sim$X) * plogis(theta["psi.logit"])
    theta["tau_uu"] <- exp(theta["tau_uu.log"])
    theta["tau_vv"] <- exp(theta["tau_vv.log"])
    theta["tau_uv"] <- sqrt(theta["tau_uu"]*theta["tau_vv"])*tanh(theta["rho_uv"])
    theta["sigma2"] <- exp(theta["sigma2.log"])

    iOut <- try(logLik_mixed10(Y = df.sim$Y, t = df.sim$X,
                             beta = theta["beta"], psi = theta["psi"],
                             tau_uu = theta["tau_uu"],
                             tau_uv = theta["tau_uv"],
                             tau_vv = theta["tau_vv"],
                             sigma2 = theta["sigma2"],
                             indiv = FALSE), silent = TRUE)

    if(inherits(iOut,"try-error")){
      cat("ERROR: ",paste(theta[c("beta","psi","tau_uu","tau_uv","tau_vv",
                                "sigma2")], collapse=","))
      return(Inf) ## optim is minimizing
    }else{
      return(-iOut) ## optim is minimizing
    }
  }, control = list(reltol = 1e-3))
```

After back-transformation we obtain the following estimates:

```
res.optim$par["psi"] <- max(df.sim$X) * plogis(res.optim$par["psi.logit"])
res.optim$par["tau_uu"] <- exp(res.optim$par["tau_uu.log"])
res.optim$par["tau_vv"] <- exp(res.optim$par["tau_vv.log"])
res.optim$par["tau_uv"] <- sqrt(res.optim$par["tau_uu"]*res.optim$par["tau_vv"])*tanh(res.optim$par["rho_uv"])
```

```
res.optim$par["sigma2"] <- exp(res.optim$par["sigma2.log"])
res.optim$par[c("beta", "psi", "tau_uu", "tau_uv", "tau_vv", "sigma2")]
```

beta	psi	tau_uu	tau_uv	tau_vv	sigma2
0.989314425	1.942741606	0.008735823	0.014492654	0.039373193	0.006122932

2.5 Estimating the random effects

Knowing the model parameters and a vector of outcomes for a specific subject, we can evaluate the subject-specific random effect as the expected value of the random effect given the observations. We therefore need the joint distribution of (Y, u, v) :

$$\begin{aligned}\mathbb{E}[Y] &= \beta t + \beta \mathbb{E}[(t - \psi - v_i) \mathbb{1}_{t-\psi-v_i>0}] + \mathbb{E}[u_i(t - \psi - v_i) \mathbb{1}_{t-\psi-v_i>0}] \\ &= \beta t + \beta(t - \psi) - \mathbb{E}[v_i \mathbb{1}_{t-\psi-v_i>0}] + (t - \psi) \mathbb{E}[u_i \mathbb{1}_{t-\psi-v_i>0}] - \mathbb{E}[u_i v_i \mathbb{1}_{t-\psi-v_i>0}] \\ &= \beta t + \beta(t - \psi) - \mathbb{E}[v_i \mathbb{1}_{v_i< t-\psi}] + (t - \psi) \mathbb{E}[u_i \mathbb{1}_{v_i< t-\psi}] - \mathbb{E}[u_i v_i \mathbb{1}_{v_i< t-\psi}]\end{aligned}$$

We now focus on the expectation terms:

$$\begin{aligned}\mathbb{E}[v_i \mathbb{1}_{v_i< t-\psi}] &= \int_{-\infty}^{t-\psi} \frac{v_i}{\sqrt{2\pi\tau_{v,v}}} \exp\left(-\frac{v_i^2}{2\tau_{v,v}}\right) dv_i \\ &= -\frac{\sqrt{\tau_{v,v}}}{\sqrt{2\pi}} \left[\exp\left(-\frac{v_i^2}{2\tau_{v,v}}\right) \right]_{-\infty}^{t-\psi} = -\frac{\exp\left(-\frac{(t-\psi)^2}{2\tau_{v,v}}\right) \sqrt{\tau_{v,v}}}{\sqrt{2\pi}} \\ &= \mu_1(t, \psi, \tau_{v,v})\end{aligned}$$

$$\begin{aligned}\mathbb{E}[u_i \mathbb{1}_{v_i< t-\psi}] &= \int_{-\infty}^{+\infty} \int_{-\infty}^{t-\psi} \frac{u_i}{2\pi\sqrt{\tau_{u,u}\tau_{v,v} - \tau_{u,v}^2}} \exp\left(-\frac{w_{v,v}v_i^2 + 2w_{u,v}u_iv_i + w_{u,u}u_i^2}{2}\right) dv_i du_i \\ &= \frac{1}{\sqrt{2\pi w_{v,v}(\tau_{u,u}\tau_{v,v} - \tau_{u,v}^2)}} \int_{-\infty}^{+\infty} u_i \exp\left(-\frac{u_i^2(w_{u,u} - \frac{w_{u,v}^2}{w_{v,v}})}{2}\right) \Phi\left((t - \psi)\sqrt{w_{v,v}} + \frac{w_{u,v}}{\sqrt{w_{v,v}}}u_i\right) du_i \\ &= \mu_2(t, \psi, \tau_{u,u}, \tau_{u,v}, \tau_{v,v})\end{aligned}$$

$$\mathbb{E}[u_i v_i \mathbb{1}_{v_i< t-\psi}] = \int_{-\infty}^{+\infty} \int_{-\infty}^{t-\psi} \frac{u_i v_i}{2\pi\sqrt{\tau_{u,u}\tau_{v,v} - \tau_{u,v}^2}} \exp\left(-\frac{w_{v,v}v_i^2 + 2w_{u,v}u_iv_i + w_{u,u}u_i^2}{2}\right) dv_i du_i$$