North American Bat Monitoring Program in British Columbia

Data Summary and Activity Trend Analyses: 2017 - 2025

Camila Hurtado Taylor Hart

2026-10-19

Table of contents

1 Executive Summary **4**

2 Land Acknowledgement **4**

3 Introduction **5**
3.1 Overview of NABat and the NNW Bat Hub . . . 5
3.2 Threats to bat populations in BC . . . 5
3.2.1 White-Nose Syndrome . . . 5
3.2.2 Wind Energy Development . . . 6
3.3 Acoustic Monitoring of Bats . . . 8
3.4 Updates in Modelling Approaches . . . 8

4 Methods **9**
4.1 Data Collection . . . 9
4.1.1 Stationary Surveys . . . 10
4.1.2 Transect Surveys . . . 10
4.1.3 Equipment . . . 10
4.1.4 Landscape and Weather Covariates . . . 11
4.2 Acoustic Processing . . . 11
4.2.1 Subsetting for manual verification . . . 11
4.3 Statistical Analyses . . . 12
4.3.1 Stationary Activity Trends . . . 12
4.3.2 Transect Models . . . 14
4.3.3 Dynamic Occupancy . . . 15

5 Results **16**
5.1 Species Detection . . . 16
5.2 Activity and Abundance Trend Models . . . 18
5.3 Dynamic Occupancy Models . . . 19

6 Discussion **23**
6.1 Species Ranges . . . 23
6.2 Trend analyses . . . 23
6.3 Challenges and Lessons Learned . . . 25
6.4 Looking Forward . . . 25

7 Appendix A **27**
7.1 Pallid bat (*Antrozous pallidus*) . . . 27
7.2 Townsend's big-eared bat (*Corynorhinus townsendii*) . . . 30
7.3 Big brown bat (*Eptesicus fuscus*) . . . 34
7.4 Spotted bat (*Euderma maculatum*) . . . 38
7.5 Eastern-red bat (*Lasiurus borealis*) . . . 42
7.6 Hoary bat (*Lasiurus cinereus*) . . . 46
7.7 Silver-haired bat (*Lasionycteris noctivagans*) . . . 50
7.8 California Myotis (*Myotis californicus*) . . . 54
7.9 Western small-footed Myotis (*Myotis ciliolabrum*) . . . 58

7.10 Long-eared Myotis (*Myotis evotis*) 62
7.11 Little brown Myotis (*Myotis lucifugus*) 66
7.12 Northern Myotis (*Myotis septentrionalis*) 70
7.13 Fringed Myotis (*Myotis thysanodes*) 74
7.14 Long-legged Myotis (*Myotis volans*) 78
7.15 Yuma Myotis (*Myotis yumanensis*) 82

Appendix B: Species composition by site **87**

Literature Cited **289**

1 Executive Summary

The North American Bat Monitoring Program (NABat) is a multi-agency initiative administered by the US Geological Survey (USGS). In 2024, the North by Northwest (NNW) Bat Hub was formed by combining the former Alberta Hub and the BC and Southeast Alaska (BC-SE AK) Hub. The NNW Bat Hub leads NABat in British Columbia, with key support from provincial and state representatives and WCSC.

In 2025, 57 grid cells were monitored across British Columbia. Two complementary analyses, dynamic occupancy models and activity trend models, were used to evaluate population trends. Most species showed no discernible trend, suggesting bat populations in the province are generally stable. Little brown Myotis and western long-eared Myotis increased in both activity and occupancy, although the western long-eared Myotis increase may partly reflect improved detection since 2023. Fringed Myotis declined in occupancy, mainly in the Okanagan Region. Species ranges remained broadly stable, with northern Myotis and eastern red bat showing potential range expansions. Given growing threats to bat populations in BC, continued monitoring is essential to detect any changes in populations.

2 Land Acknowledgement

Biodiversity Pathways respectfully acknowledges that our work takes place on Treaty 8 and Douglas Treaties Territories as well as the traditional and unceded territories of First Nations and Métis Peoples across all regions of British Columbia, whose histories, languages, and cultures are deeply connected to the biodiversity we monitor. We acknowledge the traditional teachings of the lands that we work on, and that reciprocal, meaningful, and respectful relationships with Indigenous peoples make our work possible. We are deeply grateful for their stewardship of these lands, and we are committed to supporting Indigenous-led monitoring programs, while learning Indigenous ways of knowing, being, and doing.

3 Introduction

3.1 Overview of NABat and the NNW Bat Hub

North American Bat Monitoring Program (NABat) is a large-scale coordinated effort to monitor bat species to address gaps in knowledge and lack of long-term studies across North America (Loeb et al. 2015).The program is administered by the US Geological Survey (USGS) and implemented by the North by Northwest Bat (NNW) Hub in British Columbia, Alberta and S.E Alaska.

NNW Bat Hub was established in 2024 with collaboration from Government of British Columbia, Government of Alberta, and Government of Alaska. The hub was formed by combining the previous Alberta Hub and the BC and southeast Alaska (BC-SE_AK) Hub. The implementation of the program within British Columbia has key support from Wildlife Conservation Society Canada (WCSC), government staff, other wildlife biologists, Indigenous Nations, community members and naturalists across the province.

The centralization of efforts for the hub will expand exciting programs to coordinate more efficiently across jurisdictions, ensuring consistent and long-term bat monitoring through standardized data collection, and streamlined bioacoustic data management. It also seeks to enhance engagement with NABat, foster collaboration among stakeholders, and produce annual reports on regional monitoring.

3.2 Threats to bat populations in BC

Bat populations in British Columbia face a wide range of threats including forestry practices, anthropogenic expansion, and climate change among others. A comprehensive review of tese threats and their relative impacts on individual bat species can be found in the BC Bat Action Plan (Team 2024). Of these threats, White-Nose Syndrome and wind energy development have emerged as two of the highest-priority concerns.

3.2.1 White-Nose Syndrome

White-nose Syndrome (WNS) is a deadly fungal disease caused by the fungus *Pseudogymnoascus destructans* (*Pd*) that affects hibernating bats. As of 2026, WNS has been confirmed in nine Canadian provinces including Alberta, and *Pd* has been detected within British Columbia at two locations (Figure 1: CWHC-RCSF (2024)). Continued monitoring of populations in BC is critical as *Pd* and WNS continue to spread throughout the province.



Figure 1: White-nose syndrome (WNS) and Pseudogymnoascus destructans (Pd) occurrence in Canada. Source: CWHC-RCSF (2026)

3.2.2 Wind Energy Development

Wind energy farms has been linked to high mortality rates across all bat species in Canada, with long distance migratory species accounting for 73% of recorded fatalities (Zimmerling and Francis 2016). In British Columbia alone, one estimate predicts an annual mortality rate of 680 bats per wind turbine (Zimmerling and Francis 2016). There are currently close to 300 turbines active in the province across 10 different projects with the most turbines found in the Peace region (Figure 2). In late 2024, B.C announced nine new wind projects which would require the installation of over 300 new wind turbines across the province (Energy and Climate Solutions 2024). As of 2023, COSEWIC designated all three of B.C.'s long distance migratory species as endangered due in part to fatalities at wind turbine facilities (Status of Endangered Wildlife in Canada 2023).



Figure 2: Current wind turbines in British Columbia based on the Canadian Wind Turbine Database (Source: https://search.open.canada.ca/openmap/79fdad93-9025-49ad-ba16-c26d718cc070)

3.3 Acoustic Monitoring of Bats

Acoustic recording is a cost-effective method for monitoring bat populations across large geographic and temporal scales. However, several limitations affect the conclusions that can be drawn from these data:

1. **Imperfect species identification** — Unlike birds, bats rely on echolocation rather than song, and species with similar ecology and physiology often produce overlapping calls. While detectors are deployed to maximize recordings of diagnostic Search Phase calls, species identifiability will always vary depending on the local assemblage.
2. **Variable detectability** — Bat species are not equally detectable acoustically. Larger, wide-ranging species produce louder calls that travel farther, while smaller species with quieter calls or specialized habitat use are less reliably recorded. As a result, trend estimates cannot be produced for all species using acoustic methods alone.
3. **Data variability** — Many factors influence recording quality and bat activity, including wind speed, humidity, and local habitat. Additionally, stationary detectors cannot account for individual recapture, meaning acoustic data can only inform relative activity and occupancy.
4. **Processing limitations** — Auto-ID software enables efficient processing of large datasets but has known high misclassification rates for bats, making manual verification necessary to ensure accurate results.

3.4 Updates in Modelling Approaches

Bat monitoring data can be analyzed using different statistical approaches, each with distinct assumptions and outputs. Activity trend models examine changes in relative acoustic activity over time, providing a straightforward measure of whether bat detections are increasing, decreasing, or remaining stable across the monitoring network.

Dynamic occupancy models take a different approach, estimating the probability that a species occupies a given site while explicitly accounting for imperfect detection. Rather than treating a non-detection as an absence, these models distinguish between a species being truly absent and simply going undetected, an important consideration given the variable detectability of bats discussed above. They also track colonization and extinction probabilities across sites over time, providing insight into range shifts and local population dynamics.

The two approaches differ in sensitivity and therefore serve complementary roles. Activity data are inherently noisy, influenced by weather, local foraging conditions, and the number of individuals passing a detector, but they respond quickly to change and can provide an early indication of decline. Occupancy, in contrast, changes only when a species is lost from (or gains) sites, so a stronger and more sustained signal is needed before a trend becomes apparent. Occupancy models can therefore act as a second, more conservative test of patterns first suggested by activity data. When both approaches point in the same direction, the evidence for a real population change is stronger, and confidence in the results is correspondingly higher.

In 2025, both model types were run to examine differences in results and assess the complementary information each approach provides.

4 Methods

4.1 Data Collection

The BC NABat monitoring program was established in 2016 by Wildlife Conservation Society Canada (WCSC) and has since sampled 65 grid cells. Cells were drawn from the NABat Generalized Random Tessellation Stratified (GRTS) sampling frame (Loeb et al. 2015), and selected based on accessibility, the presence of bat habitat, suitable recorder sites and in GRTS priority order. Where possible, cells were also selected based on the precense of a relatively straight road of 30–45 km for a driving transect.

Table 1: Total number active grid cells and number of site nights sampled summarized by calendar year in British Columbia. Site nights are defined as the cumulative number of nights during which recorders were operational

Year	Number of Grid Cells Sampled	Number of BC Parks Grids	Site Nights
2016	22	2	532
2017	27	1	900
2018	47	15	939
2019	49	16	1489
2020	53	17	1530
2021	51	16	1448
2022	56	16	1635
2023	51	16	1436
2024	58	18	1410
2025	57	15	1826

In 2025 a total of 57 grid cells were monitored for a total of 1826 site nights (Table 1). Of the cells sampled in 2025 a total of 15 were sampled by BC Parks staff (Table 2). The network is well-distributed across the province, with a slight bias toward southern BC and population centres (Figure 3).

Table 2: Summary of yearly sampling of grid cells in BC Parks.

		Year							
GRTS Cell ID	Park	2016	2017	2018	2019	2020	2021	2022	202
243562	Kimberley								
6202	Goldstream								
986	Bowron								
38442	Churn Creek								
115322	Tweedsmuir								
145578	Premier Lake								
212906	Horne Lake								

265749	Kettle/Conkle
283586	Lakelse Lake
317428	Manning
16426	Carp
37274	Mount Robson
132650	Junction Sheep
172090	Mount Seymour
175354	Liard Hot Springs
230186	Skagit
17706	Wells Gray
26170	Yakoun
206394	Naikoon

4.1.1 Stationary Surveys

Two to four stationary detectors were deployed across at least two quadrants per grid cell for a minimum of seven nights. Sites are selected to maximize species detection and recording quality across different habitat types (see Rae and Lausen (2024) for full site selection criteria). Deployment occured between mid-May and mid-July, after bats have emerged from hibernation but before young bats are volant. Sites and timing are kept consistent between years to minimize variability in conditions affecting bat activity.

4.1.2 Transect Surveys

Where possible, driving transects were conducted following Loeb et al. (2015). Two replicates of a 30–35 km route are driven at 30–35 km/h with a rooftop detector, using straight roads that minimize switchbacks. Transects are conducted concurrently with stationary deployments.

4.1.3 Equipment

A wide range of equipment has been used across sampling locations in BC. Until 2024, stationary acoustic units were primarily Wildlife Acoustics SM4Bat-FS or SM2Bat units (Maynard, MA), alongside an increasing number of Titley Scientific Swift and Express units (Brisbane, AU). Mobile transects were conducted using either the Wildlife Acoustics EchoMeter Touch2Pro or Titley Scientific Anabat units (SD1/2 zero-cross or Walkabout models).

2025 represents a transition year for equipment standardization. All SM2Bat units are being replaced with Titley Scientific Chorus units, and Anabat units are being phased out in favour of the EchoMeter Touch2Pro for mobile transects. To account for detection differences between outgoing and incoming equipment, paired deployments were conducted at a number of grid cells this year, both stationary (SM2Bat and Chorus) and mobile (Anabat SD1 and EchoMeter Touch2Pro).

4.1.4 Landscape and Weather Covariates

We used the same covariates for the activity trend models as Rae and Lausen (2022). Where possible, temperature and relative humidity were recorded in each grid cell with onsite data loggers. Where loggers were unavailable, values came from the nearest climate station in the Pacific Climate Impacts Consortium (PCIC) Provincial Climate Data Set (PCDS), which aggregates and quality-controls observations from multiple provincial and federal networks. From these station records we also derived nightly mean wind speed (km/h) and nightly total precipitation (mm). Degree days above 10 °C since January 1 were calculated from daily temperature records for each site-night.

Sun and moon positions were calculated with the astral package (v3.2) in Python. Sunset and sunrise times defined the night period for each site-night. Moon illumination was then calculated as the proportion of the lunar disc illuminated multiplied by the proportion of the night the moon spent above the horizon. Site elevation was extracted from the Open-Meteo Elevation API. Distance to clutter and percent clutter were recorded by grid leaders at deployment, and distance to water and water body type were derived from the BC Freshwater Atlas.

For the dynamic occupancy models, we explored additional factors that might influence colonization and extinction: distance to the nearest road and distance to the nearest timber harvest, each measured from the survey location to the closest feature. Road features came from the Human Footprint Index (HFI) layer under development by the Geospatial Centre at Biodiversity Pathways (unpublished data). Harvest features came from the Province of British Columbia's consolidated cutblocks layer. All continuous covariates were standardized by centering on the mean and dividing by the standard deviation. Missing values were imputed with the global mean before standardization.

Since 2022, insect activity has also been sampled along transect routes using vehicle-mounted sticky traps. Samples are identified to order, and we will evaluate insect abundance as a potential covariate of bat activity in future years.

4.2 Acoustic Processing

In 2025 acoustic analysis mostly followed the standards set in previous years (Rae and Lausen 2022, 2024). This included running all files through two automatic classifiers, Kaleidoscope's Bats of North America 5.4.0 classifier and Sonobat 3.0's Northwestern British Columbia classifier, to remove files that do not contain bat pulses and giving a preliminary bat species identification.

Manual vetting largely follows the protocol of Reichert et al. (2018a) and previous analysis standards developed by WCSC (Rae and Lausen 2022) . The key difference in the 2024 and 2025 acoustic analysis compared to previous years is that not all automatic species identifications were manually verified. Instead, based on findings from Stratton et al. (2022) , at least 50% of recordings were selected for manual verification.

4.2.1 Subsetting for manual verification

Starting in 2024, the NNW Bat Hub implemented manual vetting of only a subset of all the bioacoustic data collected under the NABat program. Given the project's large scale, manually vetting all recordings is not feasible long-term. To address this, the NNW Bat Hub, in collaboration with WCSC and contractor Brian Paterson, developed procedures to ensure that selecting a subset

of recordings for manual review minimizes biases in analytical results. These standards prioritize the accurate identification of rare and difficult-to-recognize species to maintain the integrity of models and analyses.

To ensure data accuracy, manual verification was conducted based on the number of tagged recordings at each site. For sites with 200 or fewer tagged recordings, all non-noise recordings were manually reviewed. For sites with more than 200 tagged recordings, verification depended on the proportion of recordings assigned a species label. All tags for rare species (comprising 10% of prior years' identifications) and commonly misidentified species were fully vetted. Specifically, all Eastern Red Bat (*Lasiurus borealis*) tags were manually reviewed due to frequent misclassification as Little Brown Bats (*Myotis lucifugus*). Additionally, recordings of common species were randomly selected for verification until 50% of total recordings were reviewed. If fewer than 50% of recordings had species tags, all classifier-assigned tags were manually verified, and "NoID" recordings were randomly selected until 50% of all recordings were reviewed.

4.3 Statistical Analyses

4.3.1 Stationary Activity Trends

Stationary activity trends followed closely what was established by WCSC and Dr. Carl Schwarz in 2021 (Rae and Lausen 2022). This included conducting trend analyses examining activity patterns in the data collected to date using negative binomial GLMM models with a set of species-specific covariates (Table 3). Species-specific covariates for stationary models were selected using AIC-based stepwise selection (Table 3).

Table 3: Top-performing stationary models selected by AIC-based stepwise selection. Definitions of covariates are: Temperature = nightly temperature (°C); Wind speed = nightly mean wind speed (km/h); RH = nightly mean relative humidity (%); dd = cumulative degree days above 10 °C since January 1; Moon = proportion of the moon illuminated × proportion of the night the moon was above the horizon; Clutter = distance (m) to the nearest clutter object; Nearby water = whether the detector was within 100 m of water (1 = yes, 0 = no); I(YearS²) = quadratic year term allowing for non-linear temporal trends. A * B denotes the main effects of A and B plus their interaction.

Species	Species Code	Top Performing Model Covariates
Pallid Bat	ANPA	Nightly_Mean_Temp
Townsend's Big-eared Bat	COTO	dd,Nightly_Mean_Temp
Big Brown Bat	EPFU	Nightly_Mean_Windsp,moon*Distance_to_Clutter___m_,Nightly_Min_Temp
Spotted Bat	EUMA	Nightly_Mean_Temp
Eastern Red Bat	LABO	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Mean_Temp
Hoary Bat	LACI	moon*Distance_to_Clutter___m_,dd,Nightly_Mean_Temp,I(YearS^2)

Silver-haired Bat	LANO	Nightly_Mean_RH,Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Min_Temp
California Myotis	MYCA	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Mean_Temp
Western Small-footed Myotis	MYCI	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Mean_Temp
Long-eared Myotis	MYEV	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,dd,Nightly_Min_Temp
Little Brown Myotis	MYLU	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Mean_Temp
Northern Myotis	MYSE	Nightly_Mean_RH,moon*Distance_to_Clutter___m_,Water_Nearby
Fringed Myotis	MYTH	dd
Long-legged Myotis	MYVO	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,Nightly_Mean_Temp
Yuma Myotis	MYYU	Nightly_Mean_Windsp*Distance_to_Clutter___m_,Water_Nearby,dd

All models followed the same basic structure, modelling the total number of detected calls as a negative binomial response with log link:

```
Total detections ~ YearS + log(Sample Nights) + (1|Grid Cell ID/Quadrant)
+ species specific covariates
```

Where total detections is the total number of detected calls of a given species at a site in a given year. YearS is the standardized survey year (years since 2016) and provides the estimate of the temporal activity trend. log(Sample Nights) is the natural logarithm of the number of detector-nights and accounts for variation in sampling effort among site-years. The random intercept (1|Grid Cell ID/Quadrant) gives one random intercept per cell, plus one per specific quadrant within a cell. This allow baseline activity to vary among grid cells and the quadrants within them.

Model fit was assessed for each final model using simulation-based scaled residuals from the DHARMa package , with 1,000 simulations per model. For each model, we inspected residual quantile–quantile plots and plots of residuals against predicted values and against each predictor. We also tested residuals for uniformity (Kolmogorov–Smirnov test), dispersion and outliers. Because large sample sizes can make trivial deviations statistically significant, we considered the magnitude of each deviation alongside its p-value (Harrison et al. 2018). Models were considered adequately fitted if no more than one of these three tests was significant ($p < 0.05$). Where two or more tests were significant, we examined the residual plots and retained a model only if any deviations were small and showed no systematic pattern.

We conducted our statistical analyses at provincial and regional scales. For the regional analyses, we assigned all sampled grid cells to four geographic regions (Southwestern, Southcentral, Southeastern and Northern), each roughly corresponding to a different species set (Figure 3). Model fit was assessed in the same way as for the provicial scale stationary models.

4.3.1.1 Use of Automated Identification Results in Stationary Trend Analyses

Stationary trend analyses used AutoID classifications from Kaleidoscope Pro. Manual verification was applied only to flag non-bat noise; where manual review suggested a different species than the AutoID label, the original AutoID classification was retained.

This approach was first identified as necessary in 2023 (Rae and Lausen 2022), when analyses of manually verified data revealed a consistent positive trend across all species, this was interpreted as an artifact of reviewer learning over time rather than a true biological signal. As verifier experience increased across years, calls were increasingly assigned to specific species rather than left unidentified, producing an apparent increase in detections independent of actual bat activity. Because this observer effect is difficult to control for statistically, AutoID classifications are now used as the standard for all activity trend analyses, as automated classifiers apply consistent criteria across all years. While AutoID is subject to some misclassification, the error rate is consistent across years and does not confound detection of genuine temporal trends.

4.3.2 Transect Models

The province-wide transect abundance trend models used only completly manually verified detections. Transects are surveyed from a moving vehicle, so each bat is likely recorded only once as the vehicle passes. Call counts therefore serve as an index of relative abundance rather than of activity. When a detection was identified as a species couplet, it was counted toward each species in the couplet.

All models had the same basic structure: total detections were modelled as a negative binomial response with a log link:

```
Total detections ~ YearS + log(transect length) + (1|Grid Cell ID)
+ species specific covariates
```

Total detections is the number of calls of a given species detected on a transect in a given year. YearS is survey year expressed as years since 2016, and provides the estimate of the temporal activity trend. log(transect length) is the natural logarithm of the total length of the transect sampled and accounts for variation in sampling effort among transects. The random intercept (1|Grid Cell ID) allow baseline activity to vary among grid cells.

Covariates were limited to weather variables, because landscape-level covariates cannot be represented meaningfully for a survey that moves through the landscape. Species-specific covariates for transect models were selected based on those previously established in Rae and Lausen (2022) and were selected using AIC-based stepwise selection (Table 4). Model fit was assessed in the same way as for the stationary models.

Since transects are only conducted in roughly 50% of our grid cells on a typical year and each transect records fewer bats per night than stationary monitoring, we have elected not to conduct regional-scale transect trend analyses due to low sample sizes.

Table 4: Top-performing transect models selected by AIC-based stepwise selection. Definitions of covariates are: Temperature = nightly temperature (°C); Wind speed = nightly mean wind speed (km/h); RH = nightly mean relative humidity (%); Moon = proportion of the moon illuminated × proportion of the night the moon was above the horizon.

Species	Species Code	Top Performing Model Covariates
Pallid Bat	ANPA	Nightly_Mean_Temp
Big Brown Bat	EPFU	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Hoary Bat	LACI	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Silver-haired Bat	LANO	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
California Myotis	MYCA	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Long-eared Myotis	MYEV	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Little Brown Myotis	MYLU	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Long-legged Myotis	MYVO	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me
Yuma Myotis	MYYU	Nightly_Mean_Temp,Nightly_Mean_RH,Nightly_Me

4.3.3 Dynamic Occupancy

We fitted multi-season (dynamic) occupancy models (MacKenzie et al. 2003) to estimate trends in occupancy, colonization, and persistence while accounting for imperfect detection. This analysis used nightly acoustic detection/non-detection data from passive bat detectors deployed at all the sites in British Columbia (BC).

Initial occupancy (ψ) was modeled as a function of distance to water and elevation. Colonization (γ) and persistence (φ) were modeled as a function of distance to water, elevation, year-varying distance to harvest, and distance to road. Detection probability (p) was modeled as a function of percent clutter, nightly mean temperature, and Julian night.

We excluded 2016 because few sites were surveyed that year. Starting the time series in 2016 would have required the model to estimate initial occupancy ( ) for the many sites not yet sampled, and a poorly informed can propagate through the colonization and extinction estimates and produce false trends.

To reduce temporal autocorrelation between consecutive nights, we subsampled every other night, using a maximum of four nights per site-year as repeat surveys. Sites were included if they had at least two nights of data in a given year and data from at least two years. After filtering, 185 sites in 63 grid cells in British Columbia were retained. Models were fit separately for each of the 15 bat species.

Models were fit in a Bayesian framework using JAGS (Version 4.3.2) via the jagsUI package (Version 1.6.3) in r (Version 4.5.1). We ran 3 chains in parallel for 30,000 iterations each, discarding the first 15,000 as burn-in and thinning the remainder by 10, yielding 4,500 posterior samples for inference. Priors on all regression coefficients, including region-specific intercepts, were normal with mean 0 and precision 0.368 (SD 1.65 on the logit scale). Standard deviations of the year-level random effects on colonization, persistence, and detection were assigned half-Cauchy(0, 2.5) priors.

Convergence was assessed using Rhat diagnostics, with values below 1.1 taken to indicate convergence. Results are summarized using posterior means and 95% Bayesian credible intervals. We calculated

an overall occupancy trend metric ( ) as the ratio of estimated occupancy in the final year to that in the initial year; values of > 1 with 95% credible intervals entirely above 1 indicate an increase in occupancy, values <1 with intervals entirely below 1 indicate a decline, and intervals overlapping 1 indicate an uncertain or stable.

5 Results

2025 marked the tenth year of coordinated NABat sampling done in BC. A total of grid cells were sampled with the support of over 60 individuals including contractors, volunteers and staff members from BC Parks, BC WLRS, and BC Ministry of Forests. Grid leaders have also collected a total of 16 transect insect sticky trap samples in 2025, which will be added to previous years of data to explore the addition of insect abundance as a covariate in the transect models.

5.1 Species Detection

All 15 species monitored by the program were detected in 2025. Total acoustic activity varied substantially among grid cells and regions (Figure 3). Little brown myotis and silver-haired bat remain two of the most widespread species in the province. Notable changes in 2025 include:

- **Pallid bat** was detected at only one site, in each of 2023 and 2024. In 2025 it was detected at 14 sites across 6 grid cells, its highest site count since monitoring began and slightly above 2022 (13 sites across 6 grid cells). Pallid bat was first recorded in the Kootenay Region in 2022 and was not detected there in 2023 or 2024. In 2025 it was detected at 8 Kootenay sites across 4 grid cells. Two of these grid cells (111466 and 243562) had no previous detections.
- **Eastern red bat** was detected at 35 sites across 17 grid cells in 2025. That is its highest site count to date, although it was found in more grid cells in 2022 (21). Four grid cells had no previous detections: 111466 and 243562 (Kootenay), 194666 (Okanagan) and 186330 (Northern). Eastern red bat was not detected in the South Coastal Region in 2025.
- **Northern myotis** was detected at 15 sites across 10 grid cells in 2025. In the southern Kootenay Region it was detected for the first time at 4 sites across 3 grid cells (20330, 89962 and 218282), near the southern edge of its range.
- **Long-legged myotis** was detected at 76 sites across 39 grid cells in 2025, down from 92 sites across 45 grid cells in 2023 (the highest year of detection for the species). Most of these losses were in the Okanagan Region (8 sites, 5 grid cells) and the South Coastal Region (5 sites, 2 grid cells). The remaining losses were in the Kootenay and Northern regions (3 sites and 1 grid cell each).

Figure 3 is an interactive map. Readers can zoom in to see the individual stationary monitoring sites (blue circles), and click a site to open a pop-up with a summary pie chart of total detections by species and year. Readers can also search for specific grid cells using the icon.



Figure 3: Acoustic bat monitoring sites in British Columbia, 2016–2025. Regions were defined by Wildlife Conservation Society Canada (WCSC) from species range maps to group areas with similar species composition.

5.2 Activity and Abundance Trend Models

2025 was the first year in which provincial-scale trends in stationary acoustic activity could be modelled for all 15 bat species (**?@tbl-provtrend**) . Model diagnostics indicated that all provincial stationary models fit adequately for interpreting the direction of activity trends, although minor deviations from model assumptions occurred in most models. Because these assumptions were not fully met, the magnitudes of the estimated trends should be interpreted with caution, and we therefore focus on trend direction throughout. For mobile transects, low numbers of recordings limited reliable trend estimation to 8 of the 15 species.

Table 5: Provincial stationary and transect trend estimates from 2017-2025 NABat monitoring in BC adjusted for covariates. Trends estimated on a logarithmic scale, thus a trend estimate of 0.05 corresponds to an annual change of 5%. Significant results (p value < 0.05) are highlighted in bold.

	Stationary Models		Transect Models	
Species	Trend Estimate ± 95%CI	P-value	Trend Estimate ± 95%CI	P-value
Pallid Bat	**0.13 ± 0.07**	**<0.001**		
Townsend's Big-eared Bat	**0.10 ± 0.05**	**<0.001**		
Big Brown Bat	**0.05 ± 0.04**	**0.007**	0.06 ± 0.08	0.142
Spotted Bat	0.01 ± 0.12	0.869		
Eastern Red Bat	-0.03 ± 0.04	0.264		
Hoary Bat	-0.02 ± 0.15	0.820	-0.05 ± 0.14	0.461
Silver-haired Bat	0.02 ± 0.04	0.294	0.06 ± 0.08	0.116
California Myotis	0.01 ± 0.03	0.464	0.05 ± 0.12	0.415
Western Small-footed Myotis	0.00 ± 0.05	0.978		
Long-eared Myotis	**0.10 ± 0.04**	**<0.001**	0.06 ± 0.13	0.412
Little Brown Myotis	0.01 ± 0.03	0.472	**0.19 ± 0.08**	**<0.001**
Northern Myotis	-0.02 ± 0.07	0.527		
Fringed Myotis	0.01 ± 0.04	0.713		
Long-legged Myotis	-0.02 ± 0.03	0.373	**0.32 ± 0.12**	**<0.001**
Yuma Myotis	**0.06 ± 0.04**	**0.002**	0.09 ± 0.16	0.294
Canyon Bat				
Mexican Free-tailed Bat				

At the provincial scale, no species showed a significant negative trend in either the stationary or the transect models. Stationary activity increased significantly for five species: pallid bat, Townsend's big-eared bat, big brown bat, long-eared Myotis and Yuma Myotis (Table 5) . In the transect models, only little brown Myotis and long-legged Myotis showed significant positive trends.

All regional trend models converged; however, after inspecting model fit, we judged seven region-specific results to be unreliable and excluded them. Most of the remaining regional trends were not significant (Table 6). The Kootenay and South Coastal regions had the most significant trends, all of them positive. In the Kootenay region, activity increased for five species (pallid bat, Townsend's big-eared bat, western small-footed Myotis, long-eared Myotis and little brown Myotis). In the

South Coastal region, activity increased for five species (Townsend's big-eared bat, big brown bat, California Myotis, long-eared Myotis and Yuma Myotis). Big brown bat also increased in the Okanagan region. The only significant negative regional trend was for Yuma Myotis in the Northern region.

Table 6: Regional trend estimates from 2017-2024 NABat monitoring in BC adjusted for covariates. Trends estimated on a logarithmic scale, thus a trend estimate of 0.05 corresponds to an annual change of 5%. Significant results (p value < 0.05) are highlighted in bold.

	Kootenay Region		Northern Region	
Species	Estimate ± 95%CI	P-value	Estimate ± 95%CI	P-value
Pallid Bat	**0.23 ± 0.13**	**<0.001**		
Townsend's Big-eared Bat	**0.08 ± 0.07**	**0.017**		
Big Brown Bat	0.03 ± 0.07	0.396	-0.00 ± 0.13	0.964
Spotted Bat				
Eastern Red Bat	0.05 ± 0.09	0.269	-0.03 ± 0.10	0.503
Hoary Bat				
Silver-haired Bat	0.03 ± 0.06	0.316	0.06 ± 0.11	0.298
California Myotis	0.01 ± 0.06	0.760	-0.10 ± 0.10	0.064
Western Small-footed Myotis	**0.10 ± 0.07**	**0.009**		
Long-eared Myotis	**0.14 ± 0.06**	**<0.001**	0.04 ± 0.10	0.413
Little Brown Myotis	**0.10 ± 0.06**	**<0.001**	-0.04 ± 0.08	0.381
Northern Myotis	-0.10 ± 0.12	0.100	0.04 ± 0.08	0.324
Fringed Myotis	0.07 ± 0.08	0.103		
Long-legged Myotis	0.02 ± 0.06	0.487		
Yuma Myotis	0.05 ± 0.08	0.217	**-0.17 ± 0.17**	**0.048**

5.3 Dynamic Occupancy Models

We assessed occupancy trends in two ways: the mean yearly change in occupancy across 2017–2025, and the posterior probability that occupancy in 2025 was higher than in 2017. We considered a posterior probability above 0.95 to be strong evidence of an increase, and one below 0.05 to be strong evidence of a decline.

Overall, occupancy increased across British Columbia between 2017 and 2025 for four of the 15 species modelled (Figure 4). Western long-eared Myotis, little brown Myotis, northern Myotis and silver-haired bat all showed a positive mean yearly trend and a high posterior probability of an increase in occupancy from 2017 to 2025, indicating strong evidence of an overall increase. Fringed Myotis was the only species with a consistent decrease in occupancy: its mean yearly change was negative, and the posterior probability that 2025 occupancy exceeded 2017 occupancy was very low, indicating strong evidence of a decline.

Yuma Myotis, western small-footed Myotis, spotted bat, pallid bat and hoary bat showed no detectable occupancy trend. Their mean yearly change was not significant, and the posterior probability of an increase from 2017 to 2025 gave no clear evidence of either an increase or a decline.

Big brown bat showed no significant mean yearly change, but the posterior probability of an increase between 2017 and 2025 was low, which points to a decline over that period. This was driven by a high posterior mean occupancy in the first year and a slightly below-average posterior mean in the last year (Appendix A), so it more likely reflects a stable population with an unusually high starting point than a true decline.

Eastern red bat, long-legged Myotis, Townsend's big-eared bat and California Myotis had positive mean yearly trends. However, the posterior probability that occupancy was higher in 2025 than in 2017 was not high enough to conclude that occupancy increased overall. This pattern reflects nonlinear change across years (see species specific results in Appendix A). Occupancy fluctuated enough to produce a positive mean yearly trend, but the net difference between the first and last years was small, suggesting these species remained relatively stable over the monitoring period.

Among the regional models, the Kootenay Region showed the most positive trends (Figure 5). Eastern red bat, silver-haired bat, western small-footed Myotis, western long-eared Myotis, little brown Myotis and long-legged Myotis all had a positive mean yearly trend and a high posterior probability of an increase in occupancy from 2017 to 2025.

In the Northern Region, big brown bat had a negative mean yearly trend and a low posterior probability of an increase, indicating a decline. Eastern red bat also had a low posterior probability of an increase, although its mean yearly change was not significant.

In the Okanagan Region, Townsend's big-eared bat and eastern red bat had positive mean yearly trends and high posterior probabilities of an increase. Fringed Myotis declined, with both a negative mean yearly trend and a low posterior probability of an increase. Yuma Myotis also had a low posterior probability of an increase, although its mean yearly change was not significant.

In the South Coastal Region, hoary bat had a negative mean yearly trend and a low posterior probability of an increase. Silver-haired bat, California Myotis, western long-eared Myotis and Yuma Myotis had positive mean yearly trends and high posterior probabilities of an increase.

To show how occupancy varied across space and time, we mapped the model-estimated probability of occupancy at each site for every year from 2017 to 2025 (**?@fig-occ-slider**). The map lets readers select a species and move through the years to see where occupancy has changed. Each point's colour shows the model's estimate of for yearly probability of occupancy, while its shape shows whether the species was detected that year (circle) or not (triangle). Because the occupancy model accounts for imperfect detection, a site where a species was not detected can still have a high estimated probability of occupancy. This is especially likely for species that are hard to detect acoustically, or at sites with few survey nights in a given year. The map therefore shows where a species was likely present, not just where it was recorded.

Because the dynamic occupancy model estimates detection probability separately for each year, it also shows how detectability changed over the monitoring period for each species. This lets us account for changes in sampling and processing, such as the equipment change in 2025, so that shifts in detectability are not mistaken for changes in occupancy. The species-specific results (Appendix A) show that the probability of detecting pallid bat and eastern red bat increased in 2025, whereas detection probability for all other species stayed within its normal year-to-year variation.



Figure 4: Estimated occupancy trends for 15 bat species in British Columbia (2017–2025) fit across all monitoring sites. Points and bars show the posterior mean and 95% credible interval of the linear change in occupancy per year. The dashed line marks zero, where there is no change in mean yearly opccupancy. Colour shows the posterior probability (at 0.95) that the occupancy in 2025 was higher (green) or lower (red) than in 2017 . Grey means the probability is uncertain. Slope and colour can disagree when trends are non-linear.



Figure 5: Regional estimated occupancy trends for 15 bat species in British Columbia (2017–2025). Points and bars show the posterior mean and 95% credible interval of the linear change in occupancy per year. The dashed line marks zero, where there is no change in mean yearly occupancy. Colour shows the posterior probability (at 0.95) that the occupancy in 2025 was higher (green) or lower (red) than in 2017 . Grey means the probability is uncertain. Slope and colour can disagree when trends are non-linear.

6 Discussion

6.1 Species Ranges

Species ranges across the project period (2016–2025) remain broadly stable, with no grid cells showing conspicuous or sustained absences of previously detected species. In 2025 there was a normal variation in the disctribution of species detections across the province, but three species are worth highlighting:

- **Northern Myotis** was detected for the first time in three grid cells (20330, 218282, and 89962) in the southern Kootenay region. These cells lie at the southern edge of the species' known range, and the detections may indicate a small range expansion. Capture or genetic (guano) surveys will be needed to confirm its presence.
- **Long-legged myotis** was detected at fewer sites in 2025 (76 sites across 39 grid cells) than in 2024 (85 sites across 42 grid cells). Most of the decline was in the South Coastal Region: among sites surveyed in both years, detections fell from 18 sites to 12, with 7 sites losing the species and 1 gaining it. Year-to-year turnover in site-level detections is common for most species in the program, so this decline likely reflects natural variation rather than a population change. However, the proportion of sites with detections has now declined for two consecutive years, so this species should be monitored closely in future years.
- **Eastern red bat** detections increased in the Kootenay and Okanagan regions in 2025. The species was detected in 17 grid cells, the most to date, including four with no previous detections (111466, 243562, 194666, and 186330). Although many knowledge gaps remain for this species, these results continue to support a westward range expansion (Solick et al. 2020). In contrast, eastern red bat was not detected in the South Coastal Region in 2025 after being recorded there every year from 2021 to 2024. Its calls overlap substantially with those of little brown Myotis, making diagnostic recordings difficult to obtain, so this apparent absence may partly reflect the difficulty of confirming the species acoustically rather than a true loss.

6.2 Trend analyses

In 2025, we used two complementary analyses to assess population trends. Dynamic occupancy models estimate how the proportion of sites occupied by each species changes over time, while accounting for imperfect detection (the chance that a species is present at a site but not recorded). Activity trend analyses estimate changes in the number of bat passes recorded per night, which reflects how heavily sites are used. Because the two analyses capture different aspects of population change, agreement between them strengthens confidence in the results.

Acoustic data suggest that bat populations in BC are generally stable. Across both analyses, most species showed no discernible trend. Four species (pallid bat, Townsend's big-eared bat, big brown bat, and Yuma Myotis) showed positive trends in activity but no trend in occupancy (Table 7). This pattern suggests that activity is increasing at sites where these species already occur, rather than the species occupying new sites. These species may be increasing, but confirming this will require positive activity trends sustained over several more years, ideally accompanied by a positive occupancy trend.

Table 7: Summary of provincial-scale population trends for bat species in British Columbia. Activity and transect trends are based on the year coefficient (2016-2025) from models of nightly stationary and mobile transect activity, with increases or decreases identified at $p < 0.05$. Dynamic occupancy trends are based on the change in occupancy probability between the first and last survey year across all sites (2017-2025), with increases or decreases identified when the 95% credible interval excluded zero. "Not assessed" indicates species that were excluded from, or not modelled in, a given analysis.

```
# A tibble: 15 x 5
   species common_name              act_trend tran_trend   occ_trend
   <chr>   <chr>                    <fct>     <fct>        <fct>
 1 ANPA    Pallid bat               Increase  Not assessed No trend
 2 COTO    Townsend's big-eared bat Increase  Not assessed No trend
 3 EPFU    Big brown bat            Increase  No trend     No trend
 4 EUMA    Spotted bat              No trend  Not assessed No trend
 5 LABO    Eastern red bat          No trend  Not assessed No trend
 6 LACI    Hoary bat                No trend  No trend     No trend
 7 LANO    Silver-haired bat        No trend  No trend     Increase
 8 MYCA    California Myotis        No trend  No trend     No trend
 9 MYCI    Small-footed Myotis      No trend  Not assessed No trend
10 MYEV    Long-eared Myotis        Increase  No trend     Increase
11 MYLU    Little brown Myotis      No trend  Increase     Increase
12 MYSE    Northern Myotis          No trend  Not assessed No trend
13 MYTH    Fringed Myotis           No trend  Not assessed Decrease
14 MYVO    Long-legged Myotis       No trend  Increase     No trend
15 MYYU    Yuma Myotis              Increase  No trend     No trend
```

Little brown Myotis showed a positive trend in transect activity (Table 5) and an overall increase in occupancy (Figure 4; Appendix A), suggesting the species is increasing across the province. However, *Pseudogymnoascus destructans* (Pd), the fungus that causes white-nose syndrome (WNS), was confirmed in the South Coastal Region in 2026 (Kellner, pers. comm.), so this trend is likely to change in coming years. Continued monitoring will be essential for detecting the effects of WNS on this species in BC.

Long-eared Myotis increased in both stationary acoustic activity and occupancy (Table 5;Figure 4), with increases in both analyses concentrated in the Kootenay and South Coastal regions. The reason for this increase is unclear, and it may be partly an artifact of detection. Detection probability for this species was higher from 2023 to 2025 than from 2017 to 2022 (Appendix A), which suggests that something changed in 2023 that made the species easier to detect. Equipment updates are the most likely cause. If equipment changes can be ruled out, a closer look at changes in the Kootenay and South Coastal regions in this period is needed to help explain the increase.

Silver-haired bat may also be increasing in the Kootenay and South Coastal regions, although this is based on occupancy results alone (Table 5; Figure 4). A similar increase in occupancy has been reported for this species in Southeast Alaska (Hurtado and Hart 2026). This trend warrants close monitoring, particularly given planned increases in wind energy development in northern BC.

Fringed Myotis showed no trend in activity but declined in occupancy, driven primarily by a decrease in the Okanagan Region. Continued monitoring will be needed to confirm this decline and to help identify potential causes. Lile et al. (2026) found that the number of Myotis species captured has declined since the arrival of WNS in South Dakota, and fringed Myotis was among the species with significantly fewer captures. A decline in fringed Myotis is therefore a concern: if the species is already decreasing before WNS becomes established in BC, the disease could accelerate that

6.3 Challenges and Lessons Learned

6.3.0.1 Timing of Data Retrieval and Submission

Field crews across British Columbia continue to reliably collect recordings for their grid cells. However, organizing and submitting these recordings, along with associated metadata, remains a challenge. Delays in data submission from grid leaders extend processing and analysis times, making it difficult to complete reports by the end of the fiscal year. The transition in project management this year further complicated data submission. We anticipate these challenges will be resolved in future years as teams adjust to the new process. To improve efficiency, we plan to increase reminders to grid leaders and provide additional support to address ongoing challenges in data organization and submission.

6.4 Looking Forward

The BC NABat program has yielded a large and robust acoustic dataset that provides solid baseline information on distribution of 15 resident BC bat species. The data that has been collected so far are an invaluable reference point for BC's bat distributions and activity that improve with each additional year of monitoring. Key considerations for the future of the program include:

- **Manual Verification Efficiency.** For many years, the program manually verified 100% of acoustic recordings. As the program continues to expand, this level of effort is no longer sustainable, and manual vetting was reduced to 50% of recordings following the results of (**stratton2022?**). Since 2023, manual verification has been used only to confirm species presence, while activity metrics are based on automated Kaleidoscope classifications. For the occupancy analyses, manual verification is only needed to confirm species presence at each site and to remove false positive IDs. The additional verification the program currently completes is valuable for maintaining a clean dataset that can be used to evaluate how vetting effort affects results, as it already has been in (**stratton2022?**). However, it is not required for the analyses presented in this report. If the program needs to find efficiencies, reducing manual verification to the level required for occupancy analyses should be considered
- **Efficiencies in monitoring.** Grid leaders should continue to be encouraged to use Survey123 to streamline site deployment; in 2024 and 2025 this platform was not used consistently across all grids. Using it also helps ensure that metadata are consistent and complete. In 2026 we can also examine efficiencies in weather data collection, specifically evaluating whether continued use of on-site temperature and humidity loggers yields results comparable to weather data drawn from the nearest weather stations.

- **Adding and maintaining grid cells.** Changes in priorities and budgets have reduced repeat visits at some grid cells across the province. Supporting continued sampling at established grid cells is key to tracking population changes over time, especially given the imminent threat of WNS. Collaboration with other agencies and organizations can help sustain monitoring at existing grid cells and expand it to new ones. One approach being explored by the NNW Bat Hub is engaging industry partners, such as those in the mining, forestry, and wind energy sectors, in coordinated monitoring within their operating areas, with data contributed both to the hub and to the broader NABat effort. These partnerships may be easier to establish where regulatory frameworks support them, for example by requiring that bat monitoring data collected for industry projects be submitted to the hub and NABat.

7 Appendix A

7.1 Pallid bat (*Antrozous pallidus*)






ANPA — Pallid bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


ANPA — Pallid bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-4
-2
0
2
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.2 Townsend's big-eared bat (*Corynorhinus townsendii*)




COTO — Townsend's big-eared bat: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 30
150 detection nights
Northern Region
60 sites
NOT DETECTED
Okanagan Region
33 sites, detected at 23
109 detection nights
South Coastal Region
56 sites, detected at 37
156 detection nights
Not detected
(assumed absent)
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


COTO — Townsend's big-eared bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


COTO — Townsend's big-eared bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
-1
0
1
2
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.3 Big brown bat (*Eptesicus fuscus*)




EPFU — Big brown bat: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 38
396 detection nights
Northern Region
60 sites, detected at 38
234 detection nights
Okanagan Region
33 sites, detected at 32
443 detection nights
South Coastal Region
56 sites, detected at 48
479 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


EPFU — Big brown bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


EPFU — Big brown bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1.5
-1.0
-0.5
0.0
0.5
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.4 Spotted bat (*Euderma maculatum*)



EUMA — Spotted bat: dynamics by region

Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.




EUMA — Spotted bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


EUMA — Spotted bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
0
2
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.5 Eastern-red bat (*Lasiurus borealis*)




LABO — Eastern red bat: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 21
65 detection nights
Northern Region
60 sites, detected at 31
112 detection nights
Okanagan Region
33 sites, detected at 16
58 detection nights
South Coastal Region
56 sites, detected at 12
16 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


LABO — Eastern red bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


LABO — Eastern red bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1
0
1
2
3
4
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.6 Hoary bat (*Lasiurus cinereus*)




LACI — Hoary bat: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 37
358 detection nights
Northern Region
60 sites, detected at 25
88 detection nights
Okanagan Region
33 sites, detected at 31
377 detection nights
South Coastal Region
56 sites, detected at 49
461 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year

LACI — Hoary bat: P(occupancy) by site and year

Surveyed site-years only. Triangles = not detected that year; their colour is model inference.




LACI — Hoary bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1.0
-0.5
0.0
0.5
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.7 Silver-haired bat (*Lasionycteris noctivagans*)




LANO — Silver-haired bat: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 38
532 detection nights
Northern Region
60 sites, detected at 55
599 detection nights
Okanagan Region
33 sites, detected at 33
622 detection nights
South Coastal Region
56 sites, detected at 55
872 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


LANO — Silver-haired bat: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


LANO — Silver-haired bat: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1.0
-0.5
0.0
0.5
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.8 California Myotis (*Myotis californicus*)




MYCA — California myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 37
435 detection nights
Northern Region
60 sites, detected at 28
242 detection nights
Okanagan Region
33 sites, detected at 31
436 detection nights
South Coastal Region
56 sites, detected at 56
683 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


MYCA — California myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYCA — California myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1.5
-1.0
-0.5
0.0
0.5
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.9 Western small-footed Myotis (*Myotis ciliolabrum*)




MYCI — Western small-footed myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 13
66 detection nights
Northern Region
60 sites
NOT DETECTED
Okanagan Region
33 sites, detected at 31
467 detection nights
South Coastal Region
56 sites, detected at 4
8 detection nights
Not detected
(assumed absent)
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


MYCI — Western small-footed myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYCI — Western small-footed myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
-1
0
1
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.10 Long-eared Myotis (*Myotis evotis*)




MYEV — Western long-eared myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 39
498 detection nights
Northern Region
60 sites, detected at 58
468 detection nights
Okanagan Region
33 sites, detected at 33
375 detection nights
South Coastal Region
56 sites, detected at 53
456 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


MYEV — Western long-eared myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYEV — Western long-eared myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
-1
0
1
2
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.11 Little brown Myotis (*Myotis lucifugus*)




MYLU — Little brown myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 39
675 detection nights
Northern Region
60 sites, detected at 60
850 detection nights
Okanagan Region
33 sites, detected at 33
554 detection nights
South Coastal Region
56 sites, detected at 54
871 detection nights
Occupancy
Colonization
Persistence
Probability
Year
0.00
0.25
0.50
0.75
1.00
2017
2019
2021
2023
2025


MYLU — Little brown myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYLU — Little brown myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1
0
1
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.12 Northern Myotis (*Myotis septentrionalis*)




MYSE — Northern myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 12
19 detection nights
Northern Region
60 sites, detected at 31
137 detection nights
Okanagan Region
33 sites
NOT DETECTED
South Coastal Region
56 sites
NOT DETECTED
Not detected
(assumed absent)
Not detected
(assumed absent)
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


MYSE — Northern myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYSE — Northern myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-1
0
1
2
3
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.13 Fringed Myotis (*Myotis thysanodes*)




MYTH — Fringed myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 14
47 detection nights
Northern Region
60 sites
NOT DETECTED
Okanagan Region
33 sites, detected at 22
166 detection nights
South Coastal Region
56 sites, detected at 3
8 detection nights
Not detected
(assumed absent)
Occupancy
Colonization
Persistence
Probability
Year
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025


MYTH — Fringed myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYTH — Fringed myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
-1
0
1
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.14 Long-legged Myotis (*Myotis volans*)




MYVO — Long-legged myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 38
402 detection nights
Northern Region
60 sites, detected at 50
506 detection nights
Okanagan Region
33 sites, detected at 32
298 detection nights
South Coastal Region
56 sites, detected at 53
451 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year

MYVO — Long-legged myotis: P(occupancy) by site and year

Surveyed site-years only. Triangles = not detected that year; their colour is model inference.




MYVO — Long-legged myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-0.5
0.0
0.5
1.0
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0



7.15 Yuma Myotis (*Myotis yumanensis*)




MYYU — Yuma myotis: dynamics by region
Posterior mean (95% CI). Colonization and persistence are realised rates; persistence is blank where too few sites were occupied to estimate it.
Kootenay Region
39 sites, detected at 26
232 detection nights
Northern Region
60 sites, detected at 13
125 detection nights
Okanagan Region
33 sites, detected at 26
226 detection nights
South Coastal Region
56 sites, detected at 52
600 detection nights
Occupancy
Colonization
Persistence
Probability
1.00
0.75
0.50
0.25
0.00
2017
2019
2021
2023
2025
Year


MYYU — Yuma myotis: P(occupancy) by site and year
Surveyed site-years only. Triangles = not detected that year; their colour is model inference.
2017
2018
2019
2020
2021
2022
2023
2024
2025
57
54
51
48
-130
-125
-120
-115
P(occupied)
0.00 0.25 0.50 0.75 1.00
Detected
Not detected


MYYU — Yuma myotis: covariate effects
Posterior mean and 95% credible interval. BC-wide slopes, shared across regions.
Elevation
Distance to water
Initial occupancy
Elevation
Distance to water
Distance to road
Distance to harvest
Colonization
Elevation
Distance to water
Distance to road
Distance to harvest
Persistence
Clutter
Temperature
Julian date
Detection
-2
-1
0
1
Effect on logit scale (per 1 SD of covariate)
Positive (95% CI above 0)
Negative (95% CI below 0)
95% CI includes 0


MYYU — Yuma myotis: detection probability by year
Per-visit detection at average covariate values (95% CI). Dashed line = overall mean.
Detection probability (p)
1.00
0.75
0.50
0.25
0.00
2017
2018
2019
2020
2021
2022
2023
2024
2025
Year

Appendix B: Species composition by site

Pie charts show the proportion of identified bat passes by species for each year a site was surveyed (most recent first); totals are given under each year. Tables give the underlying counts.

110378 NE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	17	93	57	189	190	161	170	677	1,554
MYLU	6	30	31	235	382	157	161	217	1,219
MYCI	9	3	5	105	93	36	32	62	345
LACI	0	4	2	25	130	49	9	101	320
EPFU	6	13	6	37	42	14	23	70	211
MYCA	2	4	5	33	33	9	20	11	117
MYEV	1	6	2	18	18	15	22	34	116
LABO	0	0	1	0	16	10	1	35	63
EUMA	0	3	1	12	30	3	7	4	60
ANPA	0	4	0	9	8	8	4	25	58
MYVO	2	0	0	19	6	11	9	8	55
MYYU	0	1	0	7	12	8	7	7	42
MYTH	0	2	3	0	2	0	0	1	8
COTO	0	0	0	2	1	0	0	1	4
MYSE	0	0	0	0	0	0	0	1	1

110378 NW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	293	4,675	833	615	1,483	2,206	3,149	922	1,484	908	16,568
MYLU	177	5,007	438	326	176	916	786	276	188	506	8,796
EPFU	138	137	56	10	123	252	177	33	202	120	1,248
LACI	13	497	2	11	5	73	157	35	8	99	900
MYCI	5	277	170	8	12	80	66	27	23	50	718
MYCA	11	229	59	28	38	83	53	24	24	55	604
MYEV	9	75	9	37	23	95	79	62	21	127	537
MYVO	0	101	12	6	9	20	9	20	2	15	194
MYYU	15	54	6	4	5	16	9	9	0	5	123
LABO	0	0	0	0	0	3	23	24	5	31	86
MYTH	3	4	6	0	10	12	9	13	0	0	57
ANPA	0	0	1	1	0	28	6	6	0	10	52
COTO	0	5	0	1	0	10	10	3	2	1	32
EUMA	0	3	7	0	3	0	5	4	1	0	23
PAHE	0	0	0	0	0	0	0	0	1	0	1

110378 SE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	9	94	1,513	656	3,623	144	254	51	6,344
LACI	3	201	3	24	37	49	0	6	323
MYCI	0	35	2	7	28	73	15	40	200
MYEV	1	12	2	15	11	11	6	17	75
MYLU	8	7	0	12	22	8	5	12	74
EPFU	0	0	0	5	30	1	11	5	52
MYVO	1	3	0	1	4	7	5	10	31
MYCA	1	3	0	1	2	2	5	4	18
ANPA	0	3	0	4	2	2	1	1	13
LABO	0	0	0	0	3	4	0	6	13
COTO	0	5	1	0	1	0	0	3	10
MYTH	0	4	0	1	1	0	0	0	6
EUMA	0	0	0	0	2	1	0	0	3
MYYU	0	1	0	0	0	0	0	0	1
MYSE	0	0	0	0	0	0	0	1	1

110378 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
LACI	3	806	1	2	16	101	1,047	12	10	2	2,000
LANO	19	586	20	41	22	172	121	101	244	326	1,652
MYCI	1	316	46	11	14	219	190	71	108	151	1,127
MYCA	24	253	12	14	4	70	54	22	36	22	511
EPFU	38	65	4	1	3	43	18	15	16	61	264
MYLU	10	133	4	2	1	19	26	19	15	29	258
MYEV	5	87	8	27	31	20	26	16	13	13	246
MYVO	0	23	1	1	8	9	32	7	16	27	124
MYYU	2	16	2	0	0	7	3	0	2	3	35
ANPA	0	2	0	0	0	9	5	2	1	12	31
MYTH	0	4	2	1	1	1	3	2	4	0	18
COTO	0	1	0	1	1	2	2	1	2	1	11
LABO	0	0	0	0	0	0	2	2	1	3	8
EUMA	0	0	0	1	1	0	3	0	1	0	6
PAHE	0	2	0	0	0	0	0	0	0	0	2
MYSE	0	0	0	0	0	0	0	0	0	1	1

111466 NE



Species	2023	2025	Total
MYLU	462	997	1,459
MYYU	32	288	320
EPFU	228	57	285
LACI	104	99	203
MYCA	58	138	196
LANO	68	23	91
MYVO	25	55	80
MYCI	23	37	60
MYEV	17	10	27
LABO	1	22	23
MYTH	3	7	10
COTO	3	0	3
PAHE	0	3	3
MYSE	0	1	1

111466 NW



Species	2023	2025	Total
MYLU	12	24	36
LACI	32	0	32
MYTH	14	9	23
LANO	7	0	7
EPFU	3	3	6
MYVO	2	3	5
LABO	0	5	5
MYEV	1	3	4
MYCA	1	2	3
MYYU	2	1	3
ANPA	0	3	3
MYCI	0	2	2
MYSE	0	2	2
COTO	1	0	1

114602 NE



Species	2018	2019	2020	2022	2024	2025	Total
MYLU	6	20	5	726	80	96	933
EPFU	35	8	8	336	91	104	582
LANO	4	13	30	54	41	182	324
LACI	25	43	0	3	51	71	193
MYVO	8	34	9	63	17	30	161
MYCA	11	7	6	20	16	35	95
MYEV	4	19	2	14	10	25	74
COTO	1	3	6	6	10	8	34
MYYU	2	5	1	14	4	8	34
MYCI	0	0	0	0	0	1	1

114602 NW



Species	2018	2019	2020	2021	2022	2024	2025	Total
LANO	10	25	217	117	33	25	71	498
MYCA	3	3	48	20	4	34	50	162
LACI	0	3	10	28	1	0	34	76
MYEV	1	3	10	13	6	15	9	57
MYLU	0	3	7	9	5	25	2	51
EPFU	0	5	3	20	2	3	10	43
MYVO	2	2	2	8	2	7	3	26
MYYU	0	0	2	3	0	0	2	7
COTO	0	0	0	1	1	2	1	5
TABR	0	0	0	0	1	0	0	1
MYCI	0	0	0	0	0	0	1	1

114602 SE



Species	2018	2019	2020	2021	2022	2024	2025	Total
LANO	8	137	44	718	66	33	109	1,115
MYLU	1	172	93	396	112	95	170	1,039
LACI	3	29	165	120	0	19	242	578
MYVO	0	21	40	91	11	6	9	178
MYCA	0	13	17	49	15	5	13	112
EPFU	0	3	4	17	8	17	43	92
COTO	2	11	13	14	10	5	26	81
MYEV	0	15	13	12	10	4	21	75
MYYU	0	5	7	35	1	2	15	65

114602 SW



Species	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	68	1,041	556	532	157	249	331	2,934
LANO	199	232	736	913	136	334	109	2,659
LACI	12	3	24	22	1	1	165	228
MYYU	12	27	23	22	18	7	19	128
EPFU	4	1	41	22	1	37	14	120
MYEV	5	4	11	11	6	17	10	64
MYCA	2	9	14	13	10	8	4	60
MYVO	1	7	1	2	1	0	2	14
COTO	1	2	1	2	0	2	0	8
TABR	1	0	1	0	0	0	0	2
LABO	0	0	0	0	0	0	2	2

115322 NE



Species	2018	2020	2021	2022	2023	2025	Total
MYYU	871	273	88	7	3	615	1,857
MYEV	209	8	194	9	15	328	763
LANO	89	51	35	2	22	562	761
MYVO	72	2	98	5	8	271	456
EPFU	63	44	34	1	9	169	320
MYLU	26	20	7	7	12	149	221
MYCA	22	18	18	0	72	82	212
LACI	0	0	21	0	0	99	120
MYSE	0	0	0	0	0	38	38

115322 NW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYYU	55	1,105	68	258	5	19	186	445	2,141
LANO	88	206	34	58	331	105	44	297	1,163
MYEV	70	167	156	85	65	164	282	109	1,098
MYCA	12	75	25	93	29	281	50	82	647
LACI	0	1	0	11	0	1	207	285	505
MYVO	12	33	35	55	9	58	32	133	367
MYLU	22	44	6	57	31	18	31	43	252
EPFU	13	19	4	8	2	21	26	88	181
MYSE	0	0	0	0	0	0	2	8	10

115322 SE



Species	2018	2019	2022	2023	2024	2025	Total
MYCA	98	133	54	840	107	61	1,293
MYYU	248	14	218	67	139	347	1,033
MYEV	101	59	195	115	139	262	871
LACI	0	0	400	0	0	46	446
MYLU	39	4	20	266	59	20	408
LANO	16	66	113	101	24	72	392
MYVO	50	20	23	22	45	50	210
EPFU	0	0	30	5	30	70	135
MYSE	0	0	0	0	7	30	37

115322 SW



Species	2018	2019	2020	2021	2023	2024	2025	Total
MYVO	64	13	0	1	6	5	6	95
MYCA	13	5	10	3	20	23	9	83
MYEV	10	3	0	2	10	13	10	48
MYYU	9	2	1	2	0	6	4	24
MYLU	5	1	0	0	3	1	0	10
LANO	2	0	0	0	0	1	1	4
MYSE	0	0	0	0	0	2	0	2
EPFU	0	1	0	0	0	0	0	1

1194 NW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	54	16	8	12	12	50	64	11	40	27	294
LANO	6	7	12	16	1	8	13	1	29	12	105
MYCA	16	5	4	5	3	15	25	4	8	3	88
MYYU	7	4	1	1	1	4	8	3	5	0	34
LACI	7	2	2	0	3	4	4	6	4	0	32
MYEV	7	0	0	1	1	2	7	0	7	6	31
MYVO	5	0	2	0	0	2	3	1	1	1	15
EPFU	0	0	0	0	0	0	2	0	1	2	5

1194 SW



Species	2016	2017	2018	2019	2020	2021	Total
MYLU	486	633	0	28	29	133	1,309
MYYU	1,108	120	1	5	1	26	1,261
MYCA	155	262	27	35	24	159	662
LANO	68	387	5	17	37	10	524
LACI	18	26	0	0	19	25	88
MYVO	13	31	0	7	2	33	86
MYEV	3	33	0	6	1	11	54
EPFU	0	35	0	1	1	3	40
COTO	0	1	0	0	0	0	1

1194 SW2



Species	2020	2022	2023	2024	2025	Total
MYCA	1	149	189	1,283	101	1,723
MYLU	1	484	27	390	119	1,021
LANO	33	447	15	152	325	972
MYYU	0	156	25	207	94	482
MYEV	0	31	10	55	9	105
EPFU	0	18	1	6	17	42
LACI	2	2	4	1	30	39
MYVO	0	4	0	8	2	14
LABO	0	4	0	3	2	9
MYSE	0	0	0	2	0	2
ANPA	0	0	0	0	2	2
COTO	0	1	0	0	0	1

12522 NE



Species	2016	2017	2019	2020	2021	2022	Total
LANO	260	1	379	59	407	256	1,362
MYCA	147	1	32	0	139	64	383
MYLU	40	1	85	2	87	44	259
MYYU	2	0	7	1	138	27	175
EPFU	15	0	2	2	16	58	93
LACI	24	0	50	0	1	16	91
MYVO	29	0	0	0	4	2	35
MYEV	2	0	4	1	8	2	17
COTO	3	0	0	1	0	0	4
TABR	0	0	1	1	0	1	3

12522 NW



Species	2016	2017	2018	2019	2020	2024	Total
MYYU	78	3	1	0	2	1,060	1,144
LACI	95	7	3	26	1	127	259
LANO	122	7	0	21	9	24	183
MYCA	52	9	1	0	0	44	106
EPFU	39	2	4	0	0	11	56
MYLU	8	0	0	0	0	13	21
MYVO	0	0	0	0	0	17	17
COTO	0	0	1	0	0	4	5
MYEV	1	0	0	0	0	2	3
MYSE	0	0	0	0	0	2	2
TABR	0	0	1	0	0	0	1

12522 SE



Species	2016	2017	2018	2020	2021	2023	2024	Total
LANO	292	66	18	5	2	58	10	451
MYCA	65	43	1	0	0	3	7	119
MYLU	72	5	2	0	0	0	0	79
EPFU	58	12	2	0	2	4	0	78
MYYU	38	0	0	0	0	0	0	38
LACI	8	4	0	0	0	7	3	22
MYVO	17	1	0	0	0	2	0	20
MYEV	5	2	0	1	1	4	0	13
COTO	1	0	0	0	0	0	0	1

12522 SW



Species	2016	2018	2019	2020	2021	2022	Total
MYCA	1,161	0	230	1	245	105	1,742
MYVO	682	0	1	0	12	32	727
MYLU	223	0	14	2	2	7	248
MYYU	179	1	22	1	5	8	216
MYEV	47	0	64	3	32	39	185
LANO	7	0	7	10	7	41	72
COTO	3	0	8	0	6	20	37
EPFU	10	0	0	0	4	2	16
LACI	0	2	0	1	0	0	3

127850 NE



Species	2017	2018	2019	2020	2021	Total
MYLU	5	0	67	52	90	214
MYEV	3	3	10	19	55	90
LACI	0	2	1	33	18	54
LANO	1	9	6	5	9	30
MYCA	0	1	17	0	11	29
MYVO	0	0	3	2	15	20
EPFU	2	0	2	1	12	17
MYYU	0	0	16	0	1	17
MYTH	0	0	3	1	0	4
COTO	0	0	0	3	0	3
LABO	0	0	1	0	0	1

127850 NW



Species	2016	2017	2018	2019	2020	2021	2022	2024	Total
LACI	220	127	4	49	243	350	70	0	1,063
EPFU	257	167	6	16	37	34	310	24	851
MYLU	66	158	3	15	28	274	44	17	605
LANO	62	27	0	103	42	130	26	0	390
MYEV	48	69	6	2	5	31	58	11	230
MYCA	28	32	0	0	3	32	10	0	105
MYVO	24	14	1	1	5	26	5	1	77
COTO	0	4	1	0	11	11	2	0	29
MYYU	3	4	0	0	1	2	4	0	14
MYTH	0	0	0	0	0	1	0	0	1
LABO	0	0	0	0	0	0	1	0	1

127850 SE



Species	2016	2019	2020	2021	2022	2023	Total
MYLU	131	7	997	183	39	137	1,494
LANO	26	43	193	150	32	113	557
LACI	3	334	44	35	10	19	445
MYEV	49	9	124	24	33	136	375
MYYU	7	3	128	77	5	24	244
MYCA	18	3	54	43	5	37	160
MYVO	18	3	24	16	6	27	94
EPFU	23	14	9	10	15	22	93
COTO	0	0	6	1	0	10	17
MYTH	1	3	1	0	2	1	8
LABO	0	0	0	0	0	3	3

127850 SW



Species	2017	2018	2019	2020	2021	2022	2023	2024	Total
LACI	855	1	2	1,063	136	465	225	1,013	3,760
LANO	33	0	0	82	23	12	16	131	297
MYEV	42	4	0	44	21	23	100	55	289
MYLU	46	3	1	27	48	12	59	50	246
MYCA	40	13	0	11	10	14	56	30	174
MYVO	9	1	0	16	23	2	32	1	84
EPFU	15	1	0	17	4	11	13	15	76
MYTH	38	0	0	4	3	0	10	4	59
COTO	6	0	0	20	5	2	0	6	39
MYYU	16	0	0	2	5	0	7	7	37
LABO	0	0	0	0	0	0	0	3	3

128058 NE



Species	2018	2019	2020	2021	2022	Total
MYCA	75	36	174	404	51	740
MYYU	2	2	48	37	104	193
MYEV	23	6	27	17	1	74
EPFU	4	1	0	34	7	46
LANO	7	4	7	21	3	42
LACI	2	0	2	22	0	26
MYLU	6	0	2	10	0	18
MYVO	2	0	0	0	0	2

128058 NW



Species	2018	2019	2020	2021	2022	Total
LANO	3,780	3,676	2,886	3,353	1,860	15,555
LACI	487	186	23	788	4	1,488
MYLU	104	31	192	184	389	900
MYVO	1	0	4	253	46	304
EPFU	22	1	124	25	129	301
MYCA	24	30	26	34	103	217

Species	2018	2019	2020	2021	2022	Total
MYYU	20	34	17	38	66	175
MYEV	0	4	4	2	3	13
COTO	1	1	2	7	0	11
TABR	0	1	2	1	0	4

128058 SW



Species	2018	2019	2020	2021	2022	Total
LANO	153	33	140	101	233	660
LACI	46	13	7	97	0	163
MYCA	16	18	22	18	26	100
MYLU	17	21	12	30	18	98
MYYU	11	16	9	17	22	75
EPFU	4	3	11	3	12	33
COTO	3	3	1	1	5	13
MYEV	1	1	1	1	3	7
MYVO	1	0	3	2	0	6
TABR	0	0	1	1	0	2

132650 SE



Species	2020	2021	2022	2023	2024	2025	Total
MYCI	61	622	20	156	153	449	1,461
MYLU	22	134	2	5	21	31	215
EPFU	20	61	0	3	19	52	155
LANO	39	31	0	1	3	58	132
LACI	34	2	0	0	0	82	118
MYVO	1	4	0	3	2	39	49
MYEV	4	10	0	4	4	13	35
EUMA	6	11	0	0	7	4	28
MYCA	0	4	0	2	3	7	16
MYYU	0	1	0	0	0	10	11
MYSE	0	0	0	0	0	3	3
MYTH	1	0	0	0	0	0	1
COTO	0	1	0	0	0	0	1

132650 SW



Species	2020	2021	2022	2023	2024	2025	Total
LANO	21	0	11	11	0	29	72
MYCI	30	3	5	1	0	21	60
EUMA	4	0	17	16	2	3	42
LACI	9	0	1	0	0	22	32
MYLU	3	2	1	0	0	5	11
EPFU	3	0	0	1	0	2	6
MYEV	0	0	1	0	0	5	6
COTO	0	0	1	0	0	0	1
LABO	0	0	0	0	0	1	1
MYVO	0	0	0	0	0	1	1

137274 NE



Species	2018	2019	2020	2022	2023	Total
MYLU	1,431	290	2,601	4,994	2,737	12,053
LANO	6	21	107	578	15	727
MYYU	66	75	50	250	209	650
EPFU	19	7	107	71	148	352
MYCA	26	25	119	76	41	287
MYVO	34	12	18	86	75	225
LACI	7	61	0	3	6	77
COTO	0	2	0	0	1	3
LABO	0	0	0	2	1	3
TABR	2	0	0	0	0	2
MYEV	0	0	0	0	2	2

137274 NW



Species	2018	2019	2020	2021	2022	2023	Total
MYLU	288	176	1,235	732	959	128	3,518
LACI	30	140	288	1,119	82	1	1,660
EPFU	304	1	70	125	41	93	634
LANO	50	18	117	62	149	12	408
MYCA	46	7	115	71	88	11	338
MYYU	60	19	25	137	56	16	313
MYVO	25	27	56	105	47	24	284
MYEV	1	7	0	7	10	5	30
TABR	5	0	0	1	0	0	6
LABO	0	0	0	1	1	0	2

137274 SE



Species	2018	2019	2020	2021	2022	2023	Total
LANO	271	150	392	310	1,004	268	2,395
MYYU	69	269	558	111	305	818	2,130
MYLU	91	116	283	246	360	674	1,770
MYCA	20	20	140	40	147	198	565
EPFU	3	0	1	1	54	61	120
MYVO	18	11	20	2	7	12	70
LACI	3	6	4	26	16	9	64
MYEV	2	9	4	0	7	3	25
COTO	0	3	1	0	4	3	11
LABO	0	0	0	1	0	1	2
TABR	0	0	0	0	0	1	1

143274 NE



Species	2019	2020	2021	2022	2023	2024	2025	Total
LANO	47	1	331	345	145	286	1,489	2,644
MYEV	36	5	370	621	447	459	573	2,511
MYLU	5	0	510	115	199	117	738	1,684
MYCA	8	1	303	360	291	257	371	1,591
EPFU	42	9	101	163	258	236	456	1,265
MYVO	2	0	335	102	71	92	603	1,205
LACI	12	0	169	73	21	26	204	505
MYYU	1	0	88	66	40	21	237	453
COTO	1	0	11	8	21	5	10	56
MYTH	1	0	9	3	0	1	2	16
EUMA	0	0	0	2	0	0	0	2
MYCI	0	0	0	0	0	0	1	1

143274 NW



Species	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	24	2	36	9	893	27	2,098	3,089
LANO	68	12	130	63	115	300	571	1,259
MYCA	8	2	5	4	139	33	120	311
EPFU	0	2	5	0	28	52	124	211
MYYU	12	2	4	2	36	28	58	142
LACI	2	0	42	0	6	5	43	98
MYEV	4	5	10	5	11	11	7	53
MYVO	3	1	2	0	20	6	18	50
COTO	0	1	6	0	1	0	0	8
MYCI	0	0	0	0	0	0	2	2
MYTH	0	0	0	1	0	0	0	1

143274 SE



Species	2019	2020	2021	2022	2023	2024	2025	Total
LANO	29	81	62	56	165	86	221	700
MYLU	73	28	89	53	105	102	122	572
MYCA	18	8	50	11	105	20	35	247
LACI	5	20	51	0	16	6	143	241
COTO	1	12	19	45	23	70	34	204
MYYU	10	12	53	10	29	15	14	143
EPFU	7	0	3	1	9	26	20	66
MYEV	7	0	13	1	6	9	17	53
MYVO	3	1	2	3	11	3	3	26
LABO	0	0	0	0	0	1	0	1

143274 SW



Species	2019	2020	2021	2022	2023	2024	2025	Total
EPFU	25	0	172	1,023	643	212	394	2,469
MYEV	130	1	270	350	290	403	485	1,929
LANO	11	0	238	218	155	38	235	895
LACI	11	0	184	19	117	3	121	455
MYCA	8	0	23	7	38	30	51	157
COTO	6	0	17	3	62	5	9	102
MYVO	6	0	7	4	24	6	31	78
MYLU	9	0	7	2	9	7	21	55
MYTH	0	0	9	5	15	0	4	33
MYYU	1	0	6	0	5	2	8	22

143338 NE



Species	2016	Total
MYLU	42	42
LACI	4	4
EPFU	3	3
LANO	3	3
MYVO	2	2
TABR	2	2
MYEV	1	1

143338 NW



Species	2016	Total
MYLU	19	19
MYCA	14	14
MYVO	5	5
LANO	3	3
MYEV	2	2

Species	2016	Total
EPFU	1	1

143338 SE



Species	2016	Total
EPFU	3	3
LACI	1	1
LANO	1	1
MYLU	1	1
TABR	1	1

145578 NE



Species	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	68	23	79	45	109	127	81	532
LANO	56	32	61	17	86	4	19	275
MYEV	7	7	16	11	7	25	20	93
EPFU	27	5	3	8	6	20	14	83
COTO	3	11	6	2	17	21	17	77
LACI	7	2	3	6	1	0	14	33
MYCA	3	0	2	3	8	9	1	26
ANPA	0	0	0	0	6	0	13	19
LABO	0	0	0	0	1	1	15	17
MYVO	0	1	1	3	3	0	3	11
MYCI	0	1	3	0	0	0	0	4
MYYU	0	0	0	0	3	1	0	4
MYSE	0	0	0	0	0	0	2	2
MYTH	0	0	0	0	0	0	2	2
PAHE	0	0	0	0	1	0	0	1

145578 NW



Species	2019	2021	2022	2023	2024	2025	Total
LACI	0	2	0	4	0	1,034	1,040
MYLU	1	29	9	43	43	76	201
MYEV	6	51	3	29	10	29	128
LANO	0	2	1	0	17	43	63
MYCA	3	5	0	5	5	22	40
COTO	0	3	4	6	15	7	35

Species	2019	2021	2022	2023	2024	2025	Total
MYVO	3	4	0	8	5	7	27
EPFU	1	2	0	5	5	10	23
LABO	0	0	0	0	0	8	8
ANPA	0	0	0	0	0	5	5
MYTH	0	2	0	1	0	0	3
MYCI	0	0	0	0	0	1	1
MYSE	0	0	0	0	0	1	1

145578 SE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	246	287	811	922	587	1,088	52	2,635	6,628
EPFU	8	4	165	290	39	527	38	582	1,653
LANO	25	91	216	223	215	186	2	549	1,507
LACI	0	12	40	154	61	250	1	538	1,056
MYEV	8	12	245	64	27	113	24	160	653
LABO	0	1	2	7	1	5	0	243	259
MYVO	16	24	33	39	10	25	1	74	222
PAHE	0	0	0	0	2	0	0	152	154
MYCA	24	6	18	7	14	13	2	37	121
COTO	0	7	10	18	6	40	0	16	97
ANPA	0	0	0	0	2	0	0	88	90
MYCI	0	16	27	2	5	11	0	18	79
MYTH	0	0	5	3	0	5	0	11	24
MYYU	0	0	0	0	1	3	0	6	10

Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYSE	0	0	0	0	0	0	0	7	7

145578 SW



Species	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	68	389	311	943	475	919	554	3,659
MYEV	183	248	35	112	67	52	33	730
LANO	38	84	100	135	36	23	167	583
LACI	18	53	72	26	20	0	143	332
EPFU	30	47	31	78	9	25	76	296
MYCA	4	9	4	36	19	51	7	130
MYVO	17	48	12	11	14	4	20	126
LABO	0	1	9	5	3	1	50	69
MYCI	1	16	0	0	24	10	12	63
ANPA	0	0	0	1	0	0	36	37
COTO	1	2	6	4	4	0	19	36
MYTH	7	0	1	5	10	4	9	36
PAHE	0	0	0	9	0	0	0	9
MYYU	0	0	0	0	1	3	4	8
MYSE	0	0	0	0	0	0	1	1

147306 NE1



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
EPFU	1,232	158	256	230	21	594	776	1,388	230	149	5,034
MYLU	177	220	569	431	362	514	341	617	182	1,473	4,886
MYYU	144	25	165	96	49	98	44	69	41	248	979
LANO	31	9	122	104	28	289	34	130	20	141	908
MYCA	12	17	80	49	8	74	43	49	28	123	483
MYEV	16	41	40	55	2	25	49	41	3	70	342
LACI	3	5	48	12	12	68	7	22	8	34	219
MYTH	4	6	23	45	31	11	8	4	7	54	193
MYVO	3	4	13	4	0	8	6	12	2	1	53
COTO	7	4	6	2	9	1	1	4	0	6	40
MYSE	0	0	0	0	0	0	0	0	0	14	14
LABO	0	0	0	0	0	0	0	0	0	4	4
ANPA	0	0	0	0	0	0	0	0	0	3	3
MYCI	0	0	0	0	0	0	0	0	1	0	1

147306 NE2



Species	2016	2017	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	264	645	0	6	1,506	442	1,049	1,551	3,239	8,702
MYYU	310	143	6	7	237	59	110	222	319	1,413
EPFU	46	17	0	0	223	187	168	253	113	1,007
LANO	24	12	0	9	273	55	82	57	134	646
MYCA	33	62	0	2	180	30	52	96	73	528
MYTH	6	24	0	0	25	8	104	32	95	294
LACI	9	9	0	0	88	4	70	28	74	282
MYEV	7	4	0	0	28	8	7	14	47	115
COTO	3	3	0	0	3	7	34	35	14	99
MYVO	39	16	0	0	6	0	3	7	1	72
MYSE	0	0	0	0	0	0	0	0	8	8
MYCI	0	0	0	0	0	0	0	7	0	7
LABO	0	0	0	0	0	0	0	0	5	5
ANPA	0	0	0	0	0	0	0	0	1	1

147306 SE1



Species	2016	2017	2019	2020	2021	2022	2023	2024	2025	Total
MYYU	185	489	325	3	60	277	465	294	685	2,783
MYLU	23	434	140	1	21	621	173	592	735	2,740
MYCA	9	238	232	9	49	157	326	333	589	1,942
LACI	168	217	28	0	29	7	288	27	211	975
EPFU	60	51	47	0	8	166	96	210	96	734
LANO	17	87	104	0	15	97	113	51	164	648
MYTH	4	9	23	0	2	1	11	41	164	255
MYEV	1	20	12	0	1	2	8	19	23	86
MYVO	2	17	4	0	3	13	9	8	8	64
COTO	6	4	3	0	0	0	10	6	2	31
MYCI	0	0	0	0	0	2	2	4	0	8
MYSE	0	0	0	0	0	0	0	2	3	5
ANPA	0	0	0	0	0	0	0	0	3	3
LABO	0	0	0	0	0	0	0	1	0	1

147306 SE2



Species	2016	2017	2020	2021	2022	2023	2024	2025	Total
MYLU	81	858	0	698	147	752	1,582	2,528	6,646
LANO	108	210	4	584	127	361	134	376	1,904
EPFU	63	225	10	133	62	479	341	478	1,791
MYYU	59	287	0	147	81	189	258	383	1,404
MYCA	8	128	0	204	165	212	139	291	1,147
LACI	8	191	0	417	5	74	16	113	824
MYVO	8	22	0	39	3	12	6	3	93
MYEV	6	14	0	12	9	10	17	14	82
MYTH	4	5	0	6	16	4	8	15	58
COTO	13	1	0	2	2	5	4	13	40
MYCI	0	0	0	0	1	2	4	0	7
MYSE	0	0	0	0	0	0	0	2	2
LABO	0	0	0	1	0	0	0	0	1

151210 NE1



Species	2019	2021	2022	2023	2024	2025	Total
MYCA	5,508	2,581	1,446	2,445	5,062	1,286	18,328
MYLU	880	621	174	1,618	1,367	2,212	6,872
MYYU	167	315	484	1,561	287	295	3,109
MYVO	882	232	55	230	9	47	1,455
MYEV	14	1	53	10	26	32	136
LANO	0	13	1	3	19	94	130
LACI	0	0	0	0	0	1	1

151210 NE2



Species	2019	Total
MYCA	13	13
MYVO	6	6
MYEV	1	1
MYYU	1	1

159722 NE



Species	2023	2025	Total
MYLU	4,966	4,212	9,178
MYYU	1,279	1,936	3,215
MYVO	64	434	498
MYCA	134	267	401
LACI	141	22	163
LANO	36	20	56
EPFU	12	25	37
MYEV	8	4	12
LABO	9	0	9
COTO	1	0	1
MYCI	0	1	1

159722 SE



Species	2023	2025	Total
MYLU	1,420	1,144	2,564
LANO	758	336	1,094
MYVO	244	35	279
LACI	103	7	110
EPFU	53	53	106
MYEV	31	72	103
MYCA	42	59	101
COTO	7	1	8
LABO	8	0	8
MYYU	4	2	6

16426 NE



Species	2018	2019	2020	2021	2022	2024	Total
MYLU	204	67	39	79	48	33	470
LANO	6	10	10	6	10	4	46
MYVO	2	6	4	1	6	7	26
MYEV	4	0	0	6	2	0	12
LABO	1	0	1	0	3	0	5
EPFU	1	0	0	0	0	1	2
MYCA	0	0	0	1	0	0	1
MYSE	0	0	0	0	1	0	1

16426 SE



Species	2018	2019	2020	2021	2022	2024	Total
MYLU	1,321	486	37	196	903	19	2,962
MYVO	591	185	27	1,004	21	37	1,865
MYCA	39	5	13	11	3	0	71
LANO	9	4	2	4	3	0	22
LACI	0	16	0	0	1	0	17
MYEV	4	5	1	1	2	0	13
MYSE	2	1	2	2	0	0	7
LABO	0	4	1	1	1	0	7

170042 NE



Species	2016	Total
TABR	2	2
LANO	1	1

170042 SE



Species	2016	Total
MYCA	83	83
LANO	21	21
MYLU	21	21
MYVO	20	20
TABR	1	1

172090 NE



Species	2018	2019	2020	2021	2022	2023	Total
LANO	42	221	52	47	48	21	431
MYLU	1	8	1	15	17	1	43
LACI	3	17	0	0	4	5	29
MYCA	7	4	2	4	11	1	29
MYVO	1	0	0	3	3	3	10
MYYU	2	4	1	0	0	0	7
MYEV	0	2	0	2	1	2	7
EPFU	3	0	1	2	0	0	6

172090 NW



Species	2018	2019	2021	2022	2023	Total
LANO	63	205	193	517	122	1,100
LACI	8	68	3	15	103	197
MYLU	19	8	25	49	14	115
MYCA	1	2	3	14	2	22
MYYU	1	0	0	4	1	6
MYEV	0	1	0	4	0	5
MYVO	1	0	1	0	1	3
EPFU	1	0	0	1	0	2

172090 SE



Species	2018	2019	2020	2021	2022	2023	Total
LANO	185	36	72	60	99	14	466
EPFU	0	0	1	138	3	2	144
LACI	52	0	0	1	13	22	88
MYLU	5	0	5	27	38	4	79
MYVO	6	0	0	6	9	0	21
MYCA	1	0	4	3	8	3	19
MYYU	1	0	1	4	3	0	9
MYEV	0	0	2	0	2	1	5
COTO	1	0	0	0	0	0	1

172090 SW



Species	2018	2019	2020	2021	2022	2023	Total
MYLU	85	42	68	296	623	158	1,272
LANO	90	107	51	54	350	17	669
MYCA	6	6	13	28	98	51	202
MYYU	10	11	19	22	28	32	122
EPFU	4	10	15	16	24	30	99
MYVO	32	0	6	6	14	4	62
LACI	11	17	0	0	1	3	32
MYEV	1	1	4	0	6	1	13

175354 SE



Species	2018	2019	2020	Total
EPFU	106	88	89	283
MYEV	51	32	25	108
MYLU	47	19	13	79
MYVO	24	21	11	56
MYSE	5	3	2	10
LANO	1	2	1	4
LABO	0	1	0	1
LACI	0	0	1	1

175354 SW



Species	2018	2019	2020	2024	2025	Total
MYLU	43	11	10	56	61	181
MYVO	35	15	15	47	30	142
MYEV	13	11	3	63	28	118
EPFU	5	4	0	20	43	72
LANO	1	1	1	1	35	39

Species	2018	2019	2020	2024	2025	Total
MYSE	3	1	2	15	6	27
LABO	1	0	0	0	7	8
LACI	0	1	0	3	3	7

17706 NW



Species	2019	2020	2024	2025	Total
MYLU	463	8	0	45	516
MYCA	49	1	1	9	60
LANO	53	0	0	0	53
MYEV	28	0	1	0	29
EPFU	11	0	0	0	11
MYVO	7	0	0	3	10

17706 SE



Species	2019	2020	2023	2024	2025	Total
LANO	36	256	25	9	19	345
MYLU	15	3	11	2	19	50
MYCA	5	2	18	3	14	42
MYEV	5	1	13	2	1	22
EPFU	16	0	4	0	1	21
MYVO	7	1	2	0	2	12
LACI	1	0	1	0	0	2
LABO	0	0	0	0	2	2

17706 SW



Species	2019	2020	2023	2025	Total
EPFU	839	8	223	99	1,169
MYLU	48	7	362	79	496
LANO	127	31	76	103	337
MYVO	21	1	148	51	221
MYEV	12	1	14	9	36
MYCA	6	1	13	8	28
LACI	6	2	3	8	19
LABO	1	1	0	4	6
MYSE	0	0	2	1	3
MYYU	2	0	0	0	2

186330 NE



Species	2024	2025	Total
MYLU	755	1,327	2,082
MYVO	176	57	233
MYEV	11	33	44
EPFU	33	1	34
LABO	0	12	12
LANO	3	5	8
MYSE	2	0	2

186330 NW



Species	2024	2025	Total
MYLU	6	5	11
MYEV	1	7	8
LANO	3	3	6
MYVO	3	3	6
LABO	0	2	2

Species	2024	2025	Total
MYSE	0	2	2
MYYU	0	1	1

186490 NW



Species	2016	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	5	31	356	146	171	486	85	263	521	2,064
MYVO	6	1	243	70	58	80	21	19	256	754
LANO	49	30	12	10	136	68	91	67	171	634
MYEV	2	4	46	35	29	8	6	6	46	182
EPFU	0	9	11	2	31	2	1	5	32	93
MYSE	5	0	28	9	16	7	0	1	23	89
LACI	0	0	4	0	1	46	0	0	17	68
LABO	0	0	0	1	0	0	2	1	19	23
MYCA	0	0	2	0	5	0	4	1	3	15
MYYU	0	0	0	1	0	0	0	0	2	3

186490 SE



Species	2016	2017	2018	2019	2020	2021	2022	Total
LANO	2	8	72	36	14	229	23	384
MYLU	9	23	75	44	62	60	22	295
MYVO	0	0	15	3	2	9	4	33
MYEV	1	0	7	3	4	2	4	21
EPFU	0	0	2	1	0	0	0	3
LACI	0	1	1	0	0	0	0	2
MYSE	0	0	0	1	0	1	0	2
MYCA	0	0	0	0	0	2	0	2
LABO	0	0	0	1	0	0	0	1

186490 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	474	416	1,463	678	527	1,122	1,374	1,711	2,439	3,444	13,648
LANO	44	123	181	109	499	162	255	316	203	251	2,143
LABO	0	5	4	10	1	5	2	37	100	424	588
MYVO	3	6	92	39	11	40	54	35	39	62	381
EPFU	0	1	25	5	7	4	1	17	1	16	77
MYCA	0	0	1	3	1	9	20	4	14	0	52
MYEV	0	0	9	24	0	2	1	1	7	2	46
LACI	0	0	7	1	0	16	2	3	3	5	37
MYSE	0	0	1	1	0	0	3	0	1	1	7
MYYU	1	0	0	3	0	0	0	0	0	0	4

194666 NE



Species	2016	2017	2018	2019	2020	2021	2023	2024	Total
MYLU	154	381	97	76	6	151	195	116	1,176
EPFU	100	111	29	9	0	12	35	20	316
MYCA	19	58	35	14	0	18	63	26	233
MYYU	15	28	57	28	5	26	39	18	216
LANO	32	61	14	30	7	19	34	10	207
MYCI	8	4	7	6	0	9	19	4	57
LACI	3	7	0	38	1	0	0	2	51
MYVO	1	7	10	1	0	6	4	5	34
MYEV	1	3	1	5	0	1	4	4	19
LABO	0	0	0	0	1	0	0	0	1

194666 SE1



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
EPFU	112	370	17	0	46	229	166	119	126	326	1,511
LANO	241	345	76	0	107	103	212	48	46	131	1,309
MYLU	101	252	54	0	6	132	7	20	124	228	924
MYCA	19	43	10	0	1	91	8	9	63	165	409
MYYU	8	36	4	0	8	107	10	7	40	77	297
MYCI	16	23	3	0	0	9	12	45	17	2	127
LACI	9	7	0	0	2	0	0	1	15	20	54
MYVO	4	7	0	0	0	2	0	1	4	3	21
LABO	0	2	0	0	1	0	0	0	0	11	14
MYEV	2	4	0	1	0	0	0	0	2	3	12
MYTH	0	0	0	0	1	0	0	0	2	0	3
PAHE	0	0	0	0	0	0	0	0	0	2	2
COTO	1	0	0	0	0	0	0	0	0	0	1

194666 SE2



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	424	83	27	141	8	208	99	32	15	1,027	2,064
EPFU	222	253	38	59	8	64	316	36	67	189	1,252
MYCA	151	59	26	137	20	80	104	22	22	321	942
MYYU	320	52	28	73	7	54	39	19	21	267	880
LANO	50	67	14	54	60	104	146	20	19	126	660
LACI	15	27	0	75	21	2	5	0	23	218	386
MYEV	20	73	10	89	16	21	25	11	10	36	311
MYCI	31	13	3	26	0	11	13	1	2	4	104
MYVO	24	6	7	35	0	10	4	3	3	4	96
COTO	1	2	0	20	0	2	7	0	0	41	73
LABO	1	0	2	0	0	0	0	0	0	37	40
MYTH	2	0	0	3	0	0	0	0	0	2	7
PAHE	0	0	0	0	0	0	0	0	0	3	3
MYSE	0	0	0	0	0	0	0	0	0	1	1

194666 SW

Species	2016	2017	2018	2019	2020	2021	2022	2023	2025	Total
LANO	66	42	0	22	5	8	36	1	430	610
MYLU	60	9	2	23	0	22	1	0	395	512
MYCA	14	1	2	12	0	9	8	0	202	248
EPFU	48	15	2	9	0	31	8	2	28	143
LACI	4	28	0	7	1	0	0	0	21	61
MYEV	9	5	1	14	2	10	1	1	18	61
MYCI	9	1	0	6	0	2	6	0	0	24
LABO	0	0	0	0	0	0	0	0	21	21
MYYU	4	2	0	1	0	3	3	0	4	17
MYVO	5	0	0	1	0	1	4	0	0	11
MYTH	2	0	0	0	0	0	0	0	1	3
TABR	3	0	0	0	0	0	0	0	0	3
COTO	0	0	0	1	0	0	0	0	0	1

19514 SW



Species	2024	2025	Total
MYYU	333	737	1,070
MYCA	161	247	408
LACI	160	131	291
LANO	119	9	128
MYLU	32	9	41
MYEV	30	10	40
EPFU	24	4	28
COTO	4	0	4
MYVO	2	2	4
LABO	3	0	3
MYCI	1	0	1

195242 SE1



Species	2019	2021	2022	2023	2024	2025	Total
MYCA	47	350	126	120	112	542	1,297
MYLU	37	45	15	65	23	121	306
MYYU	42	106	10	85	39	7	289
MYEV	29	24	32	35	32	92	244
MYVO	26	15	6	20	6	55	128
MYCI	0	0	0	0	0	4	4
EPFU	1	0	0	0	0	0	1

195242 SE2



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	182	192	240	86	90	396	1,186
MYCA	14	4	84	8	7	34	151
MYYU	12	3	27	9	5	13	69
LANO	12	0	6	7	13	14	52
MYEV	1	3	8	9	2	11	34
MYVO	3	7	14	4	2	4	34
EPFU	0	0	0	0	0	2	2

195242 SW1



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	2,415	3,959	381	3,108	2,133	3,581	15,577
LANO	482	962	114	700	997	136	3,391
MYCA	393	565	186	976	201	220	2,541
MYYU	427	98	25	515	95	370	1,530
MYVO	3	25	5	8	13	3	57
MYEV	18	4	2	3	0	5	32
EPFU	3	1	1	3	9	0	17
LACI	0	0	0	1	1	1	3
COTO	0	0	0	2	0	0	2

195242 SW2



Species	2019	2021	2022	2023	2024	2025	Total
MYCA	157	156	160	136	130	301	1,040
MYYU	75	89	29	203	54	40	490
MYEV	8	23	32	35	54	193	345
MYVO	34	34	2	35	2	139	246
MYLU	98	73	13	17	7	13	221
LANO	23	27	19	8	7	12	96
EPFU	6	0	0	0	1	0	7

195242 SW3



Species	2019	2021	2022	2023	2024	2025	Total
MYCA	114	318	365	25	170	32	1,024
MYYU	9	63	471	3	49	1	596
MYLU	5	210	62	29	13	7	326
MYVO	2	34	70	8	3	1	118
MYEV	7	6	12	24	7	12	68
LANO	4	8	0	2	7	0	21
MYCI	0	0	0	0	0	3	3

20330 NE



Species	2016	2017	2018	2019	2020	2022	2023	2024	2025	Total
MYLU	4	8	3	2	15	14	25	30	123	224
LANO	0	17	1	12	10	6	5	9	21	81
MYEV	3	6	1	0	4	4	4	20	10	52
LACI	0	18	0	0	0	0	1	2	27	48
EPFU	5	9	0	3	1	0	4	7	11	40
MYCA	1	4	0	1	5	2	0	6	14	33
MYVO	0	7	2	3	1	1	2	3	14	33
COTO	5	7	0	0	6	0	1	3	6	28
ANPA	0	0	0	0	0	1	0	0	15	16
MYYU	0	2	0	0	1	0	2	1	9	15
MYCI	0	0	0	0	0	0	2	1	6	9
MYSE	0	0	0	0	0	0	0	0	5	5
LABO	0	0	0	0	0	0	0	0	2	2

20330 SE



Species	2016	2017	2018	2020	2022	2023	2024	2025	Total
LANO	8	119	47	144	21	93	37	185	654
EPFU	32	119	19	42	121	57	39	39	468
LACI	2	124	48	8	3	23	14	68	290
MYLU	8	25	13	14	10	32	40	146	288
MYCA	4	27	8	6	3	46	9	35	138
MYEV	6	51	1	12	8	8	17	7	110
ANPA	0	0	0	0	62	0	1	2	65
MYYU	3	5	2	2	2	2	2	12	30
COTO	0	1	1	2	0	1	3	1	9
MYVO	0	1	1	1	0	0	0	4	7
LABO	0	0	0	0	0	0	1	6	7
PAHE	0	0	0	0	0	0	0	7	7
MYTH	0	0	0	6	0	0	0	0	6
MYCI	0	0	0	0	0	3	0	2	5
MYSE	0	0	0	0	0	0	0	2	2

20330 SW1



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	42	122	20	96	129	157	215	174	246	262	1,463
LANO	13	209	6	62	105	172	70	127	141	342	1,247
EPFU	59	185	12	57	28	112	38	168	101	423	1,183
LACI	1	209	5	2	10	33	24	24	8	54	370
MYCA	32	34	1	25	17	29	22	10	16	60	246
MYEV	14	19	0	8	12	15	20	4	17	9	118
MYYU	19	3	3	9	4	7	8	3	15	20	91
COTO	4	2	0	6	13	6	3	3	4	3	44
MYVO	0	0	0	0	0	1	0	1	0	11	13
PAHE	0	0	0	0	0	0	4	0	0	7	11
ANPA	0	0	0	0	0	0	3	0	0	5	8
MYSE	0	0	0	0	0	0	0	0	0	4	4
LABO	0	0	0	0	0	0	0	0	0	2	2
MYCI	0	0	0	0	0	0	0	0	0	2	2
EUMA	0	0	0	0	0	0	0	0	1	0	1
MYTH	0	0	0	0	0	0	0	0	1	0	1

20330 SW2



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	27	162	12	255	289	344	43	655	720	1,089	3,596
MYCA	7	152	5	77	184	182	5	459	825	953	2,849
LANO	22	193	0	87	132	79	12	42	79	286	932
MYYU	19	34	0	37	104	49	0	72	222	231	768
EPFU	11	39	1	49	21	24	12	20	29	39	245
LACI	0	27	3	2	5	17	4	1	4	58	121
MYCI	0	0	0	18	0	0	0	51	5	30	104
MYVO	0	1	0	8	30	22	0	3	7	22	93
PAHE	0	0	0	0	0	0	1	0	5	42	48
MYEV	1	14	0	9	12	0	0	2	3	0	41
LABO	0	0	0	0	0	2	0	0	2	6	10
COTO	0	1	0	0	5	0	0	0	2	0	8
ANPA	0	0	0	0	0	0	0	0	0	4	4
MYSE	0	0	0	0	0	0	0	0	0	3	3

206394 NE



Species	2022	2023	2024	2025	Total
MYLU	113	872	13	38	1,036
LANO	13	287	8	269	577
MYEV	7	37	0	40	84
MYCA	13	24	0	0	37
MYSE	0	0	0	3	3
MYYU	0	0	0	1	1

206394 NW



Species	2022	2023	2024	2025	Total
LANO	385	976	200	1,278	2,839
MYLU	469	70	26	240	805
MYEV	17	17	5	4	43
MYCA	1	8	5	2	16
EPFU	0	0	0	4	4
LABO	0	0	0	1	1

Species	2022	2023	2024	2025	Total
MYVO	0	0	0	1	1
MYYU	0	0	0	1	1

206394 SE



Species	2022	2023	2024	2025	Total
LANO	55	125	88	185	453
MYLU	43	26	15	170	254
MYCA	7	1	6	2	16
MYEV	2	2	2	9	15
MYYU	0	0	0	3	3
EPFU	0	0	0	2	2
MYSE	0	0	0	1	1

206394 SW



Species	2022	2023	2024	2025	Total
MYLU	124	138	78	113	453
LANO	52	103	45	188	388
MYCA	12	11	5	5	33
MYEV	7	5	6	8	26
LABO	0	0	0	4	4
EPFU	0	0	0	2	2
MYYU	0	0	0	1	1

20730 NE



Species	2022	2024	2025	Total
MYLU	366	506	890	1,762
LANO	26	15	542	583
MYVO	3	32	38	73
EPFU	0	1	58	59
MYEV	5	25	25	55
LABO	2	34	13	49
MYSE	0	4	6	10
MYYU	0	1	0	1
LACI	0	0	1	1

20730 NW



Species	2022	2024	2025	Total
MYLU	29	109	222	360
LANO	2	41	226	269
MYVO	2	4	17	23
LABO	0	5	17	22
MYEV	7	8	2	17
EPFU	0	0	6	6
MYSE	0	0	3	3

20730 SE



Species	2022	2024	2025	Total
MYLU	18	89	13	120
LABO	0	32	1	33
MYVO	6	16	5	27
LANO	1	6	4	11
MYEV	2	3	0	5

Species	2022	2024	2025	Total
MYSE	0	0	1	1

20730 SW



Species	2022	2025	Total
MYLU	42	25	67
MYVO	6	5	11
MYEV	1	4	5
LABO	0	5	5
EPFU	0	2	2
LANO	0	2	2
MYYU	0	1	1

212266 NE



Species	2016	2017	2018	2019	2021	2022	2023	2024	2025	Total
MYCI	79	269	43	13	635	468	40	2	271	1,820
MYLU	12	189	167	326	423	298	146	18	71	1,650
LANO	25	92	29	179	118	167	58	95	478	1,241
MYEV	3	29	74	29	97	59	12	6	30	339
MYCA	5	9	18	22	82	120	7	8	42	313
MYTH	4	17	52	6	16	38	36	3	31	203
LACI	1	5	37	13	4	2	42	4	61	169
EPFU	3	36	19	14	12	16	15	10	43	168
MYVO	1	0	11	4	29	30	1	0	9	85
EUMA	1	2	13	9	0	1	6	0	2	34
LABO	0	1	0	0	3	0	0	0	20	24
MYYU	1	4	2	2	1	1	0	0	0	11
COTO	0	0	1	0	2	4	1	0	2	10

212266 NW



Species	2017	2018	2019	2021	2023	2024	2025	Total
LANO	57	416	28	0	39	7	65	612
MYLU	99	278	46	5	14	0	16	458
MYCA	87	28	4	14	42	2	157	334
EPFU	19	32	18	0	104	1	22	196
MYYU	4	7	1	0	1	0	173	186
MYEV	10	27	11	1	76	3	12	140
MYCI	12	16	0	0	2	0	9	39
MYVO	1	10	2	1	14	2	9	39
LACI	1	2	0	0	1	0	20	24
LABO	0	0	0	0	0	0	23	23
MYTH	2	3	1	0	0	0	2	8
COTO	0	2	0	0	2	0	1	5
ANPA	0	0	0	0	0	0	1	1

212266 SE



Species	2016	2017	2018	2019	2022	2023	2024	2025	Total
MYLU	393	144	236	53	0	173	124	614	1,737
LANO	448	25	138	22	0	94	51	432	1,210
EUMA	1	0	26	32	0	116	369	656	1,200
MYCI	170	78	69	3	0	125	27	119	591
LACI	30	5	27	9	0	7	23	244	345
EPFU	49	1	16	18	1	19	29	206	339
MYCA	77	3	32	0	0	34	15	58	219
LABO	0	0	0	0	2	9	0	168	179
COTO	2	0	8	10	0	18	49	28	115
MYYU	34	2	14	1	2	35	1	26	115
MYVO	26	2	17	3	0	14	0	30	92
MYEV	8	0	17	1	0	9	6	16	57
MYTH	1	0	0	1	0	0	2	0	4

212266 SW



Species	2016	2018	2019	2021	2022	2023	2024	2025	Total
MYLU	2,753	2,095	281	263	414	677	944	5,197	12,624
EPFU	616	593	89	19	311	333	362	1,430	3,753
LANO	148	152	36	48	358	44	214	1,964	2,964
MYCI	727	157	5	104	257	110	157	953	2,470
MYCA	280	81	21	59	343	138	239	430	1,591
MYVO	172	112	24	66	44	68	77	289	852
LACI	190	188	6	0	8	9	2	183	586
COTO	68	21	26	15	23	47	39	55	294
MYEV	52	19	16	1	24	25	7	19	163
MYYU	19	9	10	1	18	20	8	48	133
MYTH	39	2	4	3	38	5	24	7	122
LABO	0	0	0	0	6	0	0	96	102
ANPA	0	0	0	0	0	0	0	10	10
EUMA	0	0	0	0	4	0	2	0	6

212906 NW



Species	2018	2019	2020	2022	2023	2024	2025	Total
MYCA	8	3	60	63	91	32	80	337
LANO	1	25	217	16	9	3	15	286
MYLU	12	3	21	38	65	34	76	249
MYYU	6	0	10	8	23	6	16	69
MYVO	4	2	6	9	23	3	9	56
MYEV	0	3	7	6	16	7	5	44
EPFU	1	5	3	0	0	0	1	10
COTO	1	0	1	0	6	0	0	8
LACI	1	3	3	1	0	0	0	8

212906 SE



Species	2018	2019	2020	2022	2023	2024	2025	Total
MYLU	56	171	132	378	409	188	570	1,904
MYCA	45	13	75	386	299	590	345	1,753
MYYU	47	5	34	380	227	488	300	1,481
LANO	137	136	58	248	568	93	130	1,370
MYEV	17	14	43	77	59	72	156	438
MYVO	14	21	33	49	47	41	201	406
LACI	7	29	53	6	240	0	48	383
EPFU	5	3	8	6	46	1	61	130
COTO	1	11	16	1	11	0	9	49
MYCI	0	0	0	0	0	0	1	1
MYTH	0	0	0	0	0	0	1	1

212906 SW



Species	2018	2019	2020	2023	2024	2025	Total
MYLU	5	1,041	560	45	18	62	1,731
LANO	5	229	699	3	2	13	951
MYCA	8	9	19	32	35	73	176
MYYU	2	27	26	16	16	32	119
MYVO	2	7	8	28	4	53	102
MYEV	3	4	13	12	28	30	90
EPFU	0	1	46	0	3	1	51
COTO	0	2	1	0	2	30	35
LACI	2	3	25	0	2	1	33
LABO	0	0	3	0	0	0	3
MYCI	0	0	0	0	0	1	1

216746 NW



Species	2021	2022	2024	2025	Total
MYYU	321	255	53	1	630
MYCA	158	90	132	134	514
MYLU	190	6	36	37	269
MYVO	17	0	0	119	136
LANO	0	0	79	4	83
MYEV	2	2	3	3	10
COTO	2	0	0	0	2
EPFU	0	0	1	0	1

216746 NW1



Species	2019	2023	Total
MYYU	393	388	781
MYCA	103	49	152
MYLU	27	57	84
MYVO	19	4	23

Species	2019	2023	Total
LANO	9	6	15
MYEV	1	13	14
COTO	0	2	2

216746 NW2



Species	2019	Total
MYLU	4	4
MYEV	1	1

216746 NW3



Species	2019	Total
MYLU	31	31
MYYU	12	12
MYCA	9	9

Species	2019	Total
LANO	6	6
MYEV	6	6
MYVO	2	2

216746 SE



Species	2019	2021	2022	2023	2024	2025	Total
MYEV	19	108	21	113	140	108	509
MYLU	62	59	28	48	81	44	322
LANO	1	1	4	6	43	13	68
MYCA	2	2	9	3	14	4	34
COTO	0	4	2	15	0	0	21
MYVO	1	0	1	3	0	0	5
MYYU	0	1	0	1	1	0	3
EPFU	0	0	0	0	1	1	2

216746 SW1



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	2,070	1,224	493	1,524	1,592	1,975	8,878
MYCA	524	246	179	97	1,159	943	3,148
MYYU	190	97	90	90	111	149	727
MYVO	325	59	8	29	62	175	658
MYEV	10	1	0	23	6	29	69
LANO	3	0	4	0	3	7	17
LACI	0	0	7	1	0	0	8
MYCI	0	0	0	0	0	2	2
EPFU	0	0	0	0	1	0	1
COTO	0	0	0	0	0	1	1

216746 SW2



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	15	96	7	28	25	35	206
MYVO	14	16	21	77	35	27	190
MYCA	16	36	16	13	18	6	105
MYYU	4	3	3	23	30	0	63
LANO	0	0	1	4	7	5	17
MYEV	0	4	4	2	0	4	14
LACI	0	0	1	0	0	1	2
EPFU	0	0	0	0	1	0	1

218282 SE



Species	2020	2021	2022	2024	2025	Total
LANO	28	16	11	21	114	190
MYLU	4	13	15	13	9	54
MYCA	0	3	4	7	29	43
LACI	0	0	2	1	23	26
MYEV	5	0	4	5	1	15
EPFU	0	0	1	0	2	3
COTO	1	0	0	1	0	2
MYVO	2	0	0	0	0	2
MYYU	0	0	1	0	0	1
LABO	0	0	0	0	1	1

218282 SW



Species	2020	2021	2022	2023	2024	2025	Total
MYLU	710	1,919	839	672	777	4,464	9,381
MYYU	104	188	549	43	80	1,392	2,356
MYCA	145	338	241	139	140	1,024	2,027
LANO	93	126	22	49	117	588	995
MYVO	64	117	247	65	32	176	701
MYEV	74	16	34	20	27	47	218
EPFU	22	17	3	6	8	41	97
MYSE	0	0	1	0	1	20	22
LACI	1	3	3	2	0	0	9
COTO	0	0	1	4	0	0	5
LABO	0	0	0	0	0	5	5
ANPA	0	0	0	0	0	1	1

219626 SW



Species	2024	2025	Total
MYYU	395	253	648
MYCA	234	95	329
MYLU	56	23	79
MYVO	27	19	46
MYEV	19	19	38
LANO	20	3	23
LACI	8	2	10
EPFU	1	1	2
COTO	0	2	2
LABO	0	1	1

226090 NE



Species	2019	2020	2021	2022	2023	2025	Total
MYLU	9	1	9	2	8	1	30
MYCA	3	0	3	0	0	0	6
MYVO	0	0	3	0	1	1	5
MYYU	0	1	1	0	1	0	3
MYCI	0	0	0	0	3	0	3
LACI	0	0	0	0	0	3	3
MYEV	0	0	0	0	0	3	3
LANO	1	0	1	0	0	0	2
EPFU	0	0	0	0	0	1	1
LABO	0	0	0	0	0	1	1

226090 NW



Species	2018	2019	2020	2021	2022	2023	2025	Total
MYCI	71	481	60	135	114	4	13	878
MYLU	41	121	31	34	49	19	7	302
LANO	47	15	47	10	40	2	9	170
LACI	6	13	4	10	10	7	36	86
MYEV	3	13	32	6	9	0	9	72
COTO	0	4	1	12	6	4	4	31
MYCA	0	20	2	5	4	0	0	31
MYTH	3	0	17	1	2	0	1	24
MYYU	1	0	0	3	4	0	2	10
MYVO	2	0	1	1	1	2	1	8
EPFU	1	1	1	1	0	1	0	5
LABO	0	0	0	1	1	0	1	3

226090 SE



Species	2018	2019	2020	2021	2022	2025	Total
MYLU	12	4	10	5	29	0	60
LANO	9	1	8	0	28	0	46
MYEV	4	2	2	0	5	1	14
MYCI	5	0	0	0	1	0	6
MYCA	1	0	0	0	2	0	3
LABO	1	0	0	0	1	0	2
MYVO	1	0	1	0	0	0	2
LACI	1	0	0	0	0	0	1
EPFU	0	0	0	0	1	0	1

230186 NE



Species	2018	2019	2020	2021	2024	Total
MYLU	89	443	468	817	1,125	2,942
LACI	10	144	11	469	58	692
LANO	63	19	34	135	369	620
EUMA	0	3	23	140	0	166
MYYU	6	15	14	25	36	96
MYVO	1	63	1	5	18	88
MYCA	2	13	6	11	51	83
MYEV	2	4	2	12	31	51
EPFU	4	0	0	2	40	46
LABO	0	0	0	1	19	20
COTO	0	0	1	6	2	9
PAHE	0	0	1	0	0	1

230186 NE2



Species	2023	Total
MYLU	220	220

Species	2023	Total
MYVO	44	44
LANO	20	20
LACI	11	11
EPFU	9	9
MYCA	7	7
MYYU	5	5
MYEV	4	4
LABO	1	1

230186 SE



Species	2018	2019	2020	2021	2023	2024	Total
MYLU	198	178	437	1,315	2,573	2,513	7,214
MYYU	233	232	198	596	790	2,074	4,123
LANO	133	55	263	438	239	450	1,578
MYCA	33	133	50	123	366	492	1,197
LACI	25	2	6	119	129	317	598
EPFU	22	4	25	182	11	153	397
MYEV	33	11	11	38	126	74	293
MYVO	2	6	15	20	19	59	121
LABO	0	0	5	8	0	51	64
COTO	0	0	0	4	1	6	11
MYTH	0	0	0	0	0	3	3
PAHE	0	2	0	0	0	0	2
EUMA	0	0	1	0	0	0	1

Species	2018	2019	2020	2021	2023	2024	Total

241002 NE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	5	9	7	532	85	77	105	125	945
LANO	0	0	0	10	3	17	25	10	65
MYEV	3	1	0	5	5	5	20	3	42
MYVO	0	2	0	3	3	4	5	10	27
MYCA	1	0	0	2	1	0	4	0	8
EPFU	0	0	0	1	0	3	1	2	7
LABO	0	0	0	0	1	1	1	2	5
LACI	0	0	0	0	1	1	0	3	5
MYSE	0	0	0	0	0	0	1	1	2

241002 NW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	0	9	10	81	20	99	113	236	568
LANO	10	29	12	35	18	18	175	250	547
MYEV	11	17	5	93	51	64	91	57	389
EPFU	10	43	21	28	14	45	50	142	353
MYVO	0	8	11	76	10	18	14	42	179
LACI	7	17	1	38	1	22	2	36	124
MYCA	0	1	1	11	6	44	18	24	105
MYSE	1	3	0	1	1	0	3	28	37
MYYU	0	0	0	0	0	0	4	14	18
COTO	0	0	0	0	0	0	2	6	8
LABO	0	0	0	0	2	1	0	0	3

243562 NE



Species	2016	2017	2019	2020	2021	2023	2024	2025	Total
MYEV	9	38	19	11	2	97	248	22	446
LACI	0	0	45	0	1	0	300	3	349
EPFU	1	5	12	7	4	40	25	23	117
MYVO	0	1	5	0	3	11	11	3	34
MYLU	0	6	4	3	0	11	4	5	33
LANO	1	1	3	2	1	6	6	9	29
MYCA	0	2	1	0	0	5	10	0	18
ANPA	0	0	0	0	0	0	0	18	18
MYSE	0	0	0	0	0	0	3	8	11
COTO	2	0	0	0	0	1	1	0	4
MYCI	0	1	0	0	0	1	0	0	2
LABO	0	0	0	0	0	0	0	1	1
PAHE	0	0	0	0	0	0	0	1	1

243562 SE



Species	2016	2017	2019	2021	2022	2023	2024	2025	Total
LANO	0	0	30	0	11	2	24	15	82
MYLU	0	8	14	5	2	15	2	10	56
MYEV	2	0	11	0	0	3	3	1	20
EPFU	0	1	6	1	3	5	0	4	20
MYVO	1	5	3	0	0	3	0	1	13
MYCA	0	2	1	0	2	0	1	2	8
MYCI	0	0	2	0	0	1	0	0	3
COTO	0	0	0	0	0	1	2	0	3
LACI	0	0	0	0	0	0	0	3	3
LABO	0	0	1	0	0	0	0	1	2
MYTH	0	0	0	0	2	0	0	0	2
ANPA	0	0	0	0	0	0	0	1	1

243562 SW1



Species	2016	2017	2018	2019	2020	2021	2022	2023	2025	Total
MYLU	72	32	157	43	38	22	110	105	83	662
MYEV	26	37	261	92	48	18	49	63	34	628
LANO	32	2	201	69	16	6	46	21	106	499
EPFU	21	8	104	19	11	2	29	34	126	354
MYCA	22	5	27	16	9	6	13	18	53	169
LACI	8	0	105	4	0	1	2	3	11	134
MYVO	3	1	42	8	3	1	7	5	14	84
COTO	0	2	12	6	11	2	4	2	5	44
ANPA	0	0	0	0	0	0	12	0	20	32
MYCI	12	0	0	1	0	0	5	6	3	27
LABO	0	0	0	1	0	0	9	1	8	19
MYSE	0	0	0	0	0	0	0	0	10	10
MYYU	1	1	0	2	0	0	2	0	1	7
MYTH	0	1	2	0	0	0	1	1	1	6
PAHE	0	0	0	0	0	0	2	0	1	3

243562 SW2



Species	2016	2017	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	110	52	248	141	121	54	178	6	24	934
MYEV	144	98	248	85	98	55	138	17	13	896
LANO	12	5	8	36	85	381	36	16	33	612
MYCA	15	104	121	41	43	28	72	4	14	442
EPFU	75	4	22	22	3	60	14	9	19	228
MYVO	5	9	35	24	6	10	18	1	10	118
LACI	4	0	4	1	60	1	4	0	2	76
MYCI	4	3	24	0	0	8	3	0	2	44
COTO	7	4	1	10	4	9	5	0	1	41
LABO	0	0	2	1	0	0	2	0	4	9
MYTH	0	0	9	0	0	0	0	0	0	9
ANPA	0	0	0	0	0	3	0	0	6	9
PAHE	0	0	0	0	0	0	0	0	6	6
MYYU	0	2	2	0	0	0	1	0	0	5
MYSE	0	0	0	0	0	0	0	0	1	1

249914 NW



Species	2022	Total
MYLU	281	281
LANO	173	173
MYVO	6	6
EPFU	5	5
MYYU	5	5

249914 SE



Species	2022	Total
MYLU	695	695
LANO	162	162
EPFU	115	115
MYVO	59	59
MYCA	7	7
MYYU	1	1

249914 SW



Species	2022	Total
MYLU	27	27
LANO	25	25
MYVO	4	4
MYCA	2	2
MYYU	2	2
EPFU	1	1

252394 NW



Species	2024	2025	Total
MYLU	105	1,204	1,309
MYYU	67	486	553
MYCA	32	207	239
LANO	47	12	59
MYEV	4	42	46
MYVO	3	14	17

Species	2024	2025	Total
LABO	0	3	3
LACI	0	2	2
COTO	1	0	1
EPFU	0	1	1
MYCI	0	1	1

26170 NE



Species	2022	2023	2024	2025	Total
MYLU	45	35	61	24	165
MYCA	50	71	32	2	155
MYEV	10	73	8	2	93
LANO	17	26	12	0	55
MYYU	5	0	5	2	12

26170 NW



Species	2022	2023	2024	2025	Total
MYCA	912	825	68	235	2,040
MYLU	52	75	11	1,016	1,154
LANO	18	90	6	142	256
MYEV	11	24	4	102	141
MYYU	0	0	0	75	75
MYSE	0	0	0	7	7
EPFU	0	0	0	1	1
MYVO	0	0	0	1	1

26170 SE



Species	2022	2023	2024	2025	Total
MYLU	586	339	252	540	1,717
MYCA	428	424	379	244	1,475
MYEV	374	79	124	42	619
LANO	72	125	125	64	386
MYYU	24	0	21	56	101
MYVO	0	0	24	4	28
LABO	0	0	5	0	5
MYSE	0	0	4	1	5
EPFU	0	0	2	1	3
LACI	0	0	1	0	1

26170 SW



Species	2022	2023	2024	2025	Total
MYCA	42	72	25	68	207
LANO	40	54	36	52	182
MYLU	19	22	16	50	107
MYEV	6	12	4	22	44
MYYU	2	0	2	30	34
MYSE	0	0	1	0	1

265749 NE



Species	2018	2019	2020	2021	2022	2023	Total
LACI	6	3	15	10	10	78	122
EUMA	0	0	2	0	84	0	86
LANO	5	29	3	10	5	20	72
MYEV	0	8	6	3	3	34	54
MYCA	2	1	5	8	11	7	34
MYLU	0	7	2	3	4	8	24
MYVO	1	0	0	0	0	11	12
EPFU	0	2	0	1	2	5	10
MYCI	0	0	0	1	0	4	5
MYTH	0	0	0	0	0	3	3
COTO	0	0	0	1	0	1	2
PAHE	0	1	0	0	0	0	1

265749 NW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	27	41	129	122	109	311	588	883	2,210
MYEV	8	41	80	38	67	122	205	190	751
LACI	1	2	5	10	19	142	332	147	658
MYCA	7	4	23	47	13	151	196	171	612
LANO	4	130	12	19	22	74	100	182	543
MYCI	17	0	30	12	5	113	86	0	263
MYVO	1	2	15	14	3	25	20	0	80
EPFU	1	5	0	2	0	15	22	5	50
MYYU	2	0	2	3	0	8	11	0	26

Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
COTO	0	0	0	0	0	4	3	2	9
MYTH	0	2	1	0	0	0	5	0	8
EUMA	1	0	0	0	1	0	1	3	6

265749 SE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	5	223	341	290	94	170	500	646	2,269
LACI	34	8	34	20	92	152	233	214	787
EPFU	3	40	35	30	60	58	252	185	663
MYEV	3	16	53	17	44	36	137	74	380
MYLU	0	10	18	17	15	120	131	38	349
MYTH	9	5	28	24	35	38	72	40	251
MYCA	1	1	28	2	7	14	87	11	151
MYVO	0	3	11	10	5	28	70	0	127
COTO	1	0	1	0	1	20	11	2	36
MYCI	1	1	2	0	2	11	19	0	36
MYYU	0	1	0	0	0	0	5	1	7
EUMA	0	0	0	0	2	0	3	0	5
PAHE	0	1	0	0	0	0	0	0	1

265749 SW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	42	584	213	61	198	424	885	2,214	4,621
MYLU	4	19	19	5	63	139	269	546	1,064
MYCA	2	7	116	10	181	94	242	11	663
EPFU	0	30	12	15	19	104	59	237	476
LACI	1	13	2	3	18	151	68	123	379
MYCI	0	1	11	3	11	19	36	0	81
MYVO	0	6	2	2	0	25	6	0	41
MYEV	1	7	6	1	4	2	9	10	40
MYYU	0	0	16	0	6	0	8	0	30
COTO	1	0	0	0	0	0	2	0	3
MYTH	0	0	0	0	1	0	0	0	1

268521 NE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LACI	0	38	58	47	3	86	36	245	513
MYVO	5	0	12	5	67	42	139	92	362
MYLU	6	32	44	26	20	20	49	36	233
LANO	1	14	6	4	35	13	34	8	115
EPFU	0	19	7	0	5	8	22	5	66
MYEV	1	0	5	2	7	29	10	6	60
LABO	0	2	0	0	1	0	0	15	18
MYCA	0	0	1	0	3	0	2	6	12
MYSE	0	0	0	0	0	0	0	4	4
MYYU	0	0	1	0	0	0	2	0	3
COTO	0	0	0	0	0	0	2	0	2

268521 NW



Species	2018	2019	2021	2022	2023	2024	2025	Total
MYLU	3	58	8	27	49	169	17	331
LACI	1	18	17	3	108	2	3	152
LANO	0	15	13	16	19	16	62	141
MYVO	13	13	5	48	18	11	6	114
MYEV	7	10	1	4	16	4	3	45
EPFU	3	3	0	2	8	3	4	23
LABO	0	0	0	1	0	0	4	5
MYCA	0	1	1	0	0	1	0	3
COTO	0	0	1	0	0	0	0	1

268521 SE



Species	2018	2019	2021	2022	2023	2024	2025	Total
MYLU	22	322	135	210	168	662	407	1,926
LANO	6	248	163	4	32	196	150	799
LACI	2	113	140	0	45	20	24	344
MYVO	3	10	10	92	10	8	8	141
MYEV	1	28	8	11	2	46	8	104
LABO	0	0	0	40	0	0	13	53
MYCA	1	5	2	26	3	2	2	41
MYYU	0	1	0	15	0	4	2	22
EPFU	0	0	0	3	0	8	4	15
COTO	0	0	0	0	0	1	0	1

269777 NW



Species	2017	2018	2019	2020	Total
MYEV	93	22	145	89	349
MYVO	53	0	89	133	275
MYLU	36	9	84	34	163
EPFU	7	0	0	0	7

269777 SW



Species	2017	2018	2019	2020	Total
MYLU	49	224	8	95	376
EPFU	17	183	1	1	202
MYEV	14	11	22	34	81
MYVO	0	10	0	28	38
LANO	0	2	0	1	3
MYSE	1	0	0	1	2
LACI	1	0	0	0	1

271731 SE1



Species	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYEV	12	9	56	68	11	26	64	8	38	292
MYLU	0	1	25	18	106	26	31	4	14	225
MYVO	0	1	8	29	7	19	30	9	45	148
LANO	0	4	0	3	17	11	3	11	10	59
MYCA	0	0	1	9	1	6	1	3	5	26
MYSE	3	1	2	0	0	0	0	1	8	15
LACI	0	0	0	0	2	0	9	0	0	11
EPFU	0	0	0	0	0	1	1	0	4	6
COTO	0	0	0	0	0	0	1	0	1	2
MYYU	0	0	0	0	0	0	0	0	1	1

271731 SE2



Species	2017	2019	2021	2023	2024	2025	Total
MYLU	59	4	61	1,252	369	333	2,078
MYEV	275	17	183	100	199	13	787
MYVO	108	1	36	159	21	14	339
LANO	0	2	3	81	23	1	110
EPFU	0	0	1	4	24	14	43
MYCA	10	0	5	11	10	1	37
MYSE	24	0	0	7	3	3	37
LACI	0	0	4	10	7	2	23
COTO	0	0	0	1	2	5	8
LABO	0	0	0	0	0	2	2

271731 SW



Species	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	7	12	2,360	39	54	2	77	2,551
MYEV	46	364	303	139	90	75	86	1,103
MYVO	18	7	318	63	105	4	236	751
EPFU	0	0	1	7	1	6	43	58
LACI	2	0	20	4	18	1	3	48
MYSE	5	0	0	6	5	1	30	47
LANO	3	2	1	7	7	5	5	30
COTO	0	0	0	8	7	1	9	25
MYCA	0	1	0	2	0	1	3	7
MYYU	0	0	1	0	0	0	6	7

273050 NE1



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	25	68	2,099	3	50	31	2,276
MYCA	40	16	58	1	110	119	344
MYVO	7	3	318	0	8	5	341
MYYU	4	1	9	0	30	47	91
LANO	0	12	0	0	16	1	29
MYEV	4	3	2	1	4	4	18
LACI	0	0	0	0	0	8	8
COTO	0	0	0	0	1	1	2

273050 NE2



Species	2019	2021	2022	2023	2024	2025	Total
MYCA	31	48	62	105	80	31	357
MYLU	0	64	0	1	2	4	71
MYYU	5	4	5	1	0	0	15
MYEV	6	1	4	0	1	2	14
LANO	0	4	0	2	3	0	9
MYVO	0	1	1	3	0	1	6
EPFU	0	0	0	1	2	1	4

273050 SE



Species	2024	2025	Total
MYYU	1,318	2,352	3,670
MYLU	168	1,546	1,714
MYCA	379	176	555
MYVO	3	402	405
MYEV	6	23	29
MYCI	0	27	27
LACI	1	2	3
COTO	2	0	2
LABO	1	0	1

273050 SE1



Species	2019	2021	2022	2023	Total
MYYU	2,337	264	328	1,962	4,891
MYCA	859	98	214	58	1,229
MYLU	135	168	7	524	834
MYVO	26	25	9	80	140
MYEV	60	1	0	3	64
LANO	0	1	0	0	1
EPFU	0	0	1	0	1

273050 SE2



Species	2019	2021	2022	2023	2024	2025	Total
MYLU	307	474	253	579	461	495	2,569
LANO	249	335	372	5	155	197	1,313
MYCA	6	11	26	95	19	2	159
MYVO	15	32	10	11	10	24	102
MYYU	10	23	5	9	1	8	56
MYEV	0	0	0	1	0	3	4
LACI	0	0	1	0	0	2	3
EPFU	1	0	0	0	0	0	1
LABO	0	0	0	0	0	1	1

283586 NE



Species	2018	2019	2020	2022	2023	2024	2025	Total
MYLU	5	3	65	0	135	54	72	334
MYYU	0	7	7	2	6	7	4	33
LANO	8	16	1	0	0	0	0	25
MYVO	3	8	3	0	0	0	0	14
MYCA	1	5	0	0	0	3	1	10
MYEV	0	4	0	0	0	0	1	5
LABO	0	0	0	0	1	0	0	1
MYSE	0	0	0	0	1	0	0	1

283586 NW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	7	18	31	7	152	147	980	624	1,966
LANO	82	105	140	0	498	273	154	437	1,689
MYYU	4	9	6	0	68	35	104	105	331
LABO	0	1	2	0	3	0	120	77	203
MYEV	3	10	18	0	16	21	39	26	133
MYVO	3	4	8	1	5	11	14	11	57
MYCA	8	8	0	1	0	0	11	28	56
LACI	0	0	0	0	0	0	9	15	24
EPFU	0	0	0	0	0	0	3	3	6
MYSE	0	0	0	0	0	1	0	0	1

283586 SE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	509	478	1,178	2,155	1,565	1,541	2,738	635	10,799
LANO	130	400	236	70	60	64	372	163	1,495
MYYU	7	29	22	22	47	35	1,130	36	1,328
MYVO	23	203	47	55	22	19	784	144	1,297
LABO	0	0	0	0	0	3	282	54	339
LACI	0	0	1	0	0	0	184	1	186
MYEV	34	25	15	20	7	18	14	0	133
MYCA	8	25	1	8	4	0	16	12	74
EPFU	0	0	0	0	0	0	38	6	44
MYSE	0	2	0	0	0	2	7	0	11

283586 SW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	0	34	46	131	143	87	62	167	670
MYLU	0	6	4	16	97	45	7	12	187
MYEV	1	2	4	13	14	39	6	7	86
MYVO	1	0	1	12	22	9	0	5	50
LACI	0	0	0	2	1	1	6	14	24
MYYU	0	0	0	0	1	2	0	1	4
EPFU	0	0	0	0	0	0	1	3	4
MYCA	0	1	0	0	0	0	0	0	1
MYSE	0	0	0	0	0	1	0	0	1
LABO	0	0	0	0	0	0	1	0	1

294832 NE



Species	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	1,588	1,196	989	426	540	639	131	402	5,911
EPFU	242	226	626	196	263	684	153	295	2,685
LANO	462	108	17	131	913	309	47	349	2,336
LABO	12	113	6	1	9	1	0	26	168
MYVO	24	25	13	4	13	60	3	6	148
LACI	7	18	0	9	1	0	75	34	144
MYSE	0	1	2	0	1	16	1	2	23
MYEV	1	1	4	0	0	3	0	0	9

294832 NW



Species	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	930	1,926	697	1,276	1,136	3,215	1,267	2,168	12,615
MYVO	192	776	100	594	723	200	211	173	2,969
MYSE	16	56	1	1	21	119	8	140	362
LANO	8	29	0	15	64	3	5	54	178
LABO	10	19	0	1	1	0	0	69	100
EPFU	1	10	1	1	3	3	7	2	28
MYEV	2	0	9	0	0	0	0	1	12
LACI	1	2	1	2	1	0	0	4	11

294832 SE



Species	2017	2018	2019	2020	2021	2022	2024	Total
MYLU	242	1,190	36	25	788	297	418	2,996
EPFU	140	371	24	8	191	13	66	813
LANO	51	63	10	0	326	21	36	507
LABO	15	202	3	0	4	0	40	264
MYVO	12	23	5	0	11	24	11	86
LACI	8	27	0	0	34	0	0	69
MYSE	0	2	0	0	1	6	1	10
MYEV	0	0	0	0	0	2	0	2

294832 SW



Species	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	445	57	274	1,310	624	183	880	3,178	6,951
MYVO	61	40	29	208	194	69	181	843	1,625
MYSE	9	11	2	13	55	48	38	265	441
LANO	14	0	2	11	12	3	4	8	54
LACI	0	9	0	0	1	0	13	2	25
EPFU	0	1	0	0	5	0	7	9	22
MYEV	1	0	1	0	1	0	1	13	17
LABO	5	0	1	0	0	0	1	6	13
MYYU	0	0	0	0	0	0	0	1	1

317428 NE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	5	63	88	1,469	841	368	227	663	3,724
LANO	49	146	40	1,542	253	203	106	1,364	3,703
EPFU	0	6	3	188	3	52	42	518	812
MYEV	5	60	25	90	29	129	34	132	504
LACI	0	0	1	215	2	17	15	93	343
MYVO	1	9	1	37	20	13	25	38	144
MYCA	1	21	4	33	21	13	16	16	125
MYYU	0	2	0	2	0	2	4	2	12
MYCI	0	4	0	0	0	0	2	0	6
COTO	0	0	0	2	1	0	0	0	3
MYTH	0	1	0	0	0	0	0	1	2
LABO	0	0	0	0	0	0	0	1	1

317428 NW



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	3	60	9	5	78	79	104	272	610
MYLU	1	15	13	9	106	34	69	85	332
EPFU	0	1	1	5	9	16	1	65	98
MYEV	1	6	8	0	8	16	13	45	97
LACI	1	1	5	1	2	35	5	39	89
MYVO	0	1	4	1	8	13	7	24	58
MYCA	0	0	1	0	0	4	1	3	9
MYTH	0	1	0	0	0	0	0	4	5
LABO	0	0	0	0	0	0	0	5	5
EUMA	0	0	0	0	0	0	1	2	3
COTO	0	0	0	0	0	1	1	0	2
MYCI	0	1	0	0	0	0	0	0	1
MYYU	0	0	0	0	1	0	0	0	1

317428 SE



Species	2018	2019	2020	2021	2022	2023	2024	Total
LANO	9	139	35	1,322	165	111	54	1,835
MYLU	8	82	102	607	110	149	168	1,226
LACI	2	1	13	188	16	14	16	250
EPFU	2	7	11	44	18	7	9	98
MYEV	1	3	4	21	15	21	20	85
MYCA	1	7	2	15	3	6	15	49
MYVO	0	4	5	20	2	10	5	46
MYYU	0	0	0	5	0	0	5	10
MYCI	0	5	0	0	0	0	1	6
COTO	0	0	0	2	0	1	1	4
EUMA	0	0	0	0	1	0	0	1

317428 SW



Species	2018	2019	2020	2021	2022	2023	Total
MYLU	445	309	293	3,445	237	1,993	6,722
LANO	54	152	14	1,187	4	167	1,578
LACI	166	1	1	1,328	0	19	1,515
MYCA	63	100	39	108	20	304	634
MYEV	11	34	20	234	15	227	541
EPFU	1	1	6	87	0	87	182
MYVO	7	4	7	126	3	14	161
MYCI	0	105	0	0	0	0	105
MYYU	3	6	2	16	2	14	43
LABO	0	0	1	2	1	24	28
COTO	0	0	1	4	0	0	5

326072 NE



Species	2016	2017	2018	2019	2020	2021	2022	2023	Total
MYLU	59	48	11	80	67	621	2,485	637	4,008
LANO	7	27	15	29	43	105	38	58	322
MYYU	17	2	6	1	32	26	10	72	166
MYCA	106	1	1	1	0	11	11	17	148
EPFU	7	11	0	1	3	26	32	7	87
MYVO	1	2	0	1	0	9	23	18	54
LACI	0	8	1	13	2	0	7	3	34
MYEV	0	1	0	0	1	0	2	1	5
COTO	0	1	0	0	0	0	0	0	1
TABR	0	0	0	1	0	0	0	0	1

326072 NW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	34	225	25	161	1,113	408	472	67	1,673	19	4,197
MYLU	21	108	0	34	8	32	728	151	1,319	235	2,636
MYVO	3	4	0	2	0	5	289	82	297	439	1,121
EPFU	67	1	6	6	1	4	143	11	97	65	401
LACI	1	55	55	10	12	1	7	4	73	88	306
MYCA	1	4	0	3	1	1	9	12	38	121	190
MYYU	4	5	2	26	3	4	5	4	17	13	83
MYEV	1	2	0	2	8	7	4	8	5	20	57
LABO	0	0	0	0	0	0	2	0	0	8	10
COTO	0	0	0	0	0	0	0	7	0	0	7
TABR	2	0	0	0	0	0	0	0	0	0	2
MYTH	0	0	0	0	0	0	0	0	1	1	2
MYCI	0	0	0	0	0	0	0	0	0	2	2

326072 SE



Species	2016	2017	2018	2019	2020	2021	2022	Total
LANO	4	28	1	49	129	54	11	276
MYLU	1	30	2	10	2	13	63	121
MYCA	0	11	1	1	6	26	4	49
LACI	0	12	0	10	1	3	5	31
EPFU	3	2	0	0	0	3	16	24
MYEV	0	6	0	1	5	1	3	16
MYVO	1	4	0	0	0	1	1	7
MYYU	2	0	0	1	0	0	3	6
COTO	0	2	0	0	0	0	0	2

326072 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	144	355	150	376	190	317	1,910	231	1,352	8,311	13,336
LANO	1	39	23	134	146	138	764	23	191	35	1,494
MYYU	43	23	36	30	6	13	48	11	81	449	740
LACI	0	20	15	34	9	5	105	4	137	221	550
MYCA	16	4	14	40	3	19	126	17	75	202	516
EPFU	1	4	3	0	3	5	51	2	33	108	210
MYVO	6	4	5	4	1	2	25	8	32	121	208
MYEV	0	7	1	9	20	1	47	6	14	37	142
COTO	0	7	0	0	0	0	2	0	4	11	24
MYTH	0	0	0	0	0	0	0	0	2	16	18
LABO	0	0	0	0	0	0	1	0	0	9	10
TABR	0	0	1	0	0	0	0	0	0	0	1

37274 NE



Species	2018	2019	2020	2021	2022	2024	Total
MYLU	39	12	10	8	11	6	86
MYEV	3	2	6	1	10	8	30
LANO	0	3	2	0	0	3	8
EPFU	3	2	1	0	0	0	6
MYVO	1	1	0	0	0	1	3
MYSE	1	0	0	0	0	0	1

37274 SE



Species	2018	2019	2020	2021	2022	2024	Total
MYLU	97	35	9	199	264	122	726
LANO	18	43	0	14	118	14	207
MYEV	17	4	9	20	47	17	114
MYVO	21	3	0	18	15	26	83
EPFU	8	5	0	16	0	4	33
LABO	0	0	0	2	3	0	5
MYCA	0	1	0	2	1	0	4
LACI	0	0	0	1	0	3	4

37274 SW



Species	2018	2019	2020	2021	2022	2024	Total
MYLU	390	17	150	41	24	34	656
MYVO	5	1	4	6	8	217	241
LANO	53	40	21	2	10	11	137
EPFU	18	1	1	0	2	1	23
LABO	2	0	2	5	1	0	10
MYEV	1	1	0	1	1	3	7
LACI	0	4	0	0	0	2	6
MYCA	0	0	0	1	0	0	1
MYSE	0	0	0	0	0	1	1

38442 NW



Species	2018	2020	2021	2023	2024	2025	Total
MYCI	409	781	1,058	299	2,949	986	6,482
MYVO	43	151	1,387	53	95	170	1,899
LANO	346	539	109	17	47	329	1,387
MYLU	79	99	478	94	193	195	1,138
EPFU	10	92	328	9	22	25	486
MYYU	38	26	174	74	92	33	437
MYTH	11	17	63	76	56	150	373
MYCA	10	7	44	28	74	35	198
LACI	1	12	58	1	22	27	121
EUMA	15	35	12	2	4	5	73
MYEV	1	25	7	20	3	8	64
COTO	2	1	2	3	2	1	11
LABO	0	0	1	0	0	0	1
MYSE	0	0	0	0	0	1	1

38442 SW



Species	2018	2019	2020	2021	2022	2023	2024	Total
MYCI	33	121	185	164	830	207	1,180	2,720
LANO	269	158	237	100	15	78	90	947
MYLU	17	15	9	263	124	44	125	597
MYCA	1	1	1	15	34	0	124	176
EPFU	2	7	11	83	17	10	26	156
MYYU	9	2	5	70	5	4	5	100
LACI	2	7	11	7	5	5	43	80
EUMA	4	4	5	4	1	25	11	54
MYEV	0	0	2	3	15	3	8	31
MYVO	0	0	1	6	7	0	14	28
MYTH	1	1	1	1	1	0	3	8
LABO	0	0	0	3	2	0	3	8
COTO	0	1	0	0	1	0	1	3
MYSE	0	0	0	0	0	0	1	1

42026 NE



Species	2022	2024	2025	Total
MYLU	78	100	119	297
MYVO	72	16	6	94
MYEV	13	0	2	15
LABO	0	0	11	11
EPFU	7	1	1	9
LANO	2	1	2	5
LACI	0	0	3	3
MYSE	0	0	1	1

42026 NW



Species	2022	2024	2025	Total
MYLU	43	455	479	977
MYEV	6	5	4	15
MYVO	0	9	6	15
LANO	3	7	4	14

Species	2022	2024	2025	Total
EPFU	1	4	0	5
LABO	1	2	0	3

42026 SE



Species	2022	2024	2025	Total
MYLU	382	324	344	1,050
LABO	2	4	88	94
LANO	4	74	8	86
MYVO	17	13	11	41
LACI	0	0	34	34
EPFU	1	8	9	18
MYEV	3	7	1	11
MYSE	0	2	0	2

42026 SW



Species	2022	2024	2025	Total
MYLU	869	1,814	394	3,077
MYVO	24	13	20	57
LANO	7	29	1	37
LABO	12	9	5	26
MYEV	1	4	0	5
MYSE	0	4	1	5
EPFU	0	1	0	1
LACI	0	0	1	1

47210 NE



Species	2023	Total
MYLU	465	465
MYCA	175	175
LANO	98	98
MYVO	46	46
LACI	30	30
MYEV	27	27
MYYU	26	26
MYCI	22	22
COTO	3	3
EPFU	2	2
MYTH	1	1

47210 NE2



Species	2024	2025	Total
MYCA	45	533	578
MYLU	31	240	271
LANO	21	73	94
MYCI	6	33	39
MYEV	6	32	38
LACI	13	4	17
EPFU	0	16	16
LABO	0	10	10
MYYU	1	5	6
MYVO	2	3	5
EUMA	2	1	3
ANPA	1	0	1

47210 NW



Species	2023	2024	2025	Total
MYLU	279	109	403	791
MYCA	59	31	184	274
MYEV	55	21	91	167
LANO	69	8	48	125
MYVO	52	8	20	80
MYCI	29	5	34	68
EPFU	26	0	33	59
LABO	0	8	17	25
LACI	8	2	11	21
COTO	5	0	0	5
MYYU	2	0	3	5
MYTH	2	0	0	2
ANPA	0	0	2	2
EUMA	0	0	1	1

47210 SE



Species	2023	2024	2025	Total
LANO	175	7	34	216
MYEV	63	5	36	104
EPFU	36	0	20	56
MYCA	38	1	17	56
MYLU	26	5	4	35
MYVO	33	0	0	33
LACI	17	0	8	25
MYCI	6	2	4	12
COTO	4	1	1	6
MYTH	5	0	0	5
MYYU	3	0	0	3
ANPA	0	2	0	2
LABO	0	0	1	1

47210 SW



Species	2023	2025	Total
MYEV	191	48	239
MYLU	162	20	182
MYCA	52	15	67
EPFU	48	17	65
LANO	34	21	55
MYVO	29	0	29
MYCI	20	5	25
LABO	1	13	14
LACI	11	2	13
COTO	4	1	5
MYYU	1	3	4
MYSE	0	2	2
MYTH	1	0	1

47322 NE



Species	2016	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYVO	236	85	1,289	3	108	1,094	1,347	40	252	4,454
MYLU	121	42	114	0	14	44	953	162	241	1,691
EPFU	6	1	55	1	3	10	27	150	74	327
MYEV	6	8	89	0	65	6	33	22	14	243
MYSE	23	5	17	0	4	46	6	13	9	123
LACI	0	0	11	0	0	0	2	8	59	80
LABO	2	1	23	0	0	0	0	5	22	53
LANO	1	7	11	0	1	1	1	8	15	45
MYYU	0	0	0	0	0	0	0	5	6	11

47322 NW



Species	2016	2017	2018	2019	2020	2021	2022	2024	Total
MYLU	697	811	608	2,049	0	313	244	197	4,919
MYVO	44	37	19	58	3	59	27	34	281
EPFU	175	2	0	4	2	10	6	58	257
LABO	21	18	0	3	0	7	3	66	118
MYSE	2	17	0	0	0	4	0	36	59
MYEV	2	1	0	5	0	4	14	30	56
LACI	19	2	0	1	0	0	1	30	53
LANO	3	1	0	0	0	39	8	2	53

47322 SE



Species	2016	2017	2018	2019	2020	2021	2022	Total
MYLU	145	185	1,249	404	75	194	1,537	3,789
MYVO	59	27	132	66	7	5	177	473
MYEV	11	19	38	6	5	2	22	103
LANO	0	1	3	0	0	0	31	35
MYSE	3	7	8	0	1	1	4	24
LABO	3	10	7	0	0	1	1	22
EPFU	2	1	7	1	0	2	5	18
LACI	1	1	0	0	0	0	0	2

47322 SW



Species	2016	2017	2018	2019	2020	2021	2022	2024	2025	Total
MYLU	1	12	289	52	91	128	141	53	5	772
LANO	0	0	1	0	5	49	53	1	0	109
EPFU	1	0	2	4	15	4	64	10	0	100
MYVO	0	0	29	12	2	6	10	17	2	78
MYEV	0	2	2	4	12	1	10	23	1	55
LABO	0	0	0	0	2	1	0	23	2	28
LACI	0	0	5	0	2	0	3	14	2	26
MYSE	1	0	0	0	0	0	0	2	0	3

53210 NE



Species	2025	Total
MYEV	3	3
MYLU	1	1

53210 NW



Species	2025	Total
MYEV	262	262
MYLU	179	179
LANO	155	155
EPFU	32	32
LABO	25	25
LACI	17	17
MYVO	17	17

53210 SE



Species	2025	Total
MYLU	20	20
MYVO	9	9
LABO	5	5
LANO	4	4
LACI	2	2
MYEV	2	2

53210 SW



Species	2025	Total
MYLU	434	434
LANO	254	254
MYVO	92	92
LACI	65	65
LABO	49	49
MYEV	49	49

Species	2025	Total
EPFU	25	25
MYSE	5	5

57322 NE



Species	2017	2018	2019	2020	2021	2022	2023	Total
MYLU	116	247	1,471	276	2,039	3,515	1,718	9,382
LANO	17	42	18	6	323	296	67	769
MYVO	2	2	25	1	58	49	78	215
LACI	3	3	5	3	108	3	16	141
EPFU	7	3	0	0	28	15	20	73
MYYU	1	4	4	0	27	3	16	55
MYCA	0	2	7	0	20	6	13	48
COTO	0	0	1	0	0	4	2	7
MYEV	0	0	0	0	0	1	2	3
LABO	0	0	0	0	0	0	2	2

57322 NW



Species	2017	2018	2019	2020	2021	2022	2025	Total
MYLU	3,087	721	541	219	7,150	6,387	5,918	24,023
LANO	2,399	256	645	80	1,188	995	151	5,714
LACI	50	10	10	9	9	24	40	152
MYVO	26	7	6	0	28	56	8	131
MYYU	44	18	5	8	5	21	24	125
EPFU	53	5	0	2	13	14	21	108
MYCA	31	5	12	1	7	4	16	76
MYEV	3	1	0	0	0	1	3	8
TABR	0	1	0	0	0	1	0	2
COTO	0	0	0	0	0	0	1	1
LABO	0	0	0	0	0	0	1	1

57322 SE



Species	2018	2019	2020	2021	2022	2023	2025	Total
LANO	0	124	23	235	83	25	1,470	1,960
LACI	0	15	1	336	275	185	421	1,233
MYLU	11	8	0	236	77	72	406	810
MYVO	0	2	0	31	8	10	170	221
EPFU	1	0	0	12	2	6	160	181
MYEV	0	1	0	4	0	4	21	30
MYCA	0	1	0	4	2	1	4	12
MYYU	0	0	0	2	1	3	6	12
COTO	0	0	0	0	0	1	5	6
MYTH	0	0	0	0	0	0	1	1

57322 SW



Species	2018	2019	2020	2022	2023	2025	Total
MYLU	3,235	318	221	4,540	1,522	4,524	14,360
MYVO	388	38	23	99	57	67	672
MYYU	74	8	58	37	88	170	435
MYCA	275	25	1	24	32	19	376
LANO	47	7	22	114	21	17	228
EPFU	2	1	4	19	78	10	114
MYEV	22	0	4	4	25	1	56
LACI	13	8	4	4	12	3	44
COTO	0	0	0	0	4	0	4
MYCI	0	0	0	0	0	3	3
LABO	0	0	0	0	2	0	2

58842 NW



Species	2024	2025	Total
MYLU	52	87	139
MYEV	18	13	31
LABO	1	14	15
MYVO	11	2	13
EPFU	3	4	7
LACI	0	4	4
MYSE	1	0	1
LANO	0	1	1

58842 SE



Species	2024	2025	Total
MYLU	32	8	40
MYEV	25	10	35
EPFU	10	21	31
MYVO	9	0	9

Species	2024	2025	Total
LANO	2	6	8
LABO	0	2	2
MYSE	1	0	1
LACI	0	1	1

58842 SW



Species	2024	2025	Total
MYLU	57	37	94
LABO	0	23	23
MYVO	7	7	14
LACI	1	10	11
LANO	1	9	10
EPFU	1	3	4
MYEV	1	1	2

5994 NE



Species	2020	2021	2022	2023	2024	Total
MYLU	479	37	104	338	175	1,133
EPFU	195	11	9	109	271	595
MYYU	49	2	59	159	51	320
LACI	146	18	10	10	92	276
MYEV	73	18	11	57	104	263
MYCI	144	2	7	13	0	166
MYVO	112	3	9	15	23	162
LANO	94	2	3	11	35	145
MYCA	8	25	22	12	24	91
LABO	5	0	11	1	13	30
ANPA	0	0	0	0	11	11
COTO	1	0	0	0	1	2
MYTH	2	0	0	0	0	2
MYSE	0	0	0	0	1	1

5994 SE



Species	2020	2021	2022	2023	2024	Total
MYLU	127	7	330	217	566	1,247
MYEV	149	13	249	147	364	922
LACI	131	2	136	39	77	385
EPFU	26	2	66	39	65	198
LANO	26	2	27	18	67	140
MYVO	27	0	25	16	3	71
LABO	7	0	43	0	13	63
MYCA	22	0	8	2	2	34
ANPA	0	0	22	0	5	27
COTO	11	0	3	6	2	22
MYCI	16	0	0	0	2	18
MYYU	3	0	1	3	1	8
MYTH	1	0	2	0	1	4
PAHE	0	0	2	0	0	2
MYSE	0	0	0	0	1	1

60586 NW



Species	2020	2021	2022	2023	2024	2025	Total
MYLU	225	597	498	138	840	2,730	5,028
LACI	2	653	352	32	24	29	1,092
MYVO	6	62	29	9	9	678	793
EPFU	14	115	271	9	40	19	468
MYEV	10	132	81	25	17	13	278
MYCA	1	11	7	8	15	211	253

Species	2020	2021	2022	2023	2024	2025	Total
LANO	4	45	41	4	17	22	133
LABO	6	2	45	1	9	25	88
COTO	0	9	0	1	4	0	14
MYYU	1	0	0	0	1	7	9
MYSE	0	0	0	0	2	6	8
ANPA	0	0	0	0	0	2	2

60586 SW



Species	2020	2021	2022	2023	2024	2025	Total
MYLU	81	550	253	57	427	3,751	5,119
MYVO	2	6	59	16	85	560	728
MYCA	1	40	31	2	66	177	317
LACI	5	183	24	1	4	39	256
EPFU	8	99	13	0	7	19	146
MYEV	2	5	21	1	79	25	133
LANO	0	20	8	1	4	31	64
MYSE	0	0	0	0	20	42	62
LABO	1	0	2	0	14	14	31
MYYU	0	1	2	0	0	7	10
ANPA	0	0	0	0	2	1	3
COTO	0	0	0	0	1	0	1

6202 NE



Species	2016	2018	2019	2021	2022	2023	2024	Total
LANO	1	61	59	0	4	61	366	552
EPFU	3	16	5	12	117	96	186	435
LACI	0	10	306	0	1	2	71	390
MYLU	9	17	4	14	6	107	2	159
MYCA	8	37	4	7	23	42	7	128
MYYU	0	70	2	3	5	23	0	103
MYVO	1	12	0	0	3	12	0	28
MYEV	0	0	2	4	2	3	0	11
LABO	0	0	0	0	0	1	7	8
COTO	1	0	0	0	1	1	0	3
PAHE	0	0	0	0	0	0	1	1

6202 NE2



Species	2023	2025	Total
LANO	238	380	618
EPFU	139	218	357
MYCA	12	92	104
LACI	0	34	34
MYEV	3	16	19
MYYU	3	0	3
MYLU	0	3	3
COTO	1	0	1
MYVO	1	0	1

6202 NW



Species	2018	2019	2020	2021	2022	2023	Total
LANO	81	145	30	8	71	156	491
MYCA	1	2	2	0	3	4	12
MYEV	0	4	0	2	0	4	10
MYLU	1	0	1	2	1	4	9
COTO	0	1	1	0	0	4	6
MYYU	0	0	1	0	0	2	3
MYVO	0	0	0	0	0	3	3
LACI	0	2	0	0	0	0	2

6202 SE



Species	2018	2019	2021	2022	Total
LANO	77	171	160	438	846
MYLU	5	20	26	396	447
MYYU	12	6	15	279	312
MYCA	3	2	75	52	132
EPFU	1	2	35	21	59
MYVO	0	0	13	26	39
LACI	3	16	0	7	26
MYEV	5	0	4	2	11
COTO	0	0	0	1	1

6202 SW



Species	2018	2019	2020	2021	2022	2024	2025	Total
LANO	13	59	27	9	768	108	607	1,591
MYYU	14	101	161	26	495	129	138	1,064
MYCA	6	0	26	14	161	24	29	260
MYEV	12	11	13	11	96	47	24	214
EPFU	0	0	5	1	29	5	1	41
MYLU	0	3	3	1	5	5	10	27
LACI	1	13	1	0	1	0	6	22
COTO	1	0	3	0	0	1	1	6
MYVO	0	0	0	0	3	2	0	5

70250 NW



Species	2020	2021	2022	2023	2024	2025	Total
MYLU	162	558	797	142	208	278	2,145
EPFU	106	118	934	16	115	11	1,300
LACI	39	60	381	12	53	31	576
LANO	27	53	86	10	28	15	219
MYVO	30	43	81	8	29	9	200
LABO	2	0	43	0	2	14	61
MYEV	1	4	20	2	7	2	36
MYCA	3	1	11	1	0	2	18
COTO	0	1	0	0	1	0	2
MYSE	0	0	0	0	0	1	1

70250 SW



Species	2020	2021	2022	2025	Total
MYLU	31	295	59	203	588
MYVO	8	8	5	7	28
EPFU	2	18	2	2	24
LABO	0	2	1	13	16
LANO	1	10	1	2	14
LACI	0	2	0	10	12
MYCA	1	0	0	1	2
MYEV	0	1	0	0	1
MYYU	0	0	1	0	1
MYSE	0	0	0	1	1

73002 NE



Species	2016	2017	2018	2019	2020	2021	2022	2024	2025	Total
EUMA	422	231	49	208	104	640	455	1,071	1,655	4,835
LANO	3	28	57	42	26	96	141	38	1,249	1,680
MYCI	66	43	73	116	1	109	621	346	215	1,590
EPFU	13	78	55	12	13	76	179	75	804	1,305
LACI	8	39	1	0	31	46	312	51	620	1,108
MYLU	4	6	4	4	2	13	98	70	210	411
MYVO	5	1	0	1	0	8	103	90	26	234
MYTH	36	7	1	2	2	8	4	12	18	90
MYYU	1	0	0	1	0	0	5	26	25	58
MYCA	1	1	1	0	0	2	10	11	30	56
MYEV	1	0	2	2	0	3	2	2	4	16
MYSE	0	0	0	0	0	0	0	4	1	5
LABO	0	0	0	0	0	0	0	2	1	3
COTO	1	0	0	0	0	0	0	1	0	2

73002 SE



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
EUMA	121	38	14	145	28	224	26	15	67	87	765
LANO	8	55	40	49	48	37	64	2	56	214	573
MYCI	11	35	30	18	0	31	147	63	132	12	479
LACI	1	106	5	4	55	15	16	6	98	89	395
EPFU	12	57	19	25	43	30	57	2	36	38	319
MYLU	2	4	5	0	0	13	11	0	27	51	113
MYCA	0	0	1	2	0	2	5	0	5	23	38
MYTH	0	0	3	0	0	2	3	5	1	2	16
MYEV	0	4	0	0	0	1	1	0	0	7	13
MYVO	0	2	1	5	0	0	3	0	1	1	13
COTO	0	1	2	0	0	0	1	0	0	0	4
MYYU	0	0	0	0	0	0	1	0	0	1	2
LABO	0	0	0	0	0	0	0	0	2	0	2
PAHE	0	0	0	0	0	0	0	0	1	0	1

73002 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
EPFU	221	69	93	101	4	66	300	14	205	1,062	2,135
MYCI	124	136	307	89	3	23	215	90	371	39	1,397
LANO	0	52	10	43	5	36	103	1	208	632	1,090
LACI	1	21	16	3	2	39	87	1	68	335	573
EUMA	44	5	21	56	61	63	36	57	109	19	471
MYLU	3	5	19	4	0	22	11	2	204	23	293
MYCA	9	38	17	6	3	9	55	29	58	30	254
MYVO	2	14	20	7	0	5	119	7	34	0	208
MYEV	0	5	3	39	2	7	19	0	20	20	115
MYYU	1	1	5	0	0	1	0	0	3	2	13
COTO	1	0	1	0	0	3	0	1	6	0	12
MYTH	0	0	0	0	0	0	0	0	3	7	10
ANPA	0	0	0	0	0	0	0	0	2	1	3
LABO	0	0	0	0	0	0	0	0	3	0	3
MYSE	0	0	0	0	0	0	0	0	2	0	2
PAHE	0	0	0	0	0	0	0	0	1	0	1

73786 NW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	28	53	7	7	31	32	238	233	183	72	884
LANO	10	29	9	85	65	70	36	63	189	75	631
MYYU	15	1	4	0	10	9	37	30	100	17	223
LACI	0	1	1	9	27	2	1	0	2	33	76
MYCA	4	2	0	0	1	3	14	6	6	8	44
EPFU	6	0	0	0	0	4	5	5	4	1	25
MYVO	0	0	1	0	0	0	1	0	3	0	5
LABO	0	0	0	0	0	0	0	0	1	0	1
MYCI	0	0	0	0	0	0	0	0	1	0	1
PAHE	0	0	0	0	0	0	0	0	1	0	1

73786 SE



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	148	687	20	7	287	158	1,390	1,991	1,065	682	6,435
LANO	41	248	34	162	391	109	1,319	241	181	737	3,463
EPFU	133	8	6	2	7	11	189	91	77	62	586
MYCA	1	5	2	2	5	11	19	20	6	53	124
LACI	0	53	0	2	5	1	2	0	4	16	83
MYVO	4	2	0	0	1	2	24	10	3	17	63
MYYU	4	1	1	2	2	1	10	4	1	0	26
TABR	3	0	0	0	0	0	0	0	0	0	3
MYEV	0	2	0	0	1	0	0	0	0	0	3
MYCI	0	0	0	0	0	0	0	0	0	1	1

73786 SW



Species	2016	2018	2019	2020	2021	2022	2023	2025	Total
MYLU	52	18	52	37	53	180	106	1,101	1,599
MYYU	12	9	39	29	35	30	30	401	585
LANO	29	6	35	45	58	40	36	56	305
LACI	0	5	83	58	1	2	0	105	254
MYCA	1	4	7	6	4	12	4	80	118
EPFU	7	1	0	0	0	17	6	47	78
MYVO	1	0	0	0	0	0	1	11	13
TABR	0	1	0	0	0	3	0	0	4
MYCI	0	0	0	0	0	0	0	1	1

77162 NE



Species	2016	2017	2018	2019	2020	2021	Total
MYCA	50	6	27	54	0	6	143
MYLU	1	46	8	14	5	9	83
MYEV	1	2	3	3	0	3	12
LANO	0	1	3	0	0	1	5
MYVO	0	1	1	0	0	1	3
MYSE	1	1	0	0	0	0	2
MYYU	1	0	0	0	0	0	1
EPFU	0	0	0	0	0	1	1

77162 NW1



Species	2016	2017	2018	2019	2020	2021	2022	2023	Total
MYLU	261	553	286	51	135	252	84	117	1,739
LANO	30	201	145	34	177	234	191	22	1,034
EPFU	376	110	53	4	39	22	19	1	624
LACI	4	125	75	46	3	37	5	233	528
MYVO	11	32	9	1	4	18	5	7	87
MYYU	6	14	13	3	5	12	3	3	59
MYCA	15	14	7	1	1	13	1	3	55
COTO	0	4	3	8	1	1	2	32	51
MYEV	14	13	2	3	2	1	5	1	41
MYSE	3	1	0	0	0	1	0	0	5

77162 NW2



Species	2016	2017	2018	2019	2020	2021	2022	2023	2025	Total
MYLU	98	2,344	1,047	734	499	2,178	714	1,292	554	9,460
LANO	192	106	91	31	457	739	2,268	8	508	4,400
MYEV	15	279	13	135	268	34	3	4	31	782
MYYU	8	211	60	26	26	46	38	139	21	575
MYCA	11	198	53	25	11	38	62	72	29	499
EPFU	19	18	27	1	21	16	46	0	156	304
MYVO	13	39	33	8	11	38	5	71	34	252
LACI	3	110	11	7	2	20	1	50	23	227
MYSE	6	2	1	2	4	2	1	0	31	49
COTO	0	1	1	1	1	3	2	3	1	13

77162 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2025	Total
MYLU	51	132	333	52	174	277	87	142	339	1,587
LANO	19	15	48	15	64	33	60	19	461	734
MYCA	13	29	87	66	66	63	21	306	66	717
EPFU	55	59	12	1	25	34	10	107	104	407
LACI	1	25	8	1	1	87	4	20	5	152
MYVO	1	14	26	0	7	27	15	30	19	139
MYYU	5	18	64	1	9	10	7	6	2	122
MYEV	5	17	4	3	12	7	16	23	2	89
MYSE	3	4	0	0	1	4	0	9	2	23
COTO	0	1	1	0	2	2	0	4	8	18
LABO	0	0	0	0	0	0	0	0	1	1

79018 NE



Species	2016	2017	2018	2019	2020	2022	2023	2024	2025	Total
LANO	28	91	17	16	1	40	323	106	260	882
MYCA	64	61	88	16	8	12	390	43	128	810
MYLU	54	126	18	29	1	17	54	7	27	333
MYEV	34	45	17	24	6	28	60	5	51	270
MYVO	17	77	37	47	0	6	11	1	34	230
EPFU	3	7	7	7	0	1	130	6	31	192
COTO	0	36	13	10	2	0	5	0	1	67
MYYU	5	3	0	0	0	0	51	0	1	60
LACI	1	0	2	0	1	1	4	0	0	9
MYSE	0	0	0	0	0	0	1	1	7	9
LABO	0	0	0	0	0	0	0	0	1	1

79018 NW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	121	145	2	312	29	281	30	221	147	129	1,417
MYCA	4	162	12	175	13	49	16	171	43	218	863
LACI	24	23	1	156	1	39	40	120	2	16	422
MYLU	5	89	1	92	2	20	8	56	18	67	358
MYYU	4	36	8	44	8	36	18	42	29	104	329
MYVO	2	36	6	86	0	14	2	8	4	41	199
EPFU	6	72	1	20	3	24	4	21	16	20	187
COTO	0	6	0	11	4	7	11	4	47	23	113
MYEV	0	1	1	11	4	3	7	8	2	10	47
MYSE	0	0	0	0	0	0	0	1	0	4	5
LABO	0	0	0	0	0	0	0	0	0	1	1

79018 SE



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
LANO	12	195	2	101	12	381	78	234	73	114	1,202
MYCA	3	3	2	3	1	6	21	429	1	147	616
MYVO	0	1	0	3	0	17	18	44	2	48	133
EPFU	20	31	2	33	3	5	2	15	7	8	126
MYEV	0	0	0	2	0	4	8	17	7	17	55
MYLU	2	1	0	1	0	4	1	12	2	23	46
LACI	1	3	0	0	1	1	0	22	1	3	32
COTO	0	1	0	0	0	1	4	2	10	6	24
MYYU	0	0	0	0	0	1	0	2	0	13	16
MYSE	0	0	0	0	0	0	0	2	0	3	5

79018 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	Total
MYYU	32	369	31	367	428	3,370	1,333	411	492	6,833
LANO	621	801	4	196	82	846	535	599	337	4,021
MYCA	25	165	22	1,229	32	271	270	211	485	2,710
MYLU	40	417	20	512	2	598	226	604	137	2,556
LACI	17	109	0	6	188	9	8	356	0	693
EPFU	15	73	2	19	0	159	8	35	66	377
MYEV	11	94	0	82	0	6	25	58	20	296
MYVO	8	45	4	32	1	54	34	61	18	257
COTO	1	21	0	20	5	21	8	67	7	150
MYSE	0	0	0	1	0	0	0	1	1	3

79018 SW2



Species	2025	Total
LANO	672	672
MYCA	421	421
MYYU	157	157
MYLU	129	129
LACI	37	37
MYVO	35	35
COTO	9	9
MYEV	8	8
EPFU	7	7
MYSE	3	3

86074 NE



Species	2017	2018	2020	2021	2022	2023	2024	Total
LANO	58	38	39	286	35	29	123	608
LACI	137	4	0	413	8	14	13	589
MYLU	7	0	1	270	0	10	0	288
EPFU	75	0	21	47	14	13	0	170
MYVO	3	0	1	16	0	1	0	21
MYEV	2	0	1	4	2	4	0	13
MYYU	1	0	0	4	1	5	2	13
COTO	0	0	1	6	0	0	1	8
MYCA	1	0	2	2	0	1	0	6
MYCI	0	0	0	0	0	1	0	1
MYTH	0	0	0	0	0	1	0	1

86074 NW



Species	2016	2017	2020	2022	Total
MYLU	301	239	939	2,098	3,577
LANO	409	81	246	835	1,571
MYYU	168	20	198	218	604
MYCA	130	13	83	60	286
MYVO	97	25	15	42	179
LACI	16	1	12	36	65
EPFU	5	4	10	9	28
LABO	0	0	0	19	19
MYEV	2	2	4	7	15
COTO	2	0	1	1	4
MYTH	1	0	1	1	3

86074 SE



Species	2016	2017	2020	2021	2022	Total
LACI	1	4	46	224	849	1,124
LANO	48	94	382	335	108	967

Species	2016	2017	2020	2021	2022	Total
MYLU	14	10	188	379	68	659
EPFU	23	32	2	51	171	279
MYYU	34	3	21	29	12	99
MYCA	30	8	14	28	1	81
MYVO	9	0	14	34	5	62
MYEV	1	4	8	7	1	21
COTO	2	2	1	3	1	9
LABO	0	0	0	2	1	3

86074 SW



Species	2016	2017	2019	2020	2022	2023	2024	Total
MYLU	15	23	0	45	1,346	470	130	2,029
LANO	63	21	72	82	257	69	80	644
LACI	42	1	5	1	119	65	19	252
MYYU	9	0	0	31	114	28	51	233
MYVO	1	0	0	2	138	77	8	226
MYCA	1	0	0	22	53	18	46	140
EPFU	37	4	0	10	5	5	3	64
MYEV	1	1	0	11	8	13	1	35
COTO	0	0	0	7	6	12	0	25
MYCI	0	0	0	0	7	6	6	19
LABO	0	0	0	0	0	0	5	5

89962 NE



Species	2022	2023	2025	Total
MYLU	150	719	603	1,472
LACI	56	28	352	436
LANO	81	122	196	399
EPFU	66	98	95	259
MYCA	11	81	36	128
MYYU	4	50	13	67
MYVO	0	4	18	22
MYCI	1	10	3	14
MYEV	1	7	3	11
LABO	3	0	7	10
ANPA	3	0	1	4

89962 NW



Species	2022	Total
MYLU	1,486	1,486

Species	2022	Total
LANO	216	216
EPFU	128	128
MYCA	100	100
LACI	95	95
MYYU	69	69
MYCI	23	23
MYEV	8	8
MYVO	7	7
LABO	3	3
MYTH	2	2

89962 SE



Species	2022	2024	2025	Total
LANO	447	317	281	1,045
MYLU	138	332	431	901
LACI	447	139	133	719
EPFU	464	95	46	605
MYCA	14	127	84	225
MYYU	34	31	44	109
MYEV	4	20	40	64
COTO	15	8	5	28
MYVO	2	2	9	13
MYTH	0	12	1	13
ANPA	5	0	1	6
MYCI	4	0	0	4
MYSE	0	0	3	3
LABO	1	1	0	2

89962 SW



Species	2022	2024	2025	Total
MYLU	448	1,333	1,432	3,213
LANO	146	674	724	1,544
EPFU	70	582	429	1,081
MYCA	44	181	295	520
MYYU	15	60	237	312
LACI	31	35	61	127
MYEV	66	8	22	96
MYVO	13	11	35	59
COTO	29	2	0	31
MYCI	2	5	8	15
ANPA	6	0	4	10
LABO	0	1	6	7
MYTH	1	0	0	1
MYSE	0	0	1	1

93290 NE



Species	2016	2018	2020	2021	2022	2023	2025	Total
LACI	19	152	267	536	762	21	81	1,838
LANO	42	205	158	219	253	65	147	1,089
EPFU	13	127	78	179	55	41	78	571
MYCI	48	36	141	50	120	31	51	477
MYLU	14	11	30	96	59	17	40	267
MYCA	6	3	12	4	31	4	14	74
ANPA	0	1	3	20	0	0	11	35
MYYU	0	2	9	4	10	6	3	34
MYTH	0	3	5	2	4	2	0	16
MYVO	0	0	6	7	0	0	2	15
MYEV	2	4	1	3	2	0	1	13
COTO	1	1	2	3	0	0	4	11
LABO	0	0	0	2	3	1	4	10
EUMA	0	1	0	0	0	1	0	2

93290 NW



Species	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYCI	130	151	56	408	55	1,155	128	229	1,521	3,833
EPFU	91	115	76	218	197	373	386	403	490	2,349
LANO	45	128	86	211	185	624	312	251	316	2,158
MYLU	51	34	18	34	77	317	93	142	1,041	1,807
LACI	19	77	3	85	265	235	244	196	129	1,253
MYCA	30	30	12	3	5	260	27	23	184	574
MYYU	2	11	7	1	6	128	12	18	13	198
MYVO	1	3	4	6	13	54	18	9	41	149
LABO	1	0	0	0	0	14	15	13	39	82
ANPA	0	1	0	1	15	8	7	6	25	63
MYEV	4	6	2	4	0	22	6	5	9	58
MYTH	1	0	1	4	0	13	2	8	18	47
COTO	0	1	1	7	1	13	6	9	8	46
EUMA	0	0	0	0	1	0	3	1	2	7
PAHE	0	0	0	0	0	0	0	0	1	1

93290 SE



Species	2016	2017	2018	2019	2020	2021	2023	2024	2025	Total
MYYU	1,449	889	1,438	738	1,046	1,254	5,628	11,624	6,510	30,576
MYLU	439	214	351	275	661	1,850	1,199	3,934	1,642	10,565
MYCA	377	29	91	188	132	176	1,424	877	1,104	4,398
MYCI	165	15	428	281	389	283	315	754	444	3,074
LANO	78	62	259	89	249	199	259	324	387	1,906
EPFU	35	49	161	38	214	191	191	658	237	1,774
LACI	58	40	107	3	42	175	117	176	76	794
MYVO	77	1	11	9	20	17	30	103	175	443
LABO	0	1	1	0	0	0	18	56	48	124
MYEV	4	7	5	0	9	2	13	6	11	57
ANPA	3	0	0	0	0	18	2	5	3	31
MYTH	7	1	6	0	1	1	4	3	4	27
COTO	2	0	0	0	0	0	0	2	1	5
EUMA	0	0	0	0	0	0	2	2	0	4
PAHE	0	1	0	0	0	0	0	0	0	1

93290 SW



Species	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYCI	325	793	0	134	640	26	1,707	571	411	140	4,747
LANO	14	35	1	27	111	147	289	195	244	356	1,419
EPFU	5	19	0	13	153	97	144	113	265	336	1,145
LACI	7	6	0	0	44	220	157	163	265	167	1,029
MYTH	5	6	0	1	47	57	18	31	33	49	247
MYLU	13	5	0	3	3	19	37	14	30	33	157
MYCA	1	0	0	1	15	2	53	9	7	13	101
EUMA	0	0	0	5	15	0	19	1	19	0	59
COTO	1	0	0	0	7	1	11	15	16	4	55
ANPA	1	0	0	0	0	7	8	5	2	12	35
MYVO	4	1	0	0	2	0	12	3	5	5	32
LABO	0	0	0	0	0	0	13	2	3	3	21
MYEV	2	1	1	0	1	0	5	0	1	4	15
MYYU	0	1	0	1	1	1	6	1	2	1	14

94522 NE1



Species	2017	2018	2019	Total
MYCA	1,094	128	167	1,389
MYLU	747	139	391	1,277
MYEV	62	17	17	96

94522 NE2



Species	2016	2018	2019	Total
MYLU	446	215	229	890
LANO	312	67	102	481
MYEV	20	17	126	163
MYCA	50	5	15	70

94522 NW



Species	2016	2017	2018	2019	Total
MYCA	219	2,557	1,301	182	4,259
MYLU	620	810	342	283	2,055
LANO	188	5	13	9	215
MYEV	36	4	2	1	43

94522 SE



Species	2017	2018	2019	Total
MYLU	84	34	48	166
LANO	85	13	17	115
MYEV	44	3	6	53
MYCA	2	8	7	17

9466 NE



Species	2022	2024	Total
MYLU	2	17	19
MYEV	2	3	5
MYVO	2	3	5
LANO	1	0	1

9466 NW



Species	2022	2024	2025	Total
MYLU	41	19	121	181
MYVO	8	2	58	68
MYEV	10	3	18	31
LABO	1	0	14	15
LANO	1	1	6	8
MYSE	0	0	8	8
EPFU	0	0	4	4

9466 SE



Species	2022	2024	2025	Total
MYLU	15	26	87	128
MYVO	7	2	33	42
LANO	0	2	35	37
LABO	0	0	32	32
MYEV	5	3	10	18
EPFU	0	0	3	3
LACI	0	0	3	3

9466 SW



Species	2022	2024	2025	Total
MYLU	10	166	512	688
LANO	0	16	195	211
LABO	0	2	101	103
MYVO	8	16	74	98
MYEV	4	14	23	41

Species	2022	2024	2025	Total
EPFU	0	2	38	40
MYSE	0	0	6	6
LACI	0	0	1	1

986 NE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	96	159	184	132	3,668	618	614	1,186	6,657
LABO	0	0	0	0	280	3	11	49	343
MYEV	2	49	12	1	129	23	43	32	291
MYVO	33	49	9	8	50	28	23	91	291
LANO	9	13	38	0	94	9	18	61	242
MYSE	0	0	0	0	2	11	0	26	39
LACI	0	0	1	0	1	1	2	20	25
MYCA	0	0	3	0	6	1	0	0	10
EPFU	0	1	1	0	0	0	0	3	5

986 SE



Species	2018	2019	2020	2021	2022	2023	2024	2025	Total
MYLU	9	323	2	87	4,333	6,503	1,823	1,759	14,839
LABO	0	2	0	0	107	844	260	250	1,463
MYVO	0	26	0	16	75	544	62	181	904
MYEV	0	9	0	0	150	296	39	123	617
LANO	0	23	0	3	126	104	71	240	567
MYSE	0	0	0	3	69	48	11	42	173
LACI	0	0	0	3	0	55	1	23	82
EPFU	0	0	0	1	3	12	0	14	30
MYCA	0	0	0	0	5	17	2	3	27
MYYU	0	0	0	0	0	0	0	1	1

Literature Cited

CWHC-RCSF. 2024. "Canadian Wildlife Health Cooperative - Réseau Canadien Pour La Santé de La Faune." https://www.cwhc-rcsf.ca/white_nose_syndrome_reports_and_maps.php.

Energy and Climate Solutions, Ministry of. 2024. "BC Gov News." December 9. https://news.gov.bc.ca/releases/2024ECS0048-001643.

Friedenberg, Nicholas A., and Winifred F. Frick. 2021. "Assessing Fatality Minimization for Hoary Bats Amid Continued Wind Energy Development." *Biological Conservation* 262 (October): 109309. https://doi.org/10.1016/j.biocon.2021.109309.

Harrison, Xavier A., Lynda Donaldson, Maria Eugenia Correa-Cano, et al. 2018. "A Brief Introduction to Mixed Effects Modelling and Multi-Model Inference in Ecology." *PeerJ* 6 (May): e4794. https://doi.org/10.7717/peerj.4794.

Hurtado, Camila, and Taylor Hart. 2026. *North American Bat Monitoring Program in SE Alaska – North American Bat Monitoring Program in s.e Alaska*. https://bp-sensr.github.io/SE-Alaska_NABat/NABat-AK.html.

Lile, Renee E, Ian M Abernethy, Jesse M Alston, et al. 2026. "Changes in Summer Bat Captures in the Black Hills, South Dakota, After White-Nose Syndrome Invasion." *Journal of Mammalogy*, September 28, gyag082. https://doi.org/10.1093/jmammal/gyag082.

Loeb, Susan C., Thomas J. Rodhouse, Laura E. Ellison, et al. 2015. *A Plan for the North American Bat Monitoring Program (NABat)*. SRS-GTR-208. U.S. Department of Agriculture, Forest Service, Southern Research Station. https://doi.org/10.2737/SRS-GTR-208.

MacKenzie, Darryl I., James D. Nichols, James E. Hines, Melinda G. Knutson, and Alan B. Franklin. 2003. "Estimating Site Occupancy, Colonization, and Local Extinction When a Species Is Detected Imperfectly." *Ecology* 84 (8): 2200–2207. https://doi.org/10.1890/02-3090.

Olson, Cory. n.d. "Bat Profiles." Alberta Community Bat Program. Accessed March 6, 2025. https://www.albertabats.ca/batprofiles/.

Rae, Jason, and Cori L Lausen. 2022. *North American Bat Monitoring Program in British Columbia: 2021 Data Summary and Activity Trend Analyses (2016-2021)*. Wildlife Conservation Society of Canada.

Rae, Jason, and Cori L Lausen. 2024. *North American Bat Monitoring Program in British Columbia: 2023 Data Summary and Activity Trend Analyses (2016-2023)*. Wildlife Conservation Society of Canada.

Reichert, Brian, Cori Lausen, Susan Loeb, et al. 2018a. *A Guide to Processing Bat Acoustic Data for the North American Bat Monitoring Program (NABat)*.

Reichert, Brian, Cori Lausen, Susan Loeb, et al. 2018b. *A Guide to Processing Bat Acoustic Data for the North American Bat Monitoring Program (NABat)*. Open-File Nos. 2018–1068. U.S. Geological Survey.

Slough, Brian, Cori Lausen, Brian Paterson, et al. 2022. "New Records about the Diversity, Distribution, and Seasonal Activity Patterns by Bats in Yukon and Northwestern British Columbia." *Northwestern Naturalist* 103 (August): 162–82. https://doi.org/10.1898/NWN21-10.

Solick, Donald. 2022. "Bat Acoustic Species-Pair Matrix Western US/Canada." https://www.batacousticsurveys.com/_files/ugd/a1e0ca_0ed9c3172b9145e49c50eb7a7712f909.pdf.

Solick, Donald I., Robert M. R. Barclay, Larisa Bishop-Boros, Quentin R. Hays, and Cori L. Lausen. 2020. "Distributions of Eastern and Western Red Bats in Western North America." *Western North American Naturalist* 80 (1): 90–97. https://doi.org/10.3398/064.080.0111.

Status of Endangered Wildlife in Canada, Committee on the. 2023. "Hoary Bat (Lasiurus Cinereus) Eastern Red Bat (Lasiurus Borealis) Silver-haired Bat (Lasionycteris Noctivagans): COSEWIC Assessment and Status Report 2023." https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/hoary-bat-eastern-red-bat-silver-haired-bat-2023.html.

Stratton, Christian, Kathryn M. Irvine, Katharine M. Banner, Wilson J. Wright, Cori Lausen, and Jason Rae. 2022. "Coupling Validation Effort with in Situ Bioacoustic Data Improves Estimating Relative Activity and Occupancy for Multiple Species with Cross-species Misclassifications." *Methods in Ecology and Evolution* 13 (6): 1288–303. https://doi.org/10.1111/2041-210X.13831.

Szewczak, Joe. 2018. *Acoustic Features of Western US Bats*. Humboldt State University Bat Lab.

Team, BC Bat Action. 2024. *2024-2028 BC Bat Action Plan*.

USFWS, U. S. Fish and Widlife Service. 2025. "White-Nose Syndrome." https://www.whitenosesyndrome.org/.

Zimmerling, J. Ryan, and Charles M. Francis. 2016. "Bat Mortality Due to Wind Turbines in Canada." *The Journal of Wildlife Management* 80 (8): 1360–69. https://doi.org/10.1002/jwmg.21128.