




DHARMa residuals: COTO









Levene Test for homogeneity of variance n.s.
Within−group deviation from uniformity n.s.
simulationOutput$scaledResiduals
0.00
0.25
0.50
0.75
1.00
0.0408163265306122
0.24234693877551
0.586734693877551
0.86734693877551
1
YearS (rank transformed) (catPred)



Residuals vs log(transect.length)
Quantile deviations detected (red curves)
Combined adjusted quantile test significant
DHARMa residual
1.00
0.75
0.50
0.25
0.00
0.0
0.2
0.4
0.6
0.8
1.0
log(transect.length) (rank transformed)



Residuals vs Nightly_Mean_RH
Quantile deviations detected (red curves)
Combined adjusted quantile test significant
DHARMa residual
1.00
0.75
0.50
0.25
0.00
0.0
0.2
0.4
0.6
0.8
1.0
Nightly_Mean_RH (rank transformed)



Residuals vs Nightly_Mean_Windsp
Quantile deviations detected (red curves)
Combined adjusted quantile test significant
DHARMa residual
1.00
0.75
0.50
0.25
0.00
0.0
0.2
0.4
0.6
0.8
1.0
Nightly_Mean_Windsp (rank transformed)


DHARMa nonparametric dispersion test via sd of residuals fitted vs. simulated
Frequency
0
5
10
15
20
0
1
2
3
4
Simulated values, red line = fitted model. p−value (two.sided) = 0.946


DHARMa zero−inflation test via comparison to expected zeros with simulation under H0 = fitted model
Frequency
0
50
100
150
185
190
195
Simulated values, red line = fitted model. p−value (two.sided) = 0.498


Residuals vs. time
Scaled residuals
1.0
0.8
0.6
0.4
0.2
0.0
2
4
6
8
Time
Autocorrelation
ACF
Durbin−Watson test p= 0.06822
Deviation n.s.
1.0
0.5
0.0
−0.5
−1.0
0
2
4
6
8
Lag


Levene Test for homogeneity of variance n.s.
Within−group deviation from uniformity n.s.
simulationOutput$scaledResiduals
0.00
0.25
0.50
0.75
1.00
986
6202
16426
37274
60586
86074
110378
137274
175354
212906
243562
269777
326072
grp (catPred)


Random intercept QQ: GRTS_Cell_ID (47 levels)
Sample Quantiles
Theoretical Quantiles
−1
0
1
2
3
4
5
−2
−1
0
1
2