




DHARMa residuals: LANO









Levene Test for homogeneity of variance n.s.
Within−group deviation from uniformity n.s.
simulationOutput$scaledResiduals
0.00
0.25
0.50
0.75
1.00
0.0331325301204819
0.198795180722892
0.512048192771084
0.843373493975904
1
YearS (rank transformed) (catPred)



Residuals vs log(transect.length)
No significant problems detected
DHARMa residual
1.00
0.75
0.50
0.25
0.00
0.0
0.2
0.4
0.6
0.8
1.0
log(transect.length) (rank transformed)



Residuals vs Nightly_Mean_Temp
No significant problems detected
DHARMa residual
1.00
0.75
0.50
0.25
0.00
0.0
0.2
0.4
0.6
0.8
1.0
Nightly_Mean_Temp (rank transformed)



Residuals vs Nightly_Mean_RH
No significant problems detected
DHARMa residual
0.00
0.25
0.50
0.75
1.00
0.0
0.2
0.4
0.6
0.8
1.0
Nightly_Mean_RH (rank transformed)



Residuals vs Nightly_Mean_Windsp
No significant problems detected
DHARMa residual
0.00
0.25
0.50
0.75
1.00
0.0
0.2
0.4
0.6
0.8
1.0
Nightly_Mean_Windsp (rank transformed)


DHARMa nonparametric dispersion test via sd of residuals fitted vs. simulated
Frequency
0
5
10
15
0.6
0.8
1.0
1.2
1.4
1.6
1.8
Simulated values, red line = fitted model. p−value (two.sided) = 0.098


DHARMa zero−inflation test via comparison to expected zeros with simulation under H0 = fitted model
Frequency
0
20
40
60
80
25
30
35
40
45
50
55
Simulated values, red line = fitted model. p−value (two.sided) = 1


Residuals vs. time
Scaled residuals
1.0
0.8
0.6
0.4
0.2
0.0
2
4
6
8
Time
Autocorrelation
ACF
Durbin−Watson test p= 0.11878
Deviation n.s.
1.0
0.5
0.0
−0.5
−1.0
0
2
4
6
8
Lag


Levene Test for homogeneity of variance n.s.
Within−group deviation from uniformity n.s.
simulationOutput$scaledResiduals
0.00
0.25
0.50
0.75
1.00
1194
9466
17706
38442
60586
79018
93290
127850
145578
194666
230186
265749
294832
grp (catPred)


Random intercept QQ: GRTS_Cell_ID (45 levels)
Sample Quantiles
Theoretical Quantiles
−1.0
−0.5
0.0
0.5
1.0
1.5
−2
−1
0
1
2