Free Energy Landscape Perturbation for Estimating Mutation-Induced Population Shifts from Reference Molecular Dynamics Ensembles

Tatum E. Murdock[†,‡] and Benjamin P. Brown[*,¶,‡]

†*Chemical and Physical Biology Program, Vanderbilt University, Nashville, Tennessee 37232, United States*

‡*Center for AI in Protein Dynamics, Vanderbilt University, Nashville, Tennessee 37232, United States*

¶*Department of Pharmacology, Vanderbilt University, Nashville, Tennessee 37232, United States*

E-mail: benjamin.p.brown@vanderbilt.edu

Abstract

Mutations can alter protein function by redistributing probability among conformational states, even when structural changes are modest. Directly estimating a perturbed conformational free energy landscape with molecular dynamics is expensive because each mutation requires a new converged simulation. Here, we introduce Free Energy Landscape Perturbation (FELP), a state-local reweighting framework that estimates mutation-induced population shifts from a reference ensemble. FELP partitions a reference trajectory into discrete microstates, evaluates restrained reference-to-perturbed energy differences for configurations sampled within each state, and combines spatially regularized state-local Zwanzig factors with the reference state probabilities to estimate the perturbed state distribution. FELP estimates how a perturbation changes the relative probabilities of conformational basins and reports effective-sample and overlap diagnostics that identify when the predicted shift lacks support in the sampled reference ensemble. We describe the FELP estimator, its implementation, and its performance on glycine dipeptide, chignolin, and T4 lysozyme. FELP is intended for perturbations that reshape populations within a sufficiently sampled reference landscape, providing a practical bridge between local alchemical energy evaluation and ensemble-level mechanistic interpretation.

1. Introduction

The functional repertoire of a protein is fundamentally determined by its conformational free energy landscape. Folding, stability, enzyme catalysis, signal transduction, molecular recognition, and allosteric regulation all depend on how probability is distributed over conformational states and how readily the system moves between them. Many proteins therefore function through ensembles of native-like substates, with activity controlled by changes in basin depth, transition pathways, and relative occupancy. Quantitatively describing these landscapes is central to understanding how sequence, ligand binding, protonation, and chemical modification alter biological function.

Molecular dynamics (MD) simulations provide a direct route to estimating these conformational free energy landscapes at atomistic resolution. In principle, sufficiently long trajectories recover both the relevant structural diversity and the thermodynamic probability of each state. In practice, sampling remains the cen-

tral limitation. Transitions between functional conformations can be slow, rare, and system-specific, such that converging a separate MD ensemble for every perturbation of interest is often prohibitively expensive. Parallel simulation strategies, enhanced sampling methods such as replica exchange[1] and metadynamics,[2] and Markov state models (MSMs)[3,4] have improved access to long-timescale behavior by combining many trajectories into kinetic and thermodynamic state models. However, even with these advances, each new mutation or chemical change can still demand substantial additional sampling to converge a new landscape. This expense is a principal bottleneck for systematic structure–function studies that require evaluating many perturbations against a shared conformational baseline.

To avoid the cost of full ensemble resampling, computational methods often sacrifice conformational diversity for speed. Single-structure and local relaxation methods, such as Rosetta $\Delta\Delta G$ estimates for mutation stability calculations, can test whether a particular coordinate set becomes more or less favorable after mutation, but they typically evaluate a narrow structural hypothesis.[5] Minimal relaxation avoids the cost of full resampling, yet it can convert small coordinate differences into abrupt energy changes and noisy estimates. These calculations can be useful for local stability screening, but they generally do not directly answer whether a mutation stabilizes one conformational basin over another.

Alchemical methods such as free energy perturbation (FEP) provide a formally rigorous extension of this idea. By computing free energy differences between Hamiltonians from local equilibrium samples, FEP incorporates local flexibility and avoids the abrupt energy changes of single-structure minimization.[6–11] However, practical protein mutation calculations remain structurally restricted because they are initialized from a small number of conformations and restrained to prevent structural drift. Both single-structure screening and local FEP calculations estimate the energetic effect of a perturbation in a specific structural context. Additional ensemble analysis is required to describe how that perturbation redistributes probability across a complex conformational landscape. Furthermore, exponential free energy estimators rely heavily on the assumption that the reference and target states share sufficient phase-space overlap.[12,13] Because these estimators are highly sensitive to rare and high-weight samples, inadequate overlap can produce unstable estimates that appear numerically converged but are driven by too few configurations.

Recent population-shift strategies address this ensemble problem by separating conformational sampling from perturbation evaluation. This approach originally emerged in the context of ligand binding to quantify conformational selection and induced-fit mechanisms. Implicit-ligand theory, for example, provides a rigorous analytical framework for estimating how a binding event reweights host conformations without requiring explicit simulation of the bound complex.[14–17] While traditional implicit-ligand theory typically evaluates these shifts using exact volume integrals or analytical interaction potentials over a continuous ensemble, recent methods adapt this logic to discrete kinetic models and empirical scoring. PopShift, for instance, evaluates discrete ligand docking scores against the states of a ligand-free MSM to predict binding-induced population shifts.[18] The unifying insight of these approaches is that expensive sampling can be performed once for a reference system. Once the reference state probabilities are established, the effects of many different ligands can be evaluated against the same conformational baseline. This framework moves beyond single-structure scoring by explicitly linking the energetic preference of a perturbation to the thermodynamic weight of each conformational state.

Here, we introduce Free Energy Landscape Perturbation (FELP) to apply this population-shift framework to amino acid mutations. FELP combines a sampled reference ensemble, discrete state models, restrained local perturbation evaluation, and FEP-style reweighting. The method partitions a reference ensemble into discrete microstates, evaluates the energetic effect of a mutation within each state, and reweights the reference probabilities to estimate how the mutation redistributes probability over the pre-sampled landscape. By treating a global population shift

as a collection of tractable local reweighting problems, FELP bridges the gap between single-structure scoring and full-ensemble resampling. To address the inherent phase-space overlap limitations of exponential estimators, FELP explicitly audits its predictions using effective sample size and weight-concentration diagnostics.[19,20] These diagnostics ensure that the method not only predicts how a mutation reshapes a landscape but also identifies when a proposed perturbation requires conformations that were not adequately sampled by the reference ensemble.

2. Theory

Consider a reference system with potential energy $U_0(x)$ and a perturbed system with potential energy $U_1(x)$, where x denotes a molecular configuration and $\beta = (k_B T)^{-1}$. The Boltzmann distribution maps these potential energy functions to continuous probability densities over configurations. FELP starts from a sampled reference ensemble, such as an MD trajectory or another state model with a meaningful reference prior, that represents configurations drawn from the reference distribution $p_0(x) \propto \exp[-\beta U_0(x)]$. FELP partitions this configuration space into discrete microstates $i \in \{1, \ldots, N\}$ using a chosen collective variable or feature representation. Let $\chi_i(x)$ be the indicator function for state i. The restricted partition function for state i under Hamiltonian a is

$$Z_a(i) = \int \chi_i(x) \exp[-\beta U_a(x)]\, dx, \qquad (1)$$

and the corresponding thermodynamic probability of the microstate is $\pi_a(i) = Z_a(i)/\sum_k Z_a(k)$. For the reference system, this is equivalent to integrating the continuous reference density over the microstate to obtain the reference prior $\pi_0(i)$. FELP aims to estimate the perturbed microstate probabilities $\pi_1(i)$ using only samples drawn from the reference ensemble **(Figure 1)**.

To bridge the reference and perturbed ensembles, FELP relies on the principle that global conformational shifts can be decomposed into local reweighting problems. Within a specific microstate, the ratio of restricted partition functions follows the two-state Zwanzig identity,

$$\frac{Z_1(i)}{Z_0(i)} = \langle \exp[-\beta \Delta U(x)] \rangle_{0,i}\,, \qquad (2)$$

where $\Delta U(x) = U_1(x) - U_0(x)$ and $\langle \cdot \rangle_{0,i}$ denotes an ensemble average over reference configurations conditioned on membership in state i. Normalizing the restricted partition functions over all microstates yields the central FELP estimator:

$$\pi_1(i) = \frac{\pi_0(i)\, \langle \exp[-\beta \Delta U(x)] \rangle_{0,i}}{\sum_k \pi_0(k)\, \langle \exp[-\beta \Delta U(x)] \rangle_{0,k}}. \qquad (3)$$

Equation 3 preserves the reference state decomposition and updates the probability of each state according to the perturbation's energetic effect on configurations sampled within that specific state.

The central hypothesis behind this formulation is that discretization creates domains of local phase-space overlap. By conditioning the exponential average on microstate membership, FELP assumes that reference configurations assigned to state i adequately represent the structural ensemble of the perturbed system within that same state. FELP evaluates the perturbation directly from the reference ensemble without simulating the target state or intermediate alchemical windows. If a mutation is unfavorable in one conformational basin but favorable in another, the state-local reweighting factors will predict a population shift between those basins.

In a finite trajectory, state i contains evaluated configurations x_{ij} for $j = 1, \ldots, n_i$. FELP estimates the local Zwanzig factor by computing

$$f_i = \frac{1}{n_i} \sum_{j=1}^{n_i} w_{ij}, \qquad w_{ij} = \exp[-\beta(\Delta U_{ij} - s)], \qquad (4)$$

where $s = \min(\Delta U)$ over all frames included in reweighting is a uniform stability shift that prevents numerical overflow. Because this shift is applied globally, it cancels during normalization. The sample estimator of eq 3 substitutes the empirical reference prior $\hat{\pi}_0(i)$ and the finite-sample




Figure 1. Free Energy Landscape Perturbation workflow. FELP begins from a sampled reference ensemble and a microstate atlas, assigns configurations to obtain the reference prior $\hat{\pi}_0(i)$, and evaluates symmetric reference and perturbed branches under reference-frame restraints. The default OpenMM reweighting Hamiltonian uses ff14SB-only side-chain parameters with GBNeck2 implicit solvent to compute $\Delta U = U_1 - U_0$ for each evaluated configuration. State-local Zwanzig factors are combined with $\hat{\pi}_0(i)$ to estimate $\hat{\pi}_1(i)$ and $F_1(i)$; optional neighbor pooling can regularize noisy local estimates before normalization.

factor f_i for their theoretical counterparts:

$$\hat{\pi}_1(i) = \frac{\hat{\pi}_0(i) f_i}{\sum_k \hat{\pi}_0(k) f_k}. \tag{5}$$

The corresponding relative free energy of each state is

$$F_1(i) = -k_B T \ln \hat{\pi}_1(i) + C, \tag{6}$$

where C is chosen so that the lowest finite state free energy is set to zero.

Now, for finite samples, f_i can be dominated by a small number of high-weight frames even when n_i is large. The local effective sample size,

$$\mathrm{ESS}_i = \frac{\left(\sum_j w_{ij}\right)^2}{\sum_j w_{ij}^2}, \tag{7}$$

quantifies this weight degeneracy.[21,22] Low ESS_i indicates that the local Zwanzig factor depends on too few configurations to be reliable.

Configuration space overlap requirements in unidirectional reweighting are fundamentally asymmetric. The reference ensemble must encompass the relevant configurations of the target ensemble, but the reverse is not true. It is expected that a broad reference trajectory will explore conformational states that become energetically forbidden or highly unfavorable upon perturbation. In these states, the local perturbation factor may have poor statistical support (low ESS_i), but this lack of support is inconsequential if the target probability $\hat{\pi}_1(i)$ in those states is vanishingly small. The reweighted landscape remains credible as long as the states that actually carry significant target probability are well-supported by the reference samples. However, because it is rarely known *a priori* whether the reference trajectory is sufficiently broad to cover all energetically relevant regions of the target ensemble, it is critical to explicitly quantify when the estimate relies on unsupported configurations. FELP therefore summarizes the validity of the reweighting operation by measuring the target probability mass assigned to unreliable states,

$$\mathcal{U} = \sum_i \hat{\pi}_1(i)\, \mathbf{1}\{\mathrm{ESS}_i < \epsilon_i\}, \tag{8}$$

where ϵ_i is an effective-sample floor for state i. Small $\mathcal{U}$ means the predicted landscape rests on states with adequate local sampling; large $\mathcal{U}$ is a direct reason for caution.

A separate practical issue arises from discretization. The potential energy functions U_0 and U_1 are continuous, so the true restricted partition functions of neighboring microstates should vary smoothly within a basin. Treating every microstate independently forces states with low ESS_i to produce local factors that fluctuate sharply relative to their neighbors. FELP can optionally regularize these estimates by pooling information across adjacent states in the feature-space topology. For state i, the pooled factor over eligible neighboring states $\mathcal{N}(i)$ is

$$\bar{f}_i = \frac{\sum_{j\in\mathcal{N}(i)} \mathrm{ESS}_j\, f_j}{\sum_{j\in\mathcal{N}(i)} \mathrm{ESS}_j}, \tag{9}$$

with $\bar{f}_i = f_i$ when no eligible neighbors are available. The blended factor used for reweighting is

$$f_i^{\text{blend}} = \alpha_i f_i + (1-\alpha_i)\bar{f}_i, \qquad \alpha_i = \frac{\text{ESS}_i}{\text{ESS}_i + \tau}. \tag{10}$$

States with high ESS_i retain their local factors ($\alpha_i \approx 1$); states with low ESS_i borrow strength from better-sampled neighbors. eq 5 can be evaluated with f_i^{blend} in place of the raw f_i. The blending rule is therefore a shrinkage estimator wherein ESS_i determines how strongly the state-local estimate is trusted, while τ sets the effective-sample scale at which neighborhood information is discounted. This preserves well-sampled local perturbation factors while damping isolated fluctuations in poorly sampled states.

When τ is not specified, FELP selects it by an L-curve criterion,[23] balancing fidelity to the state-local estimates against smoothness over neighboring states. The fidelity term is evaluated in log-factor space,

$$\mathcal{M}(\tau) = \sum_i \text{ESS}_i \left[\ln f_i - \ln f_i^{\text{blend}}(\tau)\right]^2, \tag{11}$$

so that well-sampled states anchor the fit and deviations are measured on the scale of local free energy corrections. The corresponding graph-smoothness penalty is

$$\mathcal{R}(\tau) = \frac{1}{2}\sum_i \sum_{j\in\mathcal{N}(i)} \left[\ln f_i^{\text{blend}}(\tau) - \ln f_j^{\text{blend}}(\tau)\right]^2. \tag{12}$$

The selected τ is the corner of the $\mathcal{M}$–$\mathcal{R}$ curve in log–log space, defined as the point farthest from the chord connecting the two extremes. Importantly, while this regularization stabilizes noisy local factors on a discretized landscape, it does not compensate for reference conformations that were not sampled; $\mathcal{U}$ remains the primary validity criterion.

3. Results

3.1. Glycine dipeptide

We first evaluated FELP on capped dipeptides, where the relevant conformational changes are visible directly in backbone Ramachandran (ϕ/ψ) space. Glycine provides the most permissive reference landscape of the canonical amino acids, making it a useful test of whether reweighting can recover landscapes that occupy a subset of the conformations sampled by the reference **(Figure 2A)**.

As a numerical control, GLY→GLY returns the reference landscape, with no visible redistribution and negligible Kullback–Leibler (KL) divergence from the independently analyzed glycine MD landscape **(Figure 2B–D)**. This confirms that the discretization, state assignment, and reweighting steps do not introduce appreciable structure when the perturbation energy is negligible.

We next considered GLY→ALA. Addition of the alanine methyl group removes much of the positive-ϕ population sampled by glycine while preserving the dominant α_R and β basins. The FELP-reweighted landscape captures this restriction and closely matches the independently sampled alanine MD surface **(Figure 2E–G)**. The remaining KL signal is localized and small, consistent with modest differences in basin depths rather than failure to identify the accessible regions of Ramachandran space.

GLY→PRO is a more pronounced perturbation because the pyrrolidine ring constrains ϕ near $-60°$ and narrows the allowed ψ range. Even in this case, FELP concentrates probability into the proline-like region and removes the broad glycine accessibilities **(Figure 2H–J)**. The KL differences are slightly larger than for alanine, as expected for the stronger perturbation, but again they reflect imperfect relative weighting among the same basins observed by MD.

These examples highlight the distinction between sampling and energetic accuracy. Reference MD was generated with explicit-solvent ff19SB/OPC, whereas FELP evaluates perturbation energies with an implicit-solvent

ff14SBonlysc/GBNeck2 model. The comparison therefore includes both the statistical reweighting problem and a force field/solvent mismatch. Despite this approximation, the topology of the alanine and proline target landscapes is recovered when the target distribution is a restriction of the glycine reference ensemble.

The reverse direction exposes the expected failure mode. When alanine is used as the reference, ALA→GLY requires probability in broad positive-ϕ regions that are absent or poorly represented in the alanine trajectory **(Figure 3B–D)**. FELP cannot create conformations that were not sampled; it can only redistribute probability over the alanine reference states. The reweighted surface therefore remains confined to the alanine-sampled region rather than recovering the full glycine landscape. In contrast, ALA→ALA returns the identity **(Figure 3E–G)**, and ALA→PRO succeeds because the proline landscape lies within the conformations sampled by alanine **(Figure 3H–J)**. Notably, however, ALA→PRO shows larger KL differences from direct proline MD than GLY→PRO, consistent with glycine providing the broader and better-sampled reference for this restricted target landscape.

3.2. Overlap diagnostics quantify when reweighting fails

We next asked whether these success and failure modes could be identified from the reweighting statistics alone, without comparing against an independently sampled target landscape. For each GLY, ALA, and PRO reference ensemble, we reweighted to all 20 canonical amino acids and computed two diagnostics: (i) the target probability assigned to low-ESS states, nominally defined here as microstates with $\mathrm{ESS}_i < 10$, and (ii) the maximum normalized frame weight **(Figure 4)**.

The low-ESS mass remains small for all glycine-reference perturbations, consistent with the broad Ramachandran sampling of glycine **(Figure 4A)**. The alanine reference shows the same pattern for most targets but fails sharply for ALA→GLY, where nearly all target probability is assigned to low-ESS states. This diagnostic identifies the same missing-conformation failure observed visually in **(Figure 3B–D)**. By contrast, the proline reference produces elevated low-ESS mass for most targets, reflecting the difficulty of using a conformationally restricted reference to predict broader amino-acid landscapes. ¡why is PRO –¿ GLY so good in the low-ESS mass estimate??¿

The maximum normalized frame weight provides a complementary view of the same overlap problem **(Figure 4B)**. Glycine-reference perturbations are distributed over many frames, with maximum weights generally near 10^{-4}–10^{-3}. Alanine-reference perturbations are also well-behaved except for targets such as glycine, where a small number of tail frames carry a disproportionate fraction of the reweighted probability. Proline-reference calculations show the largest frame-weight concentrations, in several cases approaching single-frame control of the local estimator. ¡again, why is GLY the best other than PRO? is there a bug? was this plotted correctly? The relatively low frame-weight concentration for PRO→GLY should not be interpreted as recovery of the glycine landscape; deleting the proline constraint is energetically smooth within the conformations proline already sampled, even though the broad glycine-only regions remain unavailable to the estimator???¿

These data are consistent with the central overlap criterion of FELP: reweighting is reliable when the target landscape redistributes probability among conformations already sampled by the reference, and unreliable when the target requires conformations that the reference rarely or never visits. Importantly, both diagnostics are computed from the reference calculation itself and can therefore be used prospectively to decide whether a predicted landscape should be interpreted or whether additional sampling is required.

3.3. Chignolin

We next tested FELP on chignolin, a 10-residue β-hairpin miniprotein whose folding equilibrium depends on coordinated tertiary contacts rather than a single local backbone angle.[24] Wild-type chignolin is only marginally stable ($T_\mathrm{m} \approx 315$ K).




Figure 2. FELP reweighting from a glycine dipeptide reference. (A) Reference MD free energy landscape in ϕ/ψ space. For GLY→GLY (B–D), GLY→ALA (E–G), and GLY→PRO (H–J): FELP-reweighted landscape (left), independently sampled MD landscape (center), and KL divergence (right).

In contrast, the G1Y/G10Y variant of chignolin, CLN025, was designed to increase thermodynamic stability of the folded conformation ($T_{\mathrm{m}} \approx 343$ K) while retaining the same hairpin topology.[25] Circular dichroism and NMR place chignolin near half-folded at physiologically relevant temperatures: a two-state fit anchored to the experimental melting data gives $p_{\mathrm{fold}} \approx 0.54$ for wild type at 310 K, and NMR reports roughly 60% folded for chignolin and 90% folded for CLN025 at 300 K.[24,26] We projected the landscape onto two Cα distances: Cα_1–Cα_{10}, which reports on termini proximity, and Cα_3–Cα_6, which reports on turn formation. Together, these coordinates separate the compact folded basin from partially folded and extended conformations.

Our 5 μs ff19SB/OPC reference trajectories substantially underfold both variants relative to experiment. Wild-type chignolin has $p_{\mathrm{fold}} \approx$ 0.11–0.12 at 310 K, and the MD landscape spans both the compact folded basin and a broad set of extended conformations **(Figure 5A)**. This underfolding is consistent with recent benchmark simulations: Fischer *et al.* report $16 \pm 1\%$ natively folded for chignolin in substantially longer ff19SB/OPC trajectories analyzed with Markov state models.[26] CLN025 shifts probability toward the folded basin while still sampling extended conformations **(Figure 5B)**, but it too remains less folded in MD ($31 \pm 12\%$ natively folded at 300 K in the benchmark study) than in NMR.[26] Reweighting the CLN025 reference to wild type recovers the compact basin and redistributes probability toward extended conformations, although the predicted extended ensemble is sparser and less continuous than in the directly sampled wild-type MD landscape **(Figure 5C)**.

The reverse calculation, WT→CLN025, starts from the broader wild-type reference. In this direction FELP strongly enriches the folded basin and depletes much of the extended landscape, consistent with the stabilizing role of the CLN025 mutations **(Figure 5D)**. The remaining extended density indicates that the reweighted landscape is not a simple binary folded/unfolded switch, but the dominant redistribution is toward compact conformations.

We quantified these shifts by defining the




Figure 3. FELP reweighting from an alanine dipeptide reference. Layout as in Figure 2. ALA→GLY (B–D) fails to recover the broad glycine landscape because the required conformations are absent from the reference trajectory. ALA→ALA (E–G) returns the identity. ALA→PRO (H–J) succeeds because the proline landscape lies within the conformations sampled by alanine.

folded basin as Cα_1–Cα_{10} < 8 Å and Cα_3–Cα_6 < 7 Å (dashed boxes in **Figure 5A–D**). Despite the absolute underfolding of the reference MD ensembles, the simulations recover the experimental sign and magnitude of the CLN025 stabilization: the two-state experimental estimate gives $\Delta\Delta G_{\mathrm{fold}} \approx -1.0$ kcal mol^{-1}, and the MD simulations give a similar value **(Figure 5E, F)**. FELP recovers the sign of this stabilization but overestimates its magnitude, predicting CLN025 to be too stable by roughly 1.5 kcal mol^{-1}. Interpreting this result requires separating force-field limitations in the reference ensemble from the additional approximation introduced by implicit-solvent mutation scoring. The former already compresses folded populations well below experiment in ff19SB/OPC, whereas the latter likely amplifies the terminal tyrosine stabilization because solvation contributions dominate folding thermodynamics in this small β-hairpin. Nevertheless, FELP recovers the direction of the experimental folding shift from reweighting alone.

3.4. T4 lysozyme

Finally, we applied FELP to T4 lysozyme, a larger protein system in which mutation alters the conformational landscape through hydrophobic core packing. The L99A substitution creates an internal cavity of approximately 150 Å^3 and is among the most extensively characterized cavity-creating mutations in structural biology, with conformational exchange between ground and excited states observed by NMR relaxation dispersion and crystallography.[27,28] We used L99A as the reference ensemble and reweighted toward wild type (A99L), projecting the result onto ψ_{114}, a Phe114 backbone dihedral reporting on helix F dynamics near the cavity.

The L99A reference distribution is dominated by a basin near $\psi_{114} \approx 55°$, but it also contains lower-probability excursions near $\psi_{114} \approx -40°$ and near the periodic boundary **(Figure 6A)**. Reweighting toward wild type further concentrates probability into the $\psi_{114} \approx 55°$ basin and strongly suppresses the secondary L99A populations **(Figure 6B)**. The corresponding free energy projection shows the same trend: the wild-type landscape retains the same global min-




Figure 4. Overlap diagnostics across reference ensembles. (A) Fraction of target probability mass assigned to low-ESS states ($\mathrm{ESS}_i < 10$; $\mathcal{U}$, eq 8) for all canonical amino acid targets from GLY, ALA, and PRO references. (B) Maximum normalized frame weight across the same perturbations. Error bars indicate bootstrap uncertainty across resampled evaluated frames.

imum but raises the alternative regions of ψ_{114} space relative to L99A **(Figure 6C)**.

This behavior is physically consistent with cavity filling. Removing Leu99 creates space for alternative helix F packing arrangements, whereas restoring the leucine side chain favors a narrower set of packed conformations. The T4 calculation therefore tests a different regime than the dipeptide failures: the reference mutant samples the minor conformations that are subsequently disfavored by the wild-type perturbation. Under these conditions, FELP predicts a redistribution among observed conformations rather than the appearance of a new basin outside the reference trajectory.

4. Discussion

Understanding how mutations in proteins alter their conformational free energy landscapes is central to interpreting allosteric mechanisms, predicting resistance phenotypes, and guiding protein engineering. However, converging a separate MD ensemble for every mutation of interest is prohibitively expensive. FELP addresses this by asking a narrower question: given a well-sampled reference landscape, how does a chemical perturbation redistribute probability among the states already observed?

FELP belongs to an emerging class of population-shift methods that reuse a reference conformational ensemble rather than resampling each perturbation from scratch. ¡implicit ligand theory? something else before PopShift? or is this it?¿ PopShift similarly reweights MSM states using empirical docking scores to predict ligand-induced population shifts.[18] FELP adapts this logic to amino-acid mutations by replacing empirical state scores with restrained FEP-style energy differences and by reporting overlap diagnostics alongside the predicted population shift. ¡need a good transition, or we just get rid of this; Abella did a coarser macrostate reweighting, similar to our folded vs. unfolded chignolin study¿, Abella et al. applied an analogous strategy to evaluate peptide mutations from a wild-type Markov state model, reweighting metastable macrostate populations to predict binding shifts.[29]

Our dipeptide benchmarks demonstrate that




Figure 5. FELP predicts mutation-induced folding shifts in chignolin. (A, B) MD free energy landscapes for wild-type chignolin and CLN025 projected onto Cα_1–Cα_{10} and Cα_3–Cα_6 distances. Dashed boxes indicate the folded basin. (C) FELP CLN025→WT reweighted landscape. (D) FELP WT→CLN025 reweighted landscape. (E) Folding free energies ΔG_{fold} from experiment, MD, and FELP. (F) $\Delta\Delta G_{\text{fold}}$ (CLN025 − WT).

FELP reliably recovers the perturbed landscape when the target conformational ensemble lies within the region sampled by the reference **(Figures 2, 3)**. The self-perturbation controls confirm that the pipeline is stable, and the GLY→ALA and GLY→PRO results show that FELP can correctly predict the restriction of accessible Ramachandran space upon side-chain addition. The reverse transformation, ALA→GLY, fails precisely because the target requires conformational basins absent from the reference ensemble. This failure is expected from the asymmetric overlap requirements of unidirectional reweighting and is correctly identified by the $\mathcal{U}$ diagnostic without access to a target simulation **(Figure 4)**.

The chignolin benchmark extends these overlap tests to a system where the conformational equilibrium depends on coordinated intramolecular contacts. Here the reference MD ensemble itself is not quantitatively faithful to experiment. When simulated with ff19SB/OPC, our MD simulations underfold both chignolin and CLN025 relative to NMR and calorimetry, consistent with recent long-timescale benchmark simulations.[26] FELP nevertheless captures the experimental direction of the CLN025 stabilization, though it overestimates the magnitude **(Figure 5F)**. This case therefore illustrates a distinct limitation from the dipeptide benchmarks: reweighting can remain informative for relative population shifts even when the reference force field misestimates absolute basin depths, provided the relevant conformations are sampled and the perturbation is evaluated consistently.

A related practical limitation arises from the solvent model mismatch. Reference MD simulations in our benchmarks were performed in explicit OPC water, whereas FELP evaluates ΔU using implicit GBNeck2 solvent on stripped protein coordinates. This choice avoids the need




Figure 6. FELP predicts helix F conformational redistribution upon cavity filling in T4 lysozyme. (A) Reference L99A probability distribution over ψ_{114}. (B) FELP-reweighted wild-type (L99A→WT) distribution. (C) Free energy landscapes for L99A (purple) and FELP-reweighted wild type (red).

for explicit solvent relaxation at each evaluated frame, which would substantially increase computational cost and reduce the practical advantage of the reweighting approach. However, it introduces a systematic approximation: the ΔU values reflect the mutation's effect under implicit solvation rather than in the explicit-solvent environment of the reference trajectory. For small perturbations in well-packed protein cores, this mismatch may be modest; for perturbations that change solvent exposure or disrupt hydrogen-bonding networks, the implicit-solvent ΔU may diverge from the explicit-solvent value. The overestimation of $\Delta\Delta G_{\text{fold}}$ in chignolin **(Figure 5F)** is consistent with this limitation, as solvation effects contribute directly to the folding thermodynamics of a 10-residue miniprotein. For membrane proteins, this issue is compounded: evaluating stripped coordinates in bulk implicit solvent discards protein–lipid interactions, asymmetric electrostatics, and hydrophobic mismatch effects that were present in the explicit bilayer simulation. Extending FELP to mutations that engage lipids will therefore require either explicit-environment perturbation evaluation or benchmarked approximations that preserve the relevant membrane contributions.

The local Zwanzig estimator is inherited from FEP and carries the same phase-space overlap requirements as any exponential reweighting calculation. The global landscape estimate is obtained by applying these local perturbation factors across a reference state model. Our diagnostics (ESS, maximum frame weight, $\mathcal{U}$) formalize this requirement as a computable criterion that accompanies every prediction. When diagnostics are favorable, FELP can estimate mutation-induced population shifts at far lower cost than independently converging each perturbed ensemble. When diagnostics are unfavorable, the method provides direct evidence that the reference ensemble did not adequately sample the conformations needed for the proposed perturbation and that new sampling is required.

The use of a unidirectional exponential average is not the only possible estimator for this state-local problem. Bidirectional estimators such as BAR or multistate estimators such as MBAR can reduce variance when samples from both end states, or from intermediate alchemical states, are available.[8,10] Nonequilibrium switching or staged perturbation paths could also improve overlap within difficult microstates. These alternatives would trade additional simulation or energy-evaluation cost for reduced estimator variance. FELP uses the unidirectional form because it preserves the central screening use case: many perturbations can be evaluated from a single reference ensemble without generating target trajectories. Future implementations could introduce intermediate-state evaluation selectively for microstates flagged by the overlap diagnostics.

The most important conceptual limitation is that FELP estimates the perturbed distribution over the reference state space. It cannot discover conformational basins that the reference trajectory did not visit. In the present implementation, energies are computed after local mutation and restrained relaxation without extended equilibration or explicit treatment of large-scale solvent reorganization. FELP is therefore best understood as an estimator of how a perturbation biases existing conformations, not as a substitute for free energy calculations in cases where the mutation induces substantial structural reorganization beyond the scope of the reference ensemble.

These limitations define FELP's intended utility. A predicted redistribution is most useful when it occurs on a functional free energy surface, retains low unreliable target probability mass, and is supported by distributed frame weights. Under those conditions, FELP can prioritize mutation-induced population shifts for mechanistic follow-up at substantially lower cost than direct resimulation of every variant. Conversely, high $\mathcal{U}$, concentrated frame weights, or known environmental contributions that are absent from the perturbation model provide direct evidence that the result should be treated as a sampling or model-fidelity problem rather than as a mechanistic conclusion. Future development should address the solvent mismatch through explicit-solvent evaluation protocols or hybrid strategies that reintroduce solvent contributions selectively, and broader benchmarks on proteins with experimentally characterized conformational equilibria will further clarify the practical scope of the reweighting approximation.

5. Methods

5.1. FELP workflow

The FELP workflow has four stages **(Figure 1)**. First, a sampled reference ensemble is obtained for the system expected to contain the relevant conformational ensemble. In the present benchmarks this ensemble comes from explicit solvent MD simulations, but in general the requirement is a meaningful prior over reference states. The sampled configurations are summarized into microstates defined in a chosen collective variable space, such as a uniform ϕ/ψ grid, distances or other user-specified coordinates in an explicit JSON/YAML state file, or states imported from an exported MSM bundle.

Second, each reference frame is assigned to a microstate. When raw-count priors are used, $\hat{\pi}_0(i)$ is the empirical fraction of assigned reference frames in state i. When a state file or MSM bundle contains trusted reference probabilities, FELP can instead normalize those loaded priors. For the purposes of our benchmark systems in this study, we made use of raw counts from our MD simulations as our priors.

Third, FELP evaluates selected configurations within each state. The purpose of this stage is to estimate how a mutation changes the energy of each reference frame while keeping the comparison local to the microstate in which that frame was observed. For every frame, FELP prepares two branches from the same starting pose: the reference branch enforces the original residue identity, and the perturbed branch introduces the target amino acid(s). We measure the configured collective variables in the reference pose and convert those values into restraint targets. Strong harmonic restraints in the CV definition space are applied identically to both branches, so that each evaluated structure remains pinned to the assigned conformational coordinates while the mutation is introduced and scored.

Local structure preparation was performed with PyRosetta Cartesian FastRelax using the constrained ref2015 score function. Backbone-dihedral CVs are enforced as Rosetta dihedral constraints and distance CVs as atom-pair constraints with narrow harmonic widths. In addition to these CV restraints, FELP exposes separate controls over which internal degrees of freedom may relax. By default, FastRelax was run for five cycles with Cartesian minimization, rigid-body jumps disabled, and design disabled. Side-chain repacking is restricted to a mobile residue set selected by the mobility policy; all other side chains are held fixed through the task factory. Unless a benchmark specified otherwise,

only the mutated residue side chain is mobile. These mobility settings determine how much additional relaxation is permitted around the pinned CV coordinates and therefore trade structural flexibility for computational cost; they do not replace the CV restraints themselves.

After Rosetta preparation, both branches are converted for evaluation in OpenMM with the ff14SB sidechain-only protein force field and GB-Neck2 implicit solvent. The base CV restraints are re-imposed as harmonic torsion and distance potentials, atoms outside the mobile set are fixed by zero mass, and each structure is energy-minimized until convergence. FELP records the energy difference $\Delta U_{ij} = U_1(x_{ij}) - U_0(x_{ij})$ for each evaluated frame and excludes failed or non-finite frames from reweighting.

Fourth, FELP aggregates frame-level Boltzmann factors into state-level reweighting factors. Perturbed probabilities follow eq 5, with local factors f_i computed from the in-state average of $w_{ij} = \exp[-\beta(\Delta U_{ij} - s)]$. To regularize noisy local estimates before normalization, we applied neighbor pooling across cardinal directions on the assignment grid, with the blending strength τ selected by the L-curve criterion described in **Theory**.

Dipeptide FELP settings. Dipeptide mutations used residue 2 as the perturbation site and a 45×45 ϕ/ψ grid for assignment. The ϕ and ψ angles of residue 2 were restrained in both Rosetta and OpenMM using a force constant of 100,000. No distance restraints or extra Rosetta-only backbone restraints were used. The default target-only mobility policy was used, so only the target side chain was repacked and minimized.

Chignolin FELP settings. Chignolin CLN025→WT calculations mutated residues 1 and 10 simultaneously (Y1G/Y10G) and used automatic microstates built from two distance CVs: Cα_1–Cα_{10} and Cα_3–Cα_6. These distances were the base restraints in both stages. In Rosetta they were applied as atom-pair harmonic constraints with $\sigma = 0.001$ Å; in OpenMM they were applied as harmonic bond restraints with force constant 1,000,000. The Rosetta preparation stage also included Rosetta-only ϕ/ψ restraints on residues 1, 2, 9, and 10, using force constant 25,000 and a 15° tolerance for validation. Mobility used a sequence window of one residue around each mutation site with backbone and side-chain motion enabled, so residues 1–2 and 9–10 were mobile. An identical protocol was used for the WT→CLN025 (G1Y/G10Y) calculation.

T4 lysozyme FELP settings. T4 lysozyme A99L calculations used automatic microstates along ψ_{114}. The ψ_{114} dihedral was the base restraint in both Rosetta and OpenMM, with force constant 10,000 during Rosetta preparation and 1,000,000 during OpenMM minimization. Rosetta preparation also applied Rosetta-only ϕ/ψ restraints to residue 99 with force constant 25,000 and a 15° validation tolerance. The mobile set was limited to the mutated residue, so only the residue 99 sidechain was mobile.

5.2. Reference simulations and trajectory processing

Molecular dynamics simulations. Except where specified otherwise below, reference ensembles were generated with explicit-solvent molecular dynamics using AMBER 24[30] and the GPU-accelerated `pmemd.cuda` engine. Proteins were parameterized with the ff19SB force field[31] and solvated in orthorhombic periodic boxes of OPC water.[32] Systems were neutralized with Na^+ or Cl^- counterions using the monovalent ion parameters of Joung and Cheatham.[33] Simulations used periodic boundary conditions, the particle mesh Ewald method for long-range electrostatics with a 12.0 Å real-space cutoff, and bond constraints on hydrogen-containing bonds.[34] Hydrogen mass repartitioning (HMR) was applied to non-solvent atoms with ParmEd. We sequentially minimized solvent, solute, and the full system with 500 steps of steepest-descent minimization followed by 2,000 steps of conjugate-gradient minimization in each stage. Systems were heated from 10 K to 100 K in the canonical (NVT) ensemble for 1.0 ns and then to 310 K in the isothermal–isobaric (NPT) ensemble at 1 bar using a Monte Carlo barostat and isotropic volume scaling over an additional 1.0 ns. These two stages of heating were repeated as needed until the system was stable at 1 bar and 310 K. Production used a Langevin

thermostat with collision frequency $\gamma = 1.0$ ps^{-1} at 310 K and a 4.0 fs timestep. Trajectory snapshots were saved every 10 ps.

Dipeptides. Capped dipeptide structures (ACE–X–NME) were built with `tleap`. Each dipeptide was solvated with a 20 Å padding of OPC water. Production simulations of 2500 ns were performed for each dipeptide.

Chignolin. Initial structures for wild-type chignolin and CLN025 were built in an extended conformation from sequence using `tleap`. The wild-type sequence was specified as `NGLY TYR ASP PRO GLU THR GLY THR TRP CGLY`. CLN025 was built similarly using the corresponding G1Y/G10Y sequence. Each system was solvated with a 15 Å padding of OPC water. Reference trajectories were generated using the general MD protocol above. Production simulations of 5000 ns were performed for each system.

T4 lysozyme L99A. Five X-ray structures capturing diverse starting states of T4 L99A lysozyme (PDB IDs: 4W52, 4W53, 4W54, 4W57, 4W59) were obtained and prepared by extracting chain A, retaining the first model, resolving non-canonical modified residues, and removing experimental artifacts. Systems were prepared with `tleap` using 12 Å solvent padding. Production simulations were carried out for 5000 ns from each starting state.

5.3. Diagnostics and quality control

Before state-level factors f_i are computed, FELP applies optional by-state quality control to the in-state Boltzmann weights. For each microstate, in-state weights are built from ΔU_{ij}, and the upper-tail Pareto shape k of the weight distribution is estimated with a PSIS-style rule. A state is flagged when k exceeds a sample-size-dependent reliability threshold. When frame exclusion is enabled, flagged states enter an iterative loop: the highest-weight frame is removed, diagnostics are recomputed, and the loop continues until the state is no longer flagged or the configured per-state exclusion cap is reached. Local Zwanzig factors and ESS_i are then computed from the remaining frames only.

Validity is summarized using eq 8 and its implementation extension,

$$\mathcal{U} = \sum_i \hat{\pi}_1(i)\, \mathbf{1}\{\mathrm{ESS}_i < \epsilon_i \text{ or } q_i = 1\}, \quad (13)$$

where $\epsilon_i = \max(5,\, 0.1\, n_i)$ with n_i the number of evaluated frames in state i, and $q_i = 1$ when by-state diagnostics identify dominant-frame control of the local average (including states that remain flagged after frame exclusion). The run summary also records per-state ESS_i, maximum normalized frame weights, and the target probability carried by QC-flagged states.

6. Code and Data Availability

The FELP source code and associated sample runtime scripts are freely available under an MIT license at `https://github.com/bpbrownlab/felp`.

7. Author Contributions

B.P.B. conceived the study. T.E.M. and B.P.B wrote the code, designed the experiments, performed the simulations, conducted the analyses, interpreted the results, and wrote the manuscript.

8. Disclosures

B.P.B is an advisor for Septagen Pharmaceuticals.

Acknowledgments

This work was supported by the NIH (DP1DA058349). The authors thank members of the Center for AI in Protein Dynamics and the Department of Pharmacology at Vanderbilt University for helpful discussions.

References

(1) Sugita, Y.; Okamoto, Y. Replica-exchange molecular dynamics method for protein folding. *Chemical Physics Letters* **1999**, *314*, 141–151.

(2) Laio, A.; Parrinello, M. Escaping free-energy minima. *Proceedings of the National Academy of Sciences* **2002**, *99*, 12562–12566.

(3) Noé, F.; Prinz, J.-H. Markov Models for Biomolecular Conformational Dynamics. *Current Opinion in Structural Biology* **2011**, *21*, 562–572.

(4) Husic, B. E.; Pande, V. S. Markov State Models: From an Art to a Science. *Journal of the American Chemical Society* **2018**, *140*, 2386–2396.

(5) Kellogg, E. H.; Leaver-Fay, A.; Baker, D. Role of Conformational Sampling in Computing Mutation-Induced Changes in Protein Structure and Stability. *Proteins: Structure, Function, and Bioinformatics* **2011**, *79*, 830–838.

(6) Zwanzig, R. W. High-Temperature Equation of State by a Perturbation Method. I. Nonpolar Gases. *The Journal of Chemical Physics* **1954**, *22*, 1420–1426.

(7) Lu, N.; Kofke, D. A. Accuracy of Free-Energy Perturbation Calculations in Molecular Simulation. I. Modeling. *The Journal of Chemical Physics* **2001**, *114*, 7303–7311.

(8) Shirts, M. R.; Pande, V. S. Comparison of Efficiency and Bias of Free Energies Computed by Exponential Averaging, the Bennett Acceptance Ratio, and Thermodynamic Integration. *The Journal of Chemical Physics* **2005**, *122*, 144107.

(9) Pohorille, A.; Jarzynski, C.; Chipot, C. Good Practices in Free-Energy Calculations. *The Journal of Physical Chemistry B* **2010**, *114*, 10235–10253.

(10) Shirts, M. R.; Chodera, J. D. Statistically Optimal Analysis of Samples from Multiple Equilibrium States. *The Journal of Chemical Physics* **2008**, *129*, 124105.

(11) Klimovich, P. V.; Shirts, M. R.; Mobley, D. L. Guidelines for the Analysis of Free Energy Calculations. *Journal of Computer-Aided Molecular Design* **2015**, *29*, 397–411.

(12) Wu, D.; Kofke, D. A. Phase-Space Overlap Measures. I. Fail-Safe Bias Detection in Free Energies Calculated by Molecular Simulation. *The Journal of Chemical Physics* **2005**, *123*, 054103.

(13) Wu, D.; Kofke, D. A. Phase-Space Overlap Measures. II. Design and Implementation of Staging Methods for Free-Energy Calculations. *The Journal of Chemical Physics* **2005**, *123*, 084109.

(14) Minh, D. D. L. Implicit Ligand Theory: Rigorous Binding Free Energies and Thermodynamic Expectations from Molecular Docking. *The Journal of Chemical Physics* **2012**, *137*, 104106.

(15) Nguyen, T. H.; Minh, D. D. L. Implicit Ligand Theory for Relative Binding Free Energies. *The Journal of Chemical Physics* **2018**, *148*, 104114.

(16) Schütte, C.; Nielsen, A.; Weber, M. Markov State Models and Molecular Alchemy. *Molecular Physics* **2015**, *113*, 69–78.

(17) He, X.; Zhang, C. Conformational Free Energy Changes via an Alchemical Path without Reaction Coordinates. *The Journal of Physical Chemistry Letters* **2018**, *9*, 4428–4435.

(18) Smith, L. G.; Novak, B.; Osato, M.; Mobley, D. L.; Bowman, G. R. PopShift: A Thermodynamically Sound Approach to Estimate Binding Free Energies by Accounting for Ligand-Induced Population Shifts from a Ligand-Free Markov State Model. *Journal of Chemical Theory and Computation* **2024**, *20*, 1036–1050.

(19) Lyman, E.; Zuckerman, D. M. On the Structural Convergence of Biomolecular Simulations by Determination of the Effective Sample Size. *The Journal of Physical Chemistry B* **2007**, *111*, 12876–12882.

(20) Romo, T. D.; Grossfield, A. Automated Sampling Assessment for Molecular Simulations Using the Effective Sample Size. *Journal of Chemical Theory and Computation* **2011**, *7*, 2464–2472.

(21) Kong, A. *A Note on Importance Sampling using Standardized Weights*; 1992.

(22) Elvira, V.; Martino, L.; Robert, C. P. Rethinking the Effective Sample Size. *International Statistical Review* **2022**, *90*, 525–550.

(23) Hansen, P. C.; O'Leary, D. P. The Use of the L-Curve in the Regularization of Discrete Ill-Posed Problems. *SIAM Journal on Scientific Computing* **1993**, *14*, 1487–1503.

(24) Honda, S.; Yamasaki, K.; Sawada, Y.; Morii, H. 10 Residue Folded Peptide Designed by Segment Statistics. *Structure* **2004**, *12*, 1507–1518.

(25) Honda, S.; Akiba, T.; Kato, Y. S.; Sawada, Y.; Sekijima, M.; Ishimura, M.; Ooishi, A.; Watanabe, H.; Odahara, T.; Harata, K. Crystal Structure of a Ten-Amino Acid Protein. *Journal of the American Chemical Society* **2008**, *130*, 15327–15331.

(26) Fischer, A.-L. M.; Tichy, A.; Kokot, J.; Hoerschinger, V. J.; Wild, R. F.; Riccabona, J. R.; Löffler, J. R.; Waibl, F.; Quoika, P. K.; Gschwandtner, P.; Forli, S.; Ward, A. B.; Liedl, K. R.; Zacharias, M.; Fernández-Quintero, M. L. The Role of Force Fields and Water Models in Protein Folding and Unfolding Dynamics. *Journal of Chemical Theory and Computation* **2024**, *20*, 2321–2333.

(27) Morton, A.; Baase, W. A.; Matthews, B. W. Energetic Origins of Specificity of Ligand Binding in an Interior Nonpolar Cavity of T4 Lysozyme. *Biochemistry* **1995**, *34*, 8564–8575.

(28) Vallurupalli, P.; Chakrabarti, N.; Pomès, R.; Kay, L. E. Atomistic picture of conformational exchange in a T4 lysozyme cavity mutant: an experiment-guided molecular dynamics study. *Chemical Science* **2016**, *7*, 3602–3613.

(29) Abella, J. R.; Antunes, D. A.; Clementi, C.; Kavraki, L. E. Markov state modeling reveals alternative unbinding pathways for peptide–MHC complexes. *Proceedings of the National Academy of Sciences* **2020**, *117*, 27374–27382.

(30) Case, D. A.; Cerutti, D. S.; Cruzeiro, V. W. D.; Darden, T. A.; Duke, R. E.; Ghazimirsaeed, M.; Giambasu, G. M.; Giese, T. J.; Götz, A. W.; Harris, J. A.; Kasavajhala, K.; Lee, T.-S.; Li, Z.; Lin, C.; Liu, J.; Miao, Y.; Salomon-Ferrer, R.; Shen, J.; Snyder, R.; Swails, J.; Walker, R. C.; Wang, J.; Wu, X.; Zeng, J.; Cheatham, T. E.; Roe, D. R.; Roitberg, A.; Simmerling, C.; York, D. M.; Nagan, M. C.; Merz, K. M. Recent Developments in Amber Biomolecular Simulations. *Journal of Chemical Information and Modeling* **2025**, *65*, 7835–7843.

(31) Tian, C.; Kasavajhala, K.; Belfon, K. A. A.; Raguette, L.; Huang, H.; Migues, A. N.; Bickel, J.; Wang, Y.; Pincay, J.; Wu, Q.; Simmerling, C. ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. *Journal of Chemical Theory and Computation* **2020**, *16*, 528–552.

(32) Izadi, S.; Anandakrishnan, R.; Onufriev, A. V. Building Water Models: A Different Approach. *The Journal of Physical Chemistry Letters* **2014**, *5*, 3863–3871.

(33) Joung, I. S.; Cheatham, T. E. Determination of Alkali and Halide Monovalent Ion

Parameters for Use in Explicitly Solvated Biomolecular Simulations. *The Journal of Physical Chemistry B* **2008**, *112*, 9020–9041.

(34) Miyamoto, S.; Kollman, P. A. Settle: An analytical version of the SHAKE and RATTLE algorithm for rigid water models. *Journal of Computational Chemistry* **1992**, *13*, 952–962.