Free Energy Landscape Perturbation for Estimating Mutation-Induced Population Shifts from Reference Molecular Dynamics Ensembles

Tatum E. Murdock and Benjamin P. Brown

Department of Pharmacology, Center for AI in Protein Dynamics, Vanderbilt University, Nashville, TN 37232, United States

Abstract

Protein function is governed by an ensemble of conformational states, and mutations can alter function by redistributing probability across that ensemble even when structural changes are modest. Directly estimating a perturbed conformational free energy landscape with molecular dynamics (MD) is expensive because each mutation normally requires a new converged simulation. Here, we introduce Free Energy Landscape Perturbation (FELP), a state-local reweighting framework that estimates mutation-induced population shifts from a reference MD ensemble. FELP partitions a reference trajectory into discrete microstates, evaluates restrained reference-to-perturbed energy differences for configurations sampled within each state, and combines spatially regularized state-local Zwanzig factors with the reference state probabilities to estimate the perturbed state distribution. FELP estimates how a perturbation changes the relative probabilities of conformational basins and reports effective-sample and overlap diagnostics that identify when the predicted shift lacks support in the sampled reference ensemble. We describe the FELP estimator, its implementation, and its performance on glycine dipeptide, chignolin, and T4 lysozyme. FELP is intended for perturbations that reshape populations within a sufficiently sampled reference landscape, providing a practical bridge between local alchemical energy evaluation and ensemble-level mechanistic interpretation.

Introduction

Protein function is largely governed by conformational free energies. Folding, stability, enzyme catalysis, signal transduction, molecular recognition, and allosteric regulation all depend on how probability is distributed over conformational states and how readily the system moves between them. Many proteins therefore function through ensembles of native-like substates, with activity controlled by changes in basin depth, transition pathways, and relative occupancy. Quantitatively describing these landscapes is central to understanding how sequence, ligand binding, protonation, and chemical modification alter biological function.

Molecular dynamics (MD) simulations provide a direct route to estimating these conformational free energy landscapes at atomistic resolution. In principle, sufficiently long trajectories recover both the relevant structural diversity and the thermodynamic probability of each state. In practice, sampling remains the central limitation. Transitions between functional conformations can be slow, rare, and system-specific, such that converging a separate MD ensemble for every perturbation of interest is often prohibitively expensive. Parallel simulation strategies, enhanced sampling, and Markov state models (MSMs) have improved access to long-timescale behavior by combining many trajectories into kinetic and thermodynamic state models <I probably also should cite a review of enhanced sampling strategies, including replica exchange, weighted ensemble, metadynamics, and then MSMs (which we are already citing)> [1, 2]. However, even with these advances, each new mutation or chemical change can still demand substantial additional sampling to converge a new landscape. <we want to say something about how this expense is a bottleneck for detailed structure-function studies>

To avoid the cost of full ensemble resampling, computational methods often sacrifice conformational diversity

for speed. Single-structure and local relaxation methods, such as Rosetta $\Delta\Delta G$ estimates for mutation stability calculations, can test whether a particular coordinate set becomes more or less favorable after mutation, but they typically evaluate a narrow structural hypothesis [3]. Minimal relaxation avoids the cost of full resampling, yet it can convert small coordinate differences into abrupt energy changes and noisy estimates. These calculations can be useful for local stability screening, but they generally do not directly answer whether a mutation stabilizes one conformational basin over another.

Alchemical methods like free energy perturbation (FEP) provide a formally rigorous extension of this idea. By computing free energy differences between Hamiltonians from local equilibrium samples, FEP incorporates local flexibility and avoids the abrupt energy changes of single-structure minimization [4–9]. However, practical calculations remain structurally restricted because they are initialized from a small number of conformations and restrained to prevent structural drift. Both single-structure screening and local FEP calculations estimate the energetic effect of a perturbation in a specific structural context. Additional ensemble analysis is required to describe how that perturbation redistributes probability across a complex conformational landscape. Furthermore, exponential free energy estimators rely heavily on the assumption that the reference and target states share sufficient phase-space overlap [10, 11]. Because these estimators are highly sensitive to rare and high-weight samples, inadequate overlap can produce unstable estimates that appear numerically converged but remain physically unsupported.

Recent population-shift strategies address this ensemble problem by separating conformational sampling from perturbation evaluation. This approach originally emerged in the context of ligand binding to quantify conformational selection and induced-fit mechanisms. Implicit-ligand theory, for example, provides a rigorous analytical framework for estimating how a binding event reweights host conformations without requiring explicit simulation of the bound complex [12–15]. While traditional implicit-ligand theory typically evaluates these shifts using exact volume integrals or analytical interaction potentials over a continuous ensemble, recent methods adapt this logic to discrete kinetic models and empirical scoring. PopShift, for instance, evaluates discrete ligand docking scores against the states of a ligand-free MSM to predict binding-induced population shifts [16]. The unifying insight of these approaches is that expensive sampling can be performed once for a reference system. Once the reference state probabilities are established, the effects of many different ligands can be evaluated against the same conformational baseline. This framework moves beyond single-structure scoring by explicitly linking the energetic preference of a perturbation to the thermodynamic weight of each conformational state.

Here, we introduce Free Energy Landscape Perturbation (FELP) to apply this population-shift framework to amino acid mutations. FELP combines reference MD sampling, discrete state models, restrained local perturbation evaluation, and FEP-style reweighting. The method partitions a reference ensemble into discrete microstates, evaluates the energetic effect of a mutation within each state, and reweights the reference probabilities to estimate the perturbed landscape. By treating a global population shift as a collection of tractable local reweighting problems, FELP bridges the gap between single-structure scoring and full-ensemble resampling. To address the inherent phase-space overlap limitations of exponential estimators, FELP explicitly audits its predictions using effective sample size and weight-concentration diagnostics [17, 18]. This internal measure of physical credibility ensures that the method not only predicts how a mutation reshapes a landscape but also identifies when a proposed perturbation falls outside the regime supported by the reference ensemble.

Theory

Consider a reference system with potential energy $U_0(x)$ and a perturbed system with potential energy $U_1(x)$, where x denotes a molecular configuration and $\beta = (k_B T)^{-1}$. The Boltzmann distribution maps these potential energy functions to continuous probability densities over configurations. A reference MD trajectory samples configurations from the reference density $p_0(x) \propto \exp[-\beta U_0(x)]$. FELP partitions this configuration space into discrete microstates $i \in \{1, \ldots, N\}$ using a chosen collective-variable or feature representation. Let $\chi_i(x)$ be the indicator function for state i. The restricted partition function for state i under Hamiltonian a is

$$Z_a(i) = \int \chi_i(x) \exp[-\beta U_a(x)]\, dx, \tag{1}$$

and the corresponding thermodynamic probability of the microstate is $\pi_a(i) = Z_a(i)/\sum_k Z_a(k)$. For the reference system, this is equivalent to integrating the continuous reference density over the microstate to obtain the reference prior $\pi_0(i)$. FELP aims to estimate the perturbed microstate probabilities $\pi_1(i)$ using only samples drawn from the reference ensemble.

To bridge the reference and perturbed ensembles, FELP relies on the principle that global conformational shifts can be decomposed into local reweighting problems. Within a specific microstate, the ratio of restricted partition functions follows the two-state Zwanzig identity,

$$\frac{Z_1(i)}{Z_0(i)} = \langle \exp[-\beta \Delta U(x)] \rangle_{0,i}\,, \tag{2}$$

where $\Delta U(x) = U_1(x) - U_0(x)$ and $\langle \cdot \rangle_{0,i}$ denotes an ensemble average over reference configurations conditioned on membership in state i. Normalizing the restricted partition functions over all microstates yields the central FELP estimator:

$$\pi_1(i) = \frac{\pi_0(i)\, \langle \exp[-\beta \Delta U(x)] \rangle_{0,i}}{\sum_k \pi_0(k)\, \langle \exp[-\beta \Delta U(x)] \rangle_{0,k}}. \tag{3}$$

Equation 3 preserves the reference state decomposition and updates the probability of each state according to the perturbation's energetic effect on configurations sampled within that specific state.

The central hypothesis behind this formulation is that discretization creates domains of local phase-space overlap. By conditioning the exponential average on microstate membership, FELP assumes that reference configurations assigned to state i adequately represent the structural ensemble of the perturbed system within that same state. This local two-state Zwanzig construction avoids the need for a multistate estimator like MBAR, which would be necessary if samples were drawn from multiple thermodynamic states or intermediate alchemical windows. Instead, FELP addresses the practical regime where expansive sampling is available only for the reference system. If a mutation is unfavorable in one conformational basin but favorable in another, the state-local reweighting factors will predict a population shift between those basins rather than yielding a single global stability score.

In a finite trajectory, state i contains evaluated configurations x_{ij} for $j = 1, \ldots, n_i$. FELP estimates the local Zwanzig factor by computing

$$f_i = \frac{1}{n_i} \sum_{j=1}^{n_i} w_{ij}, \qquad w_{ij} = \exp[-\beta(\Delta U_{ij} - s)], \tag{4}$$

where $s = \min(\Delta U)$ over all eligible frames is a uniform stability shift that prevents numerical overflow. Because this shift is applied globally, it cancels during normalization. The implemented reweighted probability is obtained by substituting the empirical reference prior $\hat{\pi}_0(i)$ and the finite-sample factor f_i into Equation 3. This step is the sample estimate of the theoretical expression above rather than a separate estimator. The corresponding relative free energy of each state is

$$F_1(i) = -k_B T \ln \hat{\pi}_1(i) + C, \tag{5}$$

where C is chosen so that the lowest finite state free energy is set to zero.

For finite samples, f_i can be dominated by a small number of high-weight frames even when n_i is large. The local effective sample size,

$$\mathrm{ESS}_i = \frac{\left(\sum_j w_{ij}\right)^2}{\sum_j w_{ij}^2}, \tag{6}$$

quantifies this weight degeneracy [19, 20]. Low ESS_i indicates that the local Zwanzig factor depends on too few configurations to be reliable. Additional frame-level diagnostics—dominant frame weight fraction, leave-one-out free-energy shift, and upper-tail Pareto shape estimate—flag states where single frames control the average.

In the intended use case the reference ensemble is chosen to be broader than the target, so many reference states may receive little perturbed probability. Global measures of distributional similarity between $\hat{\pi}_0$ and $\hat{\pi}_1$ can be low without indicating a problem; what matters is whether the states that carry significant $\hat{\pi}_1$ have adequate local support. FELP reports the perturbed probability mass in unreliable states,

$$\mathcal{U} = \sum_i \hat{\pi}_1(i)\, \mathbf{1}\{\mathrm{ESS}_i < \epsilon_i \text{ or } q_i = 1\}, \tag{7}$$

where q_i is a state-local quality-control flag and ϵ_i is a chosen effective-sample floor. The implementation uses advisory floors of $\mathrm{ESS}_i < 10$ and $\mathrm{ESS}_i < 50$ as well as a scale-aware floor $\epsilon_i = \max(\mathrm{ESS}_{\min}, 0.1\, n_i)$ with default $\mathrm{ESS}_{\min} = 5$. Small $\mathcal{U}$ indicates that the predicted landscape rests on well-supported local factors; large $\mathcal{U}$ indicates that the prediction depends on states where the local Zwanzig average is unreliable.

A separate practical issue arises from discretization. The potential energy functions U_0 and U_1 are continuous, so the true restricted partition functions of neighboring microstates should vary smoothly within a basin. Treating every microstate independently forces states with low ESS_i to produce local factors that fluctuate sharply relative to their neighbors. FELP regularizes these estimates by pooling information across adjacent states in the feature-space topology. For state i with neighbors $\mathcal{N}(i)$ having positive support, the pooled factor is

$$\bar{f}_i = \frac{\sum_{j\in\mathcal{N}(i)} \mathrm{ESS}_j\, f_j}{\sum_{j\in\mathcal{N}(i)} \mathrm{ESS}_j}, \tag{8}$$

and the blended factor used for reweighting is

$$f_i^{\mathrm{blend}} = \alpha_i f_i + (1-\alpha_i)\bar{f}_i, \qquad \alpha_i = \frac{\mathrm{ESS}_i}{\mathrm{ESS}_i + \tau}. \tag{9}$$

States with high ESS_i retain their local factors ($\alpha_i \approx 1$); states with low ESS_i borrow strength from well-supported neighbors. Equation 3 is then evaluated with f_i^{blend} in place of the raw f_i.

FELP selects τ by an L-curve criterion [21], balancing a misfit term,

$$\mathcal{M}(\tau) = \sum_i \mathrm{ESS}_i \left[\ln f_i - \ln f_i^{\mathrm{blend}}(\tau)\right]^2, \tag{10}$$

against a roughness term over neighboring states,

$$\mathcal{R}(\tau) = \frac{1}{2}\sum_i \sum_{j\in\mathcal{N}(i)} \left[\ln f_i^{\mathrm{blend}}(\tau) - \ln f_j^{\mathrm{blend}}(\tau)\right]^2. \tag{11}$$

The selected τ is the corner of the $\mathcal{M}$–$\mathcal{R}$ curve in log–log space, defined as the point farthest from the chord connecting the two extremes. This regularization stabilizes noisy local factors on a discretized landscape but does not compensate for missing reference support; $\mathcal{U}$ remains the primary validity criterion.

Results

FELP estimates perturbed landscapes through a four-stage local reweighting pipeline

The FELP workflow has four stages **(Figure 1)**. First, a reference MD trajectory is generated for the system expected to contain the broader or more permissive conformational ensemble. The trajectory is summarized into microstates using a uniform ϕ/ψ grid, an explicit JSON/YAML state file, or an exported MSM bundle. Microstate construction for each benchmark is specified in Methods. These microstates define both the aggregation units for reweighting and the coordinates used for plotting.

Second, each reference frame is assigned to a microstate. When raw-count priors are used, $\hat{\pi}_0(i)$ is the empirical fraction of assigned reference frames in state i. When a state file or MSM bundle contains trusted reference probabilities, FELP can instead normalize those loaded priors. The state identifiers used during assignment become the identifiers used throughout reweighting, summary tables, and plotting.

Third, FELP evaluates selected configurations within each state. Amino acid substitutions are introduced with PyRosetta. Reference-frame restraints are built from the original configuration and then applied to both the reference and perturbed systems so that the energy comparison remains local to the assigned state. The current implementation supports Rosetta preparation by FastRelax or side-chain repacking, target-only or radius-based mobility, and energy evaluation with OpenMM/AMBER or Rosetta scoring. The output of this stage is a frame-level table containing U_0, U_1, ΔU, assignment metadata, restraint deviations, and collective-variable measurements.

Fourth, FELP aggregates frame-level Boltzmann factors into state-level reweighting factors and computes the perturbed state probabilities. Local factors are neighbor-pooled on the assignment grid with an L-curve-selected blending strength before normalization. The resulting landscape can be plotted over discrete microstates or projected into the collective-variable space used for assignment. For custom collective variables, FELP stores state ranges, state centers, and frame-level measurements as metadata so that downstream visualization remains tied to the same coordinates used for reweighting.




Figure 1: Free Energy Landscape Perturbation workflow. A reference MD ensemble (e.g., glycine dipeptide) is partitioned into microstates in a chosen collective-variable space. Reference-to-perturbed energy differences are evaluated for sampled structures within each state. State-specific exponential averages reweight the reference probabilities to estimate the perturbed landscape (e.g., proline). Overlap diagnostics audit whether the predicted population shift is supported by the reference ensemble.

Glycine dipeptide

Glycine dipeptide provides a minimal test of FELP in a well-understood two-dimensional conformational landscape. The reference ensemble is projected into backbone Ramachandran space and partitioned into ϕ/ψ microstates. The current benchmark compares a glycine reference ensemble to an alanine-like perturbed ensemble, a useful first case because glycine samples a broader Ramachandran landscape than alanine. The expected result is recovery of the dominant population redistribution rather than every high-energy contour: alanine should down-weight regions that are favorable only for glycine and retain probability in the subset of Ramachandran space compatible with the perturbed side chain.

This benchmark is also the clearest setting for diagnosing the distinction between phase-space support and energetic preference. If a Ramachandran basin exists in the alanine MD reference but was absent from the

glycine-to-alanine FELP support, FELP should not be expected to recover it. Conversely, if the alanine landscape is contained within the glycine support, FELP should reproduce the major basin-level shifts with favorable effective-sample diagnostics. The glycine system will therefore serve as the primary validation of the estimator and plotting semantics.

Placeholder for glycine dipeptide reference, MD-perturbed, and FELP-reweighted landscapes.

Figure 2: FELP benchmark on a Ramachandran landscape. TODO: show the reference landscape, the independently sampled perturbed landscape, the FELP-reweighted landscape, and the difference map. Include overlap diagnostics and basin population comparisons.

Chignolin

Chignolin tests FELP on a small protein whose conformational preferences depend on coordinated intramolecular contacts. Rather than projecting only the target residue's backbone angles, we define collective variables based on intramolecular distances that report on global compaction and turn geometry. This benchmark is intended to test whether FELP can predict mutation-induced shifts in a more protein-like landscape while retaining a compact and interpretable state representation.

In this setting, naive use of a default Ramachandran grid would be misleading: the prior would be defined over target-residue ϕ/ψ bins rather than the distance coordinates used to interpret the landscape. FELP therefore treats the distance variables as first-class collective variables. The reference trajectory is binned in the joint distance space, raw-count priors are computed over those distance-defined microstates, and the perturbed landscape is visualized in the same coordinates.

The expected chignolin result is a mutation-dependent redistribution between compact, turn-like, and extended-like regions of the distance-coordinate landscape. A successful FELP run should preserve the broad topology of the independently sampled mutant landscape while identifying which microstates gain or lose probability. The analysis should report both the continuous projected surfaces and the discrete state summaries, because discrepancies between these views can reveal whether a visually smooth projected landscape is supported by enough state-local samples.

Placeholder for chignolin distance-CV landscapes.

Figure 3: Distance-coordinate FELP benchmark in chignolin. TODO: show the reference distance landscape, independently sampled mutant landscape, FELP-reweighted landscape, and population shifts for folded, turn, and extended-like states. Report the automatically generated binning or the explicit microstate JSON used for the analysis.

T4 lysozyme

T4 lysozyme provides a natural larger benchmark because it has been used extensively to study ligand binding, cavity mutants, and population-shift methods, including the PopShift validation on L99A [16, 22]. For FELP, the T4 lysozyme benchmark should be framed around mutations or chemical perturbations that redistribute probability among pre-existing conformational substates rather than requiring discovery of an entirely new fold or pocket geometry.

In this setting, the state definition should be chosen to reflect the mechanistic question. Possible collective variables include cavity-opening distances, helix or loop distances surrounding the engineered binding pocket, side-chain rotameric states of pocket residues, or MSM states derived from reference simulations. The expected result is a more stringent test than glycine or chignolin: FELP should identify mutation-induced shifts among protein substates when the reference trajectory samples those substates, while the diagnostics should flag cases in which the mutation stabilizes configurations absent from the reference simulation. This benchmark will therefore test both the scalability of FELP and the practical value of its overlap audit.

Overlap diagnostics and failure modes

FELP is not intended to replace perturbed MD when the perturbation creates a new basin absent from the reference simulation. Its value lies in rapidly estimating population shifts within a sampled landscape and in identifying when that estimate is unsupported. The most important failure mode is poor phase-space overlap. In practice, this appears as a small effective sample size, target probability concentrated in low-support states, one or a few frames dominating the reweighted distribution, or positive-prior states with no evaluated support.

The support diagnostics introduced in the Theory section distinguish FELP from purely empirical population-shift tools. A population shift can always be computed mathematically; physical credibility depends on reference support. By reporting overlap statistics alongside the reweighted landscape, FELP converts this into an auditable claim: the user can ask what shift is predicted and whether the reference ensemble contains the configurations required for that prediction to be meaningful. When both directions are computationally feasible, forward and reverse FELP calculations provide a stronger overlap test.

Discussion

FELP occupies an intermediate point between high-fidelity perturbed MD and local mutation scoring. It does not attempt to resample every mutant. Instead, it leverages extensive sampling of a reference system and asks how chemical perturbations reweight the probabilities of already observed states. There is precedent for this kind of thermodynamic reweighting in biological applications. For example, Abella et al. applied an analogous strategy to evaluate peptide mutations from a wild-type Markov state model, reweighting the transition between bound and unbound metastable macrostates [23]. FELP is a generalization of this and related approaches.

The method clarifies the relationship between physical rigor and computational throughput. The local estimator is inherited from FEP and therefore carries the same overlap requirements as any exponential reweighting calculation. The global landscape estimate is obtained by applying these local perturbation factors across a reference state model. When the reference state model is well supported and overlap diagnostics are favorable, FELP can estimate mutation-induced population shifts at far lower cost than independently converging each perturbed ensemble. When diagnostics are unfavorable, the method provides evidence that new sampling is required.

The most important conceptual limitation is that FELP estimates the perturbed distribution over the reference state space. It cannot discover a target-only state, and it should not be described as a replacement for perturbed MD when large conformational changes are expected. The practical strength of the method is that this limitation is explicit. FELP makes a prediction and a validity statement together: the predicted shift, the state support for that shift, and the extent to which the estimate is dominated by rare high-weight frames.

Several implementation details are important for practical use. First, the collective variables used for microstate assignment must match the biological question and the plotted landscape. Second, state identifiers in the microstate definition are the identifiers used for raw-count priors, reweighting, and plotting. Third, custom collective variables must be propagated through frame metadata and summary metadata so that visualization does not silently fall back to default ϕ/ψ coordinates. Finally, restraint strengths and coordinate-preservation protocols must be interpreted in the units and conventions of the engine being used. In the current implementation, Rosetta and OpenMM use different restraint parameterizations, so backend-specific restraint settings must be reported in Methods.

The planned benchmarks are designed to expose these strengths and limitations. Glycine dipeptide tests whether the estimator behaves correctly in a small, visually interpretable landscape. Chignolin tests whether distance-coordinate microstates can capture protein-like population shifts. T4 lysozyme tests whether FELP can scale to a biologically studied system while providing meaningful warnings when a mutation or ligand-coupled perturbation moves beyond the reference support. Together, these systems should establish FELP as an auditable reweighting framework for a defined and useful class of perturbations.

Free Energy Landscape Perturbation provides a framework for estimating how mutations and other chemical perturbations reshape conformational free energy landscapes without requiring a new converged MD simulation for every perturbed system. By combining reference ensemble sampling, microstate-based reweighting, local energy-difference evaluation, and overlap diagnostics, FELP connects the efficiency of perturbative calculations with the ensemble perspective required for protein function. The resulting approach is most powerful when the reference simulation captures the relevant conformational space and when diagnostics confirm that the predicted population shift is supported. Under these conditions, expansive sampling performed once for a reference model can be reused to test a family of perturbations while preserving a direct connection to thermodynamic state populations.

Methods

Reference simulations and trajectory processing

Reference trajectories are supplied to FELP as one or more topology/trajectory groups. Files passed within one trajectory group are treated as contiguous fragments of the same simulation, whereas repeated trajectory groups are treated as independent trajectories for MSM construction and provenance. For each benchmark, the final manuscript should report the MD engine, force field, solvent model, temperature, pressure treatment, time step, thermostat and barostat settings, equilibration protocol, production length, number of independent replicas, and frame stride used for FELP analysis.

Trajectory frames are loaded with a user-specified topology, optional frame start/stop indices, and stride. For each selected frame, FELP clones the reference topology with the frame coordinates, converts the structure to a PyRosetta pose when needed, and records the trajectory identifier and source frame index. These identifiers are propagated to the frame-level output so that failed frames, high-weight frames, and representative microstate samples can be traced back to the original simulation data.

Microstate construction

For Ramachandran analyses, microstates are defined on a uniform ϕ/ψ grid over $[-180°, 180°)$ for each axis. For glycine dipeptide and other ϕ/ψ benchmarks, this grid supplies both assignment and plotting coordinates.

For custom collective-variable analyses, microstates are usually defined by explicit JSON or YAML files containing state identifiers and "cv_ranges." State definitions may include plotting coordinates and loaded reference probabilities. If no probabilities are provided, FELP can compute raw-count priors from the assigned reference trajectory. Angular intervals are treated circularly so that states crossing the $-180°/180°$ boundary are handled correctly.

For the chignolin benchmark, we use two intramolecular distance coordinates without a pre-specified grid. FELP's automatic microstate generator scans the reference trajectory, measures the configured variables, initializes marginal bin widths using the Freedman–Diaconis rule, and caps the joint grid with a support

criterion targeting approximately 25 assigned frames per state and at most 144 total states. The run writes a reusable microstate JSON file recording bin edges, observed ranges, quantiles, state centers, and policy metadata; all chignolin reference and mutant analyses reuse that file so mutations are compared on an identical state decomposition.

MSM-backed analyses use a previously exported PyEMMA bundle. The bundle supplies precomputed assignments, cluster or metastable state definitions, and optionally loaded stationary probabilities. The FELP reweighting equation is unchanged; the MSM provides the state space and reference prior.

Perturbation and energy evaluation

Mutations are introduced with PyRosetta at a user-specified sequential pose residue index. FELP optionally enforces the original residue identity before scoring, which is useful for glycine-to-alanine or protonation-state studies where the reference topology must be chemically explicit. Canonical amino acids and common protonation or disulfide variants are supported when the corresponding residue types are available to PyRosetta and the selected energy backend.

For each frame, FELP first measures the configured collective variables and assigns the frame to a microstate. It then builds reference-frame restraints from the same pose. Default restraints are the target residue ϕ and ψ angles, but users may supply arbitrary backbone dihedral CVs and distance CVs. The same restraint targets are applied to the reference and perturbed systems. This makes the comparison local to the assigned reference configuration and reduces drift into a different basin before the energy difference is evaluated.

The PyRosetta preparation step can use FastRelax or side-chain repacking. The default mobility policy allows the target residue side chain to move while keeping the rest of the structure fixed; a radius-based mobility option can permit neighboring side chains, and optionally backbone atoms, to relax. For Rosetta preparation, the default score function is ref2015 with Cartesian relaxation enabled when supported. Backbone CVs are explicitly reset and checked after mutation and local preparation. Distance restraints are implemented as Rosetta atom-pair constraints with a reported width parameter, whereas OpenMM distance restraints use harmonic spring constants; both settings should be reported for production analyses.

Energy evaluation can be performed with OpenMM or Rosetta. The default OpenMM backend uses AMBER ff14SB-only side-chain protein parameters with GBNeck2 implicit solvent, evaluates both reference and perturbed structures under the same restraint targets, freezes atoms outside the configured mobility region during minimization, and reports energies in $\mathrm{kJ\,mol^{-1}}$. The optional Rosetta backend scores PyRosetta poses with the configured score function and constraints. In either backend, FELP records

$$\Delta U_{ij} = U_1(x_{ij}) - U_0(x_{ij}) \tag{12}$$

for each evaluated frame and excludes failed or non-finite frames from reweighting.

Reweighting and diagnostics

State probabilities are reweighted according to Equation 3 using neighbor-pooled blend factors f_i^{blend} with τ chosen by the L-curve criterion described in Theory. The independent per-state factors f_i are retained in the output for comparison and for overlap diagnostics. The reference prior is either loaded from the microstate definitions or estimated from raw assigned frame counts. When microstate-stratified sampling is used to reduce expensive energy evaluations, FELP assigns all frames cheaply, stores a deterministic sampling manifest, computes raw-count priors from the full assigned support, and estimates state-local Boltzmann averages from uniformly selected frames within each state. The default evaluates all assigned frames.

Diagnostics emphasize the perturbed probability carried by low-effective-sample-size or quality-control-flagged states. FELP also reports state-level effective sample sizes, maximum normalized frame weights, top-weight mass fractions, global weight-concentration summaries, and unsupported positive-prior states. The current advisory thresholds are described above and written to the run summary. Final analyses should report these diagnostics alongside each reweighted landscape rather than presenting the landscape as an unconditional prediction.

Visualization and comparison to perturbed MD

FELP writes frame-level results, microstate summaries, reweighted surfaces, run configuration, sampling manifests, and JSON summaries. Discrete microstate plots use the state probabilities and relative free energies from the microstate summary. Projected free energy plots use state-consistent frame weights to construct weighted histograms in the plotted collective-variable coordinates. Free energies are shifted so that the lowest finite value is zero; only differences between states or plotted bins are meaningful.

For each benchmark, the FELP-reweighted landscape will be compared to an independently sampled perturbed MD landscape. Planned comparisons include the identity and location of dominant basins, relative free energies between basins, per-state occupancy changes, target-versus-reference probability scatter plots, KL-divergence or occupancy-shift summaries, and whether states with the largest predicted changes are supported by adequate effective sample sizes.

Code and data availability

The FELP implementation, analysis scripts, run commands, generated microstate definitions, processed summaries, and plotting inputs will be released in a public repository and archived in a DOI-bearing repository before submission.

Acknowledgements

This work was supported by the NIH (DP1DA058349). The authors thank members of the Center for AI in Protein Dynamics and the Department of Pharmacology at Vanderbilt University for helpful discussions.

Author contributions

T.E.M. and B.P.B. conceived the study. T.E.M. performed simulations and analyses. B.P.B. developed the FELP software framework. T.E.M. and B.P.B. interpreted the results and wrote the manuscript.

Disclosures

The authors declare no competing financial interest.

References

(1) Noé, F.; Prinz, J.-H. *Current Opinion in Structural Biology* **2011**, *21*, 562–572.
(2) Husic, B. E.; Pande, V. S. *Journal of the American Chemical Society* **2018**, *140*, 2386–2396.
(3) Kellogg, E. H.; Leaver-Fay, A.; Baker, D. *Proteins: Structure, Function, and Bioinformatics* **2011**, *79*, 830–838.
(4) Zwanzig, R. W. *The Journal of Chemical Physics* **1954**, *22*, 1420–1426.
(5) Lu, N.; Kofke, D. A. *The Journal of Chemical Physics* **2001**, *114*, 7303–7311.
(6) Shirts, M. R.; Pande, V. S. *The Journal of Chemical Physics* **2005**, *122*, 144107.
(7) Pohorille, A.; Jarzynski, C.; Chipot, C. *The Journal of Physical Chemistry B* **2010**, *114*, 10235–10253.
(8) Shirts, M. R.; Chodera, J. D. *The Journal of Chemical Physics* **2008**, *129*, 124105.
(9) Klimovich, P. V.; Shirts, M. R.; Mobley, D. L. *Journal of Computer-Aided Molecular Design* **2015**, *29*, 397–411.
(10) Wu, D.; Kofke, D. A. *The Journal of Chemical Physics* **2005**, *123*, 054103.
(11) Wu, D.; Kofke, D. A. *The Journal of Chemical Physics* **2005**, *123*, 084109.
(12) Minh, D. D. L. *The Journal of Chemical Physics* **2012**, *137*, 104106.
(13) Nguyen, T. H.; Minh, D. D. L. *The Journal of Chemical Physics* **2018**, *148*, 104114.
(14) Schütte, C.; Nielsen, A.; Weber, M. *Molecular Physics* **2015**, *113*, 69–78.
(15) He, X.; Zhang, C. *The Journal of Physical Chemistry Letters* **2018**, *9*, 4428–4435.
(16) Smith, L. G.; Novak, B.; Osato, M.; Mobley, D. L.; Bowman, G. R. *Journal of Chemical Theory and Computation* **2024**, *20*, 1036–1050.
(17) Lyman, E.; Zuckerman, D. M. *The Journal of Physical Chemistry B* **2007**, *111*, 12876–12882.
(18) Romo, T. D.; Grossfield, A. *Journal of Chemical Theory and Computation* **2011**, *7*, 2464–2472.
(19) Kong, A. *A Note on Importance Sampling using Standardized Weights*; tech. rep. 348; Department of Statistics, University of Chicago, 1992.
(20) Elvira, V.; Martino, L.; Robert, C. P. *International Statistical Review* **2022**, *90*, 525–550.
(21) Hansen, P. C.; O'Leary, D. P. *SIAM Journal on Scientific Computing* **1993**, *14*, 1487–1503.
(22) Morton, A.; Baase, W. A.; Matthews, B. W. *Biochemistry* **1995**, *34*, 8564–8575.
(23) Abella, J. R.; Antunes, D. A.; Clementi, C.; Kavraki, L. E. *Proceedings of the National Academy of Sciences* **2020**, *117*, 27374–27382.