

# Informing IPCC accounting of forest carbon using the global forest carbon database (ForC v4.0)

Madison Williams<sup>1</sup>, Valentine Herrmann<sup>1</sup>, Rebecca Banbury Morgan<sup>1,2</sup>, Ben Bond-Lamberty<sup>3</sup>, Susan Cook-Patton<sup>4</sup>, Helene Muller-Landau<sup>5</sup>, Camille Piponiot<sup>6</sup>, Teagan Rogers<sup>1</sup>, and Kristina J. Anderson-Teixeira<sup>1,5 \*</sup>

<sup>1</sup>Center for Conservation Ecology, Smithsonian Conservation Biology Institute, Front Royal, VA, United States

<sup>2</sup>School of Geography, University of Leeds, Leeds, UK

<sup>3</sup>Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, United States

<sup>4</sup>The Nature Conservancy, Arlington VA 22203, USA

<sup>5</sup>Forest Global Earth Observatory, Smithsonian Tropical Research Institute, Panama, Republic of Panama

<sup>6</sup>CIRAD, Montpellier, France

**Correspondence:** Kristina J. Anderson-Teixeira (teixeirak@si.edu)

**Abstract.** Forests are critical for climate change mitigation and constitute a substantial portion of planned emissions reductions under the 2015 Paris Agreement. Yet, the efficacy of greenhouse gas (GHG) mitigation planning and reporting is dependent upon the quality of available emission factors data, including forest carbon (C) stocks and changes therein. Tens of thousands of relevant forest C estimates have been published, yet are not readily accessible to the practitioners compiling national greenhouse gas inventories. Many of these data have, however, been compiled in the Global Forest C database (ForC; <https://forc-db.github.io/>) and stand to be of value to greenhouse gas accounting if made available through the Emission Factor Database (EFDB) of the International Panel on Climate Change (IPCC). Here, we develop and document a process for semi-automated transfer of data from ForC into the EFDB, assess the data available and transferred to date, and provide recommendations for improving forest data collection, analysis, and reporting to improve accounting of forest-sector GHG emissions and removals. We begin by reconciling terminology and mapping ForC fields into EFDB. This process required some updates to the ForC database structure, leading to the release of a new version of ForC (v4.0; described here). At the time of writing, ForC contained ## values that would qualify for inclusion in the EFDB, ## of which have been transferred to date. (Some analysis of representation/ gaps.) In the future, forest C estimates in EFDB can be improved through targeted research to fill critical gaps, reporting of information required by IPCC, and continued submission of data from scientific publications to the EFDB.

*Copyright statement.* The author's copyright for this publication is transferred to institution/company.

## 1 Introduction

Forests are critical to our management of greenhouse gases. The influence the forests have on atmospheric carbon has made forest data increasingly valuable.

20 **Forests play a central role in international plans for climate change mitigation (UNFCCC, 2015).** As of 2016, forest-based mitigation accounted for 26% of total planned greenhouse gas mitigation within Nationally Determined Contributions under the Paris Agreement (Grassi et al., 2017). Effective forest-based climate change mitigation requires solid scientific data (Anderson-Teixeira and Belair, 2022).

25 **The International Panel on Climate Change (IPCC) provides guidance for national greenhouse gas (GHG) inventories for reporting to the United Nations Framework Convention on Climate Change (UNFCCC, (REFS for older guidelines), IPCC, 2019a).** Under this guidance, GHG inventories include all managed land, including most of the world's forest land (Ogle, 2018). The IPCC inventory guidelines include specific instructions for accounting for forest land . . . (IPCC, 2006, 2019b). This guidance has improved as more of the relevant underlying data has become available (Requena Suarez et al., 2019), but there remains room for continuous improvement as the science advances. For example, Cook-Patton et al. (2020) (*summarize IPCC comparison results*). Another example is Cuni-Sanchez et al. (2021). *Moreover, it is useful for those compiling national greenhouse gas inventories to have access to locally-specific information, when available.* To improve the data accessible for C accounting, the IPCC created the Emission Factor Database (EFDB; <https://www.ipcc-nggip.iges.or.jp/EFDB/main.php>), which is intended as a recognized library of emission factors and other parameters that can be used for estimating greenhouse gas emissions and removals.

35 **To ensure that planned emissions reductions are realistic, high-quality estimates of forest C stocks and fluxes must be publicly accessible.** The EFDB is intended as a recognized library, and can be used both for efforts to tally a nation's intended or accomplished GHG reductions, or as a basis of comparison for external parties to evaluate these inventories.

**The Global Forest Carbon Database, ForC, is the largest collection of published estimates of forest carbon stocks, increments, and annual fluxes (Anderson-Teixeira et al., 2018, 2021).** (*add stats/ details, maybe record of how ForC has grown over time*) As such, ForC is positioned to improve forest C accounting through the transfer of data to EFDB. The purpose of this publication is to document that process and provide recommendations for future improvements.

40 **Here, we** (1) review IPCC definitions of relevant carbon stocks and increments (2) describe mapping of ForC to IPCC's EFDB, (3) describe updates to ForC (ForC v4.0), (4) summarize the data in ForC that's relevant to EFDB, identifying gaps, and (5) provide recommendations for improving data collection, analysis, database, and accounting.

## 2 IPCC definitions of carbon stocks and incremenets

45 For quantifying forest role in global C cycle, we ultimately care about: (1) C stocks – stores of C that would be vulnerable to release to the atmosphere upon land use change (2) C increments – changes in those C stocks.

### 2.1 Carbon stocks

Forest ecosystem C stocks may be parsed into pools in various ways. IPCC parses into biomass (aboveground and below-ground), dead organic matter (dead wood and litter), and soil organic matter (Table @ref(table\_variables)). Quantifying these requires a one-time measurement.

**Table 1. Variables with definitions and measurement methods.** Definitions from IPCC Table 1.1. (See Table 1.1 in IPCC guidance).

pool	definition	major sources of estimate variation	IPCC guidance
aboveground biomass	all biomass of living vegetation, both woody and herbaceous, above the soil	allometry, min dbh	acceptable to exclude understory
belowground biomass	all biomass of live roots	allometry, min dbh, assumed ratio of belowground to aboveground biomass (IPCC table 4.4)	fine roots may be excluded when they cannot be distinguished empirically from soil organic matter or litter
dead wood	all non-living woody biomass not contained in the litter, either standing, lying on the ground, or in the soil	min dbh, ...	default min dbh = 10cm, but may be chosen by country
litter	all non-living biomass with a size greater than the limit for soil organic matter and less than the minimum diameter chosen for dead wood, lying dead, in various states of decomposition above or within the mineral or organic soil	min dbh for dead wood , ..	includes entire O horizon
soil organic matter	organic carbon in mineral soils to a specified depth	sampling depth	default sampling depth = 30cm, but may be chosen by country

**2.1.1 Biomass**

Biomass includes living vegetation, above- and below-ground.

The IPCC defines aboveground biomass as “all biomass of living vegetation, both woody and herbaceous, above the soil including stems, stumps, branches, bark, seeds, and foliage” [].

55 Belowground biomass is defined as “all biomass of live roots” [].

**2.1.2 Dead Organic Matter**

Dead organic matter includes all non-living biomass that is not within the mineral soil layer and smaller than the litter size threshold.

Dead wood is defined as. . .

60 Litter is defined as including " all non-living biomass with a diameter less than a minimum diameter chosen by the country  
(for example 10 cm), lying dead, in various states of decomposition above the mineral or organic soil. This includes litter (OL),  
funic (OF), and humic (OH) layers. Live fine roots (of less than the suggested diameter limit for belowground biomass) are  
included in litter where they cannot be distinguished from it empirically." (2003 IPCC GPG for LULUCF ([https://www.ipcc-](https://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf_files/Glossary_Acronyms_BasicInfo/Glossary.pdf)  
nggip.iges.or.jp/public/gpglulucf/gpglulucf\_files/Glossary\_Acronyms\_BasicInfo/Glossary.pdf))

### 65 **2.1.3 Soil Organic Matter**


Soil organic matter is defined as...

## **2.2 Carbon increments**

C increments are defined as the change over time, in annual increments, in each C pool. These may be estimated as the  
difference between C stocks at two time points, or as the difference between inputs and outputs to the pool (i.e., fluxes).

70 Quantifying these requires at least two measurements. (*But, Can carbon increments be inferred from a single measure and a  
known age (i.e., the approach we used in GROA)?*) Fluxes are the inputs and outputs to each pool.

### 3 Mapping ForC to EFDB



**Figure 1. Schematic illustrating the carbon pools quantified under IPCC accounting; ForC variables corresponding to the stock, increment, influx and outflux; and relationships among them.** In many cases, the match of ForC variables to IPCC criteria depends upon measurement protocols (e.g., minimum DBH). Additional caveats are as follows: 1- assumes that change in foliage biomass is negligible (see note 7); 2- incomplete: excludes large branch fall; also, under IPCC definitions, outflux from aboveground biomass should include all sizes, influx to deadwood should include only above the minimum diameter chosen for dead wood; 3- incomplete: excludes belowground components; 4-incomplete: excludes breakage into pieces less than dead wood threshold size; 5-incomplete: excludes woody mortality of stems <10 cm DBH, decomposition of dead wood (aboveground and coarse roots) into sizes classified as litter, may exclude branch fall; 6- measurements often limited to decomposition of relatively fine litter and may exclude branches and stems below the dead wood size threshold and/or the more decomposed layers of the O horizon; 7 - foliage production is generally measured by collecting leaf-fall, a method that assumes that the influx = outflux (foliage biomass is roughly constant year-to-year); 8 - excludes branch fall, which is necessary for a full accounting of woody productivity but is typically assumed negligible for calculations of net biomass change.

Table 2: **Mapping of ForC fields to EFDB.** See footnotes at end of table (still need to be properly inserted).

<b>ForC table</b>	<b>ForC field</b>	<b>EFDB field</b>	<b>Usage</b>	<b>Required</b>
Measurements	measurement.ID	Other Properties	direct mapping	(no)
	dominant.life.form	1996 Source/Sink Categories, 2006 Source/Sink Categories	used to determine land subcategories (see defin- ing_land_subcategory.md)	yes
	stand.age	1996 Source/Sink Categories, 2006 Source/Sink Categories, Parameters/ Conditions	used to determine land subcategories (see defin- ing_land_subcategory.md), directly listed in Parameters/ Conditions	(yes)
	dominant.veg, veg.notes, min.dbh	Parameters/ Conditions	direct mapping/ linking to dominant.veg description	no
	variable.name	-	link to variable info in ForC_variables table	yes
	date / start.date, end.date	Other Properties	direct mapping	no
	mean	Value	direct mapping	yes
	mean.in.original.units	Value in Common Units	direct mapping	yes
	original.units	Common Unit	direct mapping	yes
	lower95%CI, upper 95%CI, se, sd and n	Lower Confidence Limit, Upper Confidence Limit	direct or calculated	(yes)
	depth, covariate_1, cov_1.value, covariate_2, cov_2.value	Other Properties	direct mapping	no
	allometry_1, allometry_2	Comments from Data Provider	link to biomass allometry source, when provided	no

Table 2: **Mapping of ForC fields to EFDB.** See footnotes at end of table (still need to be properly inserted). (*continued*)

ForC table	ForC field	EFDB field	Usage	Required
	data.location.within.souree		confirm that data weren't digitized, facilitate finding data in original publication	yes
	ForC.investigator	Data Provider, Data Provider Contact	link to Data Provider, Data Provider Contact info	yes
Sites	site.ID, sites.sitename	Other Properties	direct mapping	(no)
	lat, lon	Region/Regional conditions	direct mapping; used to extract continent, Koeppen, and <u>FAO.ecozone</u>	(no)
	country, state, city, masl, mat, map	Region/Regional conditions	direct mapping	no
	continent, Koeppen	Region/Regional conditions	direct mapping	auto
	soil.texture, sand, silt, clay, <u>soil.classification</u>	Parameters/ Conditions	direct mapping	no
	FAO.ecozone	Parameters/ Conditions	direct mapping	auto
History	date, hist.cat, hist.type	1996 Source/Sink Categories, 2006 Source/Sink Categories, Abatement/Control technologies	used to determine distmrs.type for Source/Sink Categories, generate list of events for Abatement/Control technologies	(yes)/no**
	plot.area	Other Properties	direct mapping	no
Plots	plot.ID, plot.name	Other Properties	direct mapping	(no)
	distmrs.type	1996 Source/Sink Categories, 2006 Source/Sink Categories	used to determine land subcategories (see defining_land_subcategory.md)	auto



Table 2: **Mapping of ForC fields to EFDB.** See footnotes at end of table (still need to be properly inserted). (*continued*)

<b>ForC table</b>	<b>ForC field</b>	<b>EFDB field</b>	<b>Usage</b>	<b>Required</b>
	distmrs.type, distmrs.year, regrowth.type, regrowth.year	Other Properties	direct mapping	auto
PFT	description	Parameters/ Conditions	direct mapping	auto
variables	variable.type	Gases	For stocks in unit of organic matter, gases include CO2, CO, CH4, NO, NO2, N2O. For increments, fluxes, and stocks in units of C, gases includes only CO2.	auto
	variable.name	C pool, Equation	link to C pool, Equation	auto
	description	Description	direct mapping	auto
	extended.description	Other Properties	direct mapping	auto
	units	Unit (ID)	link to IPCC units	auto
Citations	citation.citation	Full Technical Reference	direct mapping	yes/auto
	citation.language	Reference Language	direct mapping	yes/auto
	citation.url	URL	direct mapping	no/auto
	citation.abstract	Abstract in English	direct mapping	no/auto
	source.type	Source of Data	direct mapping	yes

‘Required’ field indicates whether the field is required by EFDB: yes = value required; (yes) = input required, missing value acceptable if not reported; auto = present within ForC infrastructure, and therefore will always be exported to EFDB ; (no) = not required for EFDB, but required for ForC and therefore will always be exported to EFDB; no = not required, but exported to EFDB when a value is present.

\*\* ‘(yes)’ for most recent severe disturbance; ‘no’ for other history events

### 3.1 Carbon cycle variables

Mapping of variables is shown in Fig. 1

### 80 3.2 Land use categories

Documented at [https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB\\_mapping/defining\\_land\\_subcategory.md](https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB_mapping/defining_land_subcategory.md), [https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB\\_mapping/IPCC\\_LandUse\\_mapping.csv](https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB_mapping/IPCC_LandUse_mapping.csv), and in issue #8.

The UNFCCC requires GHG reporting for all managed lands in a country, where management is defined as “human inter-  
ventions and practices have been applied to perform production, ecological or social functions” [IPCC 2006 full report REF].  
This definition is applied differently across countries, and is not clearly defined by the majority of governments (Ogle, 2018).  
Given this, and because the IPCC definition of management does not necessarily match that which would be reported in scientific publications and hence in ForC, we do not transfer any classification of land management status from ForC to the EFDB, but do provide auxiliary info that may be useful in making this determination (e.g., geographical location).

## 90 4 Updates to ForC (ForC v4.0)

To support export of data to EFDB, and to improve the overall quality of the ForC database, we defined ## new variables, implemented some modest restructuring, resolved duplicate records, and conducted quality control. This section describes changes relative to ForC v2.0 (Anderson-Teixeira et al., 2018).

### 4.1 Defining new variables

95 We added eleven increment variables to the set of named and defined variables (or 22, counting \_OM and \_C versions), which previously included only one (aboveground biomass increment, *delta.agb*). (<https://github.com/forc-db/IPCC-EFDB-integration/issues/6>) These are directly related to C stocks as previously defined in ForC, with “delta.” added in front of the variable name.

Although these variables currently lack records, the structure exists such that records can be populated over time.

100 To provide better definition of the previously existing variable *organic.layer*, which has a nebulous definition that reflects the varied definitions adopted by original studies, we added two clearly defined variables: *litter* (relatively undecomposed plant material/ OL horizon), and *O.horizon* (entire O-horizon, including *litter* (OL)).

### 4.2 ForC restructuring

Table 3: **Table of changes to ForC fields.**

<b>Table</b>	<b>Column</b>	<b>Description</b>	<b>Changes</b>	<b>Motivation</b>
Sites	coordinates.precision	Precision of geographic coordinates, as reported by source or estimated from maps.	field added	allow identification of records with poor coordinate precision
Measurements	data.location.within.source	Location of data within the source listed in citation.ID.	field added	facilitate review, ensure traceability
	sd, se, lower95%CI, upper 95%CI	Standard deviation, standard error, and lower and upper 95 percent confidence intervals, respectively.	replaces ‘stat’ and ‘stat.name’	cleaner format; ability to handle assymetrical 95 percent confidence intervals
	mean.in.original.units, original.units	mean value and units presented in original publication	fields added	provide IPCC with original units, reduce errors/improve reproducibility
	C.conversion.factor	Assumed/ measured C content of organic matter used to convert organic matter to C.	field added	track units conversion, allow back-calculation of OM if conversion factor deemed inappropriate
PFT	description	Definition of the pftcode at the community level. Differs from individual level in that properly describes mixed plant functional types.	field added	

Table 3: **Table of changes to ForC fields.** (*continued*)

<b>Table</b>	<b>Column</b>	<b>Description</b>	<b>Changes</b>	<b>Motivation</b>
Citations	description.individual	Definition of the pftcode at the individual plant level.	field name change (previously ‘description’)	
	citation.citation	Full citation. Most of these records are automatically generated in R based upon DOI lookup.	field added	field required by IPCC
	citation.language	Language of original publication, automatically generated based on the title and abstract, with some manual entries and corrections.	field added	field required by IPCC
	citation.url	URL of original publication, generally retrieved automatically via URL lookup.	field added	field required by IPCC
	citation.abstract	Abstract, generally retrieved automatically via DOI lookup.	field added	field required by IPCC
	source.type	citation source type	field added	field required by IPCC
	pdf.in.repository	Indicates whether pdf of original study has been retrieved and saved in ForC’s reference repository	field added	housekeeping

Table 3: **Table of changes to ForC fields.** *(continued)*

Table	Column	Description	Changes	Motivation
	EFDB.ready	Indicates whether data have been checked for export to EFDB.	field added	housekeeping

**4.3 Quality control measures**

105 Prior to releasing ForC v4.0, we executed several quality control measures. First, we implemented a system of continuous integration using GitHub Actions (*sensu* Kim et al. in prep) to run some automatic checks any time the master data files are updated. Second, to improve information on geographic coordinates, we flagged and reviewed records with suspected low precision (*Issue #29*)[<https://github.com/forc-db/ForC/issues/229>]. Third, to identify erroneous climate data... (*Issue #212*)[<https://github.com/forc-db/ForC/issues/212>].

110 **4.4 Resolving duplicates**

**5 Results**

**figure: map of relevant ForC data with underlying FAO ecozones**

(summarize the data in ForC that’s relevant to EFDB, identifying gaps)

dead wood and litter comparisons will be particularly interesting, as IPCC values are based on just a handful of references  
115 for each climate zone (table 2.2 in 2019 guidelines)

**6 Recommendations**

(strongly flag both useful variables that the EFDB does not track and useful variables that papers fail to include that EFDB needs)

**6.1 Data collection and analysis needs**

120 **(Paragraph highlighting important gaps in variables / regions)**

Several variables of value to IPCC, including standing dead wood, woody mortality, delta.agb, are not calculated and presented as frequently as are AGB and ANPP\_woody, even though they can readily be derived from the same census data. We recommend that researchers calculate and report these, as specified below. Furthermore, there is an opportunity to fill data gaps by calculating these from existing census data. For example, the core census protocol of the Forest Global Earth Observatory  
125 [ForestGEO; REFS] collects the data required to calculate standing dead wood, woody mortality, and delta.agb, but these have

not been calculated and reported for all sites for which the appropriate number of censuses are available (n=1 for standing dead wood, n=2 for woody mortality and delta.agb) [but see REFS].

A universal challenge in estimating biomass (living or dead) from forest census data is applying appropriate allometries to convert DBH measurements to biomass. *(Camille/Helene can write this paragraph easily.)*

## 130 6.2 Data reporting needs

We recommend that, unless they have some specific reason to do otherwise, researchers calculate and report the values according to IPCC standards:

- adopt common standards for variables like min diameter of deadwood, select soil sampling increments to include a cutoff at 30.
- 135 – report 95% CIs, SE, or STD and n
- report C variables in article text, table, or SI table. EFDB cannot accept data digitized from figures
- present calculations of all variables that would be useful to IPCC. EFDB requires that data in the database be presented in the original article, and as such cannot accept subsequent calculations. For example, if aboveground biomass and total biomass are presented, but root biomass is not presented, root biomass cannot be subsequently calculated and sent to
- 140 EFDB. Similarly, fine and coarse root biomass can't be summed; soil carbon can't be summed across depth increments, etc.

For data synthesis projects, compilation can only be useful to the EFDB if they include all the required, along with transparent description on the methodology applied to derive emission factors (or have a brief description and a reference to the original source) and the original emission factor values are present (not modified/rounded).

- 145 **Contributing data to ForC and/or EFDB directly will ensure its broader impact.** The latter is more efficient for getting data to EFDB, but does not get the data into ForC, where it can be more broadly useful—for example, being used for basic science (e.g., Banbury Morgan et al., 2021; Anderson-Teixeira et al., 2021) or model benchmarking (Fer et al., 2021).

## 6.3 Database needs

- There are plenty of relevant, published data that are not included in ForC. Systematic review of the literature could vastly
- 150 improve data coverage. *(There are some efforts underway, including a few that Susan can specify.)*

## 6.4 IPCC

An important challenge is that forests are changing rapidly, and data collected a decade ago may no longer be relevant, particularly in the cases of C increments and fluxes.

Remote sensing biomass estimates include standing dead wood (Duncanson and MANY\_MORE, 2021).



*Code and data availability.* use this to add a statement when having data sets and software code available

*Author contributions.* (fill this in)

*Competing interests.* The authors declare no competing interests.

160 *Acknowledgements.* Thank you to all researchers who collected and published the data contained in ForC, and to all research assistants and collaborators who have helped to build the database. Funding for this study was provided by Bezos Earth Fund to The Nature Conservancy, the Institute for Global Environmental Strategies, WLS(?)



## References

- Anderson-Teixeira, K. J. and Belair, E. P.: Effective Forest-Based Climate Change Mitigation Requires Our Best Science, *Global Change Biology*, 28, 1200–1203, <https://doi.org/10.1111/gcb.16008>, 2022.
- 165 Anderson-Teixeira, K. J., Wang, M. M. H., McGarvey, J. C., Herrmann, V., Tepley, A. J., Bond-Lamberty, B., and LeBauer, D. S.: ForC : A Global Database of Forest Carbon Stocks and Fluxes, *Ecology*, 99, 1507–1507, <https://doi.org/10.1002/ecy.2229>, 2018.
- Anderson-Teixeira, K. J., Herrmann, V., Morgan, R. B., Bond-Lamberty, B., Cook-Patton, S. C., Ferson, A. E., Muller-Landau, H. C., and Wang, M. M. H.: Carbon Cycling in Mature and Regrowth Forests Globally, *Environmental Research Letters*, 16, 053 009, <https://doi.org/10.1088/1748-9326/abed01>, 2021.
- 170 Banbury Morgan, R., Herrmann, V., Kunert, N., Bond-Lamberty, B., Muller-Landau, H. C., and Anderson-Teixeira, K. J.: Global Patterns of Forest Autotrophic Carbon Fluxes, *Global Change Biology*, 27, 2840–2855, <https://doi.org/10.1111/gcb.15574>, 2021.
- Cook-Patton, S. C., Leavitt, S. M., Gibbs, D., Harris, N. L., Lister, K., Anderson-Teixeira, K. J., Briggs, R. D., Chazdon, R. L., Crowther, T. W., Ellis, P. W., Griscom, H. P., Herrmann, V., Holl, K. D., Houghton, R. A., Larrosa, C., Lomax, G., Lucas, R., Madsen, P., Malhi, Y., Paquette, A., Parker, J. D., Paul, K., Routh, D., Roxburgh, S., Saatchi, S., van den Hoogen, J., Walker, W. S., Wheeler, C. E., Wood, S. A., Xu, L., and Griscom, B. W.: Mapping Carbon Accumulation Potential from Global Natural Forest Regrowth, *Nature*, 585, 545–550, <https://doi.org/10.1038/s41586-020-2686-x>, 2020.
- 175 Cuni-Sanchez, A., Sullivan, M. J. P., Platts, P. J., Lewis, S. L., Marchant, R., Imani, G., Hubau, W., Abiem, I., Adhikari, H., Albrecht, T., Altman, J., Amani, C., Aneseyee, A. B., Avitabile, V., Banin, L., Baturike, R., Bauters, M., Beeckman, H., Begne, S. K., Bennett, A. C., Bitariho, R., Boeckx, P., Bogaert, J., Bräuning, A., Bulonvu, F., Burgess, N. D., Calders, K., Chapman, C., Chapman, H., Comiskey, J., de Haulleville, T., Decuyper, M., DeVries, B., Dolezal, J., Droissart, V., Ewango, C., Feyera, S., Gebrekirstos, A., Gereau, R., Gilpin, M., Hakizimana, D., Hall, J., Hamilton, A., Hardy, O., Hart, T., Heiskanen, J., Hemp, A., Herold, M., Hiltner, U., Horak, D., Kamdem, M.-N., Kayijamahe, C., Kenfack, D., Kinyanjui, M. J., Klein, J., Lisingo, J., Lovett, J., Lung, M., Makana, J.-R., Malhi, Y., Marshall, A., Martin, E. H., Mitchard, E. T. A., Morel, A., Mukendi, J. T., Muller, T., Nchu, F., Nyirambangutse, B., Okello, J., Peh, K. S.-H., Pellikka, P., Phillips, O. L., Plumptre, A., Qie, L., Rovero, F., Sainge, M. N., Schmitt, C. B., Sedlacek, O., Ngute, A. S. K., Sheil, D., Sheleme, D., Simegn, T. Y., Simo-Droissart, M., Sonké, B., Soromessa, T., Sunderland, T., Svoboda, M., Taedoumg, H., Taplin, J., Taylor, D., Thomas, S. C., Timberlake, J., Tuagben, D., Umunay, P., Uzabaho, E., Verbeeck, H., Vleminckx, J., Wallin, G., Wheeler, C., Willcock, S., Woods, J. T., and Zibera, E.: High Aboveground Carbon Stock of African Tropical Montane Forests, *Nature*, 596, 536–542, <https://doi.org/10.1038/s41586-021-03728-4>, 2021.
- 180 Duncanson, L. and MANY\_MORE: Aboveground Woody Biomass Product Validation Good Practices Protocol, <https://doi.org/10.5067/DOC/CEOSWGCV/LPV/AGB.001>, 2021.
- 190 Fer, I., Gardella, A. K., Shiklomanov, A. N., Campbell, E. E., Cowdery, E. M., Kauwe, M. G. D., Desai, A., Duveneck, M. J., Fisher, J. B., Haynes, K. D., Hoffman, F. M., Johnston, M. R., Kooper, R., LeBauer, D. S., Mantooth, J., Parton, W. J., Poulter, B., Quaife, T., Raiho, A., Schaefer, K., Serbin, S. P., Simkins, J., Wilcox, K. R., Viskari, T., and Dietze, M. C.: Beyond Ecosystem Modeling: A Roadmap to Community Cyberinfrastructure for Ecological Data-Model Integration, *Global Change Biology*, 27, 13–26, <https://doi.org/10.1111/gcb.15409>, 2021.
- 195 Grassi, G., House, J., Dentener, F., Federici, S., den Elzen, M., and Penman, J.: The Key Role of Forests in Meeting Climate Targets Requires Science for Credible Mitigation, *Nature Climate Change*, 7, 220–226, <https://doi.org/10.1038/nclimate3227>, 2017.

- IPCC: Agriculture, Forestry, and Other Land Use, in: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, edited by Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K., Institute for Global Environmental Strategies, Hayama, Japan, 2006.
- 200 IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, IPCC, Switzerland, 2019a.
- IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Tech. rep., 2019b.
- Ogle, S. M.: Delineating Managed Land for Reporting National Greenhouse Gas Emissions and Removals to the United Nations Framework Convention on Climate Change, p. 13, 2018.
- Requena Suarez, D., Rozendaal, D. M. A., Sy, V. D., Phillips, O. L., Alvarez-Dávila, E., Anderson-Teixeira, K., Araujo-Murakami, A.,
- 205 Arroyo, L., Baker, T. R., Bongers, F., Brien, R. J. W., Carter, S., Cook-Patton, S. C., Feldpausch, T. R., Griscom, B. W., Harris, N., Hérault, B., Coronado, E. N. H., Leavitt, S. M., Lewis, S. L., Marimon, B. S., Mendoza, A. M., N'dja, J. K., N'Guessan, A. E., Poorter, L., Qie, L., Rutishauser, E., Sist, P., Sonké, B., Sullivan, M. J. P., Vilanova, E., Wang, M. M. H., Martius, C., and Herold, M.: Estimating Aboveground Net Biomass Change for Tropical and Subtropical Forests: Refinement of IPCC Default Rates Using Forest Plot Data, *Global Change Biology*, 25, 3609–3624, <https://doi.org/10.1111/gcb.14767>, 2019.
- 210 UNFCCC: Adoption of the Paris Agreement, 2015.