Informing IPCC accounting of forest carbon using the global forest carbon database (ForC v4.0)

Madison Williams¹, Valentine Herrmann¹, Rebecca Banbury Morgan^{1,2}, Ben Bond-Lamberty³, Susan Cook-Patton⁴, Helene Muller-Landau⁵, Camille Piponiot⁶, Teagan Rogers¹, and Kristina J. Anderson-Teixeira^{1,5}*

Correspondence: Kristina J. Anderson-Teixeira (teixeirak@si.edu)

Abstract. The abstract goes here. It can also be on *multiple lines*.

Copyright statement. The author's copyright for this publication is transferred to institution/company.

Important: Always double-check with the official manuscript preparation guidelines at https://publications.copernicus.org/for_authors/manuscript_preparation.html, especially the sections "Technical instructions for LaTeX" and "Manuscript composition". Please contact Daniel Nüst, daniel.nuest@uni-muenster.de, with any problems.

1 Introduction

As we seek to mitigate climate change (UNFCCC, 2015), forests are critical. (See intros to other papers for relevant refs) Under Paris Agreement, ##% of Nationally Determined Contributions relate to forests (Grassi et al., 2017).

The International Panel on Climate Change (IPCC) provides guidance for national greenhouse gas (GHG) inventories for reporting to the United Nations Framework Convention on Climate Change (UNFCCC, (REFS for older guidelines), IPCC, 2019a). Under this guidance, GHG inventories include all managed land, including most of the world's forest land (Ogle, 2018). The IPCC inventory guidelines include specific instructions for accounting for forest land ... (IPCC, 2006, 2019b). This guidance has improved as more of the relevant underlying data has become available (Requena Suarez et al., 2019), but there remains room for continuous improvement as the science advances. For example, Cook-Patton et al. (2020) (summarize IPCC comparison results). Moreover, it is useful for those compiling national greenhouse gas inventories to have access to locally-specific information, when available. To improve the data accessible for C accounting, the IPCC created the Emission Factor Database (EFDB; https://www.ipcc-nggip.iges.or.jp/EFDB/main.php), which is intended as a recognized library of emission factors and other parameters that can be used for estimating greenhouse gas emissions and removals.

¹Center for Conservation Ecology, Smithsonian Conservation Biology Institute, Front Royal, VA, United States

²School of Geography, University of Leeds, Leeds, UK

³Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, United States

⁴The Nature Conservancy; Arlington VA 22203, USA

⁵Forest Global Earth Observatory, Smithsonian Tropical Research Institute, Panama, Republic of Panama

⁶CIRAD, Montpellier, France

The Global Forest Carbon Database, ForC, is the largest collection of published estimates of forest carbon stocks, increments, and annual fluxes (Anderson-Teixeira et al., 2018, 2021). (add stats/ details, maybe record of how ForC has grown over time) As such, ForC is positioned to improve forest C accounting through the transfer of data to EFDB. The purpose of this publication is to document that process and provide recommendations for future improvements.

Here, we (1) review IPCC definitions of relevant carbon stocks and increments (2) describe mapping of ForC to IPCC's EFDB, (3) describe updates to ForC (ForC v4.0), (4) summarize the data in ForC that's relevant to EFDB, identifying gaps, and (5) provide recommendations for improving data collection, analysis, database, and accounting.

2 IPCC definitions of carbon stocks and incremenets

For quantifying forest role in global C cycle, we ultimately care about: (1) C stocks – stores of C that would be vulnerable to release to the atmosphere upon land use change (2) C increments – changes in those C stocks.

2.1 Carbon stocks

Forest ecosystem C stocks may be parsed into pools in various ways. IPCC parses into biomass (aboveground and below-ground), dead organic matter (dead wood and litter), and soil organic matter (Table @ref(table_variables)). Quantifying these requires a one-time measurement.

2.1.1 Biomass

Biomass includes living vegetation, above- and below-ground.

The IPCC defines aboveground biomass as "all biomass of living vegetation, both woody and herbaceous, above the soil including stems, stumps, branches, bark, seeds, and foliage" [].

Belowground biomass is defined as "all biomass of live roots" [].

2.1.2 Dead Organic Matter

Dead organic matter includes all non-living biomass that is not within the mineral soil layer and smaller than the litter size threshold.

Dead wood is defined as...

Litter is defined as including " all non-living biomass with a diameter less than a minimum diameter chosen by the country (for example 10 cm), lying dead, in various states of decomposition above the mineral or organic soil. This includes litter (OL), fumic (OF), and humic (OH) layers. Live fine roots (of less than the suggested diameter limit for belowground biomass) are included in litter where they cannot be distinguished from it empirically." (2003 IPCc GPG for LULUCF (https://www.ipccnggip.iges.or.jp/public/gpglulucf/gpglulucf files/Glossary Acronyms BasicInfo/Glossary.pdf)

Table 1. Variables with definitions and measurement methods. Definitions from IPCC Table 1.1. (See Table 1.1 in IPCC guidance).

pool	definition	major sources of estimate	IPCC guidance
		variation	
aboveground biomass	all biomass of living	allometry, min dbh	acceptable to exclude
	vegetation, both woody and		understory
	herbaceous, above the soil		
belowground biomass	all biomass of live roots	allometry, min dbh, assumed	fine roots may be excluded
		ratio of belowground to	when they cannot be
		aboveground biomass (IPCC	distinguished empirically from
		table 4.4)	soil organic matter or litter
dead wood	all non-living woody biomass	min dbh,	default min dbh = 10cm, but
	not contained in the litter,		may be chosen by country
	either standing, lying on the		
	ground, or in the soil		
litter	all non-living biomass with a	min dbh for dead wood,	includes entire O horizon
	size greater than the limit for		
	soil organic matter and less		
	than the minimum diameter		
	chosen for dead wood, lying		
	dead, in various states of		
	decomposition above or within		
	the mineral or organic soil		
soil organic matter	organic carbon in mineral soils	sampling depth	default sampling depth = 30cm,
	to a specified depth		but may be chosen by country

2.1.3 Soil Organic Matter

Soil organic matter is defined as...

2.2 Carbon increments

C increments are defined as the change over time, in annual increments, in each C pool. These may be estimated as the difference between C stocks at two time points, or as the difference between inputs and outputs to the pool (i.e., fluxes). Quantifying these requires at least two measurements. (*But, Can carbon increments be inferred from a single measure and a known age (i.e., the approach we used in GROA)?*) Fluxes are the inputs and outputs to each pool.

3 Mapping ForC to EFDB




Figure 1. Schematic illustrating the carbon pools quantified under IPCC accounting; ForC variables corresponding to the stock, increment, influx and outflux; and relationships among them. In many cases, the match of ForC variables to IPCC criteria depends upon measurement protocols (e.g., minimum DBH). Additional caveats are as follows: 1- assumes minimal non-respiratory C fluxes,... (Chapin et al. 2006); 2- assumes that change in foliage biomass is negligible (see note 7); 3- incomplete: excludes large branch fall; also, under IPCC definitions, outflux from aboveground biomass should include all sizes, influx to deadwood should include only above the minimum diameter chosen for dead wood; 4- incomplete: excludes belowground components; 5-incomplete: excludes breakage into pieces less than dead wood threshold size; 6-incomplete: excludes woody mortality of stems <10 cm DBH, decomposition of dead wood (aboveground and coarse roots) into sizes classified as litter; 7- measurements often limited to decomposition of relatively fine litter and may exclude branches and stems below the dead wood size threshold and/or the more decomposed layers of the O horizon; 8 - foliage production is generally measured by collecting leaf-fall, a method that assumes that the influx = outflux (foliage biomass is roughly constant year-to-year).

Table 2: Mapping of ForC fields to EFDB. See footnotes at end of table (still need to be properly inserted).

ForC table	ForC field	EFDB field	Usage	Required
Measurements	measurement.ID	Other Properties	direct mapping	(no)
	dominant.life.form	1996 Source/Sink	used to determine	yes
		Categories, 2006	land subcategories	
		Source/Sink	(see defin-	
		Categories	ing_land_subcategory.r	nd)
	stand.age	1996 Source/Sink	used to determine	(yes)
		Categories, 2006	land subcategories	
		Source/Sink	(see defin-	
		Categories,	ing_land_subcategory.r	nd),
		Parameters/	directly listed in	,,
		Conditions	Parameters/	
			Conditions	
	dominant.veg,	Parameters/	direct mapping/	no
	veg.notes, min.dbh	Conditions	linking to	
	-		dominant.veg	
			description	
	variable.name	-	link to variable info in	yes
			ForC_variables table	
	date / start.date,	Other Properties	direct mapping	no
	end.date			
	mean	Value	direct mapping	yes
	mean.in.original.units	Value in Common	direct mapping	yes
		Units		
	original.units	Common Unit	direct mapping	yes
	lower95%CI, upper	Lower Confidence	direct or calculated	(yes)
	95%CI, se, sd and n	Limit, Upper		
		Confidence Limit		
	depth, covariate_1,	Other Properties	direct mapping	no
	cov_1.value,			
	covariate_2,			
	cov_2.value	G	1: 1 . 1:	
	allometry_1,	Comments from Data	link to biomass	no
	allometry_2	Provider	allometry source,	
			when provided	

Table 2: Mapping of ForC fields to EFDB. See footnotes at end of table (still need to be properly inserted). (continued)

ForC table	ForC field	EFDB field	Usage	Required
	data.location.within.source		confirm that data	yes
			weren't digitized,	
			facilitate finding data	
			in original publication	
	ForC.investigator	Data Provider, Data	link to Data Provider,	yes
		Provider Contact	Data Provider	
			Contact info	
Sites	site.ID, sites.sitename	Other Properties	direct mapping	(no)
	lat, lon	Region/Regional	direct mapping; used	(no)
		conditions	to extract continent,	
			Koeppen, and	
			FAO.ecozone	
	country, state, city,	Region/Regional	direct mapping	no
	masl, mat, map	conditions		
	continent, Koeppen	Region/Regional	direct mapping	auto
		conditions	1	
	soil.texture, sand, silt,	Parameters/	direct mapping	no
	clay,	Conditions		
	soil.classification FAO.ecozone	Parameters/	direct mapping	auto
	TAO.CCOZOIIC	Conditions	uncet mapping	auto
History	date, hist.cat,	1996 Source/Sink	used to determine	(yes)/no**
•	hist.type	Categories, 2006	distmrs.type for	
	• •	Source/Sink	Source/Sink	
		Categories,	Categories, generate	
		Abatement/Control	list of events for	
		technologies	Abatement/Control	
			technologies	
	plot.area	Other Properties	direct mapping	no
Plots	plot.ID, plot.name	Other Properties	direct mapping	(no)
	distmrs.type	1996 Source/Sink	used to determine	auto
		Categories, 2006	land subcategories	
		Source/Sink	(see defin-	
			`	nd)
		Categories	ing_land_subcategory.r	nd)

Table 2: Mapping of ForC fields to EFDB. See footnotes at end of table (still need to be properly inserted). (continued)

ForC table	ForC field	EFDB field	Usage	Required
	distmrs.type,	Other Properties	direct mapping	auto
	distmrs.year,			
	regrowth.type,			
	regrowth.year			
PFT	description	Parameters/	direct mapping	auto
		Conditions		
variables	variable.type	Gases	For stocks in unit of	auto
			organic matter, gases	
			include CO2, CO,	
			CH4, NO, NO2,	
			N2O. For increments,	
			fluxes, and stocks in	
			units of C, gases	
			includes only CO2.	
	variable.name	C pool, Equation	link to C pool,	auto
			Equation	
	description	Description	direct mapping	auto
	extended.description	Other Properties	direct mapping	auto
	units	Unit (ID)	link to IPCC units	auto
Citations	citation.citation	Full Technical	direct mapping	yes/auto
		Reference		
	citation.language	Reference Language	direct mapping	yes/auto
	citation.url	URL	direct mapping	no/auto
	citation.abstract	Abstract in English	direct mapping	no/auto
	source.type	Source of Data	direct mapping	yes

^{&#}x27;Required' field indicates whether the field is required by EFDB: yes = value required; (yes) = input required, missing value acceptable if not reported; auto = present within ForC infrasructure, and therefore will always be exported to EFDB; (no) = not required for EFDB, but required for ForC and therefore will always be exported to EFDB; no = not required, but exported to EFDB when a value is present.

55

^{** &#}x27;(yes)' for most recent severe disturbance; 'no' for other history events

60 3.1 Carbon cycle variables

Mapping of variables is shown in Fig. 1

Define relationship among NEE, NEP, and delta.C., especially noting role of harvest (Chapin et al., 2006).

3.2 Land use categories

Documented at https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB_mapping/defining_land_subcategory.md, https://github.com/forc-db/IPCC-EFDB-integration/blob/main/doc/ForC-EFDB_mapping/IPCC_LandUse_mapping.csv, and in issue #8.

The UNFCCC requires GHG reporting for all managed lands in a country, where management is defined as "human interventions and practices have been applied to perform production, ecological or social functions" [IPCC 2006 full report REF]. This definition is applied differently across countries, and is not clearly defined by the majority of governments (Ogle, 2018). Given this, and because the IPCC definition of management does not necessarily match that which would be reported in scientific publications and hence in ForC, we do not transfer any classification of land management status from ForC to the EFDB, but do provide auxiliary info that may be useful in making this determination (e.g., geographical location).

4 Updates to ForC (ForC v4.0)

To support export of data to EFDB, and to improve the overall quality of the ForC database, we defined ## new variables, implemented some modest restructuring, resolved duplicate records, and conducted quality control. This section describes changes relative to ForC v2.0 (Anderson-Teixeira et al., 2018).

4.1 Defining new variables

We added eleven increment variables to the set of named and defined variables (or 22, counting _OM and _C versions), which previously included only one (aboveground biomass increment, *delta.agb*). (https://github.com/forc-db/IPCC-EFDB-integration/issues/6) These are directly related to C stocks as previously defined in ForC, with "delta." added in front of the variable name.

Although these variables currently lack records, the structure exists such that records can be populated over time.

To provide better definition of the previously existing variable *organic.layer*, which has a nebulous definition that reflects the varied definitions adopted by original studies, we added two clearly defined variables: *litter* (relatively undecomposed plant material/ OL horizon), and *O.horizon* (entire O-horizon, including *litter* (OL)).

4.2 ForC restructuring

Table 3: Table of changes to ForC fields.

Table	Column	Description	Changes	Motivation
Sites	coordinates.precision	Precision of	field added	allow identification of
		geographic		records with poor
		coordinates, as		coordinate precision
		reported by source or		
		estimated from maps.		
Measurements	data.location.within.sou	rdeocation of data	field added	facilitate review,
		within the source		ensure traceability
		listed in citation.ID.		
	sd, se, lower95%CI,	Standard deviation,	replaces 'stat' and	cleaner format; ability
	upper 95%CI	standard error, and	'stat.name'	to handle
		lower and upper 95		assymetrical 95
		percent confidence		percent confidence
		intvervals,		intervals
		respectively.		
	mean.in.original.units,	mean value and units	fields added	provide IPCC with
	original.units	presented in original		original units, reduce
		publication		errors/improve
				reproducibility
	C.conversion.factor	Assumed/ measured	field added	track units
		C content of organic		conversion, allow
		matter used to convert		back-calculation of
		organic matter to C.		OM if conversion
				factor deemed
				inappropriate
PFT	description	Definition of the	field added	
		pftcode at the		
		community level.		
		Differs from		
		individual level in		
		that properly		
		describes mixed plant		
		functional types.		

Table 3: Table of changes to ForC fields. (continued)

Table	Column	Description	Changes	Motivation
	description.individual	Definition of the	field name change	
		pftcode at the	(previously	
		individual plant level.	'description')	
Citations	citation.citation	Full citation. Most of	field added	field required by
		these records are		IPCC
		automatically		
		generated in R based		
		upon DOI lookup.		
	citation.language	Language of original	field added	field required by
		publication,		IPCC
		automatically		
		generated based on		
		the title and abstract,		
		with some manual		
		entries and		
		corrections.		
	citation.url	URL of original	field added	field required by
		publication, generally		IPCC
		retrieved		
		automatically via		
		URL lookup.		
	citation.abstract	Abstract, generally	field added	field required by
		retrieved		IPCC
		automatically via		
		DOI lookup.		
	source.type	citation source type	field added	field required by
				IPCC
	pdf.in.repository	Indicates whether pdf	field added	housekeeping
		of original study has		
		been retrieved and		
		saved in ForC's		
		reference repository		

Table 3: Table of changes to ForC fields. (continued)

Table	Column	Description	Changes	Motivation
	EFDB.ready	Indicates whether	field added	housekeeping
		data have been		
		checked for export to)	
		EFDB.		

4.3 Quality control measures

Prior to releasing ForC v4.0, we executed several quality control measures. First, we implemented a system of continuous integration using GitHub Actions (*sensu* Kim et al. in prep) to run some automatic checks any time the master data files are updated. Second, to improve information on geographic coordinates, we flagged and reviewed records with suspected low precision (*Issue #29*)[https://github.com/forc-db/ForC/issues/229]. Third, to identify erroneous climate data... (*Issue #212*)[https://github.com/forc-db/ForC/issues/212].

4.4 Resolving duplicates

5 Results

95 figure: map of relevant ForC data with underlying FAO ecozones

(summarize the data in ForC that's relevant to EFDB, identifying gaps)

dead wood and litter comparisons will be particularly interesting, as IPCC values are based on just a handful of references for each climate zone (table 2.2 in 2019 guidelines)

6 Recommendations

100 (strongly flag both useful variables that the EFDB does not track and useful variables that papers fail to include that EFDB needs)

6.1 Data collection and analysis needs

(Paragraph highlighting important gaps in variables / regions)

Several variables of value to IPCC, including standing dead wood, woody mortality, delta.agb, are not calculated and presented as frequently as are AGB and ANPP_woody, even though they can readily be derived from the same census data. We recommend that researchers calculate and report these, as specified below. Furthermore, there is an opportunity to fill data gaps by calculating these from existing census data. For example, the core census protocol of the Forest Global Earth Observatory [ForestGEO; REFS] collects the data required to calculate standing dead wood, woody mortality, and delta.agb, but these have not been calculated and reported for all sites for which the appropriate number of censuses are available (n=1 for standing dead wood, n=2 for woody mortality and delta.agb) [but see REFS].

A universal challenge in estimating biomass (living or dead) from forest census data is applying appropriate allometries to convert DBH measurements to biomass. (Camille/Helene can write this paragraph easily.)

6.2 Data reporting needs

We recommend that, unless they have some specific reason to do otherwise, researchers calculate and report the values according to IPCC standards:

- adopt common standards for variables like min diameter of deadwood, select soil sampling increments to include a cutoff at 30.
- report 95% CIs, SE, or STD and n
- report C variables in article text, table, or SI table. EFDB cannot accept data digitized from figures
- For data synthesis projects, compilation can only be useful to the EFDB if they include all the required, along with transparent description on the methodology applied to derive emission factors (or have a brief description and a reference to the original source) and the original emission factor values are present (not modified/rounded).

Contributing data to ForC and/or EFDB directly will ensure its broader impact. The latter is more efficient for getting data to EFDB, but does not get the data into ForC, where it can be more broadly useful—for example, being used for basic science (e.g., Banbury Morgan et al., 2021; Anderson-Teixeira et al., 2021) or model benchmarking (Fer et al., 2021).

6.3 Database needs

There are plenty of relevant, published data that are not included in ForC. Systematic review of the literature could vastly improve data coverage. (*There are some efforts underway, including a few that Susan can specify.*)

6.4 IPCC

125

An important challenge is that forests are changing rapidly, and data collected a decaade ago may no longer be relevant, particularly in the cases of C increments and fluxes.

Remote sensing biomass estimates include standing dead wood (Duncanson and MANY MORE, 2021).

7 Conclusions

The conclusion goes here. You can modify the section name with \conclusions [modified heading if necessary].

135	Code and data availability. use this to add a statement when having data sets and software code available
	Author contributions. (fill this in)
	Competing interests. The authors declare no competing interests.
	Acknowledgements. Thank you to all researchers who collected and published the data contained in ForC, and to all research assistants and collaborators who have helped to build the database. Funding for this study was provided by Bezos Earth Fund to The Nature Conservancy.

 $140 \quad \text{the Institute for Global Environmental Strategies, WLS(?)} \\$

References

- Anderson-Teixeira, K. J., Wang, M. M. H., McGarvey, J. C., Herrmann, V., Tepley, A. J., Bond-Lamberty, B., and LeBauer, D. S.: For C: A Global Database of Forest Carbon Stocks and Fluxes, Ecology, 99, 1507–1507, https://doi.org/10.1002/ecy.2229, 2018.
- Anderson-Teixeira, K. J., Herrmann, V., Morgan, R. B., Bond-Lamberty, B., Cook-Patton, S. C., Ferson, A. E., Muller-Landau, H. C., and Wang, M. M. H.: Carbon Cycling in Mature and Regrowth Forests Globally, Environmental Research Letters, 16, 053 009, https://doi.org/10.1088/1748-9326/abed01, 2021.
 - Banbury Morgan, R., Herrmann, V., Kunert, N., Bond-Lamberty, B., Muller-Landau, H. C., and Anderson-Teixeira, K. J.: Global Patterns of Forest Autotrophic Carbon Fluxes, Global Change Biology, 27, 2840–2855, https://doi.org/10.1111/gcb.15574, 2021.
- Chapin, F., Woodwell, G., Randerson, J., Rastetter, E., Lovett, G., Baldocchi, D., Clark, D., Harmon, M., Schimel, D., Valentini, R., Wirth,
 C., Aber, J., Cole, J., Goulden, M., Harden, J., Heimann, M., Howarth, R., Matson, P., McGuire, A., Melillo, J., Mooney, H., Neff,
 J., Houghton, R., Pace, M., Ryan, M., Running, S., Sala, O., Schlesinger, W., and Schulze, E.: Reconciling Carbon-Cycle Concepts,
 Terminology, and Methods, Ecosystems, 9, 1041–1050, 2006.
 - Cook-Patton, S. C., Leavitt, S. M., Gibbs, D., Harris, N. L., Lister, K., Anderson-Teixeira, K. J., Briggs, R. D., Chazdon, R. L., Crowther, T. W., Ellis, P. W., Griscom, H. P., Herrmann, V., Holl, K. D., Houghton, R. A., Larrosa, C., Lomax, G., Lucas, R., Madsen, P., Malhi,
- Y., Paquette, A., Parker, J. D., Paul, K., Routh, D., Roxburgh, S., Saatchi, S., van den Hoogen, J., Walker, W. S., Wheeler, C. E., Wood, S. A., Xu, L., and Griscom, B. W.: Mapping Carbon Accumulation Potential from Global Natural Forest Regrowth, Nature, 585, 545–550, https://doi.org/10.1038/s41586-020-2686-x, 2020.
 - Duncanson, L. and MANY_MORE: Aboveground Woody Biomass Product Validation Good Practices Protocol, https://doi.org/10.5067/DOC/CEOSWGCV/LPV/AGB.001, 2021.
- Fer, I., Gardella, A. K., Shiklomanov, A. N., Campbell, E. E., Cowdery, E. M., Kauwe, M. G. D., Desai, A., Duveneck, M. J., Fisher, J. B., Haynes, K. D., Hoffman, F. M., Johnston, M. R., Kooper, R., LeBauer, D. S., Mantooth, J., Parton, W. J., Poulter, B., Quaife, T., Raiho, A., Schaefer, K., Serbin, S. P., Simkins, J., Wilcox, K. R., Viskari, T., and Dietze, M. C.: Beyond Ecosystem Modeling: A Roadmap to Community Cyberinfrastructure for Ecological Data-Model Integration, Global Change Biology, 27, 13–26, https://doi.org/10.1111/gcb.15409, 2021.
- Grassi, G., House, J., Dentener, F., Federici, S., den Elzen, M., and Penman, J.: The Key Role of Forests in Meeting Climate Targets Requires Science for Credible Mitigation, Nature Climate Change, 7, 220–226, https://doi.org/10.1038/nclimate3227, 2017.
 - IPCC: Agriculture, Forestry, and Other Land Use, in: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, edited by Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K., Institute for Global Environmental Strategies, Hayama, Japan, 2006.
 - IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Tech. rep., 2019a.
- 170 IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, IPCC, Switzerland, 2019b.
 - Ogle, S. M.: Delineating Managed Land for Reporting National Greenhouse Gas Emissions and Removals to the United Nations Framework Convention on Climate Change, p. 13, 2018.
 - Requena Suarez, D., Rozendaal, D. M. A., Sy, V. D., Phillips, O. L., Alvarez-Dávila, E., Anderson-Teixeira, K., Araujo-Murakami, A., Arroyo, L., Baker, T. R., Bongers, F., Brienen, R. J. W., Carter, S., Cook-Patton, S. C., Feldpausch, T. R., Griscom, B. W., Harris, N.,
- Hérault, B., Coronado, E. N. H., Leavitt, S. M., Lewis, S. L., Marimon, B. S., Mendoza, A. M., N'dja, J. K., N'Guessan, A. E., Poorter, L., Qie, L., Rutishauser, E., Sist, P., Sonké, B., Sullivan, M. J. P., Vilanova, E., Wang, M. M. H., Martius, C., and Herold, M.: Estimating

Aboveground Net Biomass Change for Tropical and Subtropical Forests: Refinement of IPCC Default Rates Using Forest Plot Data, Global Change Biology, 25, 3609–3624, https://doi.org/10.1111/gcb.14767, 2019.

UNFCCC: Adoption of the Paris Agreement, 2015.