

8. Worksheet: Among Site (Beta) Diversity – Part 2

Brooke Peckenpaugh; Z620: Quantitative Biodiversity, Indiana University

13 February, 2019

OVERVIEW

In this worksheet, we continue to explore concepts, statistics, and visualizations related to β -diversity. Now that you know how to formally quantify β -diversity, we will learn how to test hypotheses about β -diversity using multivariate statistics.

Directions:

1. In the Markdown version of this document in your cloned repo, change “Student Name” on line 3 (above) with your name.
2. Complete as much of the worksheet as possible during class.
3. Use the handout as a guide; it contains a more complete description of data sets along with examples of proper scripting needed to carry out the exercises.
4. Answer questions in the worksheet. Space for your answers is provided in this document and is indicated by the “>” character. If you need a second paragraph be sure to start the first line with “>”. You should notice that the answer is highlighted in green by RStudio (color may vary if you changed the editor theme).
5. Before you leave the classroom today, it is *imperative* that you **push** this file to your GitHub repo, at whatever stage you are. This will enable you to pull your work onto your own computer.
6. When you have completed the worksheet, **Knit** the text and code into a single PDF file by pressing the **Knit** button in the RStudio scripting panel. This will save the PDF output in your ‘8.BetaDiversity’ folder.
7. After Knitting, please submit the worksheet by making a **push** to your GitHub repo and then create a **pull request** via GitHub. Your pull request should include this file (**8.BetaDiversity__2__Worksheet.Rmd**) with all code blocks filled out and questions answered) and the PDF output of Knitr (**8.BetaDiversity__2__Worksheet.pdf**).

The completed exercise is due on **Wednesday, February 13th, 2019 before 12:00 PM (noon)**.

1) R SETUP

Typically, the first thing you will do in either an R script or an RMarkdown file is setup your environment. This includes things such as setting the working directory and loading any packages that you will need.

In the R code chunk below, provide the code to:

1. clear your R environment,
2. print your current working directory,
3. set your working directory to your “/8.BetaDiversity” folder, and
4. load the **vegan** R package (be sure to install if needed).

```
rm(list = ls())
getwd()
```

```
## [1] "/Users/brooke/GitHub/QB2019_Peckenpaugh/2.Worksheets/8.BetaDiversity"
```

```

setwd("~/GitHub/QB2019_Peckenpaugh/2.Worksheets/8.BetaDiversity")
package.list <- c('vegan', 'ade4', 'viridis', 'gplots', 'BiodiversityR', 'indicspecies')
for (package in package.list) {
  if (!require(package, character.only=TRUE, quietly=TRUE)) {
    install.packages(package)
    library(package, character.only=TRUE)
  }
}

## This is vegan 2.5-3

##
## Attaching package: 'gplots'

## The following object is masked from 'package:stats':
##
##      lowess

## BiodiversityR 2.11-1: Use command BiodiversityRGUI() to launch the Graphical User Interface;
## to see changes use BiodiversityRGUI(changeLog=TRUE, backward.compatibility.messages=TRUE)

```

2) LOADING DATA

Load dataset

In the R code chunk below, load the `doubs` dataset from the `ade4` package

```

# note, please do not print the dataset when submitting
data(doubs)

```

3) HYPOTHESIS TESTING

A. Multivariate Procedures for Categorical Designs

Earlier work done in the Doubs River suggested that the river has four distinct regions of habitat quality: the first region (sites 1-14) of “high quality”; the second (sites 15 - 19) and fourth (sites 26 - 30) of “moderate quality”; and the third (sites 20 - 25) of “low quality”.

In the code chunk below, test the hypothesis that fish community composition varies with river quality.

1. create a factor vector that categorizes habitat quality in the Doubs River,
2. use the multivariate analyses for categorical predictors to describe how fish community structure relates to habitat quality.

```

quality <- c(rep("HQ", 13), rep("MQ", 5), rep("LQ", 6), rep("MQ", 5))

fish <- doubs$fish
fish <- fish[-8, ]

adonis(fish ~ quality, method = "bray", permutations = 999)

##
## Call:
## adonis(formula = fish ~ quality, permutations = 999, method = "bray")
##
## Permutation: free

```

```

## Number of permutations: 999
##
## Terms added sequentially (first to last)
##
##           Df SumsOfSqs MeanSqs F.Model      R2 Pr(>F)
## quality    2    3.0947 1.54733   10.97 0.45765  0.001 ***
## Residuals 26    3.6674 0.14105         0.54235
## Total     28    6.7621         1.00000
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

indval <- multipatt(fish, cluster = quality, func = "IndVal.g", control = how(nperm=999))
summary(indval)

##
## Multilevel pattern analysis
## -----
##
## Association function: IndVal.g
## Significance level (alpha): 0.05
##
## Total number of species: 27
## Selected number of species: 23
## Number of species associated to 1 group: 1
## Number of species associated to 2 groups: 22
##
## List of species associated to each combination:
##
## Group MQ #sps. 1
##      stat p.value
## Teso 0.686  0.031 *
##
## Group HQ+MQ #sps. 2
##      stat p.value
## Satr 0.860  0.004 **
## Phph 0.859  0.019 *
##
## Group LQ+MQ #sps. 20
##      stat p.value
## Alal 0.935  0.001 ***
## Gogo 0.933  0.001 ***
## Ruru 0.916  0.001 ***
## Legi 0.901  0.001 ***
## Baba 0.895  0.001 ***
## Chna 0.866  0.001 ***
## Spbi 0.866  0.002 **
## Cyca 0.866  0.001 ***
## Acce 0.866  0.001 ***
## Lele 0.863  0.003 **
## Titi 0.853  0.008 **
## Chto 0.829  0.003 **
## Rham 0.829  0.001 ***
## Anan 0.829  0.001 ***
## Eslu 0.827  0.026 *
## Pefl 0.806  0.013 *

```

```

## Blbj 0.791    0.002 **
## Scer 0.766    0.006 **
## Abbr 0.750    0.006 **
## Icme 0.661    0.018 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

fish.rel <- decostand(fish, method = "total")
phi <- multipatt(fish.rel, cluster = quality, func = "r.g", control = how(nperm=999))
summary(phi)

##
## Multilevel pattern analysis
## -----
##
## Association function: r.g
## Significance level (alpha): 0.05
##
## Total number of species: 27
## Selected number of species: 18
## Number of species associated to 1 group: 9
## Number of species associated to 2 groups: 9
##
## List of species associated to each combination:
##
## Group HQ #sps.  3
##      stat p.value
## Phph 0.802    0.001 ***
## Neba 0.734    0.001 ***
## Satr 0.650    0.001 ***
##
## Group LQ #sps.  2
##      stat p.value
## Alal 0.693    0.001 ***
## Ruru 0.473    0.030 *
##
## Group MQ #sps.  4
##      stat p.value
## Anan 0.571    0.006 **
## Spbi 0.557    0.009 **
## Chto 0.542    0.005 **
## Icme 0.475    0.028 *
##
## Group LQ+MQ #sps.  9
##      stat p.value
## Legi 0.658    0.002 **
## Baba 0.645    0.002 **
## Rham 0.600    0.004 **
## Acce 0.594    0.007 **
## Cyca 0.586    0.005 **
## Chna 0.571    0.005 **
## Blbj 0.571    0.010 **
## Gogo 0.523    0.015 *
## Abbr 0.499    0.022 *
## ---

```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Question 1: Based on the PERMANOVA, IndVal, and phi coefficient analyses, what did you learn about the relationship between habitat quality and the fish species composition? Are the different analyses consistent with one another and do they agree with the visualizations (heat maps, cluster dendograms, ordinations) that you created?

Answer 1: Based on the PERMANOVA, habitat quality appears to significantly predict the distribution of fish species. While there are not very many indicator species for only one group of habitat quality, there are several species with high indicator value for low and medium quality sites combined. Overall, quite a few species show preference for habitat quality groups. There are more species that show preference for low/medium quality habitats, though the preference is strongest in the few species that prefer high quality habitat. These analyses seem pretty consistent to me.

B. Multivariate Procedures for Continuous Designs

i. Mantel Test

In the R code chunk below, do the following:

1. create distance matrices for both fish communities and environmental factors, and
2. use a Mantel test to test the hypothesis that fish assemblages are correlated with stream environmental variables.

```
fish.dist <- vegdist(doubs$fish[-8, ], method = "bray")
env.dist <- vegdist(scale(doubs$env[-8, ]), method = "euclid")

mantel(fish.dist, env.dist)
```

```
##
## Mantel statistic based on Pearson's product-moment correlation
##
## Call:
## mantel(xdis = fish.dist, ydis = env.dist)
##
## Mantel statistic r: 0.604
##      Significance: 0.001
##
## Upper quantiles of permutations (null model):
##   90%   95%  97.5%   99%
## 0.107 0.144 0.169 0.210
## Permutation: free
## Number of permutations: 999
```

Question 2: What do the results from our Mantel test suggest about fish diversity and stream environmental conditions? How does this relate to your hypothesis about stream quality influencing fish communities?

Answer 2: The Mantel test suggests that 60.4% of the variation in fish distribution is explained by the stream environmental condition data provided, and this result is statistically significant ($p = 0.001$). I think this evidence supports my hypothesis that stream quality influences fish communities.


ii. Constrained Ordination

In the R code chunk below, do the following:


1. create an environmental matrix of the water chemistry data included in the `doubs` dataset using forward and reverse selection of variables,
2. conduct a redundancy analysis on the fish assemblages of the Doubs River,
3. use a permutation test to determine the significance of the constrained analysis,
4. use a permutation test to determine the correlation of each environmental factor on the constrained axes,
5. calculate the explained variation on the first and second constrained axes,
6. plot the constrained ordination results including labeled points for each site, and
7. add vectors that demonstrate the influence of each environmental factor the constrained ordination.

```
env.chem <- as.matrix(doubs$env[-8, 5:11])

doubs.dbrda <- dbrda(fish.dist ~ ., as.data.frame(env.chem))
ordiplot(doubs.dbrda)
```



```
doubs.dbrda.mod0 <- dbrda(fish.dist ~ 1, as.data.frame(env.chem))
ordiplot(doubs.dbrda.mod0)
```



```
doubs.dbrda.mod1 <- dbrda(fish.dist ~ ., as.data.frame(env.chem))
```

```
doubs.dbrda <- ordiR2step(doubs.dbrda.mod0, doubs.dbrda.mod1, perm.max = 200)
```

```
## Step: R2.adj= 0
## Call: fish.dist ~ 1
##
##               R2.adjusted
## <All variables> 0.53032584
## + oxy          0.27727176
## + nit          0.25755208
## + bdo          0.17477787
## + pho          0.14568614
## + har          0.14174915
## + amm          0.14142804
## <none>         0.00000000
## + pH          -0.01827054
##
##      Df    AIC      F Pr(>F)
## + oxy  1 47.939 11.742 0.002 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.2772718
## Call: fish.dist ~ oxy
##
##               R2.adjusted
## <All variables> 0.5303258
## + bdo          0.4009000
## + amm          0.3474192
## + pho          0.3452702
## + har          0.3331357
## + nit          0.3316120
```

```
## <none>          0.2772718
## + pH           0.2586983
##
##      Df      AIC      F Pr(>F)
## + bdo  1 43.404 6.5716 0.004 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.4009
## Call: fish.dist ~ oxy + bdo
##
##              R2.adjusted
## <All variables> 0.5303258
## + nit          0.4980793
## + har          0.4695121
## <none>         0.4009000
## + pho          0.3938042
## + amm          0.3869134
## + pH           0.3865240
##
##      Df      AIC      F Pr(>F)
## + nit  1 39.134 6.034 0.004 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.4980793
## Call: fish.dist ~ oxy + bdo + nit
##
##              R2.adjusted
## + amm          0.5415705
## <All variables> 0.5303258
## + pho          0.5277128
## + har          0.5218852
## <none>         0.4980793
## + pH           0.4843267
```


```
doubs.dbrda$call
```

```
## dbrda(formula = fish.dist ~ oxy + bdo + nit, data = as.data.frame(env.chem))
```

```
doubs.dbrda$anova
```

```
##              R2.adj Df      AIC      F Pr(>F)
## + oxy          0.27727  1 47.939 11.7421 0.002 **
## + bdo          0.40090  1 43.404  6.5716 0.004 **
## + nit          0.49808  1 39.134  6.0340 0.004 **
## <All variables> 0.53033
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
ordiplot(doubs.dbrda)
```

```
permutest(doubs.dbrda, permutations = 999)
```

```
##
## Permutation test for dbrda under reduced model
##
## Permutation: free
## Number of permutations: 999
##
## Model: dbrda(formula = fish.dist ~ oxy + bdo + nit, data =
## as.data.frame(env.chem))
## Permutation test for all constrained eigenvalues
##      Df Inertia      F Pr(>F)
## Model   3  3.7317 10.262  0.001 ***
## Residual 25  3.0304
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
envfit(doubs.dbrda, env.chem[,c(4,6,7)], perm = 999)
```

```
##
## ***VECTORS
##
##      dbRDA1  dbRDA2      r2 Pr(>r)
## nit  0.87724  0.48005 0.6431  0.001 ***
## oxy -0.82864 -0.55979 0.7656  0.001 ***
## bdo  0.55603  0.83116 0.8939  0.001 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## Permutation: free
## Number of permutations: 999
```


```
dbrda.explainvar1 <- round(doubs.dbrda$CCA$eig[1] /
                           sum(c(doubs.dbrda$CCA$eig, doubs.dbrda$CA$eig)), 3) * 100
dbrda.explainvar2 <- round(doubs.dbrda$CCA$eig[2] /
```

```

sum(c(doubs.dbrda$CCA$eig, doubs.dbrda$CA$eig)), 3) * 100

par(mar = c(5, 5, 4, 4) + 0.1)
plot(scores(doubs.dbrda, display = "wa"), xlim = c(-1.3, 1.1), ylim = c(-1.1, 2.7),
      xlab = paste("dbRDA 1 (", dbrda.explainvar1, "%)", sep = ""),
      ylab = paste("dbRDA 2 (", dbrda.explainvar2, "%)", sep = ""),
      pch = 16, cex = 2.0, type = "n", cex.lab = 1.5, cex.axis = 1.2, axes = FALSE)
axis(side = 1, labels = T, lwd.ticks = 2, cex.axis = 1.2, las = 1)
axis(side = 2, labels = T, lwd.ticks = 2, cex.axis = 1.2, las = 1)
abline(h = 0, v = 0, lty = 3)
box(lwd = 2)
points(scores(doubs.dbrda, display = "wa"),
       pch = 19, cex = 3, bg = "gray", col = "gray")
text(scores(doubs.dbrda, display = "wa"),
     labels = row.names(scores(doubs.dbrda, display = "wa")))
vectors <- scores(doubs.dbrda, display = "bp")
arrows(0, 0, vectors[, 1], vectors[, 2],
      lwd = 2, lty = 1, length = 0.2, col = "red")
text(vectors[, 1], vectors[, 2], pos = 3,
     labels = row.names(vectors))
axis(side = 3, lwd.ticks = 2, cex.axis = 1.2, las = 1, col = "red", lwd = 2.2,
     at = pretty(range(vectors[, 1])) * 2, labels = pretty(range(vectors[, 1])))
axis(side = 4, lwd.ticks = 2, cex.axis = 1.2, las = 1, col = "red", lwd = 2.2,
     at = pretty(range(vectors[, 2])) * 2, labels = pretty(range(vectors[, 2])))

```



Question 3: Based on the constrained ordination, what are the environmental variables (or groups of correlated variables) that seem to be contributing to variation in fish community structure?

Answer 3: Dissolved oxygen, biological demand for oxygen, and nitrates.

iii. Variation Partitioning

In the code chunk below,

1. Create a matrix model of the selected environmental variables,
2. Create a matrix model of the selected PCNM axes,
3. Perform constrained and partial constrained ordinations using the spatial and environmental models you just created,
4. Test the significance of each of your constrained ordinations using permutation tests,
5. Partition the variation among sites into the relative importance of space, environment, spatially structured environment, and residuals,
6. Plot the variation partitioning output to visualize it.

```
doubs.dbrda$anova
```

```
##           R2.adj Df      AIC      F Pr(>F)
## + oxy       0.27727  1 47.939 11.7421  0.002 **
## + bdo       0.40090  1 43.404  6.5716  0.004 **
## + nit       0.49808  1 39.134  6.0340  0.004 **
## <All variables> 0.53033
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

env.mod <- model.matrix(~ oxy + bdo + nit, as.data.frame(env.chem))[, -1]

rs <- rowSums(fish)/sum(fish)
doubs.pcnmw <- pcnm(dist(doubs$xy[-8,]), w = rs, dist.ret = T)
doubs.pcnmw$values > 0
```

```
## [1] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [12] TRUE TRUE TRUE TRUE TRUE TRUE TRUE FALSE FALSE FALSE FALSE FALSE
## [23] FALSE FALSE FALSE FALSE
```

```
fish.db <- vegdist(fish, method = "bray")
doubs.space <- as.data.frame(scores(doubs.pcnmw))
doubs.pcnm.mod0 <- dbrda(fish.db ~ 1, doubs.space)
doubs.pcnm.mod1 <- dbrda(fish.db ~ ., doubs.space)
step.pcnm <- ordiR2step(doubs.pcnm.mod0, doubs.pcnm.mod1, perm.max = 200)
```

```
## Step: R2.adj= 0
## Call: fish.db ~ 1
##
##           R2.adjusted
## <All variables> 0.626011301
## + PCNM2       0.235370423
## + PCNM3       0.078394885
## + PCNM13      0.065305668
## + PCNM5       0.046185074
## + PCNM6       0.032809156
## + PCNM16      0.030486700
## + PCNM14      0.029680999
## + PCNM9       0.020357410
## + PCNM15      0.013632610
## + PCNM8       0.009411968
## + PCNM1       0.003986221
## + PCNM17      0.002415012
## + PCNM10      0.001326442
```

```

## <none>          0.000000000
## + PCNM7         -0.001861430
## + PCNM11        -0.006841522
## + PCNM4         -0.007089863
## + PCNM12        -0.014396973
##
##           Df      AIC      F Pr(>F)
## + PCNM2  1 49.574 9.619  0.002 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.2353704
## Call: fish.db ~ PCNM2
##
##           R2.adjusted
## <All variables>  0.6260113
## + PCNM3         0.3429270
## + PCNM5         0.3057368
## + PCNM1         0.2885396
## + PCNM16        0.2786746
## + PCNM14        0.2744520
## + PCNM15        0.2692809
## + PCNM6         0.2659866
## + PCNM13        0.2636194
## + PCNM9         0.2517847
## + PCNM8         0.2496240
## + PCNM10        0.2434688
## + PCNM7         0.2431476
## + PCNM17        0.2404343
## + PCNM11        0.2366833
## <none>          0.2353704
## + PCNM12        0.2288789
## + PCNM4         0.2189522
##
##           Df      AIC      F Pr(>F)
## + PCNM3  1 46.083 5.4196  0.004 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.342927
## Call: fish.db ~ PCNM2 + PCNM3
##
##           R2.adjusted
## <All variables>  0.6260113
## + PCNM5         0.4076020
## + PCNM1         0.3970300
## + PCNM16        0.3853210
## + PCNM15        0.3828748
## + PCNM14        0.3781827
## + PCNM13        0.3770376
## + PCNM6         0.3595644
## + PCNM8         0.3556885
## + PCNM7         0.3541631
## + PCNM10        0.3526775

```

```

## + PCNM17      0.3513683
## + PCNM9       0.3433672
## <none>        0.3429270
## + PCNM11      0.3416399
## + PCNM12      0.3396547
## + PCNM4       0.3311509
##
##           Df      AIC      F Pr(>F)
## + PCNM5  1 43.941 3.8385 0.006 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.407602
## Call: fish.db ~ PCNM2 + PCNM3 + PCNM5
##
##           R2.adjusted
## <All variables> 0.6260113
## + PCNM1        0.4721469
## + PCNM16       0.4631976
## + PCNM15       0.4589111
## + PCNM14       0.4535248
## + PCNM13       0.4511582
## + PCNM6        0.4305640
## + PCNM7        0.4261965
## + PCNM8        0.4224505
## + PCNM17       0.4181666
## + PCNM10       0.4154485
## + PCNM11       0.4112178
## + PCNM9        0.4111995
## + PCNM12       0.4087602
## <none>         0.4076020
## + PCNM4        0.3976526
##
##           Df      AIC      F Pr(>F)
## + PCNM1  1 41.411 4.057 0.01 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.4721469
## Call: fish.db ~ PCNM2 + PCNM3 + PCNM5 + PCNM1
##
##           R2.adjusted
## <All variables> 0.6260113
## + PCNM13       0.5212427
## + PCNM16       0.5208668
## + PCNM15       0.5161770
## + PCNM14       0.5147355
## + PCNM6        0.4999020
## + PCNM7        0.4936559
## + PCNM8        0.4904113
## + PCNM17       0.4856884
## + PCNM10       0.4835952
## + PCNM11       0.4760087
## + PCNM9        0.4751424

```


```

## + PCNM12          0.4747221
## <none>            0.4721469
## + PCNM4           0.4651218
##
##           Df      AIC      F Pr(>F)
## + PCNM13  1 39.346 3.4612 0.008 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.5212427
## Call: fish.db ~ PCNM2 + PCNM3 + PCNM5 + PCNM1 + PCNM13
##
##           R2.adjusted
## <All variables>  0.6260113
## + PCNM16         0.5767968
## + PCNM15         0.5715331
## + PCNM14         0.5698343
## + PCNM6          0.5475140
## + PCNM7          0.5392074
## + PCNM8          0.5379134
## + PCNM11         0.5281106
## + PCNM9          0.5267003
## + PCNM10         0.5265029
## + PCNM12         0.5255581
## <none>           0.5212427
## + PCNM17         0.5171800
## + PCNM4          0.5152311
##
##           Df      AIC      F Pr(>F)
## + PCNM16  1 36.48 4.0192 0.01 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.5767968
## Call: fish.db ~ PCNM2 + PCNM3 + PCNM5 + PCNM1 + PCNM13 + PCNM16
##
##           R2.adjusted
## <All variables>  0.6260113
## + PCNM6         0.6043089
## + PCNM8         0.5970286
## + PCNM12        0.5946888
## + PCNM7         0.5946475
## + PCNM9         0.5883735
## + PCNM10        0.5851333
## + PCNM15        0.5846468
## <none>           0.5767968
## + PCNM17        0.5748533
## + PCNM4         0.5733749
## + PCNM11        0.5711176
## + PCNM14        0.5652509
##
##           Df      AIC      F Pr(>F)
## + PCNM6  1 35.182 2.5296 0.046 *
## ---

```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Step: R2.adj= 0.6043089
## Call: fish.db ~ PCNM2 + PCNM3 + PCNM5 + PCNM1 + PCNM13 + PCNM16 + PCNM6
##
##               R2.adjusted
## <All variables> 0.6260113
## + PCNM8        0.6248697
## + PCNM12       0.6208788
## + PCNM10       0.6170988
## + PCNM7        0.6142419
## + PCNM15       0.6140369
## + PCNM9        0.6107110
## <none>         0.6043089
## + PCNM17       0.6037430
## + PCNM11       0.5978305
## + PCNM4        0.5963667
## + PCNM14       0.5932113
##
##      Df    AIC      F Pr(>F)
## + PCNM8  1 34.219 2.151  0.074 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
plot(step.pcnm)
```



```
step.pcnm$anova
```

##		R2.adj	Df	AIC	F	Pr(>F)
##	+ PCNM2	0.23537	1	49.574	9.6190	0.002 **
##	+ PCNM3	0.34293	1	46.083	5.4196	0.004 **
##	+ PCNM5	0.40760	1	43.941	3.8385	0.006 **
##	+ PCNM1	0.47215	1	41.411	4.0570	0.010 **
##	+ PCNM13	0.52124	1	39.346	3.4612	0.008 **

```

## + PCNM16          0.57680  1 36.480 4.0192  0.010 **
## + PCNM6           0.60431  1 35.182 2.5296  0.046 *
## <All variables> 0.62601
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

space.mod <- model.matrix(~ PCNM2 + PCNM3 + PCNM5 + PCNM1 + PCNM13 + PCNM16 + PCNM6, douds.space)[-1]

douds.total.env <- dbrda(fish.db ~ env.mod)
douds.total.space <- dbrda(fish.db ~ space.mod)
douds.env.cond.space <- dbrda(fish.db ~ env.mod + Condition(space.mod))
douds.space.cond.env <- dbrda(fish.db ~ space.mod + Condition(env.mod))
permutest(douds.env.cond.space, permutations = 999)

##
## Permutation test for dbrda under reduced model
##
## Permutation: free
## Number of permutations: 999
##
## Model: dbrda(formula = fish.db ~ env.mod + Condition(space.mod))
## Permutation test for all constrained eigenvalues
##           Df Inertia      F Pr(>F)
## Model      3 0.85158 4.423  0.001 ***
## Residual 18 1.15519
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

permutest(douds.space.cond.env, permutations = 999)

##
## Permutation test for dbrda under reduced model
##
## Permutation: free
## Number of permutations: 999
##
## Model: dbrda(formula = fish.db ~ space.mod + Condition(env.mod))
## Permutation test for all constrained eigenvalues
##           Df Inertia      F Pr(>F)
## Model      7 1.8752 4.1741  0.001 ***
## Residual 18 1.1552
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

permutest(douds.total.env, permutations = 999)

##
## Permutation test for dbrda under reduced model
##
## Permutation: free
## Number of permutations: 999
##
## Model: dbrda(formula = fish.db ~ env.mod)
## Permutation test for all constrained eigenvalues
##           Df Inertia      F Pr(>F)
## Model      3 3.7317 10.262  0.001 ***

```



```

## Residual 25 3.0304
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

permutest(doubs.total.space, permutations = 999)


##
## Permutation test for dbrda under reduced model
##
## Permutation: free
## Number of permutations: 999
##
## Model: dbrda(formula = fish.db ~ space.mod)
## Permutation test for all constrained eigenvalues
##           Df Inertia      F Pr(>F)
## Model      7 4.7553 7.1089 0.001 ***
## Residual 21 2.0068
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

doubs.varpart <- varpart(fish.db, env.mod, space.mod)
doubs.varpart

##
## Partition of squared Bray distance in dbRDA
##
## Call: varpart(Y = fish.db, X = env.mod, space.mod)
##
## Explanatory tables:
## X1:  env.mod
## X2:  space.mod
##
## No. of explanatory tables: 2
## Total variation (SS): 6.7621
## No. of observations: 29
##
## Partition table:
##
##           Df R.squared Adj.R.squared Testable
## [a+b] = X1      3  0.55186      0.49808      TRUE
## [b+c] = X2      7  0.70323      0.60431      TRUE
## [a+b+c] = X1+X2 10  0.82917      0.73426      TRUE
## Individual fractions
## [a] = X1|X2      3           0.12995      TRUE
## [b]              0           0.36813     FALSE
## [c] = X2|X1      7           0.23618      TRUE
## [d] = Residuals           0.26574     FALSE
## ---
## Use function 'dbrda' to test significance of fractions of interest

par(mar = c(2,2,2,2))
plot(doubs.varpart)
text(1, 0.25, "Space")
text(0, 0.25, "Env")
mtext("Variation Partitioning of \nDoubs Fish Diversity", side = 3, line = -3)

```



Question 4: Interpret the variation partitioning results.

Answer 4: 13% of the variance in doubs fish diversity is explained by environmental variables, after controlling for spatial variation. 24% of the variance is explained by spatial position, after controlling for environmental variation across space. 37% of variance in fish diversity is explained by spatially structured environmental variation. 27% of the variance is not explained by these variables.

SYNTHESIS

- 1) Using the jelly bean data from class (i.e., JellyBeans.txt), perform a PERMANOVA to test whether or not the vicariance event (random splitting of source community) had an affect on jelly bean composition. Based on your previous analyses with this data set, what are your thoughts about the importance of stochastic vs. deterministic factors on estimates of biodiversity?
- 2) Load the dataset you are using for your Team Project. Perform an ordination to visualize your dataset. Using this ordination, develop some hypotheses relevant to β -diversity. Use a statistic to test one of these hypotheses. Succinctly explain the finding and its relevance to your system.

```
setwd("~/GitHub/QB2019_Peckenpaugh/2.Worksheets/6.DiversitySampling")
jb <- read.table("JellyBeans.txt", sep = "\t", header = TRUE)

# Rearrange rows by group
jb <- jb[c(1:3, 6, 8, 4:5, 7, 9), ]

group <- c(rep("A", 5), rep("B", 4))
jb.num <- jb[, 3:30]

# PERMANOVA
adonis(jb.num ~ group, method = "bray", permutations = 999)
```

```
##
```

```
## Call:
## adonis(formula = jb.num ~ group, permutations = 999, method = "bray")
##
## Permutation: free
## Number of permutations: 999
##
## Terms added sequentially (first to last)
##
##              Df SumsOfSqs  MeanSqs F.Model      R2 Pr(>F)
## group         1   0.09247 0.092468   2.0401 0.22568  0.048 *
## Residuals     7   0.31727 0.045324           0.77432
## Total         8   0.40974           1.00000
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```


Answer 1: The p-value of this PERMANOVA is 0.057, which is not statistically significant considering a standard $\alpha = 0.05$. It is close to significance, however, this result might suggest that deterministic processes generally play a larger role in estimates of biodiversity than stochastic effects. I'm sure it varies by dataset, though.

```
setwd("~/Box Sync/2018-2019/Spring classes/Quantitative biodiversity")
bird <- read.csv("hf085-01-bird.csv", header = TRUE)

# Sorenson & PCoA
bird.num <- bird[3:51]
bird.rm <- vegdist(bird.num, method = "bray", binary = TRUE)
bird.pcoa <- cmdscale(bird.rm, eig = TRUE, k = 3)

explainvar1 <- round(bird.pcoa$eig[1] / sum(bird.pcoa$eig), 3) * 100
explainvar2 <- round(bird.pcoa$eig[2] / sum(bird.pcoa$eig), 3) * 100
explainvar3 <- round(bird.pcoa$eig[3] / sum(bird.pcoa$eig), 3) * 100
sum.eig <- sum(explainvar1, explainvar2, explainvar3)

# Plotting PCoA
palette(c('lightblue1', 'lightblue3', 'gray50', 'darkolivegreen3'))
par(mar = c(5, 5, 1, 2) + 0.1)
plot(bird.pcoa$points[,1], bird.pcoa$points[,2], ylim = c(-0.2, 0.7),
     xlab = paste("PCoA 1 (", explainvar1, "%)", sep = ""),
     ylab = paste("PCoA 2 (", explainvar2, "%)", sep = ""),
     pch = 16, cex = 2.0, type = "n", cex.lab = 1.5, cex.axis = 1.2, axes = FALSE)
axis(side = 1, labels = T, lwd.ticks = 2, cex.axis = 1.2, las = 1)
axis(side = 2, labels = T, lwd.ticks = 2, cex.axis = 1.2, las = 1)
abline(h = 0, v = 0, lty = 3)
box(lwd = 2)
points(bird.pcoa$points[,1], bird.pcoa$points[,2],
       pch = 19, cex = 3, bg = bird$mortality.class, col = bird$mortality.class)
legend("topleft", legend=c("Low", "Med", "Hi-A", "Hi-B"), fill=c('gray50', 'darkolivegreen3', 'lightblue1', 'lightblue3'))
```



```
# phi coefficient of association
library(dplyr)
```

```
##
## Attaching package: 'dplyr'
## The following objects are masked from 'package:stats':
##
##   filter, lag
## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union

bird <- arrange(bird, mortality.class)
bird.num <- bird[3:51]
mortality <- c(rep("Hi-A", 10), rep("Hi-B", 11), rep("Low", 11), rep("Med", 8))

bird.rel <- decostand(bird.num, method = "total")
phi <- multipatt(bird.rel, cluster = mortality, func = "r.g", control = how(nperm=999))
summary(phi)

##
## Multilevel pattern analysis
## -----
##
## Association function: r.g
## Significance level (alpha): 0.05
##
## Total number of species: 49
## Selected number of species: 6
```

```

## Number of species associated to 1 group: 5
## Number of species associated to 2 groups: 0
## Number of species associated to 3 groups: 1
##
## List of species associated to each combination:
##
## Group Hi-A #sps. 1
##      stat p.value
## bhco 0.541 0.003 **
##
## Group Hi-B #sps. 2
##      stat p.value
## howa 0.582 0.002 **
## eawp 0.414 0.039 *
##
## Group Low #sps. 2
##      stat p.value
## btnw 0.741 0.001 ***
## blbw 0.448 0.045 *
##
## Group Hi-A+Hi-B+Med #sps. 1
##      stat p.value
## etti 0.502 0.009 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

Answer 2: Based on my PCoA, while the bird species do not make clear clusters by mortality class of hemlock, there does appear to be a gradient from low to high in PCoA 1. However, PCoA 1 only explains 21.2% of the variance in bird species presence. I am hypothesizing that certain bird species may prefer certain habitats, based on the mortality class of hemlock. I am using the phi coefficient of association to test this. This test showed 6 species with statistically significant species-site group associations: one species (Brown creeper) is associated with Hi-A, two species (Eastern wood peewee, Hooded warbler) are associated with Hi-B, two species (Black-throated green warbler, Blackburnian warbler) are associated with Low mortality, and one species (Tufted titmouse) is associated with the groups Hi-A, Hi-B, and Medium. This result suggests that certain bird species may preferentially associate with either the presence or absence of hemlock in their habitat.