1 Equations in ρ coordinates

$$0 = \frac{1}{V'}\frac{\partial}{\partial\rho}\left[V'\langle|\nabla\rho|^2\rangle\frac{\partial}{\partial\rho}\Phi\right] + \frac{1}{\epsilon_0}\left(-en_\mathrm{e} + Z_\mathrm{i}en_\mathrm{i} + Z_\mathrm{z}en_\mathrm{z} + Z_\mathrm{b}en_\mathrm{b} + Z_\mathrm{b}egn_\mathrm{b}^\mathrm{rp}\right) \tag{1.1}$$

$$\begin{aligned}\frac{1}{c^2}\frac{\partial}{\partial t}\dot{\psi}_t = {} & V'\left\langle\frac{|\nabla\rho|^2}{R^2}\right\rangle\frac{\partial}{\partial\rho}\left[\frac{1}{V'\langle R^{-2}\rangle}\frac{\partial}{\partial\rho}\psi_t\right] \\ & + \frac{\mu_0}{\langle B^\theta\rangle}\left[-en_\mathrm{e}\left(\langle Bu_{\mathrm{e}\parallel}\rangle - I\left\langle\frac{u_{\mathrm{e}\zeta}}{R}\right\rangle\right) + Z_\mathrm{i}en_\mathrm{i}\left(\langle Bu_{\mathrm{i}\parallel}\rangle - I\left\langle\frac{u_{\mathrm{i}\zeta}}{R}\right\rangle\right)\right. \\ & \left. + Z_\mathrm{z}en_\mathrm{z}\left(\langle Bu_{\mathrm{z}\parallel}\rangle - I\left\langle\frac{u_{\mathrm{z}\zeta}}{R}\right\rangle\right) + Z_\mathrm{b}en_\mathrm{b}\left(\langle Bu_{\mathrm{b}\parallel}\rangle - I\left\langle\frac{u_{\mathrm{b}\zeta}}{R}\right\rangle\right)\right]\end{aligned} \tag{1.2}$$

$$\begin{aligned}\frac{1}{c^2}\frac{\partial}{\partial t}\dot{\psi} = {} & \frac{1}{V'\langle R^{-2}\rangle}\frac{\partial}{\partial\rho}\left[V'\left\langle\frac{|\nabla\rho|^2}{R^2}\right\rangle\frac{\partial}{\partial\rho}\psi\right] \\ & - \frac{\mu_0}{\langle R^{-2}\rangle}\left(-en_\mathrm{e}\left\langle\frac{u_{\mathrm{e}\zeta}}{R}\right\rangle + Z_\mathrm{i}en_\mathrm{i}\left\langle\frac{u_{\mathrm{i}\zeta}}{R}\right\rangle + Z_\mathrm{z}en_\mathrm{z}\left\langle\frac{u_{\mathrm{z}\zeta}}{R}\right\rangle + Z_\mathrm{b}en_\mathrm{b}\left\langle\frac{u_{\mathrm{b}\zeta}}{R}\right\rangle\right)\end{aligned} \tag{1.3}$$

$$\frac{\partial}{\partial t}\psi = \dot{\psi} \tag{1.4}$$

$$\frac{\partial}{\partial t}\psi_t = \dot{\psi}_t \tag{1.5}$$

$$\begin{aligned}\frac{1}{V'}\frac{\partial}{\partial t}(V'n_\mathrm{e}) = {} & -\frac{1}{V'}\frac{\partial}{\partial\rho}[V'n_\mathrm{e}(u_\mathrm{e}^\rho - u_g^\rho)] + \nu_\mathrm{ionize}\frac{n_\mathrm{e}}{n_{01}+n_{02}+n_{03}}(n_{01}+n_{02}+n_{03}) \\ & + Z_\mathrm{z}n_\mathrm{e}\bar{n}_z f_5 S_5 - Z_\mathrm{z}n_\mathrm{e}n_\mathrm{z}A_6 - \nu_\mathrm{L}(n_\mathrm{e} - n_\mathrm{esup}) + (1 - f_\mathrm{CX})\frac{P_\mathrm{b}}{E_\mathrm{b}}\end{aligned} \tag{1.6}$$

$$\begin{aligned}0 = {} & -(\langle B^2\rangle\langle R^2\rangle - I^2)\frac{\partial}{\partial\rho}n_\mathrm{e}T_\mathrm{e} + en_\mathrm{e}(\langle B^2\rangle\langle R^2\rangle - I^2)\frac{\partial}{\partial\rho}\Phi \\ & - e\frac{\partial\psi}{\partial\rho}In_\mathrm{e}\langle Bu_{\mathrm{e}\parallel}\rangle + e\frac{\partial\psi}{\partial\rho}\langle B^2\rangle n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle\end{aligned} \tag{1.7}$$

$$\begin{aligned}m_\mathrm{e}n_\mathrm{e}\frac{\partial}{\partial t}\langle Bu_{\mathrm{e}\parallel}\rangle = {} & -\hat{\mu}_1^\mathrm{e}\langle Bu_{\mathrm{e}\parallel}\rangle - \hat{\mu}_2^\mathrm{e}\langle B\hat{q}_{\mathrm{e}\parallel}\rangle + \hat{\mu}_1^\mathrm{e}BV_{1\mathrm{e}} + \hat{\mu}_2^\mathrm{e}BV_{2\mathrm{e}} \\ & + \ell_{11}^{\mathrm{ee}}\langle Bu_{\mathrm{e}\parallel}\rangle + \ell_{11}^{\mathrm{ei}}\langle Bu_{\mathrm{i}\parallel}\rangle + \ell_{11}^{\mathrm{ez}}\langle Bu_{\mathrm{z}\parallel}\rangle + \ell_{11}^{\mathrm{eb}}\langle Bu_{\mathrm{b}\parallel}\rangle \\ & - \ell_{12}^{\mathrm{ee}}\langle B\hat{q}_{\mathrm{e}\parallel}\rangle - \ell_{12}^{\mathrm{ei}}\langle B\hat{q}_{\mathrm{i}\parallel}\rangle - \ell_{12}^{\mathrm{ez}}\langle B\hat{q}_{\mathrm{z}\parallel}\rangle \\ & - en_\mathrm{e}\langle BE_\parallel\rangle - \nu_\mathrm{L}m_\mathrm{e}n_\mathrm{e}\langle Bu_{\mathrm{e}\parallel}\rangle - \nu_{0\mathrm{e}}m_\mathrm{e}n_\mathrm{e}\langle Bu_{\mathrm{e}\parallel}\rangle \\ & - \frac{1}{V'}\frac{\partial}{\partial\rho}V'\langle|\nabla\rho|\rangle v_{\mathrm{e}\zeta}m_\mathrm{e}n_\mathrm{e}\langle Bu_{\mathrm{e}\parallel}\rangle + \langle|\nabla\rho|\rangle v_{\mathrm{e}\zeta}m_\mathrm{e}n_\mathrm{e}\left(\frac{\partial}{\partial\rho}\frac{\langle B^2\rangle}{I}\right)\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\parallel}\rangle\end{aligned}$$

$$
\begin{aligned}
&+\frac{1}{V'}\frac{\partial}{\partial\rho}V'\langle|\nabla\rho|^2\rangle\chi_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial\rho}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\parallel}\rangle\\
&-\langle|\nabla\rho|^2\rangle\chi_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\left(\frac{\partial}{\partial\rho}\frac{\langle B^2\rangle}{I}\right)\frac{\partial}{\partial\rho}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\parallel}\rangle-\frac{\langle B^2\rangle}{I}\frac{1}{V'}\frac{\partial}{\partial\rho}V'\langle\Pi_{\mathrm{e}}^{\mathrm{res}}\rangle\\
&-\frac{1}{V'}\frac{\partial}{\partial\rho}V'\left[\langle|\nabla\rho|\rangle v_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\left(\langle R^2\rangle-\frac{I^2}{\langle B^2\rangle}\right)\frac{1}{I}BV_{1\mathrm{e}}\right.\\
&\left.-\langle|\nabla\rho|^2\rangle\chi_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\frac{\partial}{\partial\rho}\left(\langle R^2\rangle-\frac{I^2}{\langle B^2\rangle}\right)\frac{1}{I}BV_{1\mathrm{e}}\right] \qquad (1.8)
\end{aligned}
$$

$$
\begin{aligned}
\frac{1}{V'}\frac{\partial}{\partial t}(V'm_{\mathrm{e}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle)=&-\frac{1}{V'}\frac{\partial}{\partial\rho}V'\left[\langle|\nabla\rho|\rangle v_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle+(u_{\mathrm{e}}^{\rho}-u_g^{\rho})m_{\mathrm{e}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle\right.\\
&\left.-\langle|\nabla\rho|^2\rangle\chi_{\mathrm{e}\zeta}m_{\mathrm{e}}n_{\mathrm{e}}\frac{\partial}{\partial\rho}\langle Ru_{\mathrm{e}\zeta}\rangle+\langle\Pi_{\mathrm{e}}^{\mathrm{res}}\rangle\right]\\
&+\ell_{11}^{\mathrm{ee}}\langle Ru_{\mathrm{e}\zeta}\rangle+\ell_{11}^{\mathrm{ei}}\langle Ru_{\mathrm{i}\zeta}\rangle+\ell_{11}^{\mathrm{ez}}\langle Ru_{\mathrm{z}\zeta}\rangle+\ell_{11}^{\mathrm{eb}}\langle Ru_{\mathrm{b}\zeta}\rangle\\
&-\ell_{12}^{\mathrm{ee}}\langle R\hat{q}_{\mathrm{e}\zeta}\rangle-\ell_{12}^{\mathrm{ei}}\langle R\hat{q}_{\mathrm{i}\zeta}\rangle-\ell_{12}^{\mathrm{ez}}\langle R\hat{q}_{\mathrm{z}\zeta}\rangle\\
&-en_{\mathrm{e}}\langle RE_{\zeta}\rangle-e\frac{\partial\psi}{\partial\rho}n_{\mathrm{e}}u_{\mathrm{e}}^{\rho}\\
&-\frac{e\langle|\nabla\rho|^2\rangle D_{\mathrm{e}}}{T_{\mathrm{e}}}\frac{\partial\psi}{\partial\rho}\left[(1-C_{\mathrm{T}})T_{\mathrm{e}}\frac{\partial}{\partial\rho}n_{\mathrm{e}}+C_{\mathrm{T}}\frac{\partial}{\partial\rho}n_{\mathrm{e}}T_{\mathrm{e}}+n_{\mathrm{e}}T_{\mathrm{e}}\frac{\partial\psi}{\partial\rho}C_{\mathrm{p}}\right]\\
&-\nu_{\mathrm{L}}m_{\mathrm{e}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle-\nu_{0\mathrm{e}}m_{\mathrm{e}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle \qquad (1.9)
\end{aligned}
$$

$$
\begin{aligned}
\frac{3}{2}\frac{1}{V'^{5/3}}\frac{\partial}{\partial t}(V'^{5/3}n_{\mathrm{e}}T_{\mathrm{e}})=&-\frac{1}{V'}\frac{\partial}{\partial\rho}V'\left[-\chi_{\mathrm{e}}\langle|\nabla\rho|^2\rangle n_{\mathrm{e}}\frac{\partial}{\partial\rho}T_{\mathrm{e}}+V_{\mathrm{e}}^{\mathrm{hp}}\langle|\nabla\rho|\rangle n_{\mathrm{e}}T_{\mathrm{e}}+\frac{5}{2}n_{\mathrm{e}}T_{\mathrm{e}}(u_{\mathrm{e}}^{\rho}-u_g^{\rho})\right]\\
&-u_g^{\rho}\frac{\partial}{\partial\rho}n_{\mathrm{e}}T_{\mathrm{e}}-u_{\mathrm{i}}^{\rho}\frac{\partial}{\partial\rho}n_{\mathrm{i}}T_{\mathrm{i}}-u_{\mathrm{z}}^{\rho}\frac{\partial}{\partial\rho}n_{\mathrm{z}}T_{\mathrm{z}}-\hat{u}_{\mathrm{i}\theta}\langle\boldsymbol{B}\cdot\nabla\cdot\overleftrightarrow{\pi}_{\mathrm{i}}\rangle-\hat{u}_{\mathrm{z}\theta}\langle\boldsymbol{B}\cdot\nabla\cdot\overleftrightarrow{\pi}_{\mathrm{z}}\rangle\\
&-\frac{3}{2}n_{\mathrm{e}}\frac{T_{\mathrm{e}}-T_{\mathrm{i}}}{\tau_{\mathrm{ei}}}-\frac{3}{2}n_{\mathrm{e}}\frac{T_{\mathrm{e}}-T_{\mathrm{z}}}{\tau_{\mathrm{ez}}}\\
&-\dot{\psi}_t\langle B^{\theta}\rangle(-en_{\mathrm{e}}\hat{u}_{\mathrm{e}\theta}+Z_{\mathrm{i}}en_{\mathrm{i}}\hat{u}_{\mathrm{i}\theta}+Z_{\mathrm{z}}en_{\mathrm{z}}\hat{u}_{\mathrm{z}\theta})\\
&+\dot{\psi}\left(-en_{\mathrm{e}}\left\langle\frac{u_{\mathrm{e}\zeta}}{R}\right\rangle+Z_{\mathrm{i}}en_{\mathrm{i}}\left\langle\frac{u_{\mathrm{i}\zeta}}{R}\right\rangle+Z_{\mathrm{z}}en_{\mathrm{z}}\left\langle\frac{u_{\mathrm{z}\zeta}}{R}\right\rangle\right)\\
&-\nu_{\mathrm{L}}T_{\mathrm{e}}(n_{\mathrm{e}}-n_{\mathrm{esup}})-\frac{3}{2}\nu_{\mathrm{L}T_{\mathrm{e}}}n_{\mathrm{e}}(T_{\mathrm{e}}-T_{\mathrm{esup}})\\
&-E_{\mathrm{H}}\nu_{\mathrm{ionize}}\frac{n_{\mathrm{e}}}{n_{01}+n_{02}+n_{03}}(n_{01}+n_{02}+n_{03})\\
&+f_{\mathrm{ei}}P_{\mathrm{b}}+P_{\alpha\mathrm{e}}+P_{\mathrm{RFe}}-P_{\mathrm{Br}} \qquad (1.10)
\end{aligned}
$$

$$
\frac{5}{2}m_{\mathrm{e}}n_{\mathrm{e}}\frac{\partial}{\partial t}\langle B\hat{q}_{\mathrm{e}\parallel}\rangle=-\hat{\mu}_2^{\mathrm{e}}\langle Bu_{\mathrm{e}\parallel}\rangle-\hat{\mu}_3^{\mathrm{e}}\langle B\hat{q}_{\mathrm{e}\parallel}\rangle+\hat{\mu}_2^{\mathrm{e}}BV_{1\mathrm{e}}+\hat{\mu}_3^{\mathrm{e}}BV_{2\mathrm{e}}
$$

$$
\begin{aligned}
&- \ell_{21}^{\mathrm{ee}}\langle Bu_{\mathrm{e}\|}\rangle - \ell_{21}^{\mathrm{ei}}\langle Bu_{\mathrm{i}\|}\rangle - \ell_{21}^{\mathrm{ez}}\langle Bu_{\mathrm{z}\|}\rangle - \ell_{21}^{\mathrm{eb}}\langle Bu_{\mathrm{b}\|}\rangle \\
&+ \ell_{22}^{\mathrm{ee}}\langle B\hat{q}_{\mathrm{e}\|}\rangle + \ell_{22}^{\mathrm{ei}}\langle B\hat{q}_{\mathrm{i}\|}\rangle + \ell_{22}^{\mathrm{ez}}\langle B\hat{q}_{\mathrm{z}\|}\rangle \\
&- \nu_{\mathrm{L}} m_{\mathrm{e}} n_{\mathrm{e}}\langle B\hat{q}_{\mathrm{e}\|}\rangle - \nu_{0\mathrm{e}} m_{\mathrm{e}} n_{\mathrm{e}}\langle B\hat{q}_{\mathrm{e}\|}\rangle
\end{aligned} \tag{1.11}
$$

$$
\begin{aligned}
\frac{1}{V'}\frac{\partial}{\partial t}(V' n_{\mathrm{i}}) = &-\frac{1}{V'}\frac{\partial}{\partial \rho}[V' n_{\mathrm{i}}(u_{\mathrm{i}}^{\rho} - u_g^{\rho})] + \nu_{\mathrm{ionize}}\frac{n_{\mathrm{e}}}{n_{01}+n_{02}+n_{03}}(n_{01}+n_{02}+n_{03}) \\
&- \nu_{\mathrm{L}}(n_{\mathrm{i}} - n_{\mathrm{isup}}) + \nu_{\mathrm{b}} n_{\mathrm{b}} + \nu_{\mathrm{b}} g n_{\mathrm{b}}^{\mathrm{rp}} - f_{\mathrm{CX}}\frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} + \nu_{\mathrm{bL}} n_{\mathrm{b}} + S_{\mathrm{LC}} - \nu_{\mathrm{OL}} n_{\mathrm{i}}
\end{aligned} \tag{1.12}
$$

$$
\begin{aligned}
0 = &-(\langle B^2\rangle\langle R^2\rangle - I^2)\frac{\partial}{\partial \rho} n_{\mathrm{i}} T_{\mathrm{i}} - Z_{\mathrm{i}} e n_{\mathrm{i}}(\langle B^2\rangle\langle R^2\rangle - I^2)\frac{\partial}{\partial \rho}\Phi \\
&+ Z_{\mathrm{i}} e n_{\mathrm{i}}\frac{\partial \psi}{\partial \rho} I\langle Bu_{\mathrm{i}\|}\rangle - Z_{\mathrm{i}} e n_{\mathrm{i}}\frac{\partial \psi}{\partial \rho}\langle B^2\rangle\langle Ru_{\mathrm{i}\zeta}\rangle
\end{aligned} \tag{1.13}
$$

$$
\begin{aligned}
m_{\mathrm{i}} n_{\mathrm{i}}\frac{\partial}{\partial t}\langle Bu_{\mathrm{i}\|}\rangle = &-\hat{\mu}_1^{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle - \hat{\mu}_2^{\mathrm{i}}\langle B\hat{q}_{\mathrm{i}\|}\rangle + \hat{\mu}_1^{\mathrm{i}} BV_{1\mathrm{i}} + \hat{\mu}_2^{\mathrm{i}} BV_{2\mathrm{i}} \\
&+ \ell_{11}^{\mathrm{ie}}\langle Bu_{\mathrm{e}\|}\rangle + \ell_{11}^{\mathrm{ii}}\langle Bu_{\mathrm{i}\|}\rangle + \ell_{11}^{\mathrm{iz}}\langle Bu_{\mathrm{z}\|}\rangle + \ell_{11}^{\mathrm{ib}}\langle Bu_{\mathrm{b}\|}\rangle \\
&- \ell_{12}^{\mathrm{ie}}\langle B\hat{q}_{\mathrm{e}\|}\rangle - \ell_{12}^{\mathrm{ii}}\langle B\hat{q}_{\mathrm{i}\|}\rangle - \ell_{12}^{\mathrm{iz}}\langle B\hat{q}_{\mathrm{z}\|}\rangle \\
&+ Z_{\mathrm{i}} e n_{\mathrm{i}}\langle BE_{\|}\rangle - \nu_{\mathrm{L}} m_{\mathrm{i}} n_{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle + \nu_{\mathrm{b}} m_{\mathrm{b}} n_{\mathrm{b}}\langle Bu_{\mathrm{b}\|}\rangle - \nu_{0\mathrm{i}} m_{\mathrm{i}} n_{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle \\
&- \nu_{\mathrm{CX}} m_{\mathrm{i}} n_{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle - \nu_{\mathrm{OL}} m_{\mathrm{i}} n_{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle \\
&- \frac{1}{V'}\frac{\partial}{\partial \rho} V'\langle|\nabla\rho|\rangle v_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\langle Bu_{\mathrm{i}\|}\rangle + \langle|\nabla\rho|\rangle v_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\left(\frac{\partial}{\partial \rho}\frac{\langle B^2\rangle}{I}\right)\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{i}\|}\rangle \\
&+ \frac{1}{V'}\frac{\partial}{\partial \rho} V'\langle|\nabla\rho|^2\rangle\chi_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial \rho}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{i}\|}\rangle \\
&- \langle|\nabla\rho|^2\rangle\chi_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\left(\frac{\partial}{\partial \rho}\frac{\langle B^2\rangle}{I}\right)\frac{\partial}{\partial \rho}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{i}\|}\rangle - \frac{\langle B^2\rangle}{I}\frac{1}{V'}\frac{\partial}{\partial \rho} V'\langle\Pi_{\mathrm{i}}^{\mathrm{res}}\rangle \\
&- \frac{1}{V'}\frac{\partial}{\partial \rho} V'\left[\langle|\nabla\rho|\rangle v_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\left(\langle R^2\rangle - \frac{I^2}{\langle B^2\rangle}\right)\frac{1}{I} BV_{1\mathrm{i}}\right. \\
&\left.-\langle|\nabla\rho|^2\rangle\chi_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\frac{\partial}{\partial \rho}\left(\langle R^2\rangle - \frac{I^2}{\langle B^2\rangle}\right)\frac{1}{I} BV_{1\mathrm{i}}\right]
\end{aligned} \tag{1.14}
$$

$$
\begin{aligned}
\frac{1}{V'}\frac{\partial}{\partial t}(V' m_{\mathrm{i}} n_{\mathrm{i}}\langle Ru_{\mathrm{i}\zeta}\rangle) = &-\frac{1}{V'}\frac{\partial}{\partial \rho} V'\left[\langle|\nabla\rho|\rangle v_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\langle Ru_{\mathrm{i}\zeta}\rangle + (u_{\mathrm{i}}^{\rho} - u_g^{\rho}) m_{\mathrm{i}} n_{\mathrm{i}}\langle Ru_{\mathrm{i}\zeta}\rangle\right. \\
&\left.-\langle|\nabla\rho|^2\rangle\chi_{\mathrm{i}\zeta} m_{\mathrm{i}} n_{\mathrm{i}}\frac{\partial}{\partial \rho}\langle Ru_{\mathrm{i}\zeta}\rangle + \langle\Pi_{\mathrm{i}}^{\mathrm{res}}\rangle\right] \\
&+ \ell_{11}^{\mathrm{ie}}\langle Ru_{\mathrm{e}\zeta}\rangle + \ell_{11}^{\mathrm{ii}}\langle Ru_{\mathrm{i}\zeta}\rangle + \ell_{11}^{\mathrm{iz}}\langle Ru_{\mathrm{z}\zeta}\rangle + \ell_{11}^{\mathrm{ib}}\langle Ru_{\mathrm{b}\zeta}\rangle
\end{aligned}
$$

$$
\begin{aligned}
&- \ell_{12}^{\mathrm{ie}}\langle R\hat{q}_{\mathrm{e}\zeta}\rangle - \ell_{12}^{\mathrm{ii}}\langle R\hat{q}_{\mathrm{i}\zeta}\rangle - \ell_{12}^{\mathrm{iz}}\langle R\hat{q}_{\mathrm{z}\zeta}\rangle \\
&+ Z_{\mathrm{i}} e n_{\mathrm{i}}\langle RE_{\zeta}\rangle + Z_{\mathrm{i}} e \frac{\partial \psi}{\partial \rho} n_{\mathrm{i}} u_{\mathrm{i}}^{\rho} \\
&+ Z_{\mathrm{i}} \frac{e\langle |\nabla\rho|^2\rangle D_{\mathrm{e}}}{T_{\mathrm{e}}} \frac{\partial \psi}{\partial \rho} \frac{n_{\mathrm{i}} + n_{\mathrm{b}}}{n_{\mathrm{e}}} \left[(1 - C_{\mathrm{T}}) T_{\mathrm{e}} \frac{\partial}{\partial \rho} n_{\mathrm{e}} + C_{\mathrm{T}} \frac{\partial}{\partial \rho} n_{\mathrm{e}} T_{\mathrm{e}} + n_{\mathrm{e}} T_{\mathrm{e}} \frac{\partial \psi}{\partial \rho} C_{\mathrm{p}} \right] \\
&- \nu_{\mathrm{L}} m_{\mathrm{i}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta}\rangle + \nu_{\mathrm{b}} m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta}\rangle - \nu_{0\mathrm{i}} m_{\mathrm{i}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta}\rangle - \nu_{\mathrm{CX}} m_{\mathrm{i}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta}\rangle \\
&- \nu_{\mathrm{OL}} m_{\mathrm{i}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta}\rangle
\end{aligned} \tag{1.15}
$$

$$
\begin{aligned}
\frac{3}{2} \frac{1}{V'^{5/3}} \frac{\partial}{\partial t} (V'^{5/3} n_{\mathrm{i}} T_{\mathrm{i}}) = &- \frac{1}{V'} \frac{\partial}{\partial \rho} V' \left[-\chi_{\mathrm{i}} \langle |\nabla\rho|^2\rangle n_{\mathrm{i}} \frac{\partial}{\partial \rho} T_{\mathrm{i}} + V_{\mathrm{i}}^{\mathrm{hp}} \langle |\nabla\rho|\rangle n_{\mathrm{i}} T_{\mathrm{i}} + \frac{5}{2} n_{\mathrm{i}} T_{\mathrm{i}} (u_{\mathrm{i}}^{\rho} - u_g^{\rho}) \right] \\
&+ (u_{\mathrm{i}}^{\rho} - u_g^{\rho}) \frac{\partial}{\partial \rho} n_{\mathrm{i}} T_{\mathrm{i}} + \hat{u}_{\mathrm{i}\theta} \langle \boldsymbol{B} \cdot \nabla \cdot \overleftrightarrow{\pi}_{\mathrm{i}} \rangle - \frac{3}{2} n_{\mathrm{i}} \frac{T_{\mathrm{i}} - T_{\mathrm{e}}}{\tau_{\mathrm{ie}}} - \frac{3}{2} n_{\mathrm{i}} \frac{T_{\mathrm{i}} - T_{\mathrm{z}}}{\tau_{\mathrm{iz}}} \\
&- \nu_{\mathrm{L}} T_{\mathrm{i}} (n_{\mathrm{i}} - n_{\mathrm{isup}}) - \frac{3}{2} \nu_{\mathrm{L}T_{\mathrm{i}}} n_{\mathrm{i}} (T_{\mathrm{i}} - T_{\mathrm{isup}}) \\
&+ \frac{3}{2} \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} (n_{01} T_{01} + n_{02} T_{02} + n_{03} T_{03}) \\
&- \frac{3}{2} \nu_{\mathrm{CX}} n_{\mathrm{i}} (T_{\mathrm{i}} - T_{01}) + (1 - f_{\mathrm{ei}}) P_{\mathrm{b}} + m_{\mathrm{b}} \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} \langle B u_{\mathrm{b}\parallel}\rangle \frac{\langle B u_{\mathrm{i}\parallel}\rangle}{\langle B^2\rangle} + P_{\alpha\mathrm{i}} + P_{\mathrm{RFi}}
\end{aligned} \tag{1.16}
$$

$$
\begin{aligned}
\frac{5}{2} m_{\mathrm{i}} n_{\mathrm{i}} \frac{\partial}{\partial t} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle = &- \hat{\mu}_2^{\mathrm{i}} \langle B u_{\mathrm{i}\parallel}\rangle - \hat{\mu}_3^{\mathrm{i}} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle + \hat{\mu}_2^{\mathrm{i}} B V_{1\mathrm{i}} + \hat{\mu}_3^{\mathrm{i}} B V_{2\mathrm{i}} \\
&- \ell_{21}^{\mathrm{ie}} \langle B u_{\mathrm{e}\parallel}\rangle - \ell_{21}^{\mathrm{ii}} \langle B u_{\mathrm{i}\parallel}\rangle - \ell_{21}^{\mathrm{iz}} \langle B u_{\mathrm{z}\parallel}\rangle - \ell_{21}^{\mathrm{ib}} \langle B u_{\mathrm{b}\parallel}\rangle \\
&+ \ell_{22}^{\mathrm{ie}} \langle B\hat{q}_{\mathrm{e}\parallel}\rangle + \ell_{22}^{\mathrm{ii}} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle + \ell_{22}^{\mathrm{iz}} \langle B\hat{q}_{\mathrm{z}\parallel}\rangle \\
&- \nu_{\mathrm{L}} m_{\mathrm{i}} n_{\mathrm{i}} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle - \nu_{0\mathrm{i}} m_{\mathrm{i}} n_{\mathrm{i}} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle
\end{aligned} \tag{1.17}
$$

$$
\begin{aligned}
\frac{1}{V'} \frac{\partial}{\partial t} (V' n_{\mathrm{z}}) = &- \frac{1}{V'} \frac{\partial}{\partial \rho} [V' n_{\mathrm{z}} (u_{\mathrm{z}}^{\rho} - u_g^{\rho})] + n_{\mathrm{e}} \bar{n}_z f_5 S_5 - n_{\mathrm{e}} n_{\mathrm{z}} A_6 \\
&- \nu_{\mathrm{L}} (n_{\mathrm{z}} - n_{\mathrm{zsup}}) + S_{\mathrm{LC}} - \nu_{\mathrm{OL}} n_{\mathrm{z}}
\end{aligned} \tag{1.18}
$$

$$
\begin{aligned}
0 = &- (\langle B^2\rangle \langle R^2\rangle - I^2) \frac{\partial}{\partial \rho} n_{\mathrm{z}} T_{\mathrm{z}} - Z_{\mathrm{z}} e n_{\mathrm{z}} (\langle B^2\rangle \langle R^2\rangle - I^2) \frac{\partial}{\partial \rho} \Phi \\
&+ Z_{\mathrm{z}} e n_{\mathrm{z}} \frac{\partial \psi}{\partial \rho} I \langle B u_{\mathrm{z}\parallel}\rangle - Z_{\mathrm{z}} e n_{\mathrm{z}} \frac{\partial \psi}{\partial \rho} \langle B^2\rangle \langle R u_{\mathrm{z}\zeta}\rangle
\end{aligned} \tag{1.19}
$$

$$
\begin{aligned}
m_{\mathrm{z}} n_{\mathrm{z}} \frac{\partial}{\partial t} \langle B u_{\mathrm{z}\parallel}\rangle = &- \hat{\mu}_1^{\mathrm{z}} \langle B u_{\mathrm{z}\parallel}\rangle - \hat{\mu}_2^{\mathrm{z}} \langle B\hat{q}_{\mathrm{z}\parallel}\rangle + \hat{\mu}_1^{\mathrm{z}} B V_{1\mathrm{z}} + \hat{\mu}_2^{\mathrm{z}} B V_{2\mathrm{z}} \\
&+ \ell_{11}^{\mathrm{ze}} \langle B u_{\mathrm{e}\parallel}\rangle + \ell_{11}^{\mathrm{zi}} \langle B u_{\mathrm{i}\parallel}\rangle + \ell_{11}^{\mathrm{zz}} \langle B u_{\mathrm{z}\parallel}\rangle + \ell_{11}^{\mathrm{zb}} \langle B u_{\mathrm{b}\parallel}\rangle \\
&- \ell_{12}^{\mathrm{ze}} \langle B\hat{q}_{\mathrm{e}\parallel}\rangle - \ell_{12}^{\mathrm{zi}} \langle B\hat{q}_{\mathrm{i}\parallel}\rangle - \ell_{12}^{\mathrm{zz}} \langle B\hat{q}_{\mathrm{z}\parallel}\rangle
\end{aligned}
$$

$$
\begin{aligned}
&+ Z_{\mathrm{z}} e n_{\mathrm{z}} \langle B E_{\parallel} \rangle - \nu_{\mathrm{L}} m_{\mathrm{z}} n_{\mathrm{z}} \langle B u_{\mathrm{z}\parallel} \rangle - \nu_{0\mathrm{z}} m_{\mathrm{z}} n_{\mathrm{z}} \langle B u_{\mathrm{z}\parallel} \rangle - \nu_{\mathrm{OL}} m_{\mathrm{z}} n_{\mathrm{z}} \langle B u_{\mathrm{z}\parallel} \rangle \\
&- \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho| \rangle v_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \langle B u_{\mathrm{z}\parallel} \rangle + \langle |\nabla \rho| \rangle v_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \left(\frac{\partial}{\partial \rho} \frac{\langle B^2 \rangle}{I} \right) \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{z}\parallel} \rangle \\
&+ \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle \chi_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \frac{\langle B^2 \rangle}{I} \frac{\partial}{\partial \rho} \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{z}\parallel} \rangle \\
&- \langle |\nabla \rho|^2 \rangle \chi_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \left(\frac{\partial}{\partial \rho} \frac{\langle B^2 \rangle}{I} \right) \frac{\partial}{\partial \rho} \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{z}\parallel} \rangle - \frac{\langle B^2 \rangle}{I} \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle \Pi_{\mathrm{z}}^{\mathrm{res}} \rangle \\
&- \frac{1}{V'} \frac{\partial}{\partial \rho} V' \left[\langle |\nabla \rho| \rangle v_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \left(\langle R^2 \rangle - \frac{I^2}{\langle B^2 \rangle} \right) \frac{1}{I} B V_{1\mathrm{z}} \right. \\
&\left. - \langle |\nabla \rho|^2 \rangle \chi_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \frac{\partial}{\partial \rho} \left(\langle R^2 \rangle - \frac{I^2}{\langle B^2 \rangle} \right) \frac{1}{I} B V_{1\mathrm{z}} \right] \qquad (1.20)
\end{aligned}
$$

$$
\begin{aligned}
\frac{1}{V'} \frac{\partial}{\partial t} (V' m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle) = &- \frac{1}{V'} \frac{\partial}{\partial \rho} V' \left[\langle |\nabla \rho| \rangle v_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle + (u_{\mathrm{z}}^{\rho} - u_g^{\rho}) m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle \right. \\
&\left. - \langle |\nabla \rho|^2 \rangle \chi_{\mathrm{z}\zeta} m_{\mathrm{z}} n_{\mathrm{z}} \frac{\partial}{\partial \rho} \langle R u_{\mathrm{z}\zeta} \rangle + \langle \Pi_{\mathrm{z}}^{\mathrm{res}} \rangle \right] \\
&+ \ell_{11}^{\mathrm{ze}} \langle R u_{\mathrm{e}\zeta} \rangle + \ell_{11}^{\mathrm{zi}} \langle R u_{\mathrm{i}\zeta} \rangle + \ell_{11}^{\mathrm{zz}} \langle R u_{\mathrm{z}\zeta} \rangle + \ell_{11}^{\mathrm{zb}} \langle R u_{\mathrm{b}\zeta} \rangle \\
&- \ell_{12}^{\mathrm{ze}} \langle R \hat{q}_{\mathrm{e}\zeta} \rangle - \ell_{12}^{\mathrm{zi}} \langle R \hat{q}_{\mathrm{i}\zeta} \rangle - \ell_{12}^{\mathrm{zz}} \langle R \hat{q}_{\mathrm{z}\zeta} \rangle \\
&+ Z_{\mathrm{z}} e n_{\mathrm{z}} \langle R E_{\zeta} \rangle + Z_{\mathrm{z}} e \frac{\partial \psi}{\partial \rho} n_{\mathrm{z}} u_{\mathrm{z}}^{\rho} \\
&+ Z_{\mathrm{z}} \frac{e \langle |\nabla \rho|^2 \rangle D_{\mathrm{e}}}{T_{\mathrm{e}}} \frac{\partial \psi}{\partial \rho} \frac{n_{\mathrm{z}}}{n_{\mathrm{e}}} \left[(1 - C_{\mathrm{T}}) T_{\mathrm{e}} \frac{\partial}{\partial \rho} n_{\mathrm{e}} + C_{\mathrm{T}} \frac{\partial}{\partial \rho} n_{\mathrm{e}} T_{\mathrm{e}} + n_{\mathrm{e}} T_{\mathrm{e}} \frac{\partial \psi}{\partial \rho} C_{\mathrm{p}} \right] \\
&- \nu_{\mathrm{L}} m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle - \nu_{0\mathrm{z}} m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle - \nu_{\mathrm{OL}} m_{\mathrm{z}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle \qquad (1.21)
\end{aligned}
$$

$$
\begin{aligned}
\frac{3}{2} \frac{1}{V'^{5/3}} \frac{\partial}{\partial t} (V'^{5/3} n_{\mathrm{z}} T_{\mathrm{z}}) = &- \frac{1}{V'} \frac{\partial}{\partial \rho} V' \left[- \chi_{\mathrm{z}} \langle |\nabla \rho|^2 \rangle n_{\mathrm{z}} \frac{\partial}{\partial \rho} T_{\mathrm{z}} + V_{\mathrm{z}}^{\mathrm{hp}} \langle |\nabla \rho| \rangle n_{\mathrm{z}} T_{\mathrm{z}} + \frac{5}{2} n_{\mathrm{z}} T_{\mathrm{z}} (u_{\mathrm{z}}^{\rho} - u_g^{\rho}) \right] \\
&+ (u_{\mathrm{z}}^{\rho} - u_g^{\rho}) \frac{\partial}{\partial \rho} n_{\mathrm{z}} T_{\mathrm{z}} + \hat{u}_{\mathrm{z}\theta} \langle \boldsymbol{B} \cdot \nabla \cdot \overleftrightarrow{\pi}_{\mathrm{z}} \rangle - \frac{3}{2} n_{\mathrm{z}} \frac{T_{\mathrm{z}} - T_{\mathrm{e}}}{\tau_{\mathrm{ze}}} - \frac{3}{2} n_{\mathrm{z}} \frac{T_{\mathrm{z}} - T_{\mathrm{i}}}{\tau_{\mathrm{zi}}} \\
&- \nu_{\mathrm{L}} T_{\mathrm{z}} (n_{\mathrm{z}} - n_{\mathrm{zsup}}) - \frac{3}{2} \nu_{\mathrm{L}T_{\mathrm{z}}} n_{\mathrm{z}} (T_{\mathrm{z}} - T_{\mathrm{zsup}}) \\
&+ P_{\alpha \mathrm{z}} + P_{\mathrm{RFz}} \qquad (1.22)
\end{aligned}
$$

$$
\begin{aligned}
\frac{5}{2} m_{\mathrm{z}} n_{\mathrm{z}} \frac{\partial}{\partial t} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle = &- \hat{\mu}_2^{\mathrm{z}} \langle B u_{\mathrm{z}\parallel} \rangle - \hat{\mu}_3^{\mathrm{z}} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle + \hat{\mu}_2^{\mathrm{z}} B V_{1\mathrm{z}} + \hat{\mu}_3^{\mathrm{z}} B V_{2\mathrm{z}} \\
&- \ell_{21}^{\mathrm{ze}} \langle B u_{\mathrm{e}\parallel} \rangle - \ell_{21}^{\mathrm{zi}} \langle B u_{\mathrm{i}\parallel} \rangle - \ell_{21}^{\mathrm{zz}} \langle B u_{\mathrm{z}\parallel} \rangle - \ell_{21}^{\mathrm{zb}} \langle B u_{\mathrm{b}\parallel} \rangle \\
&+ \ell_{22}^{\mathrm{ze}} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle + \ell_{22}^{\mathrm{zi}} \langle B \hat{q}_{\mathrm{i}\parallel} \rangle + \ell_{22}^{\mathrm{zz}} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle
\end{aligned}
$$

$$- \nu_{\mathrm{L}} m_{\mathrm{z}} n_{\mathrm{z}} \langle B \hat{q}_{\mathrm{z}\|} \rangle - \nu_{0\mathrm{z}} m_{\mathrm{z}} n_{\mathrm{z}} \langle B \hat{q}_{\mathrm{z}\|} \rangle \tag{1.23}$$

$$\begin{aligned} \frac{1}{V'} \frac{\partial}{\partial t} (V' n_{\mathrm{b}}) = & -\frac{1}{V'} \frac{\partial}{\partial \rho} [V' n_{\mathrm{b}} (u_{\mathrm{b}}^{\rho} - u_g^{\rho})] + \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} - g \frac{P_{\mathrm{b}\perp}}{E_{\mathrm{b}}} - \nu_{\mathrm{b}} n_{\mathrm{b}} - \nu_{\mathrm{bL}} n_{\mathrm{b}} - \nu_{\mathrm{eff}} g (\sqrt{\delta} n_{\mathrm{b}} - n_{\mathrm{b}}^{\mathrm{rp}}) \\ & + \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle f_t D_{\mathrm{b}} \frac{\partial}{\partial \rho} n_{\mathrm{b}} \end{aligned} \tag{1.24}$$

$$\begin{aligned} 0 = & -(\langle B^2 \rangle \langle R^2 \rangle - I^2) \frac{\partial}{\partial \rho} n_{\mathrm{b}} T_{\mathrm{b}} - Z_{\mathrm{b}} e n_{\mathrm{b}} (\langle B^2 \rangle \langle R^2 \rangle - I^2) \frac{\partial}{\partial \rho} \Phi \\ & + Z_{\mathrm{b}} e n_{\mathrm{b}} \frac{\partial \psi}{\partial \rho} I \langle B u_{\mathrm{b}\|} \rangle - Z_{\mathrm{b}} e n_{\mathrm{b}} \frac{\partial \psi}{\partial \rho} \langle B^2 \rangle \langle R u_{\mathrm{b}\zeta} \rangle \end{aligned} \tag{1.25}$$

$$\begin{aligned} m_{\mathrm{b}} \frac{\partial}{\partial t} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle = & -(\hat{\mu}_1^{\mathrm{b}} - \ell_{11}^{\mathrm{bb}}) \langle B u_{\mathrm{b}\|} \rangle + \sum_{a \neq \mathrm{b}} \ell_{11}^{\mathrm{b}a} \langle B u_{a\|} \rangle + \hat{\mu}_1^{\mathrm{b}} B V_{1\mathrm{b}} + Z_{\mathrm{b}} e n_{\mathrm{b}} \langle B E_{\|}^{A} \rangle \\ & - \nu_{\mathrm{bL}} m_{\mathrm{b}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle - \nu_{\mathrm{b}} m_{\mathrm{b}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle - \nu_{0\mathrm{b}} m_{\mathrm{b}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle - \nu_{\mathrm{CX}} m_{\mathrm{b}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle + \langle B S_{\mathrm{m}\|\mathrm{b}} \rangle \end{aligned} \tag{1.26}$$

$$\begin{aligned} \frac{1}{V'} \frac{\partial}{\partial t} (V' m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle) = & -\frac{1}{V'} \frac{\partial}{\partial \rho} [V' (u_{\mathrm{b}}^{\rho} - u_g^{\rho}) m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle] \\ & + \ell_{11}^{\mathrm{be}} \langle R u_{\mathrm{e}\zeta} \rangle + \ell_{11}^{\mathrm{bi}} \langle R u_{\mathrm{i}\zeta} \rangle + \ell_{11}^{\mathrm{bz}} \langle R u_{\mathrm{z}\zeta} \rangle + \ell_{11}^{\mathrm{bb}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle \\ & + Z_{\mathrm{b}} e n_{\mathrm{b}} \dot{\psi} + Z_{\mathrm{b}} e \frac{\partial \psi}{\partial \rho} n_{\mathrm{b}} u_{\mathrm{b}}^{\rho} - \nu_{\mathrm{bL}} m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle - \nu_{\mathrm{b}} m_{\mathrm{b}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\zeta} \rangle \\ & - \nu_{0\mathrm{i}} m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle - \nu_{\mathrm{CX}} m_{\mathrm{b}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle + \frac{I}{\langle B^2 \rangle} \langle B S_{\mathrm{m}\|\mathrm{b}} \rangle \end{aligned} \tag{1.27}$$

$$\begin{aligned} \frac{1}{V'} \frac{\partial}{\partial t} (V' n_{01}) = & \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle D_{01} \frac{\partial}{\partial \rho} n_{01} - \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} n_{01} \\ & - \nu_{\mathrm{CX}} \frac{n_{\mathrm{i}}}{n_{01} + n_{02}} n_{01} + \gamma_0 \nu_{\mathrm{L}} (n_{\mathrm{i}} - n_{\mathrm{isup}}) + \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle \mathbf{\Gamma}_{\mathrm{i}}^{\mathrm{puff}} \cdot \nabla \rho \rangle \Big|_{\mathrm{b}} \end{aligned} \tag{1.28}$$

$$\frac{1}{V'} \frac{\partial}{\partial t} (V' n_{02}) = \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle D_{02} \frac{\partial}{\partial \rho} n_{02} - \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} n_{02} + \nu_{\mathrm{CX}} \frac{n_{\mathrm{i}}}{n_{01} + n_{02}} n_{01} \tag{1.29}$$

$$\frac{1}{V'} \frac{\partial}{\partial t} (V' n_{03}) = \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle D_{03} \frac{\partial}{\partial \rho} n_{03} - \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} n_{03} + f_{\mathrm{CX}} \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} \tag{1.30}$$

$$\begin{aligned} \frac{1}{V'} \frac{\partial}{\partial t} (V' \bar{n}_z) = & \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle D_{nz} \frac{\partial}{\partial \rho} \bar{n}_z + \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle |\nabla \rho|^2 \rangle \Delta D_{\mathrm{C}0} \frac{\partial}{\partial \rho} n_{\mathrm{C}0} - n_{\mathrm{e}} \bar{n}_z f_5 S_5 + n_{\mathrm{e}} n_{\mathrm{z}} A_6 \\ & + \gamma_{0\mathrm{z}} \nu_{\mathrm{L}} (n_{\mathrm{z}} - n_{\mathrm{zsup}}) + \gamma_{\mathrm{i,z}} \nu_{\mathrm{L}} (n_{\mathrm{i}} - n_{\mathrm{isup}}) - (1 - \gamma_{0\mathrm{z}}) \nu_{\mathrm{L}} (\bar{n}_z - n_{\mathrm{C}0}) \\ & + \frac{1}{V'} \frac{\partial}{\partial \rho} V' \langle \mathbf{\Gamma}_{\mathrm{z}}^{\mathrm{puff}} \cdot \nabla \rho \rangle \Big|_{\mathrm{b}} \end{aligned} \tag{1.31}$$

$$\frac{1}{V'} \frac{\partial}{\partial t} (V' g n_{\mathrm{b}}^{\mathrm{rp}}) = g \frac{P_{\mathrm{b}\perp}}{E_{\mathrm{b}}} + g \nu_{\mathrm{eff}} (\sqrt{\delta} n_{\mathrm{b}} - n_{\mathrm{b}}^{\mathrm{rp}}) - g \nu_{\mathrm{b}} n_{\mathrm{b}}^{\mathrm{rp}}$$

$$- \frac{1}{V'} \frac{\partial}{\partial \rho} V' g \left(\langle |\nabla \rho| \rangle u_{\mathrm{b}}^{\mathrm{rp}} n_{\mathrm{b}}^{\mathrm{rp}} - \langle |\nabla \rho|^2 \rangle D_{\mathrm{b}}^{\mathrm{rp}} \frac{\partial}{\partial \rho} n_{\mathrm{b}}^{\mathrm{rp}} \right) \tag{1.32}$$

2 Additional important relationship

$$\bar{n}_z = \sum_{i=0}^{\max(Z)-1} n_i \tag{2.1}$$

$$B_p = \frac{|\nabla \rho|}{R} \frac{\partial \psi}{\partial \rho}, \quad \langle B_p^2 \rangle = \left\langle \frac{|\nabla V|^2}{R^2} \right\rangle \left(\frac{\partial \psi}{\partial V} \right)^2 \tag{2.2}$$

$$B_t = \frac{I}{R} = \frac{1}{R} \frac{4\pi^2}{\langle R^{-2} \rangle} \frac{\partial \psi_t}{\partial V}, \quad \langle B_t^2 \rangle = \frac{16\pi^4}{\langle R^{-2} \rangle} \left(\frac{\partial \psi_t}{\partial V} \right)^2 \tag{2.3}$$

$$\langle B^2 \rangle = \langle B_t^2 \rangle + \langle B_p^2 \rangle = \frac{16\pi^4}{\langle R^{-2} \rangle} \left(\frac{\partial \psi_t}{\partial V} \right)^2 + \left\langle \frac{|\nabla V|^2}{R^2} \right\rangle \left(\frac{\partial \psi}{\partial V} \right)^2 \tag{2.4}$$

$$\langle B^\theta \rangle = \frac{4\pi^2}{V'} \frac{\partial \psi}{\partial \rho} = 4\pi^2 \frac{\partial \psi}{\partial V} = \frac{I \langle R^{-2} \rangle}{q} = 2\pi \left(\oint \frac{dl_p}{B_p} \right)^{-1} \tag{2.5}$$

$$\hat{q}^2 \equiv \frac{I^2}{2\langle B_p^2 \rangle} \left(\left\langle \frac{1}{R^2} \right\rangle - \frac{1}{\langle R^2 \rangle} \right) \tag{2.6}$$

$$\langle B^2 \rangle \langle R^2 \rangle - I^2 = \langle R^2 \rangle \left\langle \frac{|\nabla V|^2}{R^2} \right\rangle \left(\frac{\partial \psi}{\partial V} \right)^2 (1 + 2\hat{q}^2) \tag{2.7}$$

$$\langle B \hat{q}_{s\|} \rangle \equiv \frac{2 \langle B q_{s\|} \rangle}{5 p_s} \tag{2.8}$$

$$\hat{u}_{s\theta} = \frac{\langle B u_{s\|} \rangle - B V_{1s}}{\langle B^2 \rangle} \tag{2.9}$$

$$\begin{aligned} \left\langle \frac{u_{s\zeta}}{R} \right\rangle &= \frac{1}{1 + 2\hat{q}^2} \frac{1}{\langle R^2 \rangle} \langle R u_{s\zeta} \rangle + \frac{2\hat{q}^2}{1 + 2\hat{q}^2} \frac{1}{I} \langle B u_{s\|} \rangle \\ &= \frac{\langle R u_{s\zeta} \rangle}{\langle R^2 \rangle} + \frac{I}{\langle B^2 \rangle} \left(\left\langle \frac{1}{R^2} \right\rangle - \frac{1}{\langle R^2 \rangle} \right) (\langle B u_{s\|} \rangle - B V_{1s}) \\ &= \frac{\langle R u_{s\zeta} \rangle}{\langle R^2 \rangle} + I \left(\left\langle \frac{1}{R^2} \right\rangle - \frac{1}{\langle R^2 \rangle} \right) \hat{u}_{s\theta} \end{aligned} \tag{2.10}$$

$$\langle R \hat{q}_{s\zeta} \rangle = \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{1}{I} B V_{2s} + \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{s\|} \rangle \tag{2.11}$$

$$B V_{1s} = -I \left(\frac{1}{e_s n_s} \frac{\partial n_s T_s}{\partial \psi} + \frac{\partial \Phi}{\partial \psi} \right), \quad B V_{2s} = -\frac{I}{e_s} \frac{\partial T_s}{\partial \psi} \tag{2.12}$$

$$u^\rho \equiv \langle \boldsymbol{u}\cdot\nabla\rho\rangle, \quad u^V = \langle \boldsymbol{u}\cdot\nabla V\rangle \tag{2.13}$$

$$u^\rho_g = -\frac{4\pi^2}{IV'\langle R^{-2}\rangle}\dot{\psi}_t, \quad u^V_g = -\frac{4\pi^2}{I\langle R^{-2}\rangle}\dot{\psi}_t = -\frac{V'}{2\rho\psi_{ta}}\dot{\psi}_t \tag{2.14}$$

$$\frac{1}{V'}\frac{\partial V'}{\partial t} = \frac{\partial}{\partial V}\langle \boldsymbol{u}_g\cdot\nabla V\rangle = \frac{\partial u^V_g}{\partial V} \tag{2.15}$$

$$\langle E_r\rangle = -\langle|\nabla\rho|^2\rangle\frac{\partial r}{\partial\rho}\frac{\partial}{\partial\rho}\Phi \tag{2.16}$$

$$\langle BE_\parallel\rangle = I\langle R^{-2}\rangle\left(-\frac{\dot{\psi}_t}{q}+\dot{\psi}\right) \tag{2.17}$$

$$RE_\zeta = \dot{\psi} = \frac{V_{\rm loop}}{2\pi} \tag{2.18}$$

$$\langle \boldsymbol{B}\cdot\nabla\cdot\overleftrightarrow{\pi}_{\rm i}\rangle = \hat{\mu}^{\rm i}_1(\langle Bu_{\rm i\parallel}\rangle - BV_{\rm 1i}) + \hat{\mu}^{\rm i}_2\left(\frac{2\langle Bq_{\rm i\parallel}\rangle}{5p_{\rm i}} - BV_{\rm 2i}\right) \tag{2.19}$$

$$\hat{\ell}^{ab}_{ij} \equiv \frac{\ell^{ab}_{ij}}{m_a n_a}, \quad \hat{\ell}^{\rm bb}_{ij} \equiv \frac{\ell^{\rm bb}_{ij}}{m_{\rm e}n_{\rm e}} \tag{2.20}$$

$$\Gamma^{\rm puff} \simeq \frac{\langle \boldsymbol{\Gamma}^{\rm puff}\cdot\nabla\rho\rangle}{\langle|\nabla\rho|\rangle} \tag{2.21}$$

3 Equations in the coordinates suitable for implementation

$$0 = \frac{\partial}{\partial V}\left[\langle|\nabla V|^2\rangle\frac{\partial}{\partial V}\Phi\right] + \frac{1}{\epsilon_0}\left(-en_{\rm e} + Z_{\rm i}en_{\rm i} + Z_{\rm z}en_{\rm z} + Z_{\rm b}en_{\rm b} + Z_{\rm b}eg n^{\rm rp}_{\rm b}\right) \tag{3.1}$$

$$\begin{aligned}\epsilon_0\frac{\partial}{\partial t}\dot{\psi}_t = {}& \frac{\partial}{\partial V}\left[\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{1}{\langle R^{-2}\rangle}\frac{\partial}{\partial V}\frac{\psi_t}{\mu_0}\right] - \frac{1}{\langle R^{-2}\rangle}\frac{\partial}{\partial V}\left[\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\right]\frac{\partial}{\partial V}\frac{\psi_t}{\mu_0}\\ & + \frac{1}{\langle B^\theta\rangle}\left[-e\left(n_{\rm e}\langle Bu_{\rm e\parallel}\rangle - In_{\rm e}\left\langle\frac{u_{\rm e\zeta}}{R}\right\rangle\right) + Z_{\rm i}e\left(n_{\rm i}\langle Bu_{\rm i\parallel}\rangle - In_{\rm i}\left\langle\frac{u_{\rm i\zeta}}{R}\right\rangle\right)\right.\\ & \left. + Z_{\rm z}e\left(n_{\rm z}\langle Bu_{\rm z\parallel}\rangle - In_{\rm z}\left\langle\frac{u_{\rm z\zeta}}{R}\right\rangle\right) + Z_{\rm b}e\left(n_{\rm b}\langle Bu_{\rm b\parallel}\rangle - In_{\rm b}\left\langle\frac{u_{\rm b\zeta}}{R}\right\rangle\right)\right]\end{aligned} \tag{3.2}$$

$$\begin{aligned}\frac{1}{c^2}\left\langle\frac{1}{R^2}\right\rangle\frac{\partial}{\partial t}\dot{\psi} = {}& \frac{\partial}{\partial V}\left[\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{\partial}{\partial V}\psi\right]\\ & - \mu_0\left(-en_{\rm e}\left\langle\frac{u_{\rm e\zeta}}{R}\right\rangle + Z_{\rm i}en_{\rm i}\left\langle\frac{u_{\rm i\zeta}}{R}\right\rangle + Z_{\rm z}en_{\rm z}\left\langle\frac{u_{\rm z\zeta}}{R}\right\rangle + Z_{\rm b}en_{\rm b}\left\langle\frac{u_{\rm b\zeta}}{R}\right\rangle\right)\end{aligned} \tag{3.3}$$

$$\frac{\partial}{\partial t}\psi = \dot{\psi} \tag{3.4}$$

$$\frac{\partial}{\partial t}\frac{\psi_t}{\mu_0} = \frac{\dot{\psi}_t}{\mu_0} \tag{3.5}$$

$$\begin{aligned}\frac{\partial}{\partial t}n_\mathrm{e} = &-\frac{\partial}{\partial V}n_\mathrm{e}u_\mathrm{e}^V + u_g^V\frac{\partial}{\partial V}n_\mathrm{e} + \nu_\mathrm{ionize}\frac{n_\mathrm{e}}{n_{01}+n_{02}+n_{03}}(n_{01}+n_{02}+n_{03})\\ &+ Z_\mathrm{z}n_\mathrm{e}\bar{n}_z f_5 S_5 - Z_\mathrm{z}n_\mathrm{e}n_\mathrm{z}A_6 - \nu_\mathrm{L}(n_\mathrm{e}-n_\mathrm{esup}) + (1-f_\mathrm{CX})\frac{P_\mathrm{b}}{E_\mathrm{b}}\end{aligned} \tag{3.6}$$

$$0 = -\frac{e}{m_\mathrm{e}}n_\mathrm{e}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{1}{I}BV_{1\mathrm{e}} - \frac{e}{m_\mathrm{e}}In_\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle + \frac{e}{m_\mathrm{e}}\langle B^2\rangle n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle \tag{3.7}$$

$$\begin{aligned}n_\mathrm{e}\frac{\partial}{\partial t}\langle Bu_{\mathrm{e}\|}\rangle = &-\frac{1}{m_\mathrm{e}}\hat{\mu}_1^\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle - \frac{1}{m_\mathrm{e}}\hat{\mu}_2^\mathrm{e}\langle B\hat{q}_{\mathrm{e}\|}\rangle + \frac{1}{m_\mathrm{e}}\hat{\mu}_1^\mathrm{e}BV_{1\mathrm{e}}\\ &+ \frac{1}{em_\mathrm{e}}\hat{\mu}_2^\mathrm{e}I\frac{\partial V}{\partial\psi}\frac{1}{n_\mathrm{e}}\frac{\partial}{\partial V}n_\mathrm{e}T_\mathrm{e} - \frac{1}{em_\mathrm{e}}\hat{\mu}_2^\mathrm{e}I\frac{\partial V}{\partial\psi}\frac{T_\mathrm{e}}{n_\mathrm{e}}\frac{\partial}{\partial V}n_\mathrm{e}\\ &+ \hat{\ell}_{11}^{\mathrm{ee}}n_\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle + \hat{\ell}_{11}^{\mathrm{ei}}n_\mathrm{e}\langle Bu_{\mathrm{i}\|}\rangle + \hat{\ell}_{11}^{\mathrm{ez}}n_\mathrm{e}\langle Bu_{\mathrm{z}\|}\rangle + \hat{\ell}_{11}^{\mathrm{eb}}\frac{n_\mathrm{e}}{n_\mathrm{b}}n_\mathrm{b}\langle Bu_{\mathrm{b}\|}\rangle\\ &- \hat{\ell}_{12}^{\mathrm{ee}}n_\mathrm{e}\langle B\hat{q}_{\mathrm{e}\|}\rangle - \hat{\ell}_{12}^{\mathrm{ei}}n_\mathrm{e}\langle B\hat{q}_{\mathrm{i}\|}\rangle - \hat{\ell}_{12}^{\mathrm{ez}}n_\mathrm{e}\langle B\hat{q}_{\mathrm{z}\|}\rangle\\ &- e\frac{n_\mathrm{e}}{m_\mathrm{e}}I\langle R^{-2}\rangle\left(-\frac{\dot{\psi}_t}{q}+\dot{\psi}\right) - \nu_\mathrm{L}n_\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle - \nu_{0\mathrm{e}}n_\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle\\ &- \frac{\partial}{\partial V}\langle|\nabla\rho|\rangle v_{\mathrm{e}\zeta}n_\mathrm{e}\langle Bu_{\mathrm{e}\|}\rangle + \langle|\nabla V|\rangle v_{\mathrm{e}\zeta}n_\mathrm{e}\left(\frac{\partial}{\partial V}\frac{\langle B^2\rangle}{I}\right)\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\|}\rangle\\ &+ \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\mathrm{e}\zeta}n_\mathrm{e}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial V}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\|}\rangle\\ &- \langle|\nabla V|^2\rangle\chi_{\mathrm{e}\zeta}n_\mathrm{e}\left(\frac{\partial}{\partial V}\frac{\langle B^2\rangle}{I}\right)\frac{\partial}{\partial V}\frac{I}{\langle B^2\rangle}\langle Bu_{\mathrm{e}\|}\rangle - \frac{1}{m_\mathrm{e}}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial V}\langle\Pi_\mathrm{e}^{\mathrm{res}}\rangle\\ &- \frac{\partial}{\partial V}\left[\langle|\nabla V|\rangle v_{\mathrm{e}\zeta}n_\mathrm{e}\frac{\langle R^2\rangle}{\langle B^2\rangle}\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{\partial\psi}{\partial V}(1+2\hat{q}^2)\frac{1}{I}\frac{\partial\psi}{\partial V}BV_{1\mathrm{e}}\right.\\ &\left.-\langle|\nabla V|^2\rangle\chi_{\mathrm{e}\zeta}n_\mathrm{e}\frac{\partial}{\partial V}\frac{\langle R^2\rangle}{\langle B^2\rangle}\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{\partial\psi}{\partial V}(1+2\hat{q}^2)\frac{1}{I}\frac{\partial\psi}{\partial V}BV_{1\mathrm{e}}\right]\end{aligned} \tag{3.8}$$

$$\begin{aligned}\frac{\partial}{\partial t}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle = &-(u_\mathrm{e}^V-u_g^V)\frac{\partial}{\partial V}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle - \frac{\partial u_\mathrm{e}^V}{\partial V}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle - \frac{\partial}{\partial V}[\langle|\nabla V|\rangle v_{\mathrm{e}\zeta}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle]\\ &+ \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\mathrm{e}\zeta}\frac{\partial}{\partial V}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle - \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\mathrm{e}\zeta}\langle Ru_{\mathrm{e}\zeta}\rangle\frac{\partial}{\partial V}n_\mathrm{e} - \frac{1}{m_\mathrm{e}}\frac{\partial}{\partial V}\langle\Pi_\mathrm{e}^{\mathrm{res}}\rangle\\ &+ \hat{\ell}_{11}^{\mathrm{ee}}n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle + \hat{\ell}_{11}^{\mathrm{ei}}\frac{n_\mathrm{e}}{n_\mathrm{i}}n_\mathrm{i}\langle Ru_{\mathrm{i}\zeta}\rangle + \hat{\ell}_{11}^{\mathrm{ez}}\frac{n_\mathrm{e}}{n_\mathrm{z}}n_\mathrm{z}\langle Ru_{\mathrm{z}\zeta}\rangle + \hat{\ell}_{11}^{\mathrm{eb}}\frac{n_\mathrm{e}}{n_\mathrm{b}}n_\mathrm{b}\langle Ru_{\mathrm{b}\zeta}\rangle\end{aligned}$$

$$
\begin{aligned}
&- \hat{\ell}^{\mathrm{ee}}_{12} n_{\mathrm{e}} \frac{1}{e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{e}} - \hat{\ell}^{\mathrm{ee}}_{12} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle \\
&+ \hat{\ell}^{\mathrm{ei}}_{12} n_{\mathrm{e}} \frac{1}{Z_{\mathrm{i}} e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{i}} - \hat{\ell}^{\mathrm{ei}}_{12} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{i}\parallel} \rangle \\
&+ \hat{\ell}^{\mathrm{ez}}_{12} n_{\mathrm{e}} \frac{1}{Z_{\mathrm{z}} e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{z}} - \hat{\ell}^{\mathrm{ez}}_{12} n_{\mathrm{e}} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle \\
&- e \frac{n_{\mathrm{e}}}{m_{\mathrm{e}}} \dot{\psi} - e \frac{1}{m_{\mathrm{e}}} \frac{\partial \psi}{\partial V} n_{\mathrm{e}} u^V_{\mathrm{e}} \\
&- \frac{e^2 \langle |\nabla V|^2 \rangle D_{\mathrm{e}}}{m_{\mathrm{e}} T_{\mathrm{e}}} \frac{\partial \psi}{\partial V} \left[n_{\mathrm{e}} \frac{\partial \psi}{\partial V} \frac{1}{I} B V_{1\mathrm{e}} + n_{\mathrm{e}} \frac{\partial}{\partial V} \Phi + \frac{C_{\mathrm{T}} - 1}{e} \left(\frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - T_{\mathrm{e}} \frac{\partial}{\partial V} n_{\mathrm{e}} \right) + \frac{T_{\mathrm{e}}}{e} \frac{\partial \psi}{\partial V} C_{\mathrm{p}} n_{\mathrm{e}} \right] \\
&- \nu_{\mathrm{L}} n_{\mathrm{e}} \langle R u_{\mathrm{e}\zeta} \rangle - \nu_{0\mathrm{e}} n_{\mathrm{e}} \langle R u_{\mathrm{e}\zeta} \rangle
\end{aligned} \tag{3.9}
$$

$$
\begin{aligned}
\frac{3}{2} \frac{\partial}{\partial t} n_{\mathrm{e}} T_{\mathrm{e}} = &- \frac{5}{2} (u^V_{\mathrm{e}} - u^V_g) \frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - \frac{5}{2} \frac{\partial u^V_{\mathrm{e}}}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - \frac{\partial}{\partial V} [V^{\mathrm{hp}}_{\mathrm{e}} \langle |\nabla V| \rangle n_{\mathrm{e}} T_{\mathrm{e}}] \\
&+ \frac{\partial}{\partial V} \chi_{\mathrm{e}} \langle |\nabla V|^2 \rangle \frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - \frac{\partial}{\partial V} \chi_{\mathrm{e}} \langle |\nabla V|^2 \rangle T_{\mathrm{e}} \frac{\partial}{\partial V} n_{\mathrm{e}} \\
&- u^V_g \frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - u^V_{\mathrm{i}} \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} - u^V_{\mathrm{z}} \frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} - \hat{u}_{\mathrm{i}\theta} \langle \boldsymbol{B} \cdot \nabla \cdot \overleftrightarrow{\pi}_{\mathrm{i}} \rangle - \hat{u}_{\mathrm{z}\theta} \langle \boldsymbol{B} \cdot \nabla \cdot \overleftrightarrow{\pi}_{\mathrm{z}} \rangle \\
&- \frac{3}{2} \nu_{\mathrm{Tei}} \left(n_{\mathrm{e}} T_{\mathrm{e}} - \frac{n_{\mathrm{e}}}{n_{\mathrm{i}}} n_{\mathrm{i}} T_{\mathrm{i}} \right) - \frac{3}{2} \nu_{\mathrm{Tez}} \left(n_{\mathrm{e}} T_{\mathrm{e}} - \frac{n_{\mathrm{e}}}{n_{\mathrm{z}}} n_{\mathrm{z}} T_{\mathrm{z}} \right) \\
&- 4\pi^2 \frac{\partial \psi}{\partial V} \dot{\psi}_t (-e n_{\mathrm{e}} \hat{u}_{\mathrm{e}\theta} + Z_{\mathrm{i}} e n_{\mathrm{i}} \hat{u}_{\mathrm{i}\theta} + Z_{\mathrm{z}} e n_{\mathrm{z}} \hat{u}_{\mathrm{z}\theta}) \\
&+ \dot{\psi} \left(-e n_{\mathrm{e}} \left\langle \frac{u_{\mathrm{e}\zeta}}{R} \right\rangle + Z_{\mathrm{i}} e n_{\mathrm{i}} \left\langle \frac{u_{\mathrm{i}\zeta}}{R} \right\rangle + Z_{\mathrm{z}} e n_{\mathrm{z}} \left\langle \frac{u_{\mathrm{z}\zeta}}{R} \right\rangle \right) \\
&- \nu_{\mathrm{L}} (n_{\mathrm{e}} T_{\mathrm{e}} - n_{\mathrm{esup}} T_{\mathrm{e}}) - \frac{3}{2} \nu_{\mathrm{LT_e}} (n_{\mathrm{e}} T_{\mathrm{e}} - n_{\mathrm{e}} T_{\mathrm{esup}}) - E_{\mathrm{H}} \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} (n_{01} + n_{02} + n_{03}) \\
&+ f_{\mathrm{ei}} P_{\mathrm{b}} + P_{\alpha \mathrm{e}} + P_{\mathrm{RFe}} - P_{\mathrm{Br}}
\end{aligned} \tag{3.10}
$$

$$
\begin{aligned}
\frac{5}{2} n_{\mathrm{e}} \frac{\partial}{\partial t} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle = &- \frac{1}{m_{\mathrm{e}}} \hat{\mu}^{\mathrm{e}}_2 \langle B u_{\mathrm{e}\parallel} \rangle - \frac{1}{m_{\mathrm{e}}} \hat{\mu}^{\mathrm{e}}_3 \langle B \hat{q}_{\mathrm{e}\parallel} \rangle + \frac{1}{m_{\mathrm{e}}} \hat{\mu}^{\mathrm{e}}_2 B V_{1\mathrm{e}} \\
&+ \frac{1}{e m_{\mathrm{e}}} \hat{\mu}^{\mathrm{e}}_3 I \frac{\partial V}{\partial \psi} \frac{1}{n_{\mathrm{e}}} \frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - \frac{1}{e m_{\mathrm{e}}} \hat{\mu}^{\mathrm{e}}_3 I \frac{\partial V}{\partial \psi} \frac{T_{\mathrm{e}}}{n_{\mathrm{e}}} \frac{\partial}{\partial V} n_{\mathrm{e}} \\
&- \hat{\ell}^{\mathrm{ee}}_{21} n_{\mathrm{e}} \langle B u_{\mathrm{e}\parallel} \rangle - \hat{\ell}^{\mathrm{ei}}_{21} n_{\mathrm{e}} \langle B u_{\mathrm{i}\parallel} \rangle - \hat{\ell}^{\mathrm{ez}}_{21} n_{\mathrm{e}} \langle B u_{\mathrm{z}\parallel} \rangle - \hat{\ell}^{\mathrm{eb}}_{21} \frac{n_{\mathrm{e}}}{n_{\mathrm{b}}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\parallel} \rangle \\
&+ \hat{\ell}^{\mathrm{ee}}_{22} n_{\mathrm{e}} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle + \hat{\ell}^{\mathrm{ei}}_{22} n_{\mathrm{e}} \langle B \hat{q}_{\mathrm{i}\parallel} \rangle + \hat{\ell}^{\mathrm{ez}}_{22} n_{\mathrm{e}} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle \\
&- \nu_{\mathrm{L}} n_{\mathrm{e}} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle - \nu_{0\mathrm{e}} n_{\mathrm{e}} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle
\end{aligned} \tag{3.11}
$$

$$
0 = -n_{\mathrm{e}} \left\langle \frac{u_{\mathrm{e}\zeta}}{R} \right\rangle + \frac{1}{\langle R^2 \rangle} n_{\mathrm{e}} \langle R u_{\mathrm{e}\zeta} \rangle - \frac{2\hat{q}^2}{1 + 2\hat{q}^2} \frac{1}{\langle R^2 \rangle} n_{\mathrm{e}} \langle R u_{\mathrm{e}\zeta} \rangle + \frac{2\hat{q}^2}{1 + 2\hat{q}^2} \frac{1}{I} n_{\mathrm{e}} \langle B u_{\mathrm{e}\parallel} \rangle \tag{3.12}
$$

$$0 = -n_{\mathrm{e}} B V_{1\mathrm{e}} + \frac{1}{e}\frac{\partial V}{\partial \psi} I \frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - n_{\mathrm{e}} \frac{\partial V}{\partial \psi} I \frac{\partial}{\partial V} \Phi \tag{3.13}$$

$$\begin{aligned}\frac{\partial}{\partial t} n_{\mathrm{i}} = &-\frac{\partial}{\partial V} n_{\mathrm{i}} u_{\mathrm{i}}^{V} + u_g^{V} \frac{\partial}{\partial V} n_{\mathrm{i}} + \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} (n_{01} + n_{02} + n_{03}) - \nu_{\mathrm{L}} (n_{\mathrm{i}} - n_{\mathrm{isup}}) \\ &+ \nu_{\mathrm{b}} n_{\mathrm{b}} + \nu_{\mathrm{b}} g n_{\mathrm{b}}^{\mathrm{rp}} - f_{\mathrm{CX}} \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} + \nu_{\mathrm{bL}} n_{\mathrm{b}} + S_{\mathrm{LC}} - \nu_{\mathrm{OL}} n_{\mathrm{i}}\end{aligned} \tag{3.14}$$

$$0 = Z_{\mathrm{i}} \frac{e}{m_{\mathrm{i}}} n_{\mathrm{i}} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{1}{I} B V_{1\mathrm{i}} + Z_{\mathrm{i}} \frac{e}{m_{\mathrm{i}}} I n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle - Z_{\mathrm{i}} \frac{e}{m_{\mathrm{i}}} \langle B^2 \rangle n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle \tag{3.15}$$

$$\begin{aligned}n_{\mathrm{i}} \frac{\partial}{\partial t} \langle B u_{\mathrm{i}\parallel} \rangle = &-\frac{1}{m_{\mathrm{i}}} \hat{\mu}_1^{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle - \frac{1}{m_{\mathrm{i}}} \hat{\mu}_2^{\mathrm{i}} \langle B \hat{q}_{\mathrm{i}\parallel} \rangle + \frac{1}{m_{\mathrm{i}}} \hat{\mu}_1^{\mathrm{i}} B V_{1\mathrm{i}} \\ &- \frac{1}{Z_{\mathrm{i}} e m_{\mathrm{i}}} \hat{\mu}_2^{\mathrm{i}} I \frac{\partial V}{\partial \psi} \frac{1}{n_{\mathrm{i}}} \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} + \frac{1}{Z_{\mathrm{i}} e m_{\mathrm{i}}} \hat{\mu}_2^{\mathrm{i}} I \frac{\partial V}{\partial \psi} \frac{T_{\mathrm{i}}}{n_{\mathrm{i}}} \frac{\partial}{\partial V} n_{\mathrm{i}} \\ &+ \hat{\ell}_{11}^{\mathrm{ie}} n_{\mathrm{i}} \langle B u_{\mathrm{e}\parallel} \rangle + \hat{\ell}_{11}^{\mathrm{ii}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle + \hat{\ell}_{11}^{\mathrm{iz}} n_{\mathrm{i}} \langle B u_{\mathrm{z}\parallel} \rangle + \hat{\ell}_{11}^{\mathrm{ib}} \frac{n_{\mathrm{i}}}{n_{\mathrm{b}}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\parallel} \rangle \\ &- \hat{\ell}_{12}^{\mathrm{ie}} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{e}\parallel} \rangle - \hat{\ell}_{12}^{\mathrm{ii}} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{i}\parallel} \rangle - \hat{\ell}_{12}^{\mathrm{iz}} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{z}\parallel} \rangle \\ &+ Z_{\mathrm{i}} e \frac{n_{\mathrm{i}}}{m_{\mathrm{i}}} I \langle R^{-2} \rangle \left(-\frac{\dot{\psi}_t}{q} + \dot{\psi} \right) \\ &- \nu_{\mathrm{L}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle + \nu_{\mathrm{b}} \frac{m_{\mathrm{b}}}{m_{\mathrm{i}}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\parallel} \rangle - \nu_{0\mathrm{i}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle - \nu_{\mathrm{CX}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle - \nu_{\mathrm{LC}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle - \nu_{\mathrm{OL}} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle \\ &- \frac{\partial}{\partial V} \langle |\nabla \rho| \rangle v_{\mathrm{i}\zeta} n_{\mathrm{i}} \langle B u_{\mathrm{i}\parallel} \rangle + \langle |\nabla V| \rangle v_{\mathrm{i}\zeta} n_{\mathrm{i}} \left(\frac{\partial}{\partial V} \frac{\langle B^2 \rangle}{I} \right) \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{i}\parallel} \rangle \\ &+ \frac{\partial}{\partial V} \langle |\nabla V|^2 \rangle \chi_{\mathrm{i}\zeta} n_{\mathrm{i}} \frac{\langle B^2 \rangle}{I} \frac{\partial}{\partial V} \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{i}\parallel} \rangle \\ &- \langle |\nabla V|^2 \rangle \chi_{\mathrm{i}\zeta} n_{\mathrm{i}} \left(\frac{\partial}{\partial V} \frac{\langle B^2 \rangle}{I} \right) \frac{\partial}{\partial V} \frac{I}{\langle B^2 \rangle} \langle B u_{\mathrm{i}\parallel} \rangle - \frac{1}{m_{\mathrm{i}}} \frac{\langle B^2 \rangle}{I} \frac{\partial}{\partial V} \langle \Pi_{\mathrm{i}}^{\mathrm{res}} \rangle \\ &- \frac{\partial}{\partial V} \left[\langle |\nabla V| \rangle v_{\mathrm{i}\zeta} n_{\mathrm{i}} \frac{\langle R^2 \rangle}{\langle B^2 \rangle} \left\langle \frac{|\nabla V|^2}{R^2} \right\rangle \frac{\partial \psi}{\partial V} (1 + 2\hat{q}^2) \frac{1}{I} \frac{\partial \psi}{\partial V} B V_{1\mathrm{i}} \right. \\ &\left. - \langle |\nabla V|^2 \rangle \chi_{\mathrm{i}\zeta} n_{\mathrm{i}} \frac{\partial}{\partial V} \frac{\langle R^2 \rangle}{\langle B^2 \rangle} \left\langle \frac{|\nabla V|^2}{R^2} \right\rangle \frac{\partial \psi}{\partial V} (1 + 2\hat{q}^2) \frac{1}{I} \frac{\partial \psi}{\partial V} B V_{1\mathrm{i}} \right]\end{aligned} \tag{3.16}$$

$$\begin{aligned}\frac{\partial}{\partial t} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle = &-(u_{\mathrm{i}}^{V} - u_g^{V}) \frac{\partial}{\partial V} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \frac{\partial u_{\mathrm{i}}^{V}}{\partial V} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \frac{\partial}{\partial V} \langle |\nabla V| \rangle v_{\mathrm{i}\zeta} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle \\ &+ \frac{\partial}{\partial V} \langle |\nabla V|^2 \rangle \chi_{\mathrm{i}\zeta} \frac{\partial}{\partial V} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \frac{\partial}{\partial V} \langle |\nabla V|^2 \rangle \chi_{\mathrm{i}\zeta} \langle R u_{\mathrm{i}\zeta} \rangle \frac{\partial}{\partial V} n_{\mathrm{i}} - \frac{1}{m_{\mathrm{i}}} \frac{\partial}{\partial V} \langle \Pi_{\mathrm{i}}^{\mathrm{res}} \rangle \\ &+ \hat{\ell}_{11}^{\mathrm{ie}} \frac{n_{\mathrm{i}}}{n_{\mathrm{e}}} n_{\mathrm{e}} \langle R u_{\mathrm{e}\zeta} \rangle + \hat{\ell}_{11}^{\mathrm{ii}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle + \hat{\ell}_{11}^{\mathrm{iz}} \frac{n_{\mathrm{i}}}{n_{\mathrm{z}}} n_{\mathrm{z}} \langle R u_{\mathrm{z}\zeta} \rangle + \hat{\ell}_{11}^{\mathrm{ib}} \frac{n_{\mathrm{i}}}{n_{\mathrm{b}}} n_{\mathrm{b}} \langle R u_{\mathrm{b}\zeta} \rangle\end{aligned}$$

$$
\begin{aligned}
&- \hat{\ell}^{\mathrm{ie}}_{12} n_{\mathrm{i}} \frac{1}{e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{e}} - \hat{\ell}^{\mathrm{ie}}_{12} n_{\mathrm{i}} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{e}\|} \rangle \\
&+ \hat{\ell}^{\mathrm{ii}}_{12} n_{\mathrm{i}} \frac{1}{Z_{\mathrm{i}} e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{i}} - \hat{\ell}^{\mathrm{ii}}_{12} n_{\mathrm{i}} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{i}\|} \rangle \\
&+ \hat{\ell}^{\mathrm{iz}}_{12} n_{\mathrm{i}} \frac{1}{Z_{\mathrm{z}} e} \frac{1}{\langle B^2 \rangle} \left(\langle B^2 \rangle \langle R^2 \rangle - I^2 \right) \frac{\partial V}{\partial \psi} \frac{\partial}{\partial V} T_{\mathrm{z}} - \hat{\ell}^{\mathrm{iz}}_{12} n_{\mathrm{i}} \frac{I}{\langle B^2 \rangle} \langle B \hat{q}_{\mathrm{z}\|} \rangle \\
&+ Z_{\mathrm{i}} \frac{e}{m_{\mathrm{i}}} n_{\mathrm{i}} \dot{\psi} + Z_{\mathrm{i}} \frac{e}{m_{\mathrm{i}}} \frac{\partial \psi}{\partial V} n_{\mathrm{i}} u^V_{\mathrm{i}} \\
&+ Z_{\mathrm{i}} \frac{e^2 \langle |\nabla V|^2 \rangle D_{\mathrm{e}}}{m_{\mathrm{i}} T_{\mathrm{e}}} \frac{\partial \psi}{\partial V} \left[(n_{\mathrm{i}} + n_{\mathrm{b}}) \frac{\partial \psi}{\partial V} \frac{1}{I} B V_{1\mathrm{e}} + (n_{\mathrm{i}} + n_{\mathrm{b}}) \frac{\partial}{\partial V} \Phi \right. \\
&\left. + \frac{C_{\mathrm{T}} - 1}{e} \frac{n_{\mathrm{i}} + n_{\mathrm{b}}}{n_{\mathrm{e}}} \left(\frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - T_{\mathrm{e}} \frac{\partial}{\partial V} n_{\mathrm{e}} \right) + \frac{T_{\mathrm{e}}}{e} \frac{\partial \psi}{\partial V} C_{\mathrm{p}} n_{\mathrm{i}} + \frac{T_{\mathrm{e}}}{e} \frac{\partial \psi}{\partial V} C_{\mathrm{p}} n_{\mathrm{b}} \right] \\
&- \nu_{\mathrm{L}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle + \nu_{\mathrm{b}} \frac{m_{\mathrm{b}}}{m_{\mathrm{i}}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\zeta} \rangle - \nu_{0\mathrm{i}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \nu_{\mathrm{CX}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \nu_{\mathrm{OL}} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle
\end{aligned} \tag{3.17}
$$

$$
\begin{aligned}
\frac{3}{2} \frac{\partial}{\partial t} n_{\mathrm{i}} T_{\mathrm{i}} = &- \frac{5}{2} (u^V_{\mathrm{i}} - u^V_g) \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} - \frac{5}{2} \frac{\partial u^V_{\mathrm{i}}}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} - \frac{\partial}{\partial V} V^{\mathrm{hp}}_{\mathrm{i}} \langle |\nabla V| \rangle n_{\mathrm{i}} T_{\mathrm{i}} \\
&+ \frac{\partial}{\partial V} \chi_{\mathrm{i}} \langle |\nabla V|^2 \rangle \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} - \frac{\partial}{\partial V} \chi_{\mathrm{i}} \langle |\nabla V|^2 \rangle T_{\mathrm{i}} \frac{\partial}{\partial V} n_{\mathrm{i}} \\
&+ (u^V_{\mathrm{i}} - u^V_g) \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} + \hat{u}_{\mathrm{i}\theta} \langle \boldsymbol{B} \cdot \nabla \cdot \overleftrightarrow{\pi}_{\mathrm{i}} \rangle - \frac{3}{2} \nu_{\mathrm{Tei}} \left(\frac{n_{\mathrm{e}}}{n_{\mathrm{i}}} n_{\mathrm{i}} T_{\mathrm{i}} - n_{\mathrm{e}} T_{\mathrm{e}} \right) - \frac{3}{2} \nu_{\mathrm{Tiz}} \left(n_{\mathrm{i}} T_{\mathrm{i}} - \frac{n_{\mathrm{i}}}{n_{\mathrm{z}}} n_{\mathrm{z}} T_{\mathrm{z}} \right) \\
&- \nu_{\mathrm{L}} \left(n_{\mathrm{i}} T_{\mathrm{i}} - n_{\mathrm{isup}} T_{\mathrm{i}} \right) - \frac{3}{2} \nu_{\mathrm{LT_i}} (n_{\mathrm{i}} T_{\mathrm{i}} - n_{\mathrm{i}} T_{\mathrm{isup}}) \\
&+ \frac{3}{2} \nu_{\mathrm{ionize}} \frac{n_{\mathrm{e}}}{n_{01} + n_{02} + n_{03}} (n_{01} T_{01} + n_{02} T_{02} + n_{03} T_{03}) \\
&- \frac{3}{2} \nu_{\mathrm{CX}} (n_{\mathrm{i}} T_{\mathrm{i}} - n_{\mathrm{i}} T_{01}) + (1 - f_{\mathrm{ei}}) P_{\mathrm{b}} + m_{\mathrm{b}} \frac{1}{n_{\mathrm{b}}} \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} \frac{\langle B u_{\mathrm{i}\|} \rangle}{\langle B^2 \rangle} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle + P_{\alpha \mathrm{i}} + P_{\mathrm{RFi}}
\end{aligned} \tag{3.18}
$$

$$
\begin{aligned}
\frac{5}{2} n_{\mathrm{i}} \frac{\partial}{\partial t} \langle B \hat{q}_{\mathrm{i}\|} \rangle = &- \frac{1}{m_{\mathrm{i}}} \hat{\mu}^{\mathrm{i}}_2 \langle B u_{\mathrm{i}\|} \rangle - \frac{1}{m_{\mathrm{i}}} \hat{\mu}^{\mathrm{i}}_3 \langle B \hat{q}_{\mathrm{i}\|} \rangle + \frac{1}{m_{\mathrm{i}}} \hat{\mu}^{\mathrm{i}}_2 B V_{1\mathrm{i}} \\
&- \frac{1}{Z_{\mathrm{i}} e m_{\mathrm{i}}} \hat{\mu}^{\mathrm{i}}_3 I \frac{\partial V}{\partial \psi} \frac{1}{n_{\mathrm{i}}} \frac{\partial}{\partial V} n_{\mathrm{i}} T_{\mathrm{i}} + \frac{1}{Z_{\mathrm{i}} e m_{\mathrm{i}}} \hat{\mu}^{\mathrm{i}}_3 I \frac{\partial V}{\partial \psi} \frac{T_{\mathrm{i}}}{n_{\mathrm{i}}} \frac{\partial}{\partial V} n_{\mathrm{i}} \\
&- \hat{\ell}^{\mathrm{ie}}_{21} n_{\mathrm{i}} \langle B u_{\mathrm{e}\|} \rangle - \hat{\ell}^{\mathrm{ii}}_{21} n_{\mathrm{i}} \langle B u_{\mathrm{i}\|} \rangle - \hat{\ell}^{\mathrm{iz}}_{21} n_{\mathrm{i}} \langle B u_{\mathrm{z}\|} \rangle - \hat{\ell}^{\mathrm{ib}}_{21} \frac{n_{\mathrm{i}}}{n_{\mathrm{b}}} n_{\mathrm{b}} \langle B u_{\mathrm{b}\|} \rangle \\
&+ \hat{\ell}^{\mathrm{ie}}_{22} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{e}\|} \rangle + \hat{\ell}^{\mathrm{ii}}_{22} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{i}\|} \rangle + \hat{\ell}^{\mathrm{iz}}_{22} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{z}\|} \rangle \\
&- \nu_{\mathrm{L}} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{i}\|} \rangle - \nu_{0\mathrm{i}} n_{\mathrm{i}} \langle B \hat{q}_{\mathrm{i}\|} \rangle
\end{aligned} \tag{3.19}
$$

$$
0 = - n_{\mathrm{i}} \left\langle \frac{u_{\mathrm{i}\zeta}}{R} \right\rangle + \frac{1}{\langle R^2 \rangle} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle - \frac{2\hat{q}^2}{1 + 2\hat{q}^2} \frac{1}{\langle R^2 \rangle} n_{\mathrm{i}} \langle R u_{\mathrm{i}\zeta} \rangle + \frac{2\hat{q}^2}{1 + 2\hat{q}^2} \frac{1}{I} n_{\mathrm{i}} \langle B u_{\mathrm{i}\|} \rangle \tag{3.20}
$$

$$0 = -n_{\rm i} BV_{1\rm i} - \frac{1}{Z_{\rm i} e}\frac{\partial V}{\partial \psi} I \frac{\partial}{\partial V} n_{\rm i} T_{\rm i} - n_{\rm i}\frac{\partial V}{\partial \psi} I \frac{\partial}{\partial V}\Phi \tag{3.21}$$

$$\frac{\partial}{\partial t} n_{\rm z} = -\frac{\partial}{\partial V} n_{\rm z} u_{\rm z}^{V} + u_g^{V}\frac{\partial}{\partial V} n_{\rm z} + n_{\rm e}\bar{n}_z f_5 S_5 - n_{\rm e} n_{\rm z} A_6 - \nu_{\rm L}(n_{\rm z} - n_{\rm isup}) + S_{\rm LC} - \nu_{\rm OL} n_{\rm z} \tag{3.22}$$

$$0 = Z_{\rm z}\frac{e}{m_{\rm z}} n_{\rm z}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{1}{I} BV_{1\rm z} + Z_{\rm z}\frac{e}{m_{\rm z}} I n_{\rm z}\langle B u_{\rm z\parallel}\rangle - Z_{\rm z}\frac{e}{m_{\rm z}}\langle B^2\rangle n_{\rm z}\langle R u_{\rm z\zeta}\rangle \tag{3.23}$$

$$\begin{aligned}
n_{\rm z}\frac{\partial}{\partial t}\langle B u_{\rm z\parallel}\rangle =& -\frac{1}{m_{\rm i}}\hat{\mu}_1^{\rm z}\langle B u_{\rm z\parallel}\rangle - \frac{1}{m_{\rm i}}\hat{\mu}_2^{\rm z}\langle B\hat{q}_{\rm z\parallel}\rangle + \frac{1}{m_{\rm z}}\hat{\mu}_1^{\rm i} BV_{1\rm z} \\
&- \frac{1}{Z_{\rm z} e m_{\rm z}}\hat{\mu}_2^{\rm z} I\frac{\partial V}{\partial\psi}\frac{1}{n_{\rm z}}\frac{\partial}{\partial V} n_{\rm z} T_{\rm z} + \frac{1}{Z_{\rm z} e m_{\rm z}}\hat{\mu}_2^{\rm z} I\frac{\partial V}{\partial\psi}\frac{T_{\rm z}}{n_{\rm z}}\frac{\partial}{\partial V} n_{\rm z} \\
&+ \hat{\ell}_{11}^{\rm ze} n_{\rm z}\langle B u_{\rm e\parallel}\rangle + \hat{\ell}_{11}^{\rm zi} n_{\rm z}\langle B u_{\rm i\parallel}\rangle + \hat{\ell}_{11}^{\rm zz} n_{\rm z}\langle B u_{\rm z\parallel}\rangle + \hat{\ell}_{11}^{\rm zb}\frac{n_{\rm z}}{n_{\rm b}} n_{\rm b}\langle B u_{\rm b\parallel}\rangle \\
&- \hat{\ell}_{12}^{\rm ze} n_{\rm z}\langle B\hat{q}_{\rm e\parallel}\rangle - \hat{\ell}_{12}^{\rm zi} n_{\rm z}\langle B\hat{q}_{\rm i\parallel}\rangle - \hat{\ell}_{12}^{\rm zz} n_{\rm z}\langle B\hat{q}_{\rm z\parallel}\rangle \\
&+ Z_{\rm z} e\frac{n_{\rm z}}{m_{\rm z}} I\langle R^{-2}\rangle\left(-\frac{\dot{\psi}_t}{q} + \dot{\psi}\right) \\
&- \nu_{\rm L} n_{\rm z}\langle B u_{\rm z\parallel}\rangle - \nu_{0\rm z} n_{\rm z}\langle B u_{\rm z\parallel}\rangle - \nu_{\rm OL} n_{\rm z}\langle B u_{\rm z\parallel}\rangle \\
&- \frac{\partial}{\partial V}\langle|\nabla\rho|\rangle v_{\rm z\zeta} n_{\rm z}\langle B u_{\rm z\parallel}\rangle + \langle|\nabla V|\rangle v_{\rm z\zeta} n_{\rm z}\left(\frac{\partial}{\partial V}\frac{\langle B^2\rangle}{I}\right)\frac{I}{\langle B^2\rangle}\langle B u_{\rm z\parallel}\rangle \\
&+ \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\rm z\zeta} n_{\rm z}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial V}\frac{I}{\langle B^2\rangle}\langle B u_{\rm z\parallel}\rangle \\
&- \langle|\nabla V|^2\rangle\chi_{\rm z\zeta} n_{\rm z}\left(\frac{\partial}{\partial V}\frac{\langle B^2\rangle}{I}\right)\frac{\partial}{\partial V}\frac{I}{\langle B^2\rangle}\langle B u_{\rm z\parallel}\rangle - \frac{1}{m_{\rm z}}\frac{\langle B^2\rangle}{I}\frac{\partial}{\partial V}\langle\Pi_{\rm z}^{\rm res}\rangle \\
&- \frac{\partial}{\partial V}\left[\langle|\nabla V|\rangle v_{\rm z\zeta} n_{\rm z}\frac{\langle R^2\rangle}{\langle B^2\rangle}\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{\partial\psi}{\partial V}(1+2\hat{q}^2)\frac{1}{I}\frac{\partial\psi}{\partial V} BV_{1\rm z}\right. \\
&\left.- \langle|\nabla V|^2\rangle\chi_{\rm z\zeta} n_{\rm z}\frac{\partial}{\partial V}\frac{\langle R^2\rangle}{\langle B^2\rangle}\left\langle\frac{|\nabla V|^2}{R^2}\right\rangle\frac{\partial\psi}{\partial V}(1+2\hat{q}^2)\frac{1}{I}\frac{\partial\psi}{\partial V} BV_{1\rm z}\right]
\end{aligned} \tag{3.24}$$

$$\begin{aligned}
\frac{\partial}{\partial t} n_{\rm z}\langle R u_{\rm z\zeta}\rangle =& -(u_{\rm z}^{V} - u_g^{V})\frac{\partial}{\partial V} n_{\rm z}\langle R u_{\rm z\zeta}\rangle - \frac{\partial u_{\rm z}^{V}}{\partial V} n_{\rm z}\langle R u_{\rm z\zeta}\rangle - \frac{\partial}{\partial V}\langle|\nabla V|\rangle v_{\rm z\zeta} n_{\rm z}\langle R u_{\rm z\zeta}\rangle \\
&+ \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\rm z\zeta}\frac{\partial}{\partial V} n_{\rm z}\langle R u_{\rm z\zeta}\rangle - \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\chi_{\rm z\zeta}\langle R u_{\rm z\zeta}\rangle\frac{\partial}{\partial V} n_{\rm z} - \frac{1}{m_{\rm z}}\frac{\partial}{\partial V}\langle\Pi_{\rm z}^{\rm res}\rangle \\
&+ \hat{\ell}_{11}^{\rm ze}\frac{n_{\rm z}}{n_{\rm e}} n_{\rm e}\langle R u_{\rm e\zeta}\rangle + \hat{\ell}_{11}^{\rm zi}\frac{n_{\rm z}}{n_{\rm i}} n_{\rm i}\langle R u_{\rm i\zeta}\rangle + \hat{\ell}_{11}^{\rm zz} n_{\rm z}\langle R u_{\rm z\zeta}\rangle + \hat{\ell}_{11}^{\rm zb}\frac{n_{\rm z}}{n_{\rm b}} n_{\rm b}\langle R u_{\rm b\zeta}\rangle \\
&- \hat{\ell}_{12}^{\rm ze} n_{\rm z}\frac{1}{e}\frac{1}{\langle B^2\rangle}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{\partial V}{\partial\psi}\frac{\partial}{\partial V} T_{\rm e} - \hat{\ell}_{12}^{\rm ze} n_{\rm z}\frac{I}{\langle B^2\rangle}\langle B\hat{q}_{\rm e\parallel}\rangle
\end{aligned}$$

$$
\begin{aligned}
&+\hat{\ell}_{12}^{\mathrm{zi}} n_{\mathrm{z}} \frac{1}{Z_{\mathrm{i}} e} \frac{1}{\langle B^2\rangle}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{\partial V}{\partial \psi}\frac{\partial}{\partial V}T_{\mathrm{i}} - \hat{\ell}_{12}^{\mathrm{zi}} n_{\mathrm{z}} \frac{I}{\langle B^2\rangle}\langle B\hat{q}_{\mathrm{i}\|}\rangle \\
&+\hat{\ell}_{12}^{\mathrm{zz}} n_{\mathrm{z}} \frac{1}{Z_{\mathrm{z}} e} \frac{1}{\langle B^2\rangle}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{\partial V}{\partial \psi}\frac{\partial}{\partial V}T_{\mathrm{z}} - \hat{\ell}_{12}^{\mathrm{zz}} n_{\mathrm{z}} \frac{I}{\langle B^2\rangle}\langle B\hat{q}_{\mathrm{z}\|}\rangle \\
&+ Z_{\mathrm{z}}\frac{e}{m_{\mathrm{z}}} n_{\mathrm{z}}\dot{\psi} + Z_{\mathrm{z}}\frac{e}{m_{\mathrm{z}}}\frac{\partial \psi}{\partial V} n_{\mathrm{z}} u_{\mathrm{z}}^{V} \\
&+ Z_{\mathrm{z}}\frac{e\langle|\nabla V|^2\rangle D_{\mathrm{e}}}{m_{\mathrm{z}} T_{\mathrm{e}}}\frac{\partial \psi}{\partial V}\left[n_{\mathrm{z}}\frac{\partial \psi}{\partial V}\frac{1}{I} B V_{1\mathrm{e}} + n_{\mathrm{z}}\frac{\partial}{\partial V}\Phi\right. \\
&\left.+\frac{C_{\mathrm{T}}-1}{e}\frac{n_{\mathrm{z}}}{n_{\mathrm{e}}}\left(\frac{\partial}{\partial V} n_{\mathrm{e}} T_{\mathrm{e}} - T_{\mathrm{e}}\frac{\partial}{\partial V} n_{\mathrm{e}}\right) + \frac{T_{\mathrm{e}}}{e}\frac{\partial \psi}{\partial V} C_{\mathrm{p}} n_{\mathrm{z}}\right] \\
&- \nu_{\mathrm{L}} n_{\mathrm{z}}\langle R u_{\mathrm{z}\zeta}\rangle - \nu_{0\mathrm{z}} n_{\mathrm{z}}\langle R u_{\mathrm{z}\zeta}\rangle - \nu_{\mathrm{OL}} n_{\mathrm{z}}\langle R u_{\mathrm{z}\zeta}\rangle
\end{aligned} \tag{3.25}
$$

$$
\begin{aligned}
\frac{3}{2}\frac{\partial}{\partial t} n_{\mathrm{z}} T_{\mathrm{z}} &= -\frac{5}{2}(u_{\mathrm{z}}^{V} - u_{g}^{V})\frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} - \frac{5}{2}\frac{\partial u_{\mathrm{z}}^{V}}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} - \frac{\partial}{\partial V} V_{\mathrm{z}}^{\mathrm{hp}}\langle|\nabla V|\rangle n_{\mathrm{z}} T_{\mathrm{z}} \\
&+ \frac{\partial}{\partial V}\chi_{\mathrm{z}}\langle|\nabla V|^2\rangle\frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} - \frac{\partial}{\partial V}\chi_{\mathrm{z}}\langle|\nabla V|^2\rangle T_{\mathrm{z}}\frac{\partial}{\partial V} n_{\mathrm{z}} \\
&+ (u_{\mathrm{z}}^{V} - u_{g}^{V})\frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} + \hat{u}_{\mathrm{z}\theta}\langle \boldsymbol{B}\cdot\nabla\cdot\overleftrightarrow{\pi}_{\mathrm{z}}\rangle - \frac{3}{2}\nu_{\mathrm{Tez}}\left(\frac{n_{\mathrm{e}}}{n_{\mathrm{z}}} n_{\mathrm{z}} T_{\mathrm{z}} - n_{\mathrm{e}} T_{\mathrm{e}}\right) - \frac{3}{2}\nu_{\mathrm{Tiz}}\left(\frac{n_{\mathrm{i}}}{n_{\mathrm{z}}} n_{\mathrm{z}} T_{\mathrm{z}} - n_{\mathrm{i}} T_{\mathrm{i}}\right) \\
&- \nu_{\mathrm{L}}\left(n_{\mathrm{z}} T_{\mathrm{z}} - n_{\mathrm{zsup}} T_{\mathrm{z}}\right) - \frac{3}{2}\nu_{\mathrm{LT_z}}\left(n_{\mathrm{z}} T_{\mathrm{z}} - n_{\mathrm{z}} T_{\mathrm{zsup}}\right) + P_{\alpha\mathrm{z}} + P_{\mathrm{RFz}}
\end{aligned} \tag{3.26}
$$

$$
\begin{aligned}
\frac{5}{2} n_{\mathrm{z}}\frac{\partial}{\partial t}\langle B\hat{q}_{\mathrm{z}\|}\rangle &= -\frac{1}{m_{\mathrm{z}}}\hat{\mu}_2^{\mathrm{z}}\langle B u_{\mathrm{z}\|}\rangle - \frac{1}{m_{\mathrm{z}}}\hat{\mu}_3^{\mathrm{z}}\langle B\hat{q}_{\mathrm{z}\|}\rangle + \frac{1}{m_{\mathrm{z}}}\hat{\mu}_2^{\mathrm{z}} B V_{1\mathrm{z}} \\
&- \frac{1}{Z_{\mathrm{z}} e m_{\mathrm{z}}}\hat{\mu}_3^{\mathrm{z}} I\frac{\partial V}{\partial \psi}\frac{1}{n_{\mathrm{z}}}\frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} + \frac{1}{Z_{\mathrm{z}} e m_{\mathrm{z}}}\hat{\mu}_3^{\mathrm{z}} I\frac{\partial V}{\partial \psi}\frac{T_{\mathrm{z}}}{n_{\mathrm{z}}}\frac{\partial}{\partial V} n_{\mathrm{z}} \\
&- \hat{\ell}_{21}^{\mathrm{ze}} n_{\mathrm{z}}\langle B u_{\mathrm{e}\|}\rangle - \hat{\ell}_{21}^{\mathrm{zi}} n_{\mathrm{z}}\langle B u_{\mathrm{i}\|}\rangle - \hat{\ell}_{21}^{\mathrm{zz}} n_{\mathrm{z}}\langle B u_{\mathrm{z}\|}\rangle - \hat{\ell}_{21}^{\mathrm{zb}}\frac{n_{\mathrm{z}}}{n_{\mathrm{b}}} n_{\mathrm{b}}\langle B u_{\mathrm{b}\|}\rangle \\
&+ \hat{\ell}_{22}^{\mathrm{ze}} n_{\mathrm{z}}\langle B\hat{q}_{\mathrm{e}\|}\rangle + \hat{\ell}_{22}^{\mathrm{zi}} n_{\mathrm{z}}\langle B\hat{q}_{\mathrm{i}\|}\rangle + \hat{\ell}_{22}^{\mathrm{zz}} n_{\mathrm{z}}\langle B\hat{q}_{\mathrm{z}\|}\rangle \\
&- \nu_{\mathrm{L}} n_{\mathrm{z}}\langle B\hat{q}_{\mathrm{z}\|}\rangle - \nu_{0\mathrm{z}} n_{\mathrm{z}}\langle B\hat{q}_{\mathrm{z}\|}\rangle
\end{aligned} \tag{3.27}
$$

$$
0 = -n_{\mathrm{z}}\left\langle\frac{u_{\mathrm{z}\zeta}}{R}\right\rangle + \frac{1}{\langle R^2\rangle} n_{\mathrm{z}}\langle R u_{\mathrm{z}\zeta}\rangle - \frac{2\hat{q}^2}{1+2\hat{q}^2}\frac{1}{\langle R^2\rangle} n_{\mathrm{z}}\langle R u_{\mathrm{z}\zeta}\rangle + \frac{2\hat{q}^2}{1+2\hat{q}^2}\frac{1}{I} n_{\mathrm{z}}\langle B u_{\mathrm{z}\|}\rangle \tag{3.28}
$$

$$
0 = -n_{\mathrm{z}} B V_{1\mathrm{z}} - \frac{1}{Z_{\mathrm{z}} e}\frac{\partial V}{\partial \psi} I\frac{\partial}{\partial V} n_{\mathrm{z}} T_{\mathrm{z}} - n_{\mathrm{z}}\frac{\partial V}{\partial \psi} I\frac{\partial}{\partial V}\Phi \tag{3.29}
$$

$$\frac{\partial}{\partial t}n_{\mathrm{b}} = -\frac{\partial}{\partial V}n_{\mathrm{b}}u_{\mathrm{b}}^{V} + u_g^{V}\frac{\partial}{\partial V}n_{\mathrm{b}} + \frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} - g\frac{P_{\mathrm{b}\perp}}{E_{\mathrm{b}}} - \nu_{\mathrm{b}}n_{\mathrm{b}} - \nu_{\mathrm{bL}}n_{\mathrm{b}} - \nu_{\mathrm{eff}}g(\sqrt{\delta}n_{\mathrm{b}} - n_{\mathrm{b}}^{\mathrm{rp}})$$
$$+ \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle f_t D_{\mathrm{b}}\frac{\partial}{\partial V}n_{\mathrm{b}} \tag{3.30}$$

$$0 = Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}\left(\langle B^2\rangle\langle R^2\rangle - I^2\right)\frac{1}{I}n_{\mathrm{b}}BV_{1\mathrm{b}} + Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}In_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle - Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}\langle B^2\rangle n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle \tag{3.31}$$

$$\frac{\partial}{\partial t}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle = -\frac{1}{m_{\mathrm{b}}n_{\mathrm{b}}}\hat{\mu}_1^{\mathrm{b}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle + \frac{1}{m_{\mathrm{b}}n_{\mathrm{b}}}\hat{\mu}_1^{\mathrm{b}}n_{\mathrm{b}}BV_{1\mathrm{b}}$$
$$+ \hat{\ell}_{11}^{\mathrm{bb}}\frac{m_{\mathrm{e}}n_{\mathrm{e}}}{m_{\mathrm{b}}n_{\mathrm{b}}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle + \sum_{a\neq\mathrm{b}}\hat{\ell}_{11}^{\mathrm{b}a}\frac{m_{\mathrm{e}}n_{\mathrm{e}}}{m_{\mathrm{b}}}\langle Bu_{a\parallel}\rangle$$
$$+ Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}n_{\mathrm{b}}I\langle R^{-2}\rangle\left(-\frac{\dot{\psi}_t}{q} + \dot{\psi}\right) - \nu_{\mathrm{bL}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle - \nu_{\mathrm{b}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle - \nu_{0\mathrm{b}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle$$
$$- \nu_{\mathrm{CX}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle + \frac{1}{m_{\mathrm{b}}}\langle BS_{\mathrm{m}\parallel\mathrm{b}}\rangle \tag{3.32}$$

$$\frac{\partial}{\partial t}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle = u_g^{V}\frac{\partial}{\partial V}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle - \frac{\partial}{\partial V}\langle Ru_{\mathrm{b}\zeta}\rangle n_{\mathrm{b}}u_{\mathrm{b}}^{V}$$
$$+ \frac{1}{m_{\mathrm{b}}}\ell_{11}^{\mathrm{be}}\frac{1}{n_{\mathrm{e}}}n_{\mathrm{e}}\langle Ru_{\mathrm{e}\zeta}\rangle + \frac{1}{m_{\mathrm{b}}}\ell_{11}^{\mathrm{bi}}\frac{1}{n_{\mathrm{i}}}n_{\mathrm{i}}\langle Ru_{\mathrm{i}\zeta}\rangle + \frac{1}{m_{\mathrm{b}}}\ell_{11}^{\mathrm{bz}}\frac{1}{n_{\mathrm{z}}}n_{\mathrm{z}}\langle Ru_{\mathrm{z}\zeta}\rangle + \frac{1}{m_{\mathrm{b}}}\ell_{11}^{\mathrm{bb}}\frac{1}{n_{\mathrm{b}}}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle$$
$$+ Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}n_{\mathrm{b}}\dot{\psi} + Z_{\mathrm{b}}\frac{e}{m_{\mathrm{b}}}\frac{\partial\psi}{\partial V}n_{\mathrm{b}}u_{\mathrm{b}}^{V}$$
$$- \nu_{\mathrm{bL}}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle - \nu_{\mathrm{b}}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\zeta}\rangle - \nu_{0\mathrm{i}}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle - \nu_{\mathrm{CX}}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle + \frac{1}{m_{\mathrm{b}}}\frac{I}{\langle B^2\rangle}\langle BS_{\mathrm{m}\parallel\mathrm{b}}\rangle \tag{3.33}$$

$$0 = -n_{\mathrm{b}}\left\langle\frac{u_{\mathrm{b}\zeta}}{R}\right\rangle + \frac{1}{\langle R^2\rangle}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle - \frac{2\hat{q}^2}{1+2\hat{q}^2}\frac{1}{\langle R^2\rangle}n_{\mathrm{b}}\langle Ru_{\mathrm{b}\zeta}\rangle + \frac{2\hat{q}^2}{1+2\hat{q}^2}\frac{1}{I}n_{\mathrm{b}}\langle Bu_{\mathrm{b}\parallel}\rangle \tag{3.34}$$

$$0 = -n_{\mathrm{b}}BV_{1\mathrm{b}} - \frac{1}{Z_{\mathrm{b}}e}\frac{\partial V}{\partial\psi}In_{\mathrm{b}}\frac{\partial}{\partial V}T_{\mathrm{b}} - \frac{1}{Z_{\mathrm{b}}e}\frac{\partial V}{\partial\psi}IT_{\mathrm{b}}\frac{\partial}{\partial V}n_{\mathrm{b}} - n_{\mathrm{b}}\frac{\partial V}{\partial\psi}I\frac{\partial}{\partial V}\Phi \tag{3.35}$$

$$\frac{\partial}{\partial t}n_{01} = -\frac{\partial u_g^{V}}{\partial V}n_{01} + \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle D_{01}\frac{\partial}{\partial V}n_{01} - \nu_{\mathrm{ionize}}\frac{n_{\mathrm{e}}}{n_{01}+n_{02}+n_{03}}n_{01}$$
$$- \nu_{\mathrm{CX}}\frac{n_{\mathrm{i}}}{n_{01}+n_{02}}n_{01} + \gamma_0\nu_{\mathrm{L}}(n_{\mathrm{i}} - n_{\mathrm{isup}}) + \frac{\partial}{\partial V}V'\Gamma^{\mathrm{puff}}\Big|_{\mathrm{b}} \tag{3.36}$$

$$\frac{\partial}{\partial t}n_{02} = -\frac{\partial u_g^{V}}{\partial V}n_{02} + \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle D_{02}\frac{\partial}{\partial V}n_{02} - \nu_{\mathrm{ionize}}\frac{n_{\mathrm{e}}}{n_{01}+n_{02}+n_{03}}n_{02} + \nu_{\mathrm{CX}}\frac{n_{\mathrm{i}}}{n_{01}+n_{02}}n_{01} \tag{3.37}$$

$$\frac{\partial}{\partial t}n_{03} = -\frac{\partial u_g^{V}}{\partial V}n_{03} + \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle D_{03}\frac{\partial}{\partial V}n_{03} - \nu_{\mathrm{ionize}}\frac{n_{\mathrm{e}}}{n_{01}+n_{02}+n_{03}}n_{03} + f_{\mathrm{CX}}\frac{P_{\mathrm{b}}}{E_{\mathrm{b}}} \tag{3.38}$$

$$\frac{\partial}{\partial t}\bar{n}_z = -\frac{\partial u_g^V}{\partial V}\bar{n}_z + \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle D_{nz}\frac{\partial}{\partial V}\bar{n}_z + \frac{\partial}{\partial V}\langle|\nabla V|^2\rangle \Delta D_{\mathrm{C0}}\frac{\partial}{\partial V}n_{\mathrm{C0}} - n_{\mathrm{e}}\bar{n}_z f_5 S_5 + n_{\mathrm{e}} n_{\mathrm{z}} A_6$$
$$+ \gamma_{0\mathrm{z}}\nu_{\mathrm{L}}\left(n_{\mathrm{z}} - n_{\mathrm{zsup}}\right) + \gamma_{\mathrm{i,z}}\nu_{\mathrm{L}}\left(n_{\mathrm{i}} - n_{\mathrm{isup}}\right) - (1-\gamma_{0\mathrm{z}})\nu_{\mathrm{L}}\left(\bar{n}_z - n_{\mathrm{C0}}\right)$$
$$+ \frac{\partial}{\partial V}V'\Gamma_{\mathrm{z}}^{\mathrm{puff}}\Big|_{\mathrm{b}} \tag{3.39}$$

$$\frac{\partial}{\partial t}(g n_{\mathrm{b}}^{\mathrm{rp}}) = g\frac{P_{\mathrm{b}\perp}}{E_{\mathrm{b}}} + g\nu_{\mathrm{eff}}(\sqrt{\delta}n_{\mathrm{b}} - n_{\mathrm{b}}^{\mathrm{rp}}) - g\nu_{\mathrm{b}} n_{\mathrm{b}}^{\mathrm{rp}}$$
$$- \frac{\partial}{\partial V}g\langle|\nabla V|\rangle u_{\mathrm{b}}^{\mathrm{rp}} n_{\mathrm{b}}^{\mathrm{rp}} + \frac{\partial}{\partial V}g\langle|\nabla V|^2\rangle D_{\mathrm{b}}^{\mathrm{rp}}\frac{\partial}{\partial V}n_{\mathrm{b}}^{\mathrm{rp}} \tag{3.40}$$

4 Boundary Conditions

num.	name	variable	highest order derivative	center	edge
1	LQm1	Φ	2	N	0
2	LQm2	$\dot{\psi}_t$	0	×	×
3	LQm3	$\dot{\psi}$	0	×	×
4	LQm4	ψ	2	N	$2\pi\mu_0 I_\mathrm{p}$
5	LQm5	ψ_t/μ_0	2	0	$\langle\lvert\nabla V\rvert^2/R^2\rangle\lvert_{V_b} I_\mathrm{vac}/(4\pi^2)$
6	LQe1	n_e	2	N	N†
7	LQe2	$n_\mathrm{e}u_\mathrm{e}^V$	1	0	×
8	LQe3	$\langle Bu_{\mathrm{e}\parallel}\rangle$	2(0)	N(×)	N(×)
9	LQe4	$n_\mathrm{e}\langle Ru_{\mathrm{e}\zeta}\rangle$	2	N	N
10	LQe5	$n_\mathrm{e}T_\mathrm{e}$	2	N	N
11	LQe6	$\langle B\hat{q}_{\mathrm{e}\parallel}\rangle$	0	×	×
12	LQe7	$n_\mathrm{e}\langle u_{\mathrm{e}\zeta}/R\rangle$	0	×	×
13	LQe8	$BV_{1\mathrm{e}}$	0	×	×
14	LQi1	n_i	2	N	N†
15	LQi2	$n_\mathrm{i}u_\mathrm{i}^V$	1	0	×
16	LQi3	$\langle Bu_{\mathrm{i}\parallel}\rangle$	2(0)	N(×)	N(×)
17	LQi4	$n_\mathrm{i}\langle Ru_{\mathrm{i}\zeta}\rangle$	2	N	N
18	LQi5	$n_\mathrm{i}T_\mathrm{i}$	2	N	N
19	LQi6	$\langle B\hat{q}_{\mathrm{i}\parallel}\rangle$	0	×	×
20	LQi7	$n_\mathrm{i}\langle u_{\mathrm{i}\zeta}/R\rangle$	0	×	×
21	LQi8	$BV_{1\mathrm{i}}$	0	×	×
22	LQz1	n_z	2	N	N†
23	LQz2	$n_\mathrm{z}u_\mathrm{z}^V$	1	0	×
24	LQz3	$\langle Bu_{\mathrm{z}\parallel}\rangle$	2(0)	N(×)	N(×)
25	LQz4	$n_\mathrm{z}\langle Ru_{\mathrm{z}\zeta}\rangle$	2	N	N
26	LQz5	$n_\mathrm{z}T_\mathrm{z}$	2	N	N
27	LQz6	$\langle B\hat{q}_{\mathrm{z}\parallel}\rangle$	0	×	×
28	LQz7	$n_\mathrm{z}\langle u_{\mathrm{z}\zeta}/R\rangle$	0	×	×
29	LQz8	$BV_{1\mathrm{z}}$	0	×	×
30	LQb1	n_b	0(2)	×(N)	×(N)
31	LQb2	$n_\mathrm{b}u_\mathrm{b}^V$	1	0	0‡
32	LQb3	$n_\mathrm{b}\langle Bu_{\mathrm{b}\parallel}\rangle$	0	×	×
33	LQb4	$n_\mathrm{b}\langle Ru_{\mathrm{b}\zeta}\rangle$	0	×	×
34	LQb7	$n_\mathrm{b}\langle u_{\mathrm{b}\zeta}/R\rangle$	0	×	×
35	LQb8	$BV_{1\mathrm{b}}$	0	×	×
36	LQn1	n_{01}	2	N	$\langle\lvert\nabla V\rvert\rangle S_\mathrm{gas}\rvert_b$
37	LQn2	n_{02}	2	N	0
38	LQn3	n_{03}	2	N	0
39	LQnz	$\bar{n}_z$	2	N	$\langle\lvert\nabla V\rvert\rangle S_{\mathrm{gas,z}}\rvert_b$
40	LQr1	n_b^rp	0(2)	×(N)	×(N)

0: Dirichlet condition ($u = 0$), N: Neumann condition ($u' = 0$)

value: Neumann condition ($u' =$ value), ×: no condition

†: imposed as a constraint in V5.60 (`MDLWBC` = 1), see Sec. 8.2

NOTE: The dependent variable of LQm4 turns to be ψ/μ_0 when NB is applied. Subsequently, the edge boundary condition listed for that variable is also changed to $2\pi I_\mathrm{p}$.

NOTE: Rows 2 to 5 deserve a word, because the variable that carries the second-order operator is not the variable of the equation in which the operator is written. The LQm2 equation advances $\dot{\psi}_t$ and contains $\frac{\partial}{\partial V}[\langle|\nabla V|^2/R^2\rangle \langle R^{-2}\rangle^{-1} \frac{\partial}{\partial V}(\psi_t/\mu_0)]$; the LQm3 equation advances $\dot{\psi}$ and contains $\frac{\partial}{\partial V}[\langle|\nabla V|^2/R^2\rangle \frac{\partial}{\partial V}\psi]$. Counting per dependent variable over the whole system, as this section does, $\dot{\psi}_t$ and $\dot{\psi}$ are never differentiated and so take order 0 and no condition of their own, while ψ_t/μ_0 and ψ take order 2 and carry the two conditions each. The table above now reads that way. In the implementation the two conditions of each pair are applied to the LQm2 and LQm3 equation rows, since that is where the boundary term left by the integration by parts appears; this is the correct weak form and is not a discrepancy with the table. The axis condition of ψ_t/μ_0 is imposed there as $\dot{\psi}_t|_{V=0} = 0$, which holds $\psi_t(0)$ at its initial value, namely zero, and is therefore the Dirichlet condition listed.

NOTE: In the table the column labeled by the highest order derivative means the order of the highest order derivative of a dependent variable, not an equation, in the corresponding row. Please note that the term not integrated by parts is never counted. For example, the first row shows the highest order of derivatives is equal to 2 for Φ because the LQm1 equation has the diffusion term having the second derivative of Φ with respect to V. In addition, the LQe2 and LQi2 equations also have the first derivative of Φ, which is not a dependent variable of these equations. As a whole, the highest order derivative of Φ in this equation system is 2 that comes from the LQm1 equation. When we look at n_e, we find that the highest order derivative of n_e is 2 shown in the LQe5 equation, even though no derivative appears in the LQe1 equation. The Neumann conditions for n_e come from the LQe5 equation. When it comes to $n_\mathrm{e}u_\mathrm{e}^V$, the highest derivative is found as order 1 in the LQe1 equation, but this term is not integrated by parts. It means that any boundary condition is not naturally imposed on $n_\mathrm{e}u_\mathrm{e}^V$. The Dirichlet boundary condition should be imposed at axis.

5 Momentum pinch of V5.62

V5.62 changes the turbulent momentum pinch of the thermal species. The flux of toroidal angular momentum of species s is written with the density outside the gradient (the form of the scaling used below),

$$\Gamma_{\mathrm{L},s} = m_s \left(\langle |\nabla V| \rangle \, V_\mathrm{p} \, n_s \langle R u_{s\zeta} \rangle - \langle |\nabla V|^2 \rangle \, \mu_s \, n_s \, \partial_V \langle R u_{s\zeta} \rangle \right), \tag{5.1}$$

with the momentum diffusivity μ_s and the pinch velocity V_p ($V_\mathrm{p} > 0$ outward). The pinch is

$$V_\mathrm{p} = -\frac{\mu_\mathrm{i}}{R_0} \left[1.1 \, g\!\left(\frac{R_0}{L_n} \right) + 1.0 \right] T(\rho), \qquad \frac{R_0}{L_n} = -R_0 \, \frac{d \ln n_\mathrm{e}}{d\rho_V}, \tag{5.2}$$

where ρ_V is the volume radius; $d \ln n_\mathrm{e} / d\rho_V$ is smoothed by a Savitzky–Golay filter (see the item on smoothing in Sec. 8.9).

Origin and range of the scaling. The multi-machine scaling of Tala et al. [4] is $-R_0 V_\mathrm{pinch} / \chi_\phi = 1.1 \, R/L_n + 1.0$, both terms inward. It was obtained from NBI modulation in JET, DIII-D and AUG and from magnetic perturbations in NSTX, with the pinch and the Prandtl number averaged over $0.4 < \rho_\mathrm{tor} < 0.8$; the data cover R/L_n = 1.2–3.6 (0.2–3.8 with NSTX) and pinch numbers 0.9–6.2. The slopes of the single devices are 1.2 (JET, DIII-D), 1.5 (AUG) and 0.6 (NSTX); gyrokinetic calculations give a dependence about half as strong. The profiles are "not accurate in the edge region", and an edge torque added to the analysis does not change the coefficients inside $r/a < 0.8$. The analysis method [6] and the JET subset [5] write the flux in the form above; residual stress and intrinsic torque are neglected. No upper limit of R/L_n is stated. Beyond the range, the JETPEAK database [7] (R/L_n = 1–10, r/a up to 0.85) gives an R/L_n coefficient of 0.41–0.66 for the steady-state rotation gradient (pinch and residual stress together), and the regression of GKW calculations 0.44; neither shows a saturation.

The closure. Equation (5.2) is a reduced closure, not the Tala law extrapolated over the whole plasma: it is the Tala fit inside its data, it takes the slope beyond the data from the multi-machine and gyrokinetic information, and it is not applied in the edge and the SOL.

- $g(x) = x$ for $|x| \le x_1$ and

$$g(x) = \mathrm{sgn}(x) \left[x_1 + s(|x| - x_1) + (1 - s) \, w \tanh \frac{|x| - x_1}{w} \right] \quad (|x| > x_1), \tag{5.3}$$

  with $x_1 = 3.8$ (`PCHRLN`, the largest R/L_n of the fit), $s = 0.4$ (`PCHSLP`, so that the slope goes from 1.1 to 0.44) and $w = 1$ (`PCHWID`). g is continuous up to its second derivative. The pinch number is 7.7 at R/L_n = 8 and 8.6 at 10, within the measured pinch numbers (up to about 8), against 9.8 and 12 for the linear law. $s = 0$ saturates the pinch number at $1 + 1.1(x_1 + w)$; this is an experimental option without a basis in the literature.

- $T(\rho) = 1$ for $\rho \le \rho_1 = 0.8$ (`RhoPch1`, the outer edge of the data), 0 for $\rho \ge \rho_2 = 0.9$ (`RhoPch2`) and the quintic smoothstep $1 - \xi^3(10 - 15\xi + 6\xi^2)$, $\xi = (\rho - \rho_1)/(\rho_2 - \rho_1)$, between. T is continuous up to its second derivative and exactly zero outside ρ_2, so that the pinch carries no momentum through the wall. The location follows the data (0.8), the momentum boundary of the TOPICS validation (0.9 [8]) and the pedestal-top boundaries of present integrated modelling (0.8–0.94); the width is a numerical choice. Outside ρ_2 the momentum is transported by the diffusion only. Cubic, cosine and tanh tapers with the same end points give the same rotation to three digits.

- V_p is built from the ion viscosity and given to every thermal species; the beam has no turbulent momentum flux. With a common μ the fluxes sum to $V_\mathrm{p}L - \mu\sum_s m_s n_s \partial_V \langle R u_{s\zeta}\rangle$ times the metric factors, the pinch of the total thermal momentum $L = \sum_s m_s n_s \langle R u_{s\zeta}\rangle$ used in TOPICS [8], whether or not the species rotate alike. With species-dependent `rMus0` the pinch number $-R_0 V_\mathrm{p}/\mu_s$ differs between species and this equivalence is lost.

Correction of the sign. Up to V5.61 the code evaluated $V_\mathrm{p} = (\mu/R_0)(1.1R_0\, d\ln n_\mathrm{e}/d\rho_V + 1)$, i.e. $-R_0V_\mathrm{p}/\mu = 1.1R_0/L_n - 1$: the constant term pointed outward, against the scaling it cited. V5.62 uses +1.0 (verified on the output pinch velocity to three digits). The commented-out edge suppressions of the old code (a tanh at $\rho = 0.98$ and a Gaussian at $\rho = 1$) had no documented basis and are removed. Without the radial restriction the old pinch reached a pinch number of 14 at $\rho = 0.9$ and about 140 at the separatrix, and the convective flux at the wall carried momentum into the plasma (+0.04 N m in the JT-60U case).

Validation. JT-60U-like case of Sec. 7.7 (3.2 MW co-tangential and 0.4 MW perpendicular NBI, $\chi = \mu = 0.22(1 + 19\rho^2)$ m^2/s, `PNsSUP(1)` = 0.0125 as explained below), production defaults (`FSTPTM` = 3), 1.5 s:

case	$u_{\mathrm{i}\phi}(0)$ km/s	$u_{\mathrm{i}\phi}(0.82)$ km/s	$E_r(0.5)$ kV/m	$j_r(0.97)$ A/m^2	$j\times B$ torque N m	L_ϕ N m s
no ripple	510	57	+44	−0.017	+0.16	2.03
half ripple (FST)	370	27	+22	−0.103	−3.25	1.58
full ripple	198	−8	−5	−0.180	−7.28	1.05

The line-averaged density is 1.15×10^{19} m^{-3} and the thermal mass inside $\rho = 1$ is 1.54–1.62 mg. The first version of this table used `PNsSUP(1)` = 0.01 (the default) and `FSTPTM` = 2: 621, 441 and 226 km/s ($\bar n_\mathrm{e} = 1.02\times10^{19}$ m^{-3}, thermal mass 1.29 mg); `FSTPTM` = 3 gives 620 and 223 km/s without and with full ripple at 0.01, and V5.61 gave 658, 417 and 128 km/s. The ripple changes the rotation, E_r, the return current and its torque at `PNsSUP(1)` = 0.0125 as at 0.01. The rest of this paragraph refers to `PNsSUP(1)` = 0.01. The pinch number is 6.3 at $\rho = 0.5$ and 7.9 at 0.8 and vanishes outside 0.9; in V5.61 it was 14 at 0.9. The profile is more peaked in the core (the constant term) and lower at the edge ($u_{\mathrm{i}\phi}(0.9) = 46$ against 105 km/s). The volume integrals close to 2×10^{-5} of the injected torque in every row, the pinch flux at the wall is zero and the friction terms sum to 3×10^{-9} N m. With `NRMAX` = 120 the total angular momentum agrees with `NRMAX` = 60 to 0.5% up to 0.8 s (the Picard iteration is limited there by the known impurity terms at the axis, as in V5.61). On the test equilibria of Sec. 7.7 densities and temperatures change by less than 3×10^{-3} and the rotation by 13–40%; the mean Picard count rises by 1–4 and no step more reaches `ICMAX`. The Tala law with the corrected sign but without the radial restriction and the slope reduction spins the JT-60U case up to 881 km/s and stops at 1.4 s at the impurity grid mode of supersonic rotation.

Density of the benchmark. `PNsSUP(1)` is the upstream SOL density and sets the density level of the whole plasma (Sec. 8.9); the shape of the density profile does not depend on it. The benchmark input sets it to 0.0125 (10^{20} m^{-3}), which gives $\bar n_\mathrm{e} = 1.15\times10^{19}$ m^{-3} and $n_\mathrm{e}(\rho = 1) = 0.22\times10^{19}$ m^{-3}. The line-averaged density is the one estimated from the measured profile of the JT-60U L-mode discharge compared in [1] (Fig. 1 of [2], which gives no value), and the separatrix density lies at the lower end of the range expected for JT-60U L-modes (0.2–0.45$\times10^{19}$ m^{-3}). This value normalizes the density of this benchmark; it is not a universal value, and the default of `PNsSUP(1)` stays 0.01. A new device, discharge, equilibrium, heating scheme or particle-transport

model needs its own value, found by an outer loop of two or three runs to the target density (secant in log–log; there is no density feedback inside the code). The benchmark is JT-60U-like and not the discharge: $I_\mathrm{p} = 1.0$ MA against 1.23 MA, a circular cross-section against a D-shaped plasma of 77 m^3, and the computed density profile falls more steeply towards the edge ($n_\mathrm{e}(0.9)/\bar{n}_\mathrm{e} = 0.45$ against 0.74). The rotation was not used to choose `PNsSUP(1)`: it is a consistency check of the momentum model, not a fitting target.

The reduced ripple model is judged by the changes with the ripple rather than by the absolute rotation. The co-rotation without ripple is 1.7 times that of [1] (510 against 300 km/s); the states of that work, restored from its save files with the 2008 code, show why. The 2008 plasma was pure deuterium with a flat density ($n_\mathrm{e} = 1.4$–1.5×10^{19} m^{-3} up to $r/a = 0.7$), 1.4 times heavier than the present one (thermal mass 2.30 against 1.59 mg, moment of inertia 2.70 against 1.87×10^{-5} kg m^2); its torque was 7.63 N m (85 keV, the whole beam velocity toroidal at R_0) with half of it deposited off axis, it had no momentum pinch, and its wall held $u_\phi = 0$. The momentum confinement times L_ϕ/T_NBI of the thermal species are similar (0.137 and 0.143 s), so the mass-weighted rotation $R_0 L_\phi/I_\phi$ differs by 1.3 (132 and 174 km/s) and the more peaked profile gives the rest of the factor 1.7 on axis.

Limitations. The pinch number is taken from the electron density gradient alone; the dependences on q, $\hat{s}$, ε, $T_\mathrm{i}/T_\mathrm{e}$ found in the regressions are not included, and no residual stress is modelled. The radial restriction is fixed in ρ and does not follow the pedestal of an H-mode. The momentum diffusivity itself is prescribed ($\mu = \chi$ in the benchmark). A Coriolis-type alternative with a geometric dependence ($-R_0 V/\mu \propto \sqrt{r/R}$, as used for DTT [9]) rotates the JT-60U case 30% more slowly; its coefficient is a calibration, not a universal constant, and it is not implemented in the production code.

6 Anomalous transport in the SOL of V5.62

Inside the separatrix the anomalous coefficients of particles ($k = 1$), toroidal momentum ($k = 2$) and heat ($k = 3$) are

$$C_{k,s}(\rho) = C_{0,s}\left[\mathtt{FSDFIX}_k\, f_k(\rho) + B_k + \mathtt{FSANOM}_k\, D_\mathrm{turb}(\rho)\right], \tag{6.1}$$

where $C_{0,s}$ is `Dfs0`, `rMus0` or `Chis0` of species s, f_k the fixed profile $(1 + (\mathtt{PROF} - 1)\rho^{\mathtt{PROF1}}$, with the Gaussian modification `PROFD2` for $k = 1$), B_k the offset `PROFDB`, `PROFMB` or `PROFCB`, and D_turb the CDBM (`MDANOM` = 1) or CDIM (2) value. Only `Dfs(:,1)` enters the particle transport. The turbulence models describe closed flux surfaces. On the open field lines of the SOL ($\rho \geq 1$, from the separatrix node `NRA`; the mesh puts $\rho = 1$ exactly on it for any `rhoaccum`) the coefficient is built from the separatrix value $C^\mathrm{core}_{k,s}(1)$, Eq. (6.1) at $\rho = 1$ with D_turb of the node `NRA` (the `PROFD2` modification is zero there).

6.1 `MDLSOLT` = 0 (default): `FSPCL`

$\mathtt{FSPCL}_k$	SOL coefficient
1 (default)	$C^\mathrm{core}_k(1)$, constant
0	$C^\mathrm{core}_k(1)\, D_\mathrm{B}(\rho)/D_\mathrm{B}(1)$, $D_\mathrm{B} = T_\mathrm{e}/(16eB)$
other	$\mathtt{FSPCL}_k\, C_0[\mathtt{FSDFIX}_k f_k(1) + B_k]$ without `FSANOM`; $C_0\, \mathtt{FSANOM}_k\, D_\mathrm{turb}(1)$ with it

In V5.62 the first two rows take the whole separatrix coefficient, as the table states; before, they used only parts of it when `FSANOM` was on. With `FSPCL` = 1 the SOL took $C_0\, \mathtt{FSANOM}_k D_\mathrm{turb}(1)$ alone, so with `FSDFIX`

and FSANOM both on the coefficient stepped down at the separatrix (NRMAX = 60 with NBI: D from 1.51 to 0.76 and χ from 5.51 to 1.11 m^2/s at $t = 0$). With FSPCL = 0 the anomalous part was D_{turb} of each SOL node, the closed-surface model evaluated where it does not apply; the CDBM value nearly vanishes there, the SOL transport dropped with FSANOM, and the Picard iteration converged badly (NRMAX = 60, FSANOM only, 1 s: mean number of sweeps 20.3 and 756 retried steps with NBI, 69.8 and 8802 without; now 14.8 and 8.1 sweeps and no retry). FSDFIX alone gives the same results as before in both rows, and the third row is unchanged.

6.2 MDLSOLT = 1: connection to far-SOL values

With MDLSOLT = 1 the SOL coefficients of D, χ_s and μ_s are

$$C_s(\rho) = C_{\mathrm{SOL},s} + [C_s^{\mathrm{core}}(1) - C_{\mathrm{SOL},s}] \exp\left(-\frac{\rho - 1}{w}\right), \qquad \rho \geq 1, \tag{6.2}$$

with C_{SOL} = DSOLT (for Dfs(:,1)), CHISOLT(s) and RMUSOLT(s) in m^2/s and w = WSOLT in units of ρ; at the outer midplane $\Delta r \approx a\, w$ (ellipticity and Shafranov shift ignored). $C_s^{\mathrm{core}}(1)$ is the separatrix value above. The code evaluates Eq. (6.2) as the convex combination $C_s^{\mathrm{core}}(1)\, e^{-x} + C_{\mathrm{SOL},s}(1 - e^{-x})$, $x = (\rho - 1)/w$, so C equals $C_s^{\mathrm{core}}(1)$ exactly at NRA and stays positive. FSPCL is not used (a warning is printed if any FSPCL $\neq 1$), so the Bohm-like row is not taken either. FSPCL = 1 is the limit $w \to \infty$.

The connection is phenomenological. It separates what FSPCL ties together: the core model fixes the coefficient at the separatrix, but nothing in the code predicts the cross-field transport of the far SOL, which edge codes also prescribe rather than compute. The four inputs therefore have no universally recommended values: they are unset (-1) by default and MDLSOLT = 1 stops with an input error unless they are all positive. For orientation only, predictive edge calculations for ITER commonly take spatially uniform $D_\perp = 0.3$ and $\chi_{\mathrm{e}} = \chi_{\mathrm{i}} = 1.0\,\mathrm{m}^2/\mathrm{s}$ with a perpendicular viscosity of $0.2\,\mathrm{m}^2/\mathrm{s}$ [10], and the one integrated-modelling connection of a core separatrix value to such values that we found, in JINTRAC [11], is an exponential relaxation with an e-folding length of 1.3 cm at the outer midplane (about $\Delta\rho = 0.016$ for $a = 0.8$ m) when the separatrix value exceeds the far-SOL value; below a transport barrier it uses a flat extension and a tanh rise instead, which MDLSOLT = 1 does not model.

About four elements outside the separatrix within w resolve the connection (a guide, not a requirement). A warning is printed when w is less than two elements (checked once the mesh is built; with rhoaccum ≥ 0 the mesh is not refined at the separatrix, e.g. 0.0095 against 0.0052 at NRMAX = 60 with rhoaccum = 0.95), and when $e^{-(\rho_b - 1)/w} > 0.05$, i.e. more than 5% of $C^{\mathrm{core}}(1) - C_{\mathrm{SOL}}$ remains at the wall. MDLSOLT = 1 is an input error with MDANOM = 3 (MMM95 sets the SOL coefficients itself). The inputs are kept in a fifth extension record of the save file; an older file leaves them as they are, and an older build stops before the record.

Validation (NRMAX = 60, 3.2 MW NBI and CDBM unless stated, with the test values $D_{\mathrm{SOL}} = 0.3$, $\chi_{\mathrm{SOL}} = \mu_{\mathrm{SOL}} = 1\,\mathrm{m}^2/\mathrm{s}$ and $w = 0.02$, which are not defaults). MDLSOLT = 0 is bit-identical to the code without the option. Against FSPCL = 1 at 1 s the connection raises $T_{\mathrm{e}}(1)$ by 7%, $T_{\mathrm{i}}(1)$ by 22%, $u_{\mathrm{i}\phi}(0.9)$ by 10% and $u_{\mathrm{i}\phi}(1)$ by 28%, lowers T_{i} at $\rho = 1.05$ by 13%, and changes the core by less than 2%. The same changes within 1% are found with fixed transport and at NRMAX = 120; at 0.2 s NRMAX = 60, 120 and 240 give $T_{\mathrm{i}}(1)$ +19 to +20% and $u_{\mathrm{i}\phi}(1)$ +22 to +24%. Without NBI the changes are about 1% with CDBM, whose separatrix value is then close to the far-SOL values, and 11% in $T_{\mathrm{i}}(1)$ with fixed transport. NRMAX = 60 and 120 agree to 1% inside the separatrix with either treatment. The mean number of Picard sweeps changes by 2% or less, and the few retried steps (at most six in a run) are isolated and of the kinds also seen with FSPCL = 1 (carbon superstage, ion momentum near the separatrix or on the axis). Doubling one far-SOL value at a time (0.5 s): μ_{SOL} changes

only the rotation ($u_{\mathrm{i}\phi}(1)$ −9%, +45% near the wall; densities and temperatures unchanged to 0.05%), χ_{SOL} mainly the temperatures ($T_{\mathrm{i}}(1)$ −6%, +23% near the wall), and D_{SOL} the density by less than 1% (the SOL density is set by the parallel loss and the recycling source); doubling w changes $T_{\mathrm{i}}(1)$ and $u_{\mathrm{i}\phi}(1)$ by −8 and −10%. The particle, momentum and energy budgets close to 10^{-10} of the largest term. With 6.4 MW the option is not verified: there CDBM fails near the axis at 0.4–0.6 s with either SOL treatment, as it does in the code before these changes.

7 Beam and ripple model of V5.61

V5.61 changes the beam and ripple equations. The model itself (its assumptions, the ordering behind each term and the comparison with the 2008 ripple model [1]) is described in `beam_ripple.tex`; this section states what the code solves and what changed.

change	effect
ripple momentum terms of the beam removed (B-full)	`FSRP` = 1 with NBI
LQr1 written as the conservation law of $gn_{\mathrm{b}}^{\mathrm{rp}}$	`FSRP` = 1 with NBI
$\Gamma_{\mathrm{b}}(\rho_b) = 0$	`MDBEAM` = 1 with NBI
fast-ion friction $\ell_{11}^{fk} = \ell_{11}^{kf}$ restored	NBI (`MDLNEO` = 1, 2)
Z_{eff} and the fast-ion inventory weighted by g	`FSRP` = 1 with NBI
`FSRP` restricted to 0, 1, 2; `FSRP` = 2 keeps $gP_{\mathrm{b}\perp}$ in n_{b}	input
on-axis ripple trapping rate from the interior	`FSRP` = 1 with NBI

Runs without NBI are bit-identical to V5.60 (X-point and circular test equilibria, `NRMAX` = 60 and 240, `MDLNZD` = 0, `MDLNOFL` = 2, `FSRP` = 0 and 1). No default switch changes. For runs with NBI, and in particular with `FSRP` = 1, `NANDER` = −20 is recommended (Sec. 8.5; −50 at `NRMAX` = 240): the impurity radial flux at the wall converges slowly without it once the beam heats the edge.

7.1 `FSRP` and the ripple-well fraction

`FSRP` is a mode, not a factor: 0, no ripple; 1, full ripple model (ripple-trapped ions, LQr1, and ripple diffusion of n_{b}); 2, ripple diffusion of n_{b} without ripple-trapped ions. Other values are rejected. The ripple amplitude is $\delta(r,\theta) = \delta_{\min} I_0(N a_\delta / R_{\delta\min})$ with `DltRPn` = $\delta_{\min}$; `FSRP` only switches it on. $R_{\delta\min}$ (input `RDltRPn`, default 2.4 m) is the major radius on the midplane where the ripple amplitude has its minimum $\delta_{\min}$; the formula takes the surfaces δ = const as circles centred there, a property of the TF coil geometry (the coil shape and radial position), not of the plasma. 2.4 m is the JT-60U value that entered the 2008 code together with the ripple model ("specific value for JT-60U", also used by the routine that drew the ripple contours); it was fixed in the code up to V5.61.00, and other devices need their own value. The text of [1] gives 2.8 m, but its own numbers require 2.4 m: with $\delta_{\min} = 2 \times 10^{-6}$ and $N = 18$ the formula gives $\delta_a = 1.28\%$ at the outboard midplane and 0.30% at $\theta = \pi/2$ with 2.4 m, the values quoted in [1], and 0.085% and 0.043% with 2.8 m; the contours of its Fig. 1 are centred near 2.4 m. The 2.8 m of the text is a misprint. A point of a flux surface lies in a ripple well when

$$\alpha \equiv \frac{\epsilon |\sin\theta|}{N q \delta(r,\theta)} < 1, \tag{7.1}$$

and $g(\rho)$ (`rip_rat`) is the fraction of the poloidal circumference where it holds, found by searching the rims on the low- and high-field sides. g is not zero near the axis: close to the low-field-side midplane $\sin\theta \to 0$ and a

narrow arc always satisfies $\alpha < 1$ (for the JT-60U case below $g = 0.22$, 0.06 and 0.43 at $\rho = 0.04$, 0.25 and 0.96). V5.61 counts these narrow core wells as the criterion gives them; no cut-off is applied.

7.2 Ripple-trapped beam density: conservation form

n_b^rp (LQr1) is the density of the ripple-trapped beam ions inside the wells; the inventory per unit volume is $N_\mathrm{r} = g n_\mathrm{b}^\mathrm{rp}$. V5.61 solves the conservation law of N_r with n_b^rp as the unknown,

$$\frac{\partial}{\partial t}(g n_\mathrm{b}^\mathrm{rp}) + \frac{\partial}{\partial V}\left(gU n_\mathrm{b}^\mathrm{rp}\right) - \frac{\partial}{\partial V}\left(gK \frac{\partial}{\partial V} n_\mathrm{b}^\mathrm{rp}\right) = g\frac{P_{\mathrm{b}\perp}}{E_\mathrm{b}} + g\nu_\mathrm{eff}(\sqrt{\delta}\, n_\mathrm{b} - n_\mathrm{b}^\mathrm{rp}) - g\nu_\mathrm{b} n_\mathrm{b}^\mathrm{rp}, \tag{7.2}$$

$$U = \langle|\nabla V|\rangle u_\mathrm{b}^\mathrm{rp}, \qquad K = \langle|\nabla V|^2\rangle D_\mathrm{b}^\mathrm{rp}. \tag{7.3}$$

g stays inside every divergence. Written for N_r, the diffusive flux is $-gK\partial_V(N_\mathrm{r}/g) = -K\partial_V N_\mathrm{r} + K N_\mathrm{r}\partial_V \ln g$; the second part, the dg/dV term, is what makes the flux vanish where the wells end ($g \to 0$ with n_b^rp finite), so it cannot be dropped: without it the ripple-trapped ions diffuse out of the well region (a test with $-K\partial_V N_\mathrm{r}$ alone lost 80% of the inventory through the core-side rim). The form (7.2) contains no $1/g$.

Changes from V5.60, where LQr1 was $\partial_t n_\mathrm{b}^\mathrm{rp} + \partial_V(r u_\mathrm{b}^\mathrm{rp} n_\mathrm{b}^\mathrm{rp}) - \partial_V(K\partial_V n_\mathrm{b}^\mathrm{rp}) = \ldots$ without g:

- the dg/dV terms (above) were missing and $n_\mathrm{b} + g n_\mathrm{b}^\mathrm{rp}$ was not conserved: the particle inventory of $n_\mathrm{b} + g n_\mathrm{b}^\mathrm{rp}$ was off by 2–9% of the NBI source;
- the advection used $r u_\mathrm{b}^\mathrm{rp}$ (a remnant of the $\psi = r^2$ coordinate of the earlier code) instead of $\langle|\nabla V|\rangle u_\mathrm{b}^\mathrm{rp}$, a factor $4\pi^2 R$ too weak in circular geometry;
- the source had no g, so n_b^rp grew where there are no wells.

Discretization: the mass, source, exchange and thermalization terms are the element matrices $\int g\,(\ldots)\,w\,dV$; the exchange terms are the same matrices as in LQb1 with opposite sign, so $n_\mathrm{b} + g n_\mathrm{b}^\mathrm{rp}$ is conserved exactly by the discrete equations. The advection is in strong form, $-\int \partial_V(gU n_\mathrm{b}^\mathrm{rp}) w\,dV$; its sum over the test functions is the wall flux $[gU n_\mathrm{b}^\mathrm{rp}]_{V_b}$, the ripple loss out of the plasma, and no boundary condition is needed at either end ($gU = 0$ at the axis). The diffusion has the natural condition. A node with no well on either side gets the identity $\partial_t n_\mathrm{b}^\mathrm{rp} = 0$. No SUPG: the element Péclet number is 15–30 at the edge and 10^2–10^3 inside, but the solution shows no grid-scale structure and converges with the mesh (Sec. 7.7).

Every other equation already weighted n_b^rp by g (LQb1, Poisson, LQi1); V5.61 also weights Z_eff and the fast-ion inventory (`ANFINT`) by g.

7.3 Ripple momentum closure of the beam (B-full)

In V5.60 the parallel beam momentum (LQb3) had the two ripple terms of the 2008 model [1],

$$-\nu_\mathrm{eff} g\sqrt{\delta}\, m_\mathrm{b} n_\mathrm{b} \langle B u_{\mathrm{b}\parallel}\rangle, \qquad +\frac{\partial}{\partial V}\langle|\nabla V|^2\rangle\langle B u_{\mathrm{b}\parallel}\rangle\, f_t D_\mathrm{b}\, \frac{\partial}{\partial V} n_\mathrm{b}, \tag{7.4}$$

the momentum removed by ripple trapping and the momentum carried by the ripple diffusion flux; the toroidal momentum (LQb4) had neither. V5.61 removes both, and adds nothing to LQb4. The reason is the order of these terms in the velocity moments of the beam distribution, not the numerical instability that the second term caused on linear elements:

- The ions scattered into the ripple wells have $|u_\parallel|/v < \sqrt{\delta}$. The momentum they take is $\int_{|\xi|<\sqrt{\delta}} mv\xi\,\nu f\,d^3v$, bounded by $\sqrt{\delta}\,mv$ per ion and $O(\delta)\,mv$ for a smooth $f(\xi)$, against $mu_{\mathrm{b}\parallel} = O(1)\,mv$ per ion in the 2008 form. The same holds for mRv_ζ up to the drift, $O(\rho^*_{\mathrm{b}})$.

- Ripple diffusion acts on the banana ions. Their bounce-averaged distribution is even in ξ, so the parallel momentum flux $\int mv\xi\,\Gamma_r[f]\,d^3v$ vanishes at zero orbit width; the first non-zero part comes from the offset of the two legs of the banana, $O(\rho^*_{\mathrm{b}})$. For the toroidal momentum, $\langle mRv_\zeta\rangle_{\mathrm{b}} = P_\zeta + Ze\langle\psi\rangle_{\mathrm{b}}$ with $v_\parallel = 0$ at the tips and $\langle v_\parallel\rangle_{\mathrm{b}} = 0$, again $O(\rho^*_{\mathrm{b}})$.

- The beam fluid of TX keeps no finite-orbit-width physics (the beam viscosity is zero), so keeping these $O(\delta)$ and $O(\rho^*_{\mathrm{b}})$ terms alone would be inconsistent. $\rho^*_{\mathrm{b}} = 0.1$–$0.2$ at the edge of JT-60U for 80 keV D: the terms are dropped because the closure does not retain that order, not because they are negligible.

Both components vanish, so the closure also respects the radial force balance (LQb2): a ripple term in LQb3 alone forces a spurious radial beam flux through the multiplier Γ_{b}. Ripple particle transport is kept: the trapping, detrapping and diffusion of n_{b} in LQb1, and LQr1. The main effect of the ripple in the 2008 work, the torque of the return current, comes from the charge that these particle fluxes move and is unchanged (Sec. 7.7).

7.4 Radial beam flux at the wall

With `MDBEAM` = 1, $\Gamma_{\mathrm{b}} = n_{\mathrm{b}}u^V_{\mathrm{b}}$ (LQb2) is the Lagrange multiplier of the beam radial force balance; V5.60 had no condition on it at ρ_b. V5.61 imposes

$$\Gamma_{\mathrm{b}}(\rho_b) = 0. \tag{7.5}$$

This is a closure of the reduced model, not a statement that fast ions are not lost to the wall. The model has no surface or orbit-loss channel for the beam; outside the separatrix the beam is lost by the volume loss $\nu_{\mathrm{bL}}n_{\mathrm{b}}$. The wall value of the multiplier was not an outflow law: it was whatever the force balance at the wall node required, with the diamagnetic term of a beam with fixed $T_{\mathrm{b}} = 2E_{\mathrm{b}}/3$, whereas the fast-ion orbit width (10–20 cm) exceeds the gradient lengths of the SOL. Left free, that value pushed n_{b} at the wall node towards zero and made the Picard iteration cycle without converging (a period-5 cycle of Γ_{b} on the circular test equilibrium). The alternatives were rejected: the SOL beam removal used with `MDBEAM` = 0, 2 rewrites n_{b} outside the equations and contradicts the constraint rows; an outflow $\Gamma_{\mathrm{b}} = n_{\mathrm{b}}v_{\mathrm{out}}$ needs a physical v_{out} and would double-count ν_{bL}. The condition changes runs with NBI by 1.5–5% in the core rotation for perpendicular injection and by 0.5% for the JT-60U case below. It should be revisited if a finite-orbit-width or orbit-loss model is introduced.

7.5 Beam–thermal friction

The friction between the beam and the thermal species follows the fast-ion coefficients of Kikuchi and Azumi [3] (`imodel_neo(3)` = 2, isotropic fast ions in the pitch-angle-scattering regime): $\ell^{\mathrm{fe}}_{11} = \ell^{\mathrm{ef}}_{11}$, $\ell^{\mathrm{f}k}_{11} = \ell^{k\mathrm{f}}_{11}$ (k = i, z), $\ell^{\mathrm{ff}}_{11} = -\sum_k \ell^{\mathrm{f}k}_{11}$, the Maxwellian ℓ^{kk}_{11} reduced by $\ell^{k\mathrm{f}}_{11}$, and the heat friction $\ell^{\mathrm{ef}}_{21} = -\frac{3}{2}\ell^{\mathrm{ef}}_{11}$ with $\ell^{\mathrm{fe}}_{21} = 0$. The 11 block is symmetric, so the friction forces sum to zero; only the heat friction is not symmetric, as the paper states. In V5.60 the fast-ion row was obtained from the thermal row with the Braginskii ratio τ_{ee}/τ_{ii}, which assumes equal Coulomb logarithms, and converted back to physical units with the actual τ_{ee} and τ_{ii}. The result was $\ell^{\mathrm{fi}}_{11}/\ell^{\mathrm{if}}_{11} = \ln\Lambda_{ee}/\ln\Lambda_{ii}$ (0.82–0.86 for D, 1.00–1.08 for C) and a net force of 2–10% of the fast-ion friction: 3% of the NBI torque in the JT-60U case, 13% at lower T_{e}. V5.61 converts with the same collision times; the column sums of the 11 block are now 10^{-16} of the diagonal. Applicability: the coefficients assume an isotropic fast-ion

distribution and the slowing-down regime $v_{\mathrm{Ti}} \ll v_{\mathrm{b}}$, and they ignore the fast-ion heat flow; for a tangential beam the distribution is anisotropic.

7.6 Discrete momentum conservation of the friction

The friction force on species a is $\sum_b \ell^{ab} u_b$ with $\sum_a \ell^{ab} = 0$, so the forces cancel for any flows. The coefficients satisfy this at every node (to 10^{-8} of the diagonal, MI). The equations of species a are written per unit mass density, so the term of row a acting on the momentum $X_b = n_b\langle R u_b\rangle$ has the coefficient $\ell^{ab}/(m_a n_a) \times n_a/n_b$; up to V5.61.00 it was integrated as the product of the two interpolants (`fem_int(20,lab,Ratden)`, and `fem_int(20,lab,n)` in the parallel equations), while the diagonal term of the same column used the interpolated coefficient (`fem_int(2,lab)`). Per element the two differ by $-h\,\Delta a\,\Delta b\,(X_1 + X_2)/12$, where Δa, Δb are the changes of the two factors across the element, so the columns did not sum to zero and the friction left a net force of $O(h^2)$. It was measured directly: the residual falls by 2.2–2.3 from `NRMAX` = 60 to 90 and by 3.8–3.9 to 120, and in the JT-60U case it came almost entirely from the ion–impurity pair near $\rho = 0.8$, where n_i/n_z and the coefficients vary most. Every friction term of a column is now assembled from the nodal product of its row-dependent factors ($\ell^{ab} n_a/n_b$, $\ell^{ab} n_a$, and $\ell^{ab}_{12} n_a$ for the heat friction, with the row-independent factors $I/\langle B^2\rangle$ and $\partial_V T_b$ integrated alike in every row). The element matrices then sum to zero over a column whenever the nodal coefficients do, and the continuous equations are unchanged. The change of the solution is $O(h^2)$: in the JT-60U case at 1.5 s, $u_{\mathrm{i}\phi}(0)$ goes from 658 to 656 km/s, the total angular momentum by -0.2%, the impurity density by up to 1% and E_r by up to 0.7% (normalised to their maxima); without NBI, where the spurious force was comparable to the whole toroidal angular momentum, the small rotation changes by up to 5%. The Picard counts are unchanged.

7.7 Validation

Regression. Without NBI all runs are bit-identical to V5.60 (see the start of this section). With NBI and `FSRP` = 0 the solution changes by the intended $\Gamma_{\mathrm{b}}(\rho_b) = 0$ and the friction correction: on the X-point test equilibrium with 5 MW perpendicular NBI the total toroidal angular momentum at 0.5 s is 16% lower (the injected torque is zero there, so the spurious friction source was comparatively large), with 3 MW co-tangential plus 1 MW perpendicular 2% lower; bulk densities and temperatures change by less than 10^{-3}. On the circular test equilibrium the wall cycle of Γ_{b} is gone (V5.60: 25 of 5000 steps at `ICMAX`; V5.61: none).

Stability matrix. `FSRP` = 1, perpendicular NBI, `NANDER` = −20, X-point and circular, 0.3 s (0.15 s for $\Delta t = 2.5 \times 10^{-5}$):

case	X-point: steps at `ICMAX` / mean	circular
NRMAX 60, $\Delta t = 10^{-4}$, 2.5 / 5 / 10 MW	1 / 15, 0 / 21, 0 / 28	0 / 19, 0 / 28, 0 / 34
$\Delta t = 2.5 \times 10^{-5}$, 2×10^{-4}	0 / 11, 0 / 28	0 / 14, 1 / 37
NRMAX 120	13 / 42	1020 / 68

No run stops and none shows a grid-scale mode: the odd-even part of the beam parallel momentum near the edge is 0.016–0.026 of its maximum at `NRMAX` = 60, the level of the non-uniform mesh, and 0.005 at 120. The steps at `ICMAX` on the circular equilibrium at `NRMAX` = 120 are limited by the impurity diamagnetic flow (LQz8) at the axis node. They do not occur with `FSRP` = 0 and remain when the ripple terms inside $\rho = 0.3$ are removed (tested: $g = 0$ or $D_{\mathrm{b}}^{\mathrm{rp}} = 0$ there): it is the known weakness of the impurity equations at the axis, triggered by the state the ripple produces, not a defect of the ripple model.

Conservation. Volume integrals of every term of every equation (JT-60U case below, every step): the discrete time derivative equals the sum of the terms to 10^{-6} for all density and momentum equations; $n_{\mathrm{b}} + g n_{\mathrm{b}}^{\mathrm{rp}}$ closes with a residual below 10^{-5} of the NBI source. The injected torque is 6.73 N m (analytic $2PR_{\tan}/v_{\mathrm{b}}$: 6.76 N m). The friction between the beam and the thermal species conserves momentum since the correction of ℓ_{11}^{fk} (Sec. 7.5), and the friction terms of all species sum to less than 10^{-8} N m at all times since the assembly of Sec. 7.6; before that, the friction between the thermal ions and the impurity left a discrete residual that grew with the rotation to +0.037 N m (0.55% of the injected torque) at 1.5 s. The viscous flux at the wall is zero by the weak form; the rotation is damped by the SOL parallel loss and charge exchange at the edge.

Ripple benchmark. JT-60U-like circular case of [1]: $R = 3.4$ m, $a = 0.95$ m, $B = 2.6$ T, $I_{\mathrm{p}} = 1$ MA, $\kappa = 1.15$ for the ripple, $N = 18$, $\delta_{\min} = 2 \times 10^{-6}$ (edge ripple 1.26%), 3.2 MW co-tangential and 0.4 MW perpendicular NBI, $\chi = \mu = 0.22(1 + 19\rho^2)$ m^2/s, $D = 0.15(1 + 4\rho^2)$ m^2/s, 1.5 s:

case	$u_{\mathrm{i}\phi}(0)$ km/s	$u_{\mathrm{i}\phi}(0.82)$ km/s	$E_r(0.5)$ kV/m	$j_r(0.97)$ A/m^2	$j \times B$ torque N m	dL_ϕ/dt N m
no ripple	658	152	+83	−0.020	+0.18	+0.12
half ripple (FST)	417	60	+34	−0.108	−3.40	+0.05
full ripple	128	−42	−22	−0.185	−7.44	0.00

An earlier version of this table gave the state at 0.6 s. At that time the runs without ripple and with half the ripple were still transient ($dL_\phi/dt = +1.57$ and +0.87 N m against the injected 6.73 N m; $u_{\mathrm{i}\phi}(0) = 473$ and 325 km/s); the table now gives the quasi-steady state at 1.5 s, where dL_ϕ/dt is below 2% of the injected torque in all three runs. With ripple the return current and its $j \times B$ torque changed by 5–11% from 0.6 s to 1.5 s, so the ripple results are qualitatively unchanged; the co-rotation without ripple is 40% higher than at 0.6 s. The return current grows with the ripple and has the magnitude of [1] (−0.08 and −0.16 A/m^2 with and without FST), its torque is counter and grows, E_r moves negative and the edge rotates counter with the full ripple; halving the ripple weakens all of these. The qualitative physics of the 2008 work is kept by B-full and the conservative LQr1. The co-rotation without ripple is 2.2 times that of [1] (300 km/s). The injected torque agrees with its analytic value, so this is a question of the momentum damping and transport (the losses concentrate at the edge in the V5.60 SOL and neutral models), not of the ripple model.

8 Models, switches and defaults of V5.60

V5.60 changes the default SOL and neutral physics. The detailed physics, the verification and the development history are in `neutral_transport.tex`; this section states what the code solves.

8.1 SOL parallel loss

Outside the separatrix the parallel loss rate is

$$\nu_{\mathrm{L}} = k_L \frac{C_s}{L_c}\, g(\rho), \qquad L_c = 2\pi q R, \qquad g = \frac{x^2}{1 + x^2}, \quad x = \frac{r - a}{\delta}\ (r \geq a),\ x = 0\ (r < a), \tag{8.1}$$

and enters every SOL equation in Robin form, e.g. $-\nu_{\mathrm{L}}(n_s - n_{\mathrm{ssup}})$. k_L is `FSLP`. The ramp $g(\rho)$ is numerical smoothing: it keeps ν_{L} from jumping at $\rho = 1$, which the FEM cannot represent, and its continuum limit is the step. `MDLSOL` selects its width δ:

MDLSOL	δ
0	$3\sqrt{0.5\,T_i(0)\,m_i}\,q(a)/(Z_i e B_\phi(a))/\sqrt{a/R}$ (V5.52)
1	ion banana width at the separatrix, $\sqrt{\epsilon}\,\sqrt{T_i m_i}/(Z_i e B_\theta)$ (default)

With MDLSOL = 0 the width is evaluated with half the on-axis ion temperature and the edge q; for the X-point test equilibrium distributed with the code (eqdata, $R = 3.2$ m, $a = 0.8$ m) it is 1.8 times the whole SOL, so g is only 0.17 at the wall and the ramp suppresses the loss everywhere instead of smoothing its onset. MDLSOL = 1 evaluates the banana width with the local ion temperature and poloidal field at the separatrix and gives $g \simeq 0.99$ at the wall.

The default $k_L = 1$ is the value of the sheath-limited two-point model (pressure conservation, Mach number 1 at the target, upstream and target densities related by $n_\mathrm{t} = n_\mathrm{u}/2$) when $2\pi qR$ is read as the length between the two targets: a flux tube of that length loses $2n_\mathrm{t}C_s$ per unit area while holding $n_\mathrm{u}\,2\pi qR$, so $\nu_\mathrm{L} = C_s/(2\pi qR)$. V5.52 used the empirical value $k_L = 0.3$, a typical upstream Mach number of the SOL flow; with MDLSOL = 1 it does not give a stable SOL on the X-point test equilibrium.

8.2 Wall boundary condition for the densities

The table of the previous section lists a Neumann condition for the densities at the wall $\rho = \rho_b$, said to come naturally from the pressure equation. The natural boundary condition of that equation is, however, zero heat flux, not a condition on the density. Up to V5.52 the densities therefore had no condition at the wall, and the radial particle flux through ρ_b was fixed by the last element. On the X-point test equilibrium the ratio of the ion flux through the wall to that through the separatrix, $\Gamma_\mathrm{wall}/\Gamma_\mathrm{sep}$, was +0.57, −0.40 and −0.28 at NRMAX = 60, 120 and 480: depending on the mesh the wall acted as a source or a sink of particles, and refining the mesh did not make it vanish. The radial flux $n_s u_\mathrm{s}^V$ cannot take the condition instead: its equation is first order and not integrated by parts, so it already has its one condition at the axis. For the charged species ρ_b is a computational boundary at the outer edge of the SOL, not a material wall: their exhaust is carried by ν_L.

V5.60 (MDLWBC = 1) imposes $dn_s/d\rho = 0$ on n_e, n_i and n_z. In the P1 basis this is the constraint

$$n_s(N) = n_s(N-1), \qquad N = \texttt{NRMAX}. \tag{8.2}$$

Constrained degrees of freedom, in general. For a linear system $Ax = b$ with the constraint $x_N = x_{N-1}$, write $x = Ty$, where T is the identity on $y_1 \dots y_{N-1}$ and copies y_{N-1} into x_N. The constrained Galerkin system is

$$T^\mathrm{T} A T\, y = T^\mathrm{T} b. \tag{8.3}$$

T^T adds row N to row $N-1$; AT adds column N to column $N-1$. In finite elements this is the basis reduction $\phi_{N-1} \to \phi_{N-1} + \phi_N$: the trial and test functions are constant on the last element, which is the P1 form of $dn/d\rho = 0$. The code keeps the system square and solves the equivalent

$$\text{row } N-1 \leftarrow \text{row } N-1 + \text{row } N, \qquad \text{row } N \leftarrow x_N - x_{N-1} = 0, \tag{8.4}$$

with x_N still an unknown, so the other equations keep their coupling to it and the column sum is carried by the constraint row. Since $\sum_i \phi_i = 1$, the sum of all rows of a continuity equation is the global particle balance; each column of T^T sums to one, so that sum, and with it particle conservation, is unchanged. Replacing row N by the constraint without adding it to row $N-1$ would drop the particle balance of the wall half-cell, an $O(h)$ conservation error.

Implementation. `txcala.f90`, subroutine `BOUNDARY_ZEROGRAD(LQ)`, called from the boundary-condition block of `TXCALA` for `LQe1`, `LQi1` and `LQz1` after the Dirichlet eliminations. In the band storage the coefficient of term `NC` of equation `NQ` in row `NR` is `ALC`, `BLC` or `CLC(NR,NC,NQ)` for `X(NR+1)`, `X(NR)` or `X(NR-1)`, and `PLC` holds the known terms. For every term `NC = 0..NLCMAX`: `BLC(N-1) += CLC(N)`, `ALC(N-1) += BLC(N)`, `PLC(N-1) += PLC(N)`; row `N` is then cleared and set to `BLC(N,1) = 1`, `CLC(N,1) = -1`. The time term and its right-hand side are built from the same coefficient arrays, so they are folded with them. The condition converges at first order in the mesh size: $\Gamma_{\rm wall}/\Gamma_{\rm sep}$ is -1.8, -1.1, -0.5 and -0.3% at `NRMAX` = 60, 120, 240 and 480.

8.3 C^0–C^{5+} superstage with a source-slaved C^0 part

Carbon is carried by two densities: $n_{\rm z}$ for C^{6+} and the superstage $\bar n_z = \sum_{q=0}^{5} n_{C^{q+}}$, called the bundle below. The bundle exchanges particles with C^{6+} through the effective ionization $f_5 S_5$ and recombination A_6 of Sec. 1, and is fed by sputtering in the SOL: $\gamma_{0\rm z}$ (`rGamm0zz`) is the fraction of the C^{6+} parallel loss and $\gamma_{\rm i,z}$ (`rGamm0iz`) the fraction of the main-ion parallel loss that return as sputtered carbon.

In local ionization balance the C^0 fraction of the bundle is below 2×10^{-6}, so the bundle is almost entirely magnetized C^{1+}–C^{5+} ions. With `MDLNZD` = 1 it is therefore diffused with the anomalous particle diffusivity of the charged species, $D_{nz} = D_{\rm c}$ (`Dfs`). The exception is the carbon that has just been sputtered, which enters as neutral C^0. Its lifetime $1/(S_0 n_{\rm e})$ is of the order of microseconds, shorter than every other time scale here, so its density follows its local source balance:

$$n_{\rm C0} = \min\left(\frac{S_{\rm wall}}{S_0 n_{\rm e}},\ \bar n_z\right), \quad S_{\rm wall} = \gamma_{0\rm z}\nu_{\rm L}\max(n_{\rm z}-n_{\rm zsup},0) + \gamma_{\rm i,z}\nu_{\rm L}\max(n_{\rm i}-n_{\rm isup},0), \tag{8.5}$$

where S_0 is the $C^0 \to C^{1+}$ ionization rate (OpenADAS scd96, `data/Carbon-S0F0_20260921.txt`). $n_{\rm C0}$ enters the $\bar n_z$ equation as a known field through the excess flux with $\Delta D_{\rm C0} = D_{\rm C0} - D_{\rm c}$, the part of the C^0 flux $-D_{\rm C0}\nabla n_{\rm C0}$ that the $D_{\rm c}$ term does not already carry, so that the bundle flux is exactly $-D_{\rm c}\nabla(\bar n_z - n_{\rm C0}) - D_{\rm C0}\nabla n_{\rm C0}$. Here $D_{\rm C0}$ is the Larsen-limited (see below) value of $\bar v_{0z}^2/(3S_0 n_{\rm e})$, the short-mean-free-path diffusivity of C^0, and $\bar v_{0z} = \sqrt{4V_{0z}^2/\pi}$ its mean speed (V_{0z} = `V0z`), and through the parallel loss of the charged part of the bundle, $(1-\gamma_{0\rm z})\nu_{\rm L}(\bar n_z - n_{\rm C0})$, of which the fraction $\gamma_{0\rm z}$ is sputtered back (the $\bar n_z$ equations of Secs. 1 and 3). $n_{\rm C0}$ is evaluated from the previous Picard iterate, $S_{\rm wall}$, $S_0 n_{\rm e}$ and the bound $\bar n_z$ alike, so the bound $n_{\rm C0} \le \bar n_z$ applies to the bundle at the time level being solved for, and the $\bar n_z$ equation stays linear in $\bar n_z$. With `MDLNZD` = 0 (the V5.52 model), $D_{nz} = D_{01}$, the diffusivity of the slow hydrogen neutrals, and the $n_{\rm C0}$ terms are absent.

Element-based C^0 limiter. $D_{\rm C0}$ is Larsen-limited, $D \to D/(1 + D|\nabla n_{\rm C0}|/(\alpha\bar v_{0z} n_{\rm C0}))$, α = `FLlim`. The limiter is evaluated on each element from the element mean of $n_{\rm C0}$ and the element gradient. It acts on the whole $D_{\rm C0}$, because it bounds the whole C^0 flux by free streaming, and $D_{\rm c}$ is subtracted afterwards with its sign: $\Delta D_e = D_e - D_{{\rm c},e}$, where D_e is the limited $D_{\rm C0}$ and $D_{{\rm c},e}$ the element mean of $D_{\rm c}$. $\Delta D_e < 0$ is not a negative diffusion: it removes the part $D_{{\rm c},e}$ of the C^0 flux that the $D_{\rm c}$ term already carries, and since $n_{\rm C0}$ is a known field the matrix is unchanged. It occurs in the first element outside the separatrix, where $n_{\rm C0}$ rises from zero within one element and the limited $D_{\rm C0}$ falls to about $\alpha\bar v_{0z}h/2$ (h the radial width of the element): at NRMAX = 480, not at NRMAX $\le$ 240. Clipping ΔD_e at zero there would let the C^0 flux exceed free streaming by 7%. The element flux $F_e = \langle|\nabla V|^2\rangle_e \Delta D_e\, \Delta n_{\rm C0}/\Delta V_e$ enters the two rows of the element as $+F_e$ and $-F_e$, so the term is exactly conservative. The element evaluation is needed because inside the separatrix the source, and hence $n_{\rm C0}$, vanishes: a nodal ratio $|\nabla n_{\rm C0}|/n_{\rm C0}$ would be 0/0 at the separatrix node, and whether that node is limited would be left to rounding. The limiter is part of the model and is always applied with `MDLNZD` = 1.

Without it D_{C0} becomes very large in the SOL, and the divergence of the known C^0 flux, a source that does not scale with $\bar{n}_z$, drives $\bar{n}_z$ negative just outside the separatrix.

Validity of the source-slaved C^0 part. The slaving of C^0 to its source is implemented and is part of the production model. What remains open is the range of validity of this quasi-steady approximation and the lack of a formal positivity guarantee. The mean free path of C^0 is comparable to the width of the C^0 layer, so the local source balance does not resolve the inner structure of the layer. Positivity of $\bar{n}_z$ is kept by the limiter in the cases computed, but it is not guaranteed, because the divergence of the known C^0 flux is a source that does not scale with $\bar{n}_z$. The bound $n_{C0} = \bar{n}_z$ marks the limit of the model. It is never reached in the converged iterates of the production cases; it is reached only in the first sweep of a run started from the initial profiles, where $\bar{n}_z$ is still close to zero. Where it holds persistently, the lagged treatment of n_{C0} is not appropriate: the C^0 flux in the bounded nodes is then a diffusion of $\bar{n}_z$ itself with $\Delta D_{C0} \approx 7$–$15\, D_c$, taken from the previous iterate, and the iteration diverges (forced with a C^0 ionization rate reduced 100-fold; with the bound taken from the previous time step, the time stepping diverges instead). `TXLOOP` warns when the bound is still active in the last Picard sweep of a step, and with `IDIAG` ≥ 2 prints the number of bounded nodes at every step. Such conditions call for an independent C^0 transport equation, an implicit treatment of the C^0 flux, or a kinetic (Monte-Carlo) treatment, which would also delimit the validity of the approximation.

8.4 Larsen limiter on the hydrogen slow neutrals (optional)

The slow hydrogen neutrals are diffused with the short-mean-free-path diffusivity $D_{01} = \bar{v}_0^2/(3S_{\mathrm{tot}}) = \lambda\bar{v}_0/3$, S_{tot} being the ionization plus charge-exchange rate and $\lambda = \bar{v}_0/S_{\mathrm{tot}}$ the mean free path. The Larsen limiter replaces D by $D/(1+R)$ with

$$R = \frac{D|\nabla n_{01}|}{\alpha\,\bar{v}_0\,n_{01}} = \frac{\text{flux the diffusion law demands}}{\text{free-streaming flux it may not exceed}} = \frac{\lambda}{3\alpha L_n}, \qquad \alpha = \texttt{FLlim},$$

L_n being the gradient length: the limited flux $D|\nabla n|/(1+R)$ is the diffusion flux where $R \ll 1$ and $\alpha\bar{v}_0 n$ where $R \gg 1$. $\alpha = 1/4$ is the one-sided flux of an isotropic population: for particles of speed $\bar{v}$ spread uniformly over all directions, those crossing a plane in one direction carry $n\bar{v}\int_0^1 \mu\, d\mu/2 = n\bar{v}/4$. $\alpha = 1/2$ would correspond to all particles moving into one half-space, uniformly in angle, and $\alpha = 1$ to all of them moving in the same direction, the kinematic upper bound. In the V5.60 solution without the limiter R is 1–3 in the SOL, about 2 at the separatrix and up to 6 at $\rho \simeq 0.8$–0.9 (Kn $= \lambda/L_n \sim 1$): the diffusion approximation is marginal wherever the neutrals matter.

`MDLNOFL` $= 0$ (default) leaves D_{01} unlimited. `MDLNOFL` $= 2$ evaluates the limiter element by element, as for the C^0 flux, from the element mean $\bar{n}$ and the element difference Δn of the current iterate; since $|\Delta n|/\bar{n} \leq 2$ on an element, the factor is bounded by $1 + 2\lambda/(3\alpha h)$. `MDLNOFL` $= 1$ is the earlier nodal form, with $|\nabla n|/n$ from the spline derivative, kept only to reproduce earlier runs. The limited ionization front at $\rho \simeq 0.85$–0.9 is exponential with the decay length $\alpha\lambda \simeq 5$ mm, shorter than an element at `NRMAX` $= 60$ (about 1 cm). On that unresolved front, and in the tiny-n_{01} tail inside it, the nodal ratio is large and erratic, and the Picard iteration does not settle: with `MDLNOFL` $= 1$, 496 of 500 steps reach `ICMAX` at `NRMAX` $= 60$ (`EPS` $= 10^{-6}$), the neutral-only relaxation `RLXNO` $= 0.5$ (used only with `MDLNOFL` $= 1$) leaves 2–3. `MDLNOFL` $= 2$ needs no relaxation: 20.0 / 21.7 sweeps on average and no step at `ICMAX` (X-point / circular, `NRMAX` $= 60$), the same as without the limiter. The two forms agree to 0.5–2% where both converge. Against the limiter off, the limiter raises the neutral inventory by about 30% and changes n_e at the separatrix and at the wall by -3% and $+4\%$; $Z_{\mathrm{eff}}(0)$ is unchanged.

The limiter is a bound on the diffusion flux, not a substitute for kinetic neutral transport. A straight-flight reference (the same sources, attenuated at $S_{\rm tot}$) ionizes or charge-exchanges 35% of the slow neutrals inside the separatrix (X-point, `NRMAX` = 240); plain diffusion gives 49% and the limiter with $\alpha = 1/4$ 25%, with almost nothing beyond $\rho = 0.95$ against 12% in the reference. $\alpha = 1/2$ recovers the total but not the ballistic tail, and no single α recovers both. The difference between limiter on and off is of the size of the uncertainty from the neutral energy alone (the reference gives 24% and 48% at 0.5 and 4.5 eV). The limiter is therefore off by default.

8.5 Anderson acceleration of the Picard iteration (optional)

Each time step is solved by Picard iteration: the coefficients are evaluated from the previous iterate x_k and the linear system gives $g_k = G(x_k)$. On fine meshes the iteration has a slow mode localised on the first few nodes at the magnetic axis, in BV_{1s} (`LQ*8`) of all species, whose contraction factor per sweep tends to 1 as the axis element shrinks (0.70, 0.91, 0.98 at `NRMAX` = 120, 240, 480 with $\Delta t = 10^{-4}$); at `NRMAX` = 480 it takes about 500 sweeps to reach `EPS` = 10^{-6}.

With `NANDER` $\neq 0$ the next iterate is instead

$$x_{k+1} = g_k - \sum_j \gamma_j \,(\Delta x_j + \Delta f_j), \qquad \gamma = \arg\min \Big| f_k - \sum_j \gamma_j \Delta f_j \Big|_2,$$

with $f = g - x$ and Δx_j, Δf_j the differences over the last |`NANDER`| sweeps of the step (Anderson mixing with $\beta = 1$). Each variable is scaled by its profile RMS at the first sweep of the step, as in the convergence test. With `NANDER` < 0 the rows of `LQm1`–`LQm5` do not enter the least-squares fit, because near convergence their increments are rounding (for $\dot\psi_t$ the floor of Sec. 8.6); the correction is applied to all variables. The history is restarted at every time step, and dropped (the plain g_k is used) if the least-squares solve fails or returns $|\gamma| > 10^3$. Once the adaptive relaxation of the step has found the iteration not contracting, the rest of the step is left to the relaxation alone: the linear model behind the mixing does not hold there, and with the mixing kept on the first steps from the initial profiles at `NRMAX` = 960 end in a negative density. A fixed point of G is a fixed point of the accelerated iteration, so the converged solution is that of plain Picard; `NANDER` = 0 (default) is bit-identical to the plain iteration.

With `NANDER` = −20 (100 steps from $t = 0.02$, $\Delta t = 10^{-4}$, `EPS` = 10^{-6}, `ICMAX` = 100, X-point equilibrium) the average number of sweeps falls from 22.7, 36.4, 99.4 to 16.0, 28.4, 58.8 at `NRMAX` = 60, 120, 240, and the steps that reach `ICMAX` at `NRMAX` = 240 from 86 to 9; the circular equilibrium behaves alike. The solution agrees with the plain iteration to within the convergence tolerance. At `NRMAX` = 480 the `LQ*8` mode is removed as well. These figures were measured with the purely relative test of $\dot\psi_t$ (`LQm2`), under which almost every step at `NRMAX` = 480 still reached `ICMAX` on the rounding floor of $\dot\psi_t$, which the acceleration does not affect. With the floor term of Sec. 8.6 the steps at `NRMAX` = 240 and 480 end when the other variables have converged: with `NANDER` = −20 the average number of sweeps is 33.5 (240) and 67.1 (480), and 3 of 100 steps at 480 reach `ICMAX`, on the other variables.

Later in the JT-60U case at `NRMAX` = 240 with NBI (0.2–0.4 s) the slow modes are not a single axis mode but a band of BV_1 modes of all species, on the axis and at $\rho \approx 0.65$–0.9 (32–39 eigenvalues of the Picard map above 0.9), and the mixing removes them only when its depth exceeds the band, as GMRES would. From the state at 0.25 s ($\Delta t = 5 \times 10^{-5}$, 400 steps, `NRETRY` = 1, with the floor of Sec. 8.7) `NANDER` = −10, −20, −30, −40, −50 retake 400, 273, 6, 0, 1 steps, leave 400, 102, 0, 0, 0 of them unconverged at $\Delta t/2$, and take 1.36, 1, 0.46, 0.33, 0.39 times the CPU time of −20; clearing the history every 30 sweeps, or fitting only the BV_1

rows, is worse than -20. The band widens later: from the state at 0.6 s `NANDER` $= -40$ retakes 13% of the steps and -50 0.3%, in 0.66 times the CPU time; clearing the history when the residual grows, or leaving `LQm1`–`LQm5` out of the correction, is worse. The mixing costs about 3 ms per sweep at depth 40 (a sweep about 30 ms). At `NRMAX` $= 240$ with NBI, `NANDER` $= -50$, the deepest accepted, is advised together with $\Delta t = 5 \times 10^{-5}$ s and `NRETRY` $= 1$ (Sec. 8.9); the default of `NANDER` stays 0. The axis mode at `NRMAX` $= 60$ with 6.4 MW (Sec. 8.9) does not depend on the depth.

8.6 Convergence test of $\dot{\psi}_t$ (`LQm2`)

The sweeps of a time step stop when, for every variable X_q, $\max_j |\Delta X_{q,j}| \le \texttt{EPS}\,\mathrm{rms}(X_q)$, ΔX_q being the change over the sweep. For $X_2 = \dot{\psi}_t$ this relative test has a floor that no iteration removes, and the test is

$$\max_j |\Delta X_{2,j}| \le \texttt{EPS}\,\mathrm{rms}(X_2) + C_\mathrm{r}\,\epsilon_\mathrm{m}\,\texttt{NRMAX}^2\,\frac{\mu_0\,\mathrm{rms}(X_5)}{\texttt{ADV}\,\Delta t}, \qquad C_\mathrm{r} = 1,$$

with $X_5 = \psi_t/\mu_0$ (`LQm5`), ϵ_m the machine epsilon and `ADV` the weight of the implicit terms (2/3 for BDF2, 1 in the first step). `LQm2` is not removed from the test: the relative term is kept as it is for every variable, and only the floor term is added. The test of every other variable, the equations and the fixed point are unchanged; $\dot{\psi}$ (`LQm3`) has a floor of the same kind (Sec. 8.7).

The floor has two factors. ψ_t is mostly vacuum flux, and in Ampère's law (the `LQm2` row) the $\nabla \times B_\phi$ term applied to it is a cancellation of terms larger than the result, $|KX_5|/(|K||X_5|) = 3.4 \times 10^{-7}$ at `NRMAX` $=$ 60 falling as `NRMAX`$^{-2}$. A rounding of relative size ϵ_m in that row, in the assembly or in the solve, leaves ψ_t determined only to $(0.05\text{–}0.07)\,\epsilon_\mathrm{m}$`NRMAX`^{2}. Faraday's law (the `LQm5` row) then gives $\dot{\psi}_t$ as the difference quotient $\mu_0(X_5 - X_5^\mathrm{old})/(\texttt{ADV}\,\Delta t)$, which multiplies that rounding by $\psi_t/(\texttt{ADV}\,\Delta t\,|\dot{\psi}_t|)$, 7×10^6 at $\Delta t = 10^{-4}$, $t = 0.02$ s, and without bound as ψ_t approaches a steady state. The relative floor of $\dot{\psi}_t$ is therefore about 4×10^{-7}, 1.5×10^{-6}, 4.6×10^{-6}, 1.8×10^{-5} at `NRMAX` $= 60$, 120, 240, 480 ($\Delta t = 10^{-4}$, $t = 0.02$ s), grows as $1/\Delta t$, and grows in time: at `NRMAX` $= 60$ it reaches 10^{-5} by $t = 0.5$ s. At the floor the equations are satisfied to rounding (backward error of the `LQm2` and `LQm5` rows 10^{-15}–10^{-11}, normwise residual of every equation below 4×10^{-9}); the increment of $\dot{\psi}_t$ is the forward error of an ill-conditioned component, not an unconverged iteration. The second term of the test is that floor with a safety factor.

C_r is not a physical tolerance on $\dot{\psi}_t$. $C_\mathrm{r} = 0.3$ lies on the largest floor observed (the rounding of ψ_t reaches 4–6.5 times its geometric mean) and leaves no margin for rounding that depends on the compiler, the BLAS or the order of operations; $C_\mathrm{r} = 1$, three to four times it, is used. The two values give the same numbers of sweeps to within a few per cent and solutions that cannot be told apart. The test on $\dot{\psi}_t$ keeps its role while the iteration converges, since $\dot{\psi}_t$ responds to a change of the currents about 10^4 times more strongly than they change (`NRMAX` $= 60$). In the printed and returned convergence measures the value for `LQm2` is the part of the relative increment above the floor term.

Against a reference solved with the relative test and up to 300 (100 steps) or 200 (3000 steps) sweeps per step, `NANDER` $= -20$, the test with the floor term gives (`EPS` $= 10^{-6}$, `ICMAX` $= 100$, $\Delta t = 10^{-4}$)

	relative test sweeps / steps at `ICMAX`	with the floor term sweeps / steps at `ICMAX`
X-point, 240, 100 steps	58.8 / 9	33.5 / 0
circular, 240, 100 steps	69.8 / 10	37.1 / 0
X-point, 480, 100 steps	99.6 / 97	67.1 / 3
circular, 480, 100 steps	99.3 / 95	77.0 / 5
X-point, 240, 3000 steps	89.0 / 2167	29.9 / 0
circular, 240, 3000 steps	89.6 / 2093	33.6 / 0

The remaining steps at `ICMAX` are steps on which another variable has not converged. Densities, temperatures, current and Z_{eff} differ from the reference by less than 7×10^{-8}; $\dot{\psi}_t$ and the poloidal electric field derived from it by 0.2–1.5×10^{-4}, within the 0.3–2.1×10^{-4} by which the relative test with `ICMAX` = 100 differs from the same reference, that is, within the floor. The loop voltages and the Poynting-flux terms follow within 6×10^{-7} over 3000 steps, except the rate of change of the poloidal electric-field energy, which is built on $\epsilon_0 \dot{\psi}_t{}^2$ and carries the floor of $\dot{\psi}_t$ under either test. Volume-averaged densities, stored energy and current agree to 10^{-10} and the toroidal angular momentum to 2×10^{-8}. The BDF2 errors and orders of n, T, j and Z_{eff} are unchanged (`NRMAX` = 60, $\Delta t = 4 \times 10^{-4}$–$2.5 \times 10^{-5}$, `EPS` = 10^{-8}); only the self-convergence error of $\dot{\psi}_t$ between the two finest steps rises from 6–8×10^{-6} to 1.5×10^{-5}, its floor. With `NANDER` = 0 the slow `LQ*8` mode still decides most steps at `NRMAX` $\geq$ 240, and the floor term changes little.

8.7 Convergence test of $\dot{\psi}$ (`LQm3`)

$\dot{\psi}$ has a floor of the same origin. Ampère's law for ψ' (the `LQm3` row) holds ψ through a second difference whose terms are about `NRMAX`^{2} times larger than their sum, so ψ is determined only to a relative $\sim \epsilon_{\mathrm{m}}$`NRMAX`^{2}; Faraday's law (the `LQm4` row) gives $\dot{\psi} = -\mu_0(X_4 - c_1 X_4^n + c_2 X_4^{n-1})/(\mathtt{ADV}\,\Delta t)$ with $X_4 = \psi/\mu_0$ (ψ without NBI) and multiplies that rounding by $\tau_\psi/(\mathtt{ADV}\,\Delta t)$, $\tau_\psi = \psi/\dot{\psi}$. The test is

$$\max_j |\Delta X_{3,j}| \leq \mathtt{EPS}\,\mathrm{rms}(X_3) + C_{\mathrm{r}}\,\epsilon_{\mathrm{m}}\,\mathtt{NRMAX}^2\,\frac{\mu_0\,\mathrm{rms}(X_4)}{\mathtt{ADV}\,\Delta t}, \qquad C_{\mathrm{r}} = 1,$$

with `ADV` and Δt those of the current (sub)step, so it follows the variable-step weights of a retried step. X_4 itself and every other variable are tested against `EPS` as before.

Measured from the JT-60U state with NBI at 0.2 s (`NRMAX` = 240, τ_ψ = 23 s): from the second sweep on the increment of $\dot{\psi}$ is $\mu_0/(\mathtt{ADV}\,\Delta t)$ times that of X_4 to all printed digits ($\Delta t = 10^{-4}$–5×10^{-6}), and the relative increment of X_4 stops at (0.02–0.04) ϵ_{m}`NRMAX`^{2} (median; largest 0.07–0.15) at `NRMAX` = 240, 120, 60, at a node that moves from sweep to sweep. The floor term is 4.4×10^{-6}, 4.4×10^{-5}, 8.8×10^{-5} of rms($\dot{\psi}$) at $\Delta t = 10^{-4}$, 10^{-5}, 5×10^{-6}; it grows as $1/\Delta t$ and as the loop voltage relaxes, and so do the steps that the relative test alone leaves unconverged on `LQm3` at small Δt and late times (at $\Delta t = 10^{-5}$ their remaining increments are 0.13–0.32 ϵ_{m}`NRMAX`^{2} in X_4): ψ has converged and the test cannot go below its rounding.

The floor enters only the stopping test; the relaxation and the Anderson mixing use the raw increments. Where it does not stop a step earlier the run is bit-identical: `NRMAX` = 60 with and without NBI (300 steps); at `NRMAX` = 240 ($\Delta t = 10^{-4}$, 300 steps, with and without NBI) every step takes the same number of sweeps and only the printed variable of the largest increment changes. From the state at 0.2 s to 0.4 s with $\Delta t = 10^{-5}$ (`NANDER` = −20, `NRETRY` = 1) the retried steps fall from 2175 to 679 of 20000 and the steps not converged at $\Delta t/2$ from 604 to 75; their last substeps were on `LQm3` (330 of 331) and are now on the beam flux Γ_{b} (`LQb2`)

at $\rho \approx 0.35$ (36 of 37), whose increment moves with that of $\dot{\psi}$ at about 0.2 of its relative size. At 0.4 s the two solutions differ by less than 10^{-5} of the profile RMS except $\dot{\psi}_t$ (3×10^{-4}, its floor) and BV_1 near the axis ($\leq 8 \times 10^{-5}$).

8.8 New defaults and V5.52 compatibility

parameter	V5.52	V5.60	meaning
FSLP	0.3	1.0	k_L
MDLSOL	(0)	1	ramp width δ
MDLNZD	(0)	1	bundle transport with the source-slaved C^0 part
MDLWBC	(0)	1	wall density condition
MDLNOFL	(0)	0	Larsen limiter on D_{01} (2: element form; 1: legacy nodal form)
FLlim	–	0.25	limiter factor α (C^0; D_{01} with MDLNOFL = 1, 2)
RLXNO	–	0.5	neutral-only relaxation (legacy MDLNOFL = 1 only)
NANDER	(0)	0	Anderson acceleration of the Picard sweeps

A value in parentheses means that the switch did not exist in V5.52 and that the value reproduces the V5.52 behaviour. These are all the defaults that differ. `parm/txparm.v552default` sets FSLP = 0.3, MDLSOL = MDLNZD = MDLWBC = MDLNOFL = 0. With it V5.60 is not bit-identical to V5.52.20 (the optimized build schedules the changed routines differently), but one step from the same state gives double-precision states that agree to 10^{-12} (median over the variables) and 10^{-9} at worst, except $\dot{\psi}_t$ at its Picard noise floor (3×10^{-7}); the profile output agrees to its single-precision resolution. The table `data/Carbon-S0F0_20260921.txt` is read at start-up in either case.

8.9 Other changes since V5.51

- MDLPCK = 2 (default): block-tridiagonal solver with one step of iterative refinement; the Picard floor falls from about 10^{-5} to 5×10^{-7} on the worst equation.
- oldmixa (default 0.5): Picard under-relaxation applied only on time steps whose sweep is measured not to contract.
- BDF2 keeps its history across CONT; one backward-Euler step is taken where there is none (cold start, LOAD). After a change of Δt the first step takes the variable-step BDF2 weights of the item on NRETRY below (backward Euler only if Δt grows by more than a factor 2).
- Analytic axis limit of the Pfirsch–Schlüter Fourier factors, $f_m(\cdot, 0) = 0$; $\hat{q}^2$ at the axis from the interior instead of $q(0)^2$.
- D_{0z} uses the impurity neutral speed V_{0z}.
- FSTPTM = 3 (default): multi-ion composition channel for the turbulent particle flux. With FSTPTM = 2 the impurity is not diffused on its own density gradient, and in long runs a grid-scale mode of n_z (wavelength about three mesh intervals, sign-alternating) grows: the JT-60U case of Sec. 7.7 stops at 2.4 s with $T_z < 0$ at $\rho \approx 0.43$. The mode is tied to the mesh, not to a physical scale, and is not a physical impurity instability; it is the discrete form of an under-regularized closure, in which nothing damps short-wavelength perturbations of n_z. The diffusion of each thermal ion on its own gradient in FSTPTM = 3

damps them at the rate Dk^2; the same case then runs to 4 s with no grid-scale component above the curvature of the profile. In the NBI cases n_e and $u_{\mathrm{i}\phi}$ change by less than 1.5% (in the ohmic default case $u_{\mathrm{i}\phi} \approx 0.5$ km/s changes by 15%), the core impurity density by 25–55%. `FSTPTM` = 2 remains available and must be given for every species (`FSTPTM` = 2, 2, 2).

- Turbulent force of `FSTPTM` = 2, 3 in the wall row. The pressure-gradient part of the electron-driven turbulent force is assembled through the diamagnetic flow ($n\,BV_1$, LQ*8) and $n\Phi'$. LQe8 is the algebraic definition $n_\mathrm{e}BV_{1\mathrm{e}} = -(I/\psi')(p_\mathrm{e}'/(Z_\mathrm{e}e) + n_\mathrm{e}\Phi')$, so in the continuum this is the direct $\partial_V p_\mathrm{e}$ and the force is `FSVAHL` $\partial_V p_\mathrm{e}$ + (1 − `FSVAHL`) $T_\mathrm{e}\partial_V n_\mathrm{e}$. Discretely LQ*8 is solved with the SUPG weights $N_j + phN_j'$ (`iSUPG8` = 1) and the force rows with the Galerkin N_i: at an interior node the SUPG moments of the two elements around it cancel to $O(h^2)$, but the wall node has the last element only, and there the two forms differ by the order of the LQe8 residual. The particle flux at the wall is fixed by the wall row of LQ*4, so this gave a spurious wall flux (JT-60U case, `NRMAX` = 60: an electron inflow of 0.22×10^{20} s^{-1}). The wall row (test function of node `NRMAX`) now uses the direct-gradient form for every species (that of `FSTPTM` = 0); not with `MDFIXT` ≠ 0. The removed artificial part of the electron wall flux is 0.38 and 0.20 $\times 10^{20}$ s^{-1} at `NRMAX` = 60 and 120 (1.3 s) and 0.16, 0.10 and 0.045 at 60, 120 and 240 (0.05 s), so it goes with the mesh; the corrected wall flux itself converges slowly (+0.16 and +0.05 at 60 and 120; +0.18, +0.03 and +0.02 at 60, 120 and 240), being a small difference of large terms in the wall row. The non-ambipolar part of the wall flux (the fast-ion return current), j_r, the $j \times B$ torque and L_ϕ are unchanged to three digits, the profiles change by less than 0.5% (JT-60U case to 4 s, `FSRP` = 0 and 1, `FSTPTM` = 2 and 3, X-point and circular test equilibria), and the Picard counts are unchanged.

- `FSTPTM` = 4 (the default since the change below; 3 before): thermal-species self-gradient closure. With fast ions, `FSTPTM` = 3 drives the common transport by ∇n_e, which contains the fast-ion charge gradient, and lets the main ions carry the fast-ion share of the electron flux (weight $n_\mathrm{i} + n_\mathrm{b}$), so the main-ion continuity equation has a sink $\nabla \cdot (n_\mathrm{b}u)$ that does not vanish with n_i. `FSTPTM` = 4 drives the electrons by the thermal-ion charge density $n_\mathrm{th} = \sum_s Z_s n_s$, gives the reaction to the thermal ions with the weights $Z_s n_s/n_\mathrm{th}$ and uses n_th in the composition pairs, so that every thermal species diffuses on its own density gradient with the common thermodiffusion, $\langle \mathbf{\Gamma}_s \cdot \nabla V\rangle = -Dn_s(X_s + C_\mathrm{T}X_T)$, also with fast ions, and the fast ions have no turbulent flux. Momentum and ambipolarity hold element by element (the weights sum to one by construction). Without fast ions it equals 3 (JT-60U ohmic, 300 steps: 10^{-7} in the densities, 5×10^{-5} in $u_{\mathrm{i}\phi}$, the quasineutrality residual); it matches the diagnostic implementation it was developed in to 5×10^{-8}; `FSTPTM` = 3 results are bit-identical. The formulation is in twofluid.tex. It must be given for every species; `NCM` and the save file are unchanged. With fixed temperatures (`MDFIXT` ≠ 0, where `LQe5` carries T_e) the pressure-gradient parts are written with the given $\partial_V T_\mathrm{e}$ as for 2 and 3: $\alpha Z_\mathrm{b}(n_\mathrm{b}/n_\mathrm{e})(\partial_V p_\mathrm{e} - T_\mathrm{e}\partial_V n_\mathrm{e}) = \alpha Z_\mathrm{b} n_\mathrm{b}\partial_V T_\mathrm{e}$ for the electrons and $(n_s/n_\mathrm{th})\{(\alpha-1)n_\mathrm{e} - \alpha Z_\mathrm{b}n_\mathrm{b}\}\partial_V T_\mathrm{e}$ for the thermal ions. Known limits: the beam friction (NBCD) and the beam radial flux still reach the thermal ions through ambipolarity, so positivity of n_i is not guaranteed structurally; the more peaked core densities expose the axis `LQ*8` Picard stall below already at `NRMAX` = 60. JT-60U case at 1.5 s (`NRMAX` = 60), 4 against 3: $n_\mathrm{e}(0)$ +17% (`PNsSUP(1)` = 0.0125) and +21% (0.01), $n_\mathrm{i}(0)$ +26% and +36%, $n_\mathrm{b}/n_\mathrm{i}(0)$ 0.31 → 0.21 and 0.42 → 0.26, $u_{\mathrm{i}\phi}(0)$ −11% and −14%; with 6.4 MW NBI $n_\mathrm{i}(0)$ doubles ($n_\mathrm{b}/n_\mathrm{i}(0)$ 0.92 → 0.35); n_e at $\rho = 0.95$ within 3% in all cases, and $\sum_s Z_s\Gamma_s$ inside the separatrix below 1.5×10^{-7} of the electron flux. In the high-density cases (6.4 MW NBI, `Zeffin` = 2.5, gas puff 0.4) 216–722 steps of 15000 stall in short bursts (208 in the benchmark with 0.0125), none with `NRETRY` = 1, which changes the

profiles by less than 10^{-6}.

- Production defaults of the particle transport: `FSTPTM` = 4, `FSVAHL` (C_T) = −0.25, `PNsSUP(1)` = 0.026 (were 3, −0.5, 0.01; the JT-60U benchmark used 0.0125 with 3 and −0.5). With `FSTPTM` = 4 the thermal ions do not carry the fast-ion charge gradient, so a weaker thermodiffusion no longer depletes them (`FSTPTM` = 3 does for $C_\mathrm{T} \geq -0.4$; $C_\mathrm{T} = 0$ collapses with either); C_T then sets the density level at a given `PNsSUP`, the profile shape staying nearly the same, and `PNsSUP(1)` = 0.026 restores the line average. JT-60U benchmark (3.2 MW, `FSRP` = 0, `NRMAX` = 60, 4 s), old (3, −0.5, 0.0125) against new: line average 1.15 / 1.14 $\times 10^{19}$ m^{-3}, $n_\mathrm{e}(0)$ 1.79 / 1.72, $n_\mathrm{i}(0)$ 1.26 / 1.22, $n_\mathrm{b}/n_\mathrm{i}(0)$ 0.30 / 0.32, n_e at ρ = 0.8, 0.95, 1.0 0.69, 0.41, 0.22 / 0.68, 0.45, 0.33, peaking $n_\mathrm{e}(0)/\bar{n}_\mathrm{e}$ 1.55 / 1.51, $u_{\mathrm{i}\phi}(0)$ 510 / 439 km/s, no Picard stall (one retried step in the initial transient, where the source-slaved C0 bound is active), ambipolarity 10^{-6}; with `PNsSUP(1)` = 0.025 the line average is 1.12. The core is kept and the edge density rises by 10–50%, toward the measured edge (which the old benchmark underestimated); the rotation is 14% lower. The value of C_T is a model choice in the direction suggested by gyrokinetic estimates (weaker thermodiffusion), not a fit to them, and the edge density is still to be compared with the experiment. The old defaults are reproduced bit for bit by giving the three inputs. The save file now stores `FSTPTM` as well (extension record 3; it was the one turbulent-transport input missing from the layout, `FSVAHL` and `PNsSUP` being in it), so a run continued from a file keeps its closure; a file written before has no such record, and then `FSTPTM` keeps its current value (4 by default) with a message, so give `FSTPTM` = 3 or 2 to continue such a run unchanged. Older executables read the new files and skip the record. `parm/txparm.v552default` sets all inputs whose defaults changed after V5.52.20 back to the V5.52 values (also the pinch without saturation and taper, `FSTPTM` = 2, `FSVAHL` = −0.5, the V5.52 `PNsSUP` with `MDLNSUP` = 0, no main-wall recycling, `NRETRY` = 0, `imodel_neo(4)` = 0) and lists what no input restores: the inward constant term of the pinch, the discretely conserving friction, the V5.61 fast-ion fixes and the wall-row form. In the JT-60U-like case without NBI (1.5 s) the code just before the wall-row fix reproduces V5.52.20 with these inputs to 2×10^{-4} in the densities; the wall-row fix lowers the density level by 4.8% (n_z by 8%), uniformly in radius, and is the only noticeable difference; with NBI the core rotation is 40% higher than in V5.52.20 (pinch sign, V5.61 fast-ion fixes). The benchmark results quoted above for the wall row, the recycling and `PNsSUP(1)` = 0.0125 refer to the old defaults. All benchmark numbers of this document before the item on the potato closure below were computed with `imodel_neo(4)` = 0; that item gives the change by 1.

- `NRETRY` (default 1): a step whose Picard iteration reaches `ICMAX` without converging is taken again from its start as 2, 4, … substeps, down to $\Delta t/2^k$ for `NRETRY` = k (≤ 4), and the history is $X(t-\Delta t)$ again afterwards. The first substep, of size $h = \Delta t/2^j$, is the variable-step BDF2 on the history of the step, $X(t-\Delta t)$ and $X(t)$: with $\omega = h/\Delta t$,

$$y_{n+1} - \frac{(1+\omega)^2}{1+2\omega}\, y_n + \frac{\omega^2}{1+2\omega}\, y_{n-1} = \frac{1+\omega}{1+2\omega}\, h\, f(y_{n+1}),$$

  from the quadratic through the three points ($\omega = 1$ gives the weights 4/3, 1/3, 2/3); the later substeps are BDF2 at h. Backward Euler remains only where the step itself has no history (cold start, `LOAD`). Converged steps are untouched: a run in which no step stalls is bit-identical to `NRETRY` = 0. The retried steps keep the second order: with every step of the JT-60U case retried as two substeps (diagnostic build, from the state at 0.1 s over 8 ms, $\Delta t = 2 \times 10^{-4}$–$1.25 \times 10^{-5}$ s), the observed order is that of the continuous BDF2 (median over the variables 1.8, $u_{\mathrm{i}\phi}$ and T_i 2.0), and at $\Delta t = 10^{-4}$ s the result is 3×10^{-9}

($u_{\mathrm{i}\phi}$ 1.4×10^{-9}) from the continuous run at $\Delta t/2$; retried as four substeps, 8×10^{-10} from the run at $\Delta t/4$. With the backward-Euler first substep used before (up to 2026-10) the order was 1.0 and those distances 2.4×10^{-6} ($u_{\mathrm{i}\phi}$ 3×10^{-6}) and 7×10^{-7}. In practice 1–15% of the steps are retried, in quasi-steady phases. Across SAVE/LOAD it behaves as without retries. The summary of a run gives the number and share of retaken steps and the finest substep, and warns when more than 10% of the steps were retaken (a smaller base Δt is then the right fix) or when steps did not converge even at $\Delta t/2^{\texttt{NRETRY}}$.

- Convergence test of $\dot\psi$ (`LQm3`) with a rounding-floor term, as for $\dot\psi_t$ (Sec. 8.7). No input; a run in which no step reaches the floor is bit-identical.

- Known numerical limit, the axis `LQ*8`: with NBI the Picard map has a slowly contracting mode on the axis node of BV_1 (mostly the impurity) and the poloidal field, of the form $\lambda \approx A\Delta t/(1+A\Delta t)$, $A \approx 2\times10^5$ s^{-1} (a fast coupling taken from the previous sweep), with A growing with the mesh refinement and the core peaking of the beam, impurity and ion densities. Without NBI it does not appear up to `NRMAX` = 240. At `NRMAX` = 60 it shows as bursts of steps needing 70–140 sweeps (6.4 MW NBI with `FSTPTM` = 3 or 4, the benchmark with `FSTPTM` = 4), harmless for the solution ($\le 2\times10^{-6}$) and, in those runs, removed by `NRETRY` = 1. With the potato closure (`imodel_neo(4)` = 1) and `NANDER` = −20 or −40 the 6.4 MW case has a window at 0.92–0.96 s that leaves about 175 steps unconverged at $\Delta t/2$ with `NRETRY` = 1; `NRETRY` = 2 converges all but one, $\Delta t = 5\times10^{-5}$ s has no window, and the solutions agree there to 3.5×10^{-5}; for this regime `NRETRY` = 2 is advised (cheaper than halving Δt). At 120 it is sporadic to frequent (`NRETRY` = 1 retakes up to 15% of the steps with 3.2 MW, 65% with 6.4 MW); at 240 it is persistent, and without retries an odd-even mode of $BV_{1\mathrm{z}}$ and n_z grows on the axis until $n_\mathrm{z}(0) < 0$ at 0.1 s; with `NRETRY` ≥ 1 or $\Delta t \le 5\times10^{-5}$ s the run stays positive to 0.4 s. A smaller Δt removes the axis mode (with `NANDER` = −20 the steps left unconverged on it at $\Delta t/2$ over 0–0.4 s fall from 1278 at 10^{-4} s to 23 at 2.5×10^{-5} s and none at 10^{-5} s), but after about 0.2 s the stalls move to a band of slow BV_1 modes of all species (Sec. 8.5) and, at small Δt, to the floor of $\dot\psi$ (Sec. 8.7), which grows as $1/\Delta t$: at 10^{-5} s 11% of the steps from 0.2 to 0.4 s were retried and 604 left unconverged at $\Delta t/2$, so a step that small does not help. At `NRMAX` = 240 with NBI use $\Delta t = 5\times10^{-5}$ s, `NRETRY` = 1 and `NANDER` = −50 (the default of `NANDER` stays 0): with the floor of Sec. 8.7, from 0.2 to 0.8 s 0.4% of the steps are retried and 1 is left unconverged at $\Delta t/2$. Without the floor the same settings retry 4–5% of the steps from 0.5 to 0.7 s (7 unconverged, on $\dot\psi$); with `NANDER` = −40 and the floor 0.8% from 0.2 to 0.4 s but 7% from 0.4 to 0.8 s (10–14% from 0.5 to 0.65 s, 16 unconverged). $\Delta t = 2.5\times10^{-5}$ s costs 1.4–1.5 times as much; at 10^{-4} s the axis mode returns (22% retried, 44 unconverged at $\Delta t/2$, with the floor and `NANDER` = −40). The solutions at 0.4 s with $\Delta t = 10^{-4}$ to 10^{-5} s agree to about 10^{-5} of the profile RMS except $\dot\psi_t$ and BV_1, which carry their floors ($\le 3\times10^{-4}$). With 6.4 MW and `FSTPTM` = 4 at `NRMAX` = 120 the impurity density on the axis becomes negative at 0.86 s even with `NRETRY` = 1. The continuum solution is regular.

- Potato closure near the magnetic axis, `imodel_neo(4)` (default 1 since 2026-10; this element of `imodel_neo` was unused before). The matrix inversion (`MDLNEO` = 2) takes the viscosity of each velocity as $k_\mathrm{b}k_\mathrm{ps}/(k_\mathrm{b}+k_\mathrm{ps})$ of its banana and Pfirsch–Schlüter values (the smaller one dominates; half their harmonic mean). Near the axis the share of the fast tail limited by the banana value changes with ε ($f_\mathrm{t} \propto \sqrt{\varepsilon}$), and the poloidal momentum balance, which fixes the poloidal flow algebraically at each radius, carried this into BV_1 and Φ: they were not analytic in V inside $\rho \approx 0.05$, which for the thermal ions lies inside their potato width ($\rho \approx 0.03$–0.05), where the local theory does not hold. With `imodel_neo(4)` = 1, (i) the

viscosity of each velocity joins a potato branch (the coefficient of NCLASS, Shaing et al. 1996) and a potato-plateau branch with the coefficient of Shaing and Hazeltine, Phys. Plasmas **4** (1997) 4331, Eq. (6), $K_{\rm pp} = 0.0930\,x^{7/3}v_{\rm t}c_4^{4/3}/(q_0R_0)$ ($c_4 = 2q_0\rho_{\rm t}/(\kappa_0R_0)$, $x = v/v_{\rm t}$), whose plateau/PS limit is taken at the orbit width of the particles that set it; this viscosity gives `xmu`, the flows, the resistivity and the bootstrap current; (ii) for ChiNC, D and gflux the potato part carries $K(s) = s(2-s)$ for $s < 1$ (1 beyond), $s = \varepsilon^2/[(xc_4)^{4/3}(1+\nu_*)^{-2/3}]$: the flux of a random walk in the canonical momentum ψ whose step is the orbit width (Shaing, Hazeltine and Zarnstorff, Phys. Plasmas **4** (1997) 771, Eqs. (61) and (73)), reflected at the axis. There is no new input. 0 is the former matrix inversion. The value acts only with `MDLNEO` = 2 and `imodel_neo(5)` = 0; other settings ignore it without a message. The derivation, the literature and the validation are in `potato_axis_regularization.tex` (beside `twofluid.tex`). With 1 the centre of $T_{\rm i}$ is $O(V)$, and $T_{\rm i}$, Φ, $BV_{1{\rm z}}$ and the Φ part of $BV_{1{\rm e}}$ are regular outside the first two or three elements of a mesh (converged between `NRMAX` = 240 and 480); the pressure parts of $BV_{1{\rm e}}$ and $BV_{1{\rm i}}$ keep a ρ^3 component of $n_{\rm e}$ held by the local neoclassical electron fluxes, as with 0 (the electron potato width, $\rho \approx 0.002$, is far below the mesh; a regular $\rho \to 0$ limit would need the finite-orbit inner solution of the electrons). Fluxes, energy balance, densities and ambipolarity agree with 0 to 3–4 digits; in the JT-60U case with NBI at 1.5 s $T_{\rm i}(0)$ is 0.43% higher, $n_{\rm e}(0)$ 0.29% higher and the bootstrap current 0.43% lower. A Picard sweep costs 3.5–6.5% more (`NRMAX` = 60, 120). At `NRMAX` = 60 with NBI the axis `LQ*8` mode above opens a window (1.31–1.335 s in the JT-60U case) in which the steps reach `ICMAX` (with 0 they need 80–99 sweeps there); `NRETRY` = 1 takes them again (246 steps), all converge at $\Delta t/2$, and the solution agrees with a run at $\Delta t/2$ to 10^{-6}–4×10^{-5} before and after the window. `imodel_neo` is in the main record of the save file, so a file keeps the closure it was run with: a file written before this change holds 0 and continues with the former matrix inversion (with the message `Potato closure from the file: imodel_neo(4) = 0 (was 1)`); give `imodel_neo(4)` = 1 after `LOAD` to continue it with the new closure. New runs use 1. Older executables read the new files and use 0. JT-60U benchmark (3.2 MW, `FSRP` = 0, `NRMAX` = 60, 4 s), 0 against 1: line average 1.142 / 1.142 $\times10^{19}$ m^{-3}, $n_{\rm e}(0)$ 1.724 / 1.728, $n_{\rm i}(0)$ 1.218 / 1.228, $n_{\rm z}(0)$ 1.93 / 1.85 $\times10^{17}$ m^{-3}, $n_{\rm b}/n_{\rm i}(0)$ 0.32 / 0.32, $n_{\rm e}$ at $\rho = 0.8$, 0.95, 1.0 unchanged (0.68, 0.45, 0.33), $u_{{\rm i}\phi}(0)$ 439 / 438 km/s, $T_{\rm i}(0)$ 7.85 / 7.88 keV, $T_{\rm e}(0)$ 4.32 / 4.32 keV, bootstrap current 0.125 / 0.124 MA, $W_{\rm p}$ 1.088 MJ, ambipolarity 1.2×10^{-6} in both; retried steps 2 / 246 (the window at 1.310–1.335 s), one step of the initial transient not converged at $\Delta t/2$ in both. With `imodel_neo(4)` = 0 the benchmark of the item on the production defaults above is reproduced bit for bit (all sweeps of the 40000 steps).

- Effective energy of the beam ions in the neoclassical solver (nccoe, booth9). The beam friction and viscosity use the energy $E_{\rm b} = x_{\rm b}E_{\rm c}$ at which the slowing-down distribution $f \propto 1/(v^3+v_{\rm c}^3)$ on $3T_{\rm i} < E < E_{\rm b}$ has the beam temperature as its mean energy, $T(x_{\rm b}) = T_{\rm b}$ with $T(x) = E_{\rm c}[F(x)-F(x_0)]/\ln[(1+x^{3/2})/(1+x_0^{3/2})]$, $x_0 = 3T_{\rm i}/E_{\rm c}$. The integral F was a trapezoidal sum with $\Delta x = 0.1$ (about 4000 terms in the SOL, 6.5–9.4% of a Picard sweep with NBI); it is now the closed form $F(x) = x - G(x)$, $G = -\frac{2}{3}\ln(1+u) + \frac{1}{3}\ln(u^2-u+1) + \frac{2}{\sqrt{3}}\arctan\frac{2u-1}{\sqrt{3}}$, $u = \sqrt{x}$, and the equation is solved by Newton's method safeguarded by a bracket and bisection (`beam_energy_ratio`, shared by nccoe and booth9; for $T_{\rm b} \le 2T_{\rm i}$, where no root exists, $x_{\rm b} = x_0$). The continuous model is unchanged: the trapezoidal error of $E_{\rm b}$, up to $\sim 10^{-3}$, is removed, and with NBI the solution changes by less than $\sim 10^{-4}$ (JT-60U case, 1.5 s: rotation 1.5×10^{-5}, the NBCD current profile 10^{-4}, densities and temperatures $\le 2\times10^{-5}$); runs without NBI are bit-identical. The 4 s JT-60U benchmark of the item on the potato closure changes by at most 2.2×10^{-5} ($n_{\rm z}(0)$; $n_{\rm e}(0)$ 1.1×10^{-5}, $T_{\rm i}(0)$ 8×10^{-6}, $u_{{\rm i}\phi}(0)$ 1.5×10^{-5}, the NBCD current 5×10^{-5}, $W_{\rm p}$ unchanged): its printed values stay, except

the ambipolarity (1.3×10^{-6} with `imodel_neo(4)` = 1) and the step of the initial transient, which now converges at $\Delta t/2$; the Picard count and the 246 retried steps are the same. A Picard sweep with NBI costs 8–12% less (`NRMAX` = 60; 1.5 and 4 s).

- Recycling at the main wall. The radial ion outflow through the wall ρ_b, $\Gamma_s(\rho_b)^+ = \max(n_s\langle \boldsymbol{u}_s \cdot \nabla V\rangle, 0)$ at `NRMAX`, returns as a wall-side inflow of neutrals like the gas puff: `RWALL` (default 0.98) of the bulk ions to n_{01}, `RWALLZ` (0.42) of the impurity ions to the impurity neutrals, and `RWALLIZ` (0, off) of the bulk ions to the impurity neutrals by sputtering; the impurity never returns as a hydrogen neutral. This path is separate from the target recycling of the SOL parallel loss (`rGamm0`, `rGamm0zz`, `rGamm0iz`), which `tx_sol_particle_balance` computes from a divertor recycling loop (escape to the upstream SOL against the pump, $c_1/(c_1 + c_5) = 0.980$ for the bulk ions; for the impurity the reflection and self-sputtering yield 0.42 times an escape factor 0.051, 0.0213; sputtering by the bulk ions 0.0099). There is no such loop at the main wall, so `RWALL` is that of a saturated wall with a small retention and `RWALLZ` the yield itself. In the JT-60U case the main wall returns 0.40×10^{20} s^{-1} (target recycling 14.4, gas puff 28.1); n_e changes by 0.2% (0.6% with `FSRP` = 1), and even `RWALLIZ` = 0.15 (tested with `RWALLZ` = 0.02) changes the core impurity density by 0.04%, the wall source being screened by the SOL. `W` (TXSTAT) and `O 7` (`tx_sources.dat`) print the SOL parallel loss, the wall fluxes and the neutral sources by path (target, main wall, gas puff) with the ionization. The three coefficients are kept in the save file in a second extension record; older files load with the current values and newer files load in older builds.

- Finite-collisionality closure for the higher-order flows; triangularity of grid-traced equilibria regularized; G-EQDSK p' normalization corrected; `ipbtdir` = 1 by default.

- `MDLNSUP` (default 1): the SOL upstream densities of the ions and the impurity, `PNsSUP(2:3)`, are derived from `PNsSUP(1)` and `Zeffin` (quasi-neutral, $Z_{\rm eff}$ = `Zeffin`) whenever the parameters are read; before, they were set once from the defaults and a namelist value of `PNsSUP(1)` or `Zeffin` left them unchanged. `MDLNSUP` = 0 uses `PNsSUP(1:3)` as given. The SOL loss $-\nu_L(n_s - n_{{\rm SUP},s})$ holds the SOL density near $n_{{\rm SUP},s}$, so `PNsSUP(1)` acts as the upstream SOL density and sets the density level of the whole plasma.

- Diagnostic switches, off by default: `iaxproj` (near-axis regularity projection), `Dnfac` (particle diffusion in the impurity continuity equation).

- Input handling. A namelist entered after `P` at the prompt is checked and processed like the parameter file and a namelist typed at the prompt; before, it bypassed the input checks and the derivation of `PNsSUP(2:3)` by `MDLNSUP` = 1. `FSTPTM` must be an integer from 0 to 4 (a value such as 2.5 was taken by the branches of the coefficients for one of the options). The centre and width of the Gaussian modification `PROFD2` of the particle diffusivity, constants 0.99 and 0.07 before, are the inputs `PROFD2C` and `PROFD2W` with the same defaults, kept in a fourth extension record of the save file. `V` lists every namelist input (`NRETRY`, `NANDER`, `MDLNSUP`, `MDLNOFL` and others were missing, and `rNRFi` showed the value of `rNRFe`). In the first steps of a cold start the slow neutrals have not reached the wall and the edge of the n_{02} source region came from 0/0, i.e. from comparisons with NaN; it now stays at the separatrix by an explicit test. Default runs are bit-identical.

- Smoothing. The Savitzky–Golay smoothing of $d\ln n_e/d\rho_V$ (momentum pinch) and of dE_r/dr (ExB shear of CDBM and CDIM) in `TXCALC` was not applied: the number of points to the right shared its name with

the loop variable NR (Fortran does not tell the two apart), so the filter got NRMAX + 1 points to the right and returned the data unchanged. The filter itself treated the ends differently from the interior: the first nl points summed the weights that fall inside the data without renormalizing them (a constant came out as 0.67 of itself on the axis), and the last nr points were not smoothed. The end points now take the least-squares polynomial of the same order over the nl + nr + 1 points next to the end: polynomials up to that order are reproduced, and the noise gain is 0.5–0.97 there (0.95 at the end point itself) against 0.58 inside. The interior is unchanged, and the points are taken as equally spaced in the index, not in ρ. The changed points are not used by the pinch (replaced by a linear interpolation inside $\rho = 0.1$ and tapered to zero beyond RhoPch2), so runs without ExB shear (FSCBSH = 0), with FSLC = 0 and MDOSQZ ≤ 10 are bit-identical to those with the old ends. With ExB shear the axis node now takes dE_r/dr there almost unsmoothed (it was scaled down by the defect): FSCBSH = 1 changes by 10^{-4} (2×10^{-3} in the rotation) at NRMAX = 60 and by 10^{-5} (2–3 $\times 10^{-4}$ in the rotation) at 120 and 240. The orbit-loss rate (FSLC $\neq 0$) peaks at the separatrix and falls by orders of magnitude within a few elements; the filter rings there in the interior as before (down to -5% of the peak), and the end windows now continue the ringing near the wall ($\pm 0.3\%$). The same filter smooths the initial profiles read from file with MDINTT, MDINTN, MDINTC = -2: the first point of the smoothed region lost 35% of the density and 37% of the current density, and is now continuous with the unsmoothed part. There the same name conflict, with the initialized (hence saved) declarations, left the edge unsmoothed in a second initialization in the same run; a restart (R answered with "y") now reproduces the first initialization, and it no longer stops on an allocation of array_init_NRNS that was never freed.

- Pressure gradient of CDBM on the axis. α of CDBM (and CDIM) uses dp/dr over the minor radius. Its axis value came from a one-sided four-point difference, which was also the end condition of the spline; p is even in r, so $dp/dr = 0$ on the axis is now imposed, as for $d\Phi/d\rho$ (dE_r/dr keeps its one-sided axis value, E_r being odd). With a flat or hollow core pressure the one-sided value was large (6.4 MW NBI: +46 kPa/m on the axis against $|dp/dr| < 4$ kPa/m at the next node), and the CDBM coefficient of the axis node was hundreds of times that of the next node. Fixed transport is unchanged. With CDBM and 3.2 MW NBI the profiles change by less than 4×10^{-5} at 1 s ($u_{i\phi}(0)$ by -0.04%) with NRMAX = 60 and by less than 5×10^{-7} with 120 (-0.003%).

- CDBM alone at high NBI power. With FSANOM only (no FSDFIX, no offsets) the anomalous D, χ and μ are those of CDBM, proportional to $|\alpha|^{3/2}$, and vanish where the pressure gradient does. With 6.4 MW of NBI on the test case the thermal ions are depleted near the axis: the electron flux driven by the friction with the beam and the inward beam flux, which the thermal ions carry through ambipolarity, exceed the thermal-ion source inside $\rho \approx 0.05$, and a steady state then needs an inward turbulent flux, i.e. a hollow thermal-ion density, the deeper the smaller D. The core pressure flattens, the CDBM coefficients vanish near the axis, and the run stops with a negative impurity density on the axis (0.50 s with NRMAX = 60, 0.35 s with 120). The code keeps the CDBM-alone model. Fixed transport together with CDBM (FSDFIX with FSANOM), or the offsets PROFDB, PROFMB, PROFCB = 0.1 in all three channels, kept the same case stable; these are options for sensitivity studies. The value 0.1 is a test value that stabilized this case and has no physical justification, and fixed transport alongside CDBM is not a recommended standard setting. FSVAHL = -0.5 instead of -0.25 also removes the depletion; the default is kept.

- Dirichlet conditions in the matrix. BOUNDARY removes a variable fixed by a Dirichlet condition from the

other rows: its coefficients there are set to zero, and a nonzero boundary value moves with them to the constant term. The list of the terms concerned was kept from the first call of the process, and that first call skipped the elimination. The equations are the same either way, but the first sweep of the first run of a process differed from every later run in round-off, which Anderson acceleration carried to 10^{-7} (and 10^{-4} in n_{z0} with `MDOSQZ` = `MDOSQZN` = 1): a run repeated in the same process (`R` answered with "y"), or after a change of `NRMAX`, did not reproduce the first. The list was also never rebuilt, while the variable of a term depends on the input: after a change of `FSTPTM`, `MDBEAM` or `MDFIXT` between runs in one process the stale list removed coefficients of other variables (an $O(1)$ error after `FSTPTM` $4 \to 0$). The terms are now found on every call. A repeated run, a run after a change of input, and a fresh process with that input agree bit for bit; the first run of a fresh process changes by round-off (it is the second run of the former code): at 1 s, with and without NBI, with CDBM and with fixed transport, the densities and temperatures change by less than 4×10^{-7} and `LQm2` by 6×10^{-5}, and the number of sweeps by 0.1%. A run started by `LOAD` is unchanged.

- `LOAD` after a longer run. The save file holds the history of the global quantities (`GVX`, `GVY`) without its length, and `LOAD` read it with the length in memory: after `R`, `S`, `C`, `L` it stopped with an input error. `LOAD` does not keep that history (it starts the time-evolution graphs again at the loaded time), so the records are now skipped; the file format is unchanged, and files of all earlier versions load as before. The time-trace history in the file is used only for τ_E; at the first point after `LOAD` the quantities formed by a difference in time (τ_E and the magnetic Poynting fluxes `PoyntI(3:4)`) still depend on what was in memory.

- Edge transport barrier on a new run. With `MDLETB` $\neq 0$ the transport outside `RhoETB` is multiplied by a factor that falls from 1 to 0.1 over the `NTMAX` steps of a run and is then held. A new run in the same process started with the factor of the previous one, without the ramp; `R` now starts it from 1. `CONT` keeps the factor. The factor is not in the save file: `LOAD` computes it from the step count and `NTMAX` in the file unless it is already held in memory, which gives the held value for a file written at the end of a run, but the ramp of a run that stopped before its end is not restored.

References

[1] M. Honda, T. Takizuka, A. Fukuyama, M. Yoshida and T. Ozeki, Nucl. Fusion **48** (2008) 085003.

[2] M. Yoshida et al., Plasma Phys. Control. Fusion **48** (2006) 1673.

[3] M. Kikuchi and M. Azumi, Plasma Phys. Control. Fusion **37** (1995) 1215, Appendix.

[4] T. Tala et al., Proc. 24th IAEA Fusion Energy Conf. (San Diego, 2012) ITR/P1-19.

[5] T. Tala et al., Nucl. Fusion **51** (2011) 123002.

[6] P. Mantica et al., Phys. Plasmas **17** (2010) 092505.

[7] H. Weisen et al., Nucl. Fusion **52** (2012) 042001; **52** (2012) 114024.

[8] M. Honda et al., Nucl. Fusion **53** (2013) 073050.

[9] I. Casiraghi et al., Nucl. Fusion **61** (2021) 116068.

[10] Rivals et al., Nucl. Fusion **65** (2025) 026038.

[11] Tholerus et al., Nucl. Fusion **65** (2025) 036006 (arXiv:2408.01222).