

# More to teamwork than knowledge, skill and attitude

D Siassakos,<sup>a</sup> TJ Draycott,<sup>a</sup> JF Crofts,<sup>a</sup> LP Hunt,<sup>b</sup> C Winter,<sup>a</sup> R Fox<sup>c</sup>

<sup>a</sup> Southmead Hospital, Bristol, UK <sup>b</sup> Department of Clinical Sciences at South Bristol, Faculty of Medicine and Dentistry, Institute of Child Life and Health, Bristol, UK <sup>c</sup> Taunton and Somerset Hospital, Taunton, UK

Correspondence: Dr D Siassakos, Women's Health, Chilterns, Southmead Hospital, Westbury on Trym, Bristol BS10 5NB, UK.

Email jsiasakos@gmail.com

Accepted 1 June 2010. Published Online 7 July 2010.

**Objective** To assess whether team performance in simulated eclampsia is related to the knowledge, skills and attitudes of individual team members.

**Design** Cross-sectional analysis of data from the Simulation and Fire Drill Evaluation randomised controlled trial.

**Setting** Six secondary and tertiary maternity units in south-west England.

**Participants** One hundred and fourteen maternity professionals in 19 teams of six members; one senior and one junior obstetrician; two senior and two junior midwives.

**Methods** We validated a team performance ranking scheme with respect to magnesium administration (Magnesium Administration Rank, MAR) by expert consensus (face validity) and correlation with clinical measures (construct validity). We tested for correlation between MAR and measures of knowledge, skills and attitudes.

**Main outcome measures** Correlation between team performance (MAR) and scores in validated multiple-choice questionnaires (MCQs) (knowledge), a measure of individual manual skill to

manage an obstetric emergency (skill) and scores in a widely used teamwork/safety attitude questionnaire (attitude).

**Results** There was no relationship between team performance and cumulative individual MCQs, skill or teamwork/safety attitude scores.

**Conclusions** The knowledge, manual skills and attitudes of the individuals comprising each team, measured by established methods, did not correlate in this study with the team's clinical efficiency in the management of simulated eclampsia. The inference is that unidentified characteristic(s) play a crucial part in the efficiency of teams managing emergencies. Any emphasis of training programmes to promote individual knowledge, skills and attitudes alone may have to be re-examined. This highlights a need to understand what makes a team efficient in dealing with clinical emergencies.

**Keywords** Eclampsia, education, emergencies, knowledge, leadership, magnesium sulphate, multiple choice questions, obstetric labour complications, patient care team, pre-eclampsia, safety attitudes, simulation, skills and attitudes, teaching, teamwork, training.

Please cite this paper as: Siassakos D, Draycott T, Crofts J, Hunt L, Winter C, Fox R. More to teamwork than knowledge, skill and attitude. BJOG 2010;117:1262–1269.

## Introduction

Obstetric emergencies are unpredictable and sudden. Successful management requires a rapid coordinated response by *ad hoc* multiprofessional teams. The need to provide training in team coordination and communication for clinicians has been repeatedly identified as a safety priority for developed countries.<sup>1–3</sup>

Eclampsia is an emergency that can result in severe complications for both mother and baby.<sup>4,5</sup> The members of clinical teams that manage eclampsia might not have worked

together in this context previously. The teams need to perform a number of key tasks, in particular the administration of magnesium sulphate, in accordance with national guidelines derived from confidential enquiries and systematic reviews of evidence.<sup>6–8</sup> Magnesium administration for women with eclampsia is associated with a decrease in the recurrence of convulsions, a trend for lower maternal mortality and a significant reduction in serious maternal morbidity.<sup>8</sup> For the baby, its administration is associated with a significant decrease in perinatal mortality and fewer admissions to special care baby

units.<sup>8</sup> Some teams are efficient in administering magnesium in this context; other teams are less efficient or never consider giving it.<sup>4,9–12</sup>

It appears that some clinical teams possess characteristics that make them more efficient than others, and so are better able to achieve good outcomes by performing key actions in a timely manner. This association between efficiency and overall team performance, as well as outcome, has been shown not only for simulated eclampsia,<sup>12</sup> but also for simulated postpartum haemorrhage<sup>13</sup> and real-life umbilical cord prolapse.<sup>14</sup> Moreover, it has been shown that an inefficient response to other medical emergencies, in association with poor teamwork, is the root cause of many adverse outcomes, often leading to medical litigation.<sup>15</sup>

If the characteristics of efficient teams could be identified, the information could be used to inform training programmes. The aim of this study was to explore the relationship between team efficiency and team members' knowledge, skills and attitudes (KSA).

## Methods

### Design

This is a cross-sectional analysis of data from a large randomised controlled trial of training for obstetric emergencies [Simulation and Fire Drill Evaluation (SaFE) study].<sup>13</sup> 'SaFE' was a portfolio of studies commissioned by the Department of Health for England and Wales, the primary aim of which was to compare training at a simulation centre with training in local units, and clinical training alone versus clinical training with the addition of extra teamwork training. The methodology has been described in detail elsewhere.<sup>12,16–19</sup> In this report, we focus on the results prior to the training process.

### Participants

Participants were recruited to the study in 2004–5 from six large maternity units in the south-west of England.<sup>12</sup> Participants were randomly selected from staff lists and allocated to 24 simulation teams (four teams from each unit). The individual teams were made up of staff from one unit, and each team comprised one senior doctor, one junior doctor, two senior midwives and two junior midwives. For the SaFE study, there were no data of training effectiveness on which to base power calculations.

### Measurements

#### *Knowledge, skills and attitudes (KSA)*

All participants were evaluated individually for knowledge [using validated multiple-choice questionnaires (MCQs)],<sup>13,16</sup> skill at managing an obstetric emergency (using simulated shoulder dystocia with a standardised

scenario)<sup>19</sup> and attitudes to safety and teamwork (using a validated questionnaire).<sup>13,20</sup>

The MCQ questions, written by expert midwives, obstetricians and obstetric anaesthetists, were based on evidence-based guidance and published literature. The maximum number of marks for the MCQs was 185, which included 30 marks specific to eclampsia (Appendix S1, see Supporting information).

For skills, one of the main aims of training was to improve both the accoucheur's efficiency (to limit fetal hypoxia) and care (to reduce trauma). The multiprofessional research team looked at the SaFE study records and devised an ordinal score (rank) based on the skills and effectiveness of individual team members exhibited during shoulder dystocia drills. The ordinal score reflected recommendations from national confidential enquiries<sup>21</sup> and guidelines,<sup>22</sup> and correlated significantly with the maximum amount of force used by the participants (Kendall's  $\tau_b = 0.19$ ,  $P = 0.016$ ; Appendix S2, see Supporting information).

For attitudes, each team member completed a validated safety and teamwork attitudes questionnaire.<sup>20</sup> It comprised 57 items and participants answered with a five-point Likert scale. The maximum score was 100, and higher scores indicated more positive attitudes.

#### *Team performance*


Teams were evaluated for their ability to manage simulated eclampsia with a standardised clinical scenario. The team evaluations were undertaken before the teams entered a training programme. The team members were not aware of the nature of the simulation before it started. The video recordings were evaluated with checklists derived from national recommendations.<sup>6–8</sup> All evaluations were undertaken by two trained external assessors (doctor and midwife) working independently, who viewed the digital recordings in different sequences randomly generated by computer.

We regarded success in obtaining, preparing and administering magnesium sulphate as the most important observable and documented team action, and formulated a clinical efficiency ranking based on the administration of magnesium sulphate (Magnesium Administration Rank, MAR). To reach consensus and establish face validity of the ranking scheme, we used a 'Delphi' technique, modified for combined face-to-face meetings and email communication. Consensus was reached over four formal meetings by four clinical obstetricians, a research midwife and a statistician/methodologist. The teams were grouped, blind to their KSA scores to avoid bias, according to their performance as follows: (i) did not obtain magnesium; (ii) obtained but did not prepare magnesium; (iii) prepared but did not administer magnesium; (iv) administered magnesium but

≥6 minutes from the start of the drill; (v) administered magnesium <6 minutes from the start of the drill. The 6-minute cut-off was based arbitrarily on the frequency distribution of the teams' recorded timings to obtain roughly equal group sizes.

### Validation

The team MARs correlated with the ability to perform other key clinical actions for the management of eclampsia (Figure 1), confirming the construct validity of the scores. The team MARs also increased after training (Table 1,  $P = 0.035$ , two-tailed sign test), providing further supportive evidence of construct validity.



**Figure 1.** Correlation between Magnesium Administration Rank (MAR) and other measures of clinical team efficiency. (A) Lowering of head rest: higher magnesium scores (MAR) tended to be associated with shorter time intervals: Kendall's rank correlation coefficient  $\tau_{ab} = -0.32$ ,  $P = 0.09$ . (B) Administration of oxygen: higher magnesium scores (MAR) were generally associated with shorter time intervals:  $\tau_{ab} = -0.29$ ,  $P = 0.12$ .

**Table 1.** Validation of Magnesium (Mg) Administration Rank (MAR) as a team performance measure: comparison of team MARs before and after training. Statistically significant improvement after training confirmed the validity of the scheme ( $P = 0.035$ , two-tailed sign test)

MAR pre-training	MAR post-training				
	1	2	3	4	5
1 Did not obtain Mg	0***	1*	0*	0*	0*
2 Obtained but did not prepare Mg	0	0***	0*	3*	1*
3 Prepared but did not administer Mg	0	0	0***	1*	1*
4 Administered Mg but ≥6 minutes from start of drill	0	0	0	1***	5*
5 Administered Mg <6 minutes from start of drill	0	0	1**	2**	3***
Total	0	1	1	7	10

\*Higher rank:  $n = 12$ .

\*\*Lower rank:  $n = 3$ .

\*\*\*No change:  $n = 4$ .

### Analysis

This study tested for correlation between the team MARs and various aspects of individual team members' KSA as follows.

For 'knowledge', we used the following: (i) the team's minimum individual score (the weakest link in the team); (ii) the team's maximum individual score (regardless of profession or seniority); (iii) the senior doctors' MCQ scores alone; (iv) the average of all the team members; (v) the first principal component of the team members' scores, a weighted average that 'best' represented the individual staff member's scores and maximised the variability between them. As the pairs of junior doctors and junior midwives were interchangeable, the averages of their pairs of scores were used in the principal component analysis. We repeated the analysis for both total and eclampsia-specific MCQs.

For 'skills', we used the team members' shoulder dystocia ranking score, based on their ability to deliver a baby with simulated shoulder dystocia. Again, we tested for correlation between MAR and team average, team maximum and senior doctor's skill score.

For 'attitudes', we looked at the team average scores in the six domains ('teamwork climate', 'safety climate', 'job satisfaction', 'stress recognition', 'perceptions of management', 'work conditions') of the validated teamwork/safety attitudes questionnaire, calculated in accordance with the instructions of the authors.<sup>20</sup>

We were not interested in the effect of training and report analyses based on the baseline (pre-training) data

only. To study the relationships between MAR and KSA, we calculated nonparametric correlations (Kendall's tau<sub>b</sub>) because of the ordinal nature of MAR (a part qualitative measure). The SAS software package (SAS v 9.1, 2002–3, SAS Institute Inc., Cary, NC, USA) was used to obtain approximate 95% confidence intervals.

## Results

Although 24 teams underwent pre-training evaluation, one simulation was not recorded because of a fault in the recording equipment and four teams were incomplete (five members only) because of failure of individual attendance. To ensure comparability, we report here on the remaining 19 complete teams whose 114 participants had complete audiovisual, MCQ and shoulder dystocia records. Attitude questionnaires were complete for all 114 individuals and all 57 questions, apart from eight participants who had not answered isolated questions; domain scores for these were the averages of the relevant completed questions.

There was a wide spread of values for all components of the KSA categories (Table 2). There was no correlation

between team leader (senior doctor), team maximum or team minimum MCQ scores and team MAR (Table 3). Similarly, there was no correlation between the average of the team member MCQ scores and team rank (MAR), regardless of whether a simple or weighted average (first principal component) was used (Table 3). For eclampsia-specific MCQ scores, Kendall's tau<sub>b</sub> was consistently negative, albeit low (<0.3) and statistically nonsignificant ( $P > 0.05$ ) (Table 3b).

There was also no correlation between team MAR and team average, team maximum or senior doctors' manual skill scores (Table 4). Finally, there was no correlation between team average teamwork/safety attitude scores in any of the six domains and MAR scores (Table 5).

## Discussion

This study assessed the ability of a large cohort of multi-professional teams from across a large health region in England to manage a simulated complex obstetric emergency. We examined the relationship between team performance and measures of KSA for individual clinicians, using data from pre-training evaluations to represent the prevailing level of knowledge and skills within the region. The results are compatible with an absence of a relationship between the KSA of individuals and team performance.

We devised a method for ranking team performance (MAR) using the ability to administer magnesium sulphate, an internationally recommended treatment for eclampsia. This does not incorporate all aspects of care important for successful outcome, but is a measure of team efficiency. It has not been validated against true clinical outcomes, but face, content and construct validity were established, including an assessment against the performance of other key actions for eclampsia.

Some might argue that MCQs are a superficial measure of knowledge. The MCQ questions in this study focused on the factual knowledge required to manage successfully emergencies, including eclampsia, and were linked to the objectives of the training intervention of the SaFE study to ensure content validity.<sup>16</sup> After a pilot study, only items that were sufficiently discriminatory were retained to establish construct validity for that study,<sup>13</sup> further supported by evidence that MCQ scores improved after clinical training.<sup>16</sup> Further to this, one component of the skill assessment tested the ability to apply theoretical knowledge in a coordinated manner during a simulated emergency (shoulder dystocia) that relied solely on the individual rather than the team. Measuring attitudes is a difficult task. The attitudes scores in this study were derived from a validated questionnaire, specifically designed for maternity care, which has been shown to correlate with patient outcomes.<sup>23</sup> It is suitable for surveys to evaluate views on team

**Table 2.** Descriptive data

	Mean (SD)	Range
<b>Knowledge</b>		
<b>Total MCQ score</b>		
Team minimum	54 (13)	32–82
Team maximum	111 (9)	96–128
Senior doctor	109 (13)	72–128
Team average	80 (7)	68–93
Team first PC* (unstandardised)	44 (18)	15–92
<b>Eclampsia component of MCQ score</b>		
Team minimum	6 (3)	–2 to 10
Team maximum	20 (3)	14–24
Senior doctor	19 (3)	14–24
Team average	13 (2)	7–17
Team first PC (unstandardised)	18 (5)	3–24
<b>Manual skills</b>		
Team maximum	2.8 (1.3)	0–4
Senior doctors	2.2 (1.3)	0–4
Team average	1.1 (0.7)	0–2.2
<b>Teamwork/safety attitude components (team average)</b>		
Teamwork climate	72.0 (5.3)	57.6–81.3
Safety climate	68.4 (5.6)	56.5–79.2
Job satisfaction	64.8 (8.6)	50.8–75.6
Stress recognition	70.8 (8.0)	59.4–87.5
Perceptions of management	46.0 (7.5)	33.3–57.3
Work conditions	59.0 (6.4)	46.9–69.8

MCQ, multiple-choice questionnaire; SD, standard deviation.

\*PC, the first principal component of the team members' scores is a weighted average that 'best' represents the individual results and maximises the variability between them.

**Table 3.** Team knowledge (multiple-choice questionnaire, MCQ) scores and team rank (Magnesium Administration Rank, MAR): no correlation of MAR with either total (all obstetric emergencies) MCQs (a) or specific (eclampsia-only) MCQs (b)

	Team rank (MAR)					Kendall's tau <sub>b</sub> [approximate 95% CI]
	1 (n = 1)	2 (n = 4)	3 (n = 2)	4 (n = 6)	5 (n = 6)	
<b>(a) Total MCQ score</b>						
Team minimum	58*	48 (SD 5)	54 (SD 1)	50 (SD 16)	62 (SD 15)	0.17 [−0.16 to 0.49] <i>P</i> = 0.36
Team maximum**	116	115 (SD 10)	119 (SD 6)	104 (SD 4)	112 (SD 9)	−0.15 [−0.53 to 0.23] <i>P</i> = 0.42
Senior doctor	116	115 (SD 11)	119 (SD 6)	96 (SD 14)	112 (SD 10)	−0.16 [−0.50 to 0.19] <i>P</i> = 0.40
Team average	83	78 (SD 5)	82 (SD 2)	78 (SD 8)	84 (SD 9)	0.12 [−0.24 to 0.48] <i>P</i> = 0.51
Team first PC (unstandardised)	42	34 (SD 12)	37 (SD 1)	50 (SD 24)	47 (SD 21)	0.24 [−0.10 to 0.58] <i>P</i> = 0.19
<b>(b) MCQ eclampsia score</b>						
Team minimum	8*	6 (SD 3)	10 (SD 0)	6 (SD 2)	5 (SD 4)	−0.26 [−0.58 to 0.07] <i>P</i> = 0.17
Team maximum**	23	20 (SD 2)	22 (SD 0)	19 (SD 3)	20 (SD 3)	−0.15 [−0.52 to 0.21] <i>P</i> = 0.42
Senior doctor	23	20 (SD 2)	21 (SD 2)	17 (SD 3)	19 (SD 3)	−0.19 [−0.55 to 0.18] <i>P</i> = 0.32
Team average	15	12 (SD 2)	16 (SD 2)	13 (SD 2)	12 (SD 3)	−0.14 [−0.52 to 0.24] <i>P</i> = 0.44
Team first PC (unstandardised)	21	18 (SD 4)	20 (SD 6)	18 (SD 4)	17 (SD 7)	−0.03 [−0.37 to 0.32] <i>P</i> = 0.88

PC, principal component analysis.

\*Group means are shown together with standard deviations (SD) in parentheses.

\*\*In 15 of 19 cases, the team maximum total score was that of the senior doctor.

and patient safety, and as a quality improvement tool to identify residual or new challenges in units with established safety programmes.<sup>24,25</sup>

The data showed that there was a wide range of outcomes for the management of simulated eclampsia; some teams administered magnesium in a timely manner, but other teams did not even consider its use. To our surprise, there was no relationship between team performance in eclampsia and the measures of KSA that were used in this study. These are conventional means used to assess individual doctors and midwives,<sup>26–30</sup> but clearly, used alone, these measures can neither predict the effectiveness of clinical teams, nor help to understand the problems of ineffective teams. It follows that the focusing of postgraduate training on these conventional domains of learning for individuals alone might not maximise team performance, nor optimise the outcome of emergencies that rely on complex rapid multiprofessional working.

A criticism might be that the sample size was not sufficient. However, this study was based on a pragmatic sample from a very large study of training that will be very

difficult to replicate. Not only were *P*-values nonsignificant in our study, but correlation coefficients were also very low, indicating a true absence of correlation rather than a lack of power to detect a correlation. This lack of correlation was confirmed by visual inspection of scatter plot charts of all variables against MAR (DS and LPH).

Reports and enquiries into poor outcome and patient safety<sup>2,5,31</sup> have repeatedly demonstrated that failures in teamwork are more frequent than failures of individuals. As a result, training in healthcare has evolved to include practical team training. Regular practical clinical training (skills drills) has been associated with improvements in real-life perinatal outcomes in observational studies,<sup>14,32–34</sup> but even after training there remains a wide range in team performance.<sup>12,13</sup> The inference of our study is that there is some other characteristic of clinical teams or team members that governs team efficiency, and which is not directly related to conventional clinical measures of the individual members.

It is unclear why teams differ in their abilities, and why teams do not improve equally after training. It seems likely

**Table 4.** Tests for correlation between team rank (Magnesium Administration Rank, MAR) and team members' manual skills (shoulder dystocia practical management scores): no correlation with team maximum, team average or senior doctor's scores

	Team rank (MAR)					Kendall's tau <sub>b</sub> [approximate 95% CI]
	1 (n = 1)	2 (n = 4)	3 (n = 2)	4 (n = 6)	5 (n = 6)	
<b>Team maximum manual skill score</b>						
0 (did not deliver)	0	0	0	1	1	
1 (delivered. within 4–5 minutes)	0	1	0	0	0	0.21
2 (delivered within 3–4 minutes)	1	0	0	0	1	[–0.18 to 0.60]
3 (delivered within 2–3 minutes)	0	3	1	3	1	<i>P</i> = 0.30
4 (delivered within 1–2 minutes)	0	0	1	2	3	
<b>Senior doctor's manual skill score</b>						
0 (did not deliver)	0	0	0	2	1	
1 (delivered within 4–5 minutes)	1	1	0	0	0	0.16
2 (delivered within 3–4 minutes)	0	1	1	1	2	[–0.23 to 0.54]
3 (delivered within 2–3 minutes)	0	2	1	3	1	<i>P</i> = 0.42
4 (delivered within 1–2 minutes)	0	0	0	0	2	
<b>Team average of manual skill scores</b>	0.7	1.3 (SD 0.8)	1.3 (SD 0.9)	1.1 (SD 0.8)	1.1 (SD 0.6)	–0.01 [–0.36 to 0.34] <i>P</i> = 0.94

**Table 5.** No correlation between team rank (Magnesium Administration Rank, MAR) and team average teamwork/safety attitudes scores in any of the six domains of the validated questionnaire

Team average (team attitudes questionnaire domains)	Team rank (MAR)					Kendall's tau <sub>b</sub> [approximate 95% CI]
	1 (n = 1)	2 (n = 4)	3 (n = 2)	4 (n = 6)	5 (n = 6)	
Teamwork climate**	77.1*	69.7 (SD 8.2)	75.0 (SD 2.0)	72.6 (SD 4.2)	71.3 (SD 5.2)	–0.26 [–0.66 to 0.14] P = 0.15
Safety climate**	75.0	64.5 (SD 6.7)	69.0 (SD 0.8)	72.0 (SD 5.3)	66.1 (SD 3.7)	–0.12 [–0.50 to 0.26] P = 0.51
Job satisfaction**	75.6	60.1 (SD 9.6)	67.1 (SD 4.1)	69.9 (SD 5.1)	60.3 (SD 9.0)	–0.18 [–0.60 to 0.25] P = 0.34
Stress recognition***	63.5	75.5 (SD 11.7)	63.5 (SD 5.9)	67.7 (SD 6.6)	74.5 (SD 4.8)	0.19 [–0.18, 0.57] P = 0.29
Perceptions of management**	52.1	42.7 (SD 10.2)	50.5 (SD 3.7)	49.0 (SD 5.8)	42.7 (SD 7.3)	–0.18 [–0.54 to 0.19] P = 0.34
Work conditions***	65.6	55.5 (SD 8.3)	59.9 (SD 5.2)	62.8 (SD 5.7)	56.3 (SD 4.7)	–0.13 [–0.51 to 0.25] P = 0.46

\*Group means are shown together with standard deviations (SD) in parentheses.

\*\*We calculated the component team attitude questionnaire scores as the average of the relevant questions that were not missing.

\*\*\*For stress recognition and work conditions, there were no missing values.

that there are other characteristics of individuals, perhaps involving attributes such as composure, confidence and communication. One systematic review of team performance concluded that the ability to retrieve and use knowledge and skills is impaired during acute events.<sup>35</sup> It might be that variation in team performance is influenced by the

different abilities of team members to deal with the effects of anxiety. Further insight into how the behavioural characteristics of individuals influence the way teams work during acute events might provide a means of coaching individual clinicians and/or clinical teams to become more effective. By further studying clinical teams in simulation



and real life, we can hope to develop specific tools for assessment and training, rather than use methods that were developed for airplane pilots.<sup>12,36</sup>

## Conclusions

The data obtained in this study show that team efficiency amounts to more than the conventional clinical characteristics of individual team members. It is often said that a team is more than the sum of its parts, but, in truth, it seems likely that we simply do not yet know how to measure the full contribution of individual clinicians. There is an important need to understand these individual factors and how they make a team more efficient in dealing with emergencies. This improved understanding could inform the development of specific evidence-based methods for the analysis and improvement of teamwork in maternity care and other clinical specialities.

## Disclosure of interest

TJD and CW are members of the steering committee of PROMPT, a UK-based charity running training courses, but derive no financial interest from this association.

## Contribution to authorship

DS conceived the idea, coordinated the analysis, and wrote and edited the manuscript. TJD conceived the idea, coordinated the analysis, and wrote and edited the manuscript. JFC coordinated the SaFE study, helped with conceptual analysis and edited the manuscript. LPH reviewed the methodological considerations, conducted the statistical analysis, wrote parts of the manuscript related to methods and edited the manuscript. CW coordinated the SaFE study, helped with the conceptual analysis and edited the manuscript. RF conceived the idea, coordinated the analysis, and wrote and edited the manuscript. All authors participated in several multiprofessional meetings to develop the study tools and methods.

## Details of ethics approval

Ethical approval was granted by a Regional Research Ethics Committee (SOUTHWEST DEVON MREC 04/Q2103/68). This study was conducted in accordance with the Research Governance Framework for Health and Social Care and Good Clinical Practice. Data storage and protection were in accordance with the Research Governance Framework and the Data Protection Act.

## Funding

This study is an extension to the SaFE ('Simulation and Fire Drill Evaluation') study. The extension is funded by The North Bristol Research and Development Small Grant Scheme. The SaFE study was funded by the Department of

Health Patient Safety Programme. None of the sponsors were involved in the study design, collection, analysis and interpretation of the data, writing of the manuscript or the decision to submit the article for publication.

## Acknowledgements

The authors thank the following individuals for their contributions to the SaFE Study: Christine Bartlett, Karen Cloud, Maureen Harris, Sarah Fitzpatrick, Derek Tuffnell, Bryony Strachan, Stephanie Withers, Valentine Akande, Fiona Donald, Imogen Montague, Penny Watson, Anne McCrum, Sarah Read, Heather Smart, Melanie Robson, Katie Harrison, Neil Liversedge, Joanne Crocker and Simon Grant.

## Supporting information

The following supplementary materials are available for this article:

**Appendix S1.** Multiple-choice questions (MCQs) for eclampsia.

**Appendix S2.** Team members' skills' ranking based on effectiveness and efficiency during shoulder dystocia drills (time from delivery of the baby's head to delivery of the body). The ranking also correlated weakly, but significantly, with the maximum amount of force used by the participants (Kendall's tau<sub>b</sub> = 0.19,  $P = 0.016$ ).

Additional Supporting information may be found in the online version of this article.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author. ■

## References

- 1 King's Fund. *Safe Births: Everybody's Business. An Independent Enquiry into the Safety of Maternity Services in England*. London: King's Fund, 2008.
- 2 Joint Commission on Accreditation of Healthcare Organizations. Preventing infant death and injury during delivery. *Sentinel Event Alert* 2004;30:1–3.
- 3 Bates DW, Larizgoitia I, Prasopa-Plaizier N, Jha AK. On behalf of the Research Priority Setting Working Group of the W.H.O. World Alliance for Patient Safety, Global priorities for patient safety research. *Br Med J (Clin Res Ed)* 2009;338:1242–3.
- 4 Knight M. Eclampsia in the United Kingdom 2005. *BJOG* 2007;114:1072–8.
- 5 Lewis G, editor. The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer – 2003–2005. *The Seventh Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom*. London: CEMACH, 2007.
- 6 Royal College of Obstetricians and Gynaecologists (RCOG). *The Management of Severe Pre-eclampsia/Eclampsia. Guideline No. 10(A)*. London: RCOG Press, 2006.

- 7 ACOG Practice Bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. *Obstet Gynecol* 2002;99:159–67.
- 8 Duley L, Henderson-Smart D. Magnesium sulphate versus phenytoin for eclampsia. *Cochrane Database Syst Rev* 2003;4: Art. No.: CD000128. DOI: 10.1002/14651858.CD000128.
- 9 Schutte JM, Schuitemaker NW, van Roosmalen J, Steegers EA. Substandard care in maternal mortality due to hypertensive disease in pregnancy in the Netherlands. *BJOG* 2008;115:732–6.
- 10 Zwart JJ, Richters A, Ory F, de Vries JJ, Bloemenkamp KW, van Roosmalen J. Eclampsia in the Netherlands. *Obstet Gynecol* 2008;112: 820–7.
- 11 Thompson S, Neal S, Clark V. Clinical risk management in obstetrics: eclampsia drills. *Qual Saf Health Care* 2004;13:1279.
- 12 Ellis D, Crofts JF, Hunt LP, Read M, Fox R, James M. Hospital, simulation center, and teamwork training for eclampsia management: a randomized controlled trial. *Obstet Gynecol* 2008;111:723–31.
- 13 Strachan B, Crofts J, James M, Akande V, Hunt L, Ellis D, et al. *Proof of Principle Study of the Effect of Individual and Team Drill on the Ability of Labour Ward Staff to Manage Acute Obstetric Emergencies*. Edgbaston, Birmingham: PSRP, Department of Health, Public Health, Epidemiology and Biostatistics, University of Birmingham, 2008.
- 14 Siassakos D, Hasafa Z, Sibanda T, Fox R, Donald F, Winter C, et al. Retrospective cohort study of diagnosis–delivery interval with umbilical cord prolapse: the effect of team training. *BJOG* 2009;116: 1089–96.
- 15 Hanscom R. Medical simulation from an insurer's perspective. *Acad Emerg Med* 2008;15:984–7.
- 16 Crofts JF, Ellis D, Draycott TJ, Winter C, Hunt LP, Akande VA. Change in knowledge of midwives and obstetricians following obstetric emergency training: a randomised controlled trial of local hospital, simulation centre and teamwork training. *BJOG* 2007; 114:1534–41.
- 17 Crofts JF, Bartlett C, Ellis D, Winter C, Donald F, Hunt LP, et al. Patient-actor perception of care: a comparison of obstetric emergency training using manikins and patient-actors. *Qual Saf Health Care* 2008;17:20–4.
- 18 Crofts JF, Bartlett C, Ellis D, Hunt LP, Fox R, Draycott TJ. Management of shoulder dystocia: skill retention 6 and 12 months after training. *Obstet Gynecol* 2007;110:1069–74.
- 19 Crofts JF, Bartlett C, Ellis D, Hunt LP, Fox R, Draycott TJ. Training for shoulder dystocia: a trial of simulation using low-fidelity and high-fidelity mannequins. *Obstet Gynecol* 2006;108:1477–85.
- 20 Sexton JB, Helmreich RL, Neilands TB, Rowan K, Vella K, Boyden J, et al. The safety attitudes questionnaire: psychometric properties, benchmarking data, and emerging research. *BMC Health Services Res* 2006;6:44.
- 21 Draycott T, Fox R, Montague I. *Royal College of Obstetricians and Gynaecologists (RCOG). Shoulder Dystocia. Clinical Guideline No. 42*. London: RCOG Press, 2005.
- 22 Antepartum Term Stillbirths. *Confidential Enquiries into Stillbirths and Deaths in Infancy. 5th Annual Report*. London: Maternal and Child Health Consortium, 1998.
- 23 Colla JB, Bracken AC, Kinney LM, Weeks WB. Measuring patient safety climate: a review of surveys. *Qual Saf Health Care* 2005;14:364–6.
- 24 Siassakos D, Fox R, Hunt L, Farey J, Laxton C, Winter C, et al. Attitudes to safety and teamwork in a maternity unit with embedded team training. *Am J Med Qual* 2010; DOI: 10.1177/1062860610373379, in press.
- 25 Siassakos D, Fox R, Hunt L, Farey J, Laxton C, Winter C, et al. Profile of attitudes to safety, teamwork and working conditions in a maternity unit with embedded inter-professional learning. *Int J Gynecol Obstet* 2009;107 (Suppl 2):S339.
- 26 Johanson R, Menon V, Burns E, Kargramanya E, Osipov V, Israelyan M, et al. Managing Obstetric Emergencies and Trauma (MOET) structured skills training in Armenia, utilising models and reality based scenarios. *BMC Med Educ* 2002;2:5.
- 27 Freeth D, Berridge EJ, Mackintosh N. *NAMS/NPSA Evaluation of Safety Culture and MOSES Training in Four Maternity Units and Two Clinical Simulation Centres, Final Report*. London: National Patient Safety Agency and City University, 2008.
- 28 Birch L, Jones N, Doyle PM, Green P, McLaughlin A, Champney C, et al. Obstetric skills drills: evaluation of teaching methods. *Nurse Educ Today* 2007;27:915–22.
- 29 Haller G, Garnerin P, Morales MA, Pfister R, Berner M, Irion O, et al. Effect of crew resource management training in a multidisciplinary obstetrical setting. *Int J Qual Health Care* 2008;20:254–63.
- 30 Pratt S, Mann S, Salisbury M, Greenberg P, Marcus R, Stabile B, et al. Impact of CRM-based team training on obstetric outcomes and clinicians' patient safety attitudes. *J Qual Patient Saf* 2007;33:720–5.
- 31 CESDI. *Confidential Enquiries into Stillbirths and Deaths in Infancy. 7th Annual Report*. London: Maternal and Child Health Consortium, 2000.
- 32 Draycott TJ, Crofts JF, Ash JP, Wilson LV, Yard E, Sibanda T, et al. Improving neonatal outcome through practical shoulder dystocia training. *Obstet Gynecol* 2008;112:14–20.
- 33 Draycott T, Sibanda T, Owen L, Akande V, Winter C, Reading S, et al. Does training in obstetric emergencies improve neonatal outcome? *BJOG* 2006;113:177–82.
- 34 Siassakos D, Crofts J, Winter C, Weiner C, Draycott T. The active components of effective team training in obstetric emergencies. *BJOG* 2009;116:1028–32.
- 35 LeBlanc VR. The effects of acute stress on performance: implications for health professions education. *Acad Med* 2009;84:S25–33 10.1097/ACM.0b013e3181b37b8f.
- 36 Nielsen PE, Goldman MB, Mann S, Shapiro DE, Marcus RG, Pratt SD, et al. Effects of teamwork training on adverse outcomes and process of care in labor and delivery: a randomized controlled trial. *Obstet Gynecol* 2007;109:48–55.