

## High-dose oral vitamin C partially replenishes vitamin C levels in patients with Type 2 diabetes and low vitamin C levels but does not improve endothelial dysfunction or insulin resistance

Hui Chen,<sup>1</sup> Rajaram J. Karne,<sup>1</sup> Gail Hall,<sup>1</sup> Umberto Campia,<sup>2</sup> Julio A. Panza,<sup>2</sup> Richard O. Cannon III,<sup>2</sup> Yaohui Wang,<sup>3</sup> Arie Katz,<sup>1</sup> Mark Levine,<sup>3</sup> and Michael J. Quon

<sup>1</sup>Diabetes Unit, National Center for Complementary and Alternative Medicine; <sup>2</sup>Cardiology Branch, National Heart, Lung, and Blood Institute; <sup>3</sup>Molecular and Clinical Nutrition Section, Digestive Diseases Branch, National Institute of Diabetes, Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland

Submitted 19 July 2005; accepted in final form 23 August 2005

**Chen, Hui, Rajaram J. Karne, Gail Hall, Umberto Campia, Julio A. Panza, Richard O. Cannon III, Yaohui Wang, Arie Katz, Mark Levine, and Michael J. Quon.** High-dose oral vitamin C partially replenishes vitamin C levels in patients with Type 2 diabetes and low vitamin C levels but does not improve endothelial dysfunction or insulin resistance. *Am J Physiol Heart Circ Physiol* 290: H137–H145, 2006. First published August 26, 2005; doi:10.1152/ajpheart.00768.2005.—Endothelial dysfunction is a hallmark of Type 2 diabetes related to hyperglycemia and oxidative stress. Nitric oxide-dependent vasodilator actions of insulin may augment glucose disposal. Thus endothelial dysfunction may worsen insulin resistance. Intra-arterial administration of vitamin C improves endothelial dysfunction in diabetes. In the present study, we investigated effects of high-dose oral vitamin C to alter endothelial dysfunction and insulin resistance in Type 2 diabetes. Plasma vitamin C levels in 109 diabetic subjects were lower than healthy ( $36 \pm 2 \mu\text{M}$ ) levels. Thirty-two diabetic subjects with low plasma vitamin C ( $<40 \mu\text{M}$ ) were subsequently enrolled in a randomized, double-blind, placebo-controlled study of vitamin C (800 mg/day for 4 wk). Insulin sensitivity (determined by glucose clamp) and forearm blood flow in response to ACh, sodium nitroprusside (SNP), or insulin (determined by plethysmography) were assessed before and after 4 wk of treatment. In the placebo group ( $n = 17$  subjects), plasma vitamin C ( $22 \pm 3 \mu\text{M}$ ), fasting glucose ( $159 \pm 12 \text{ mg/dl}$ ), insulin ( $19 \pm 7 \mu\text{U/ml}$ ), and  $\text{SI}_{\text{Clamp}}$  [ $2.06 \pm 0.29 \times 10^{-4} \text{ dl} \cdot \text{kg}^{-1} \cdot \text{min}^{-1} / (\mu\text{U/ml})$ ] did not change significantly after placebo treatment. In the vitamin C group ( $n = 15$  subjects), basal plasma vitamin C ( $23 \pm 2 \mu\text{M}$ ) increased to  $48 \pm 6 \mu\text{M}$  ( $P < 0.01$ ) after treatment, but this was significantly less than that expected for healthy subjects ( $>80 \mu\text{M}$ ). No significant changes in fasting glucose ( $156 \pm 11 \text{ mg/dl}$ ), insulin ( $14 \pm 2 \mu\text{U/ml}$ ),  $\text{SI}_{\text{Clamp}}$  [ $2.71 \pm 0.46 \times 10^{-4} \text{ dl} \cdot \text{kg}^{-1} \cdot \text{min}^{-1} / (\mu\text{U/ml})$ ], or forearm blood flow in response to ACh, SNP, or insulin were observed after vitamin C treatment. We conclude that high-dose oral vitamin C therapy, resulting in incomplete replenishment of vitamin C levels, is ineffective at improving endothelial dysfunction and insulin resistance in Type 2 diabetes.

hypertension; hyperglycemia; hypercholesterolemia; insulin sensitivity; sodium nitroprusside

TYPE 2 DIABETES is characterized by both endothelial dysfunction and insulin resistance (14, 20). Vascular actions of insulin stimulating production of nitric oxide (NO) from endothelium (81) lead to vasodilation and increased blood flow to skeletal muscle that significantly augment insulin-mediated glucose

disposal (3–5, 7, 64). Thus endothelial dysfunction may directly contribute to insulin resistance. Conversely, insulin resistance may contribute to endothelial dysfunction because signaling pathways mediating metabolic actions of insulin in skeletal muscle and adipose tissue are strikingly similar to those regulating insulin-stimulated production of NO in vascular endothelium (34, 49–51, 78, 81, 82). Indeed, impaired vasodilator actions of insulin in obese and diabetic subjects are positively correlated with metabolic insulin resistance (40, 72).

The hyperglycemia of diabetes worsens both endothelial dysfunction and insulin resistance by multiple independent mechanisms including increased oxidative stress that accelerates the inactivation of endothelium-derived NO (8, 13, 29, 55, 80). Patients with diabetes or the metabolic syndrome have low levels of the antioxidant vitamin C (24, 67, 79). Acute intra-arterial administration of pharmacological doses of vitamin C improves endothelial dysfunction in subjects with diabetes, obesity, hypertension, hypercholesterolemia, or acute hyperglycemia (9, 54, 61, 65, 73–76). Intravenous infusion of pharmacological doses of vitamin C in diabetic subjects increases renal perfusion (21). However, in hypertensive subjects, improvement of brachial artery endothelial function with intra-arterial vitamin C is not associated with improvement in insulin-mediated glucose uptake in the forearm (54). In addition to direct antioxidant effects, vitamin C may ameliorate hyperglycemia-induced oxidative stress in the endothelium by regulating prooxidant enzyme activity (77). However, neither intra-arterial nor intravenous administration of vitamin C is a practical clinical therapy.

It is controversial whether oral vitamin C supplementation in humans is beneficial. In several epidemiological studies, an inverse correlation between vitamin C intake and mortality is observed (33, 34). Although acute oral vitamin C therapy improves vascular function in patients with coronary artery disease (42) and chronic oral vitamin C therapy improves endothelial function in children with hyperlipidemia (23), chronic supplementation with oral vitamin C is not associated with decreases in 5-yr mortality, coronary artery disease, carotid intima-media thickness, blood pressure, or diabetic hyperlipidemia (2, 12, 29a, 31, 35). In some studies, chronic oral administration of vitamin C to patients with Type 2 diabetes causes a decline in plasma free radicals that is associated with

Address for reprint requests and other correspondence: M. J. Quon, Diabetes Unit, NCCAM, NIH, Bldg. 10, Rm. 6C-205, 10 Center Dr. MSC 1632, Bethesda, MD 20892-1632 (e-mail: quonm@nih.gov).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

improved whole body glucose disposal (60), decreased blood pressure (53), and improved endothelial function (62). However, other studies (10, 19) are unable to document the beneficial effects of oral vitamin C supplementation on blood pressure, oxidative stress, and endothelial function in Type 2 diabetes. One recent report (41) suggests that supplemental vitamin C may actually increase the risk of cardiovascular disease in women with diabetes. Because there is no consensus on the beneficial effects of vitamin C supplementation in Type 2 diabetes and because endothelial dysfunction is coupled with insulin resistance, we conducted a randomized, double-blind, placebo-controlled trial of high-dose oral vitamin C (800 mg/day for 4 wk) to evaluate whether this simple therapy may be beneficial in subjects with Type 2 diabetes with respect to vascular and metabolic functions.

## METHODS

**Study subjects.** One hundred nine subjects with Type 2 diabetes between the ages of 23 and 65 yr underwent screening for inclusion in our clinical study (*study 1*). All subjects met the American Diabetes Association criteria for Type 2 diabetes (1) and had their blood drawn for measurement of plasma vitamin C levels. Subjects were excluded from further study if they were on insulin therapy or if they had severe systemic diseases, known atherosclerosis, disease predisposing to vasculitis or Raynaud's phenomenon, bleeding disorders, history of kidney stones, glucose-6-phosphate dehydrogenase deficiency, infection with HIV or hepatitis B, or were pregnant. Diabetic subjects with hypertension whose blood pressure exceeded 170/109 mmHg off of their antihypertensive medication were also excluded from further study. This cutoff of 170/109 mmHg was chosen for safety reasons. Screened subjects from *study 1* whose plasma vitamin C levels were lower than 40  $\mu$ M and whose fasting blood glucose did not exceed 300 mg/dl off of their antidiabetic medication were invited to enroll in the complete study (*study 2*). Out of 109 screened subjects, 37 subjects enrolled and 32 subjects completed all phases of *study 2*. In addition to having diabetes, 9 out of the 32 subjects who completed *study 2* had essential hypertension. The enrolled subjects were then randomized to receive daily oral doses of either vitamin C (800 mg/25 ml of an aqueous solution) or placebo (500 mg citric acid/25 ml, pH 7.0) for 4 wk in a double-blind fashion. Citric acid was used in the placebo because its taste is indistinguishable from that of ascorbic acid. Subjects were advised to avoid dietary or medicinal intake of vitamin C during the study. All antidiabetic and antihypertensive medications were stopped for 1 wk before subjects were assessed at the beginning and the end of treatment with either vitamin C or placebo. Thus subjects had their medication withdrawn for 1 wk before each glucose clamp and forearm blood flow (FBF) study. Each subject underwent FBF studies and hyperinsulinemic isoglycemic glucose clamp on the same day before and after 4 wk of treatment with either placebo or vitamin C. Informed consent was obtained from each subject. The study protocol was approved by the Institutional Review Board of the National Heart, Lung, and Blood Institute, and all procedures followed were in accordance with institutional guidelines. All studies were conducted in the Clinical Center at the National Institutes of Health.

**Plasma vitamin C measurement.** Plasma vitamin C levels were measured at the initial screening. Fasting plasma vitamin C levels were measured again immediately before and after a 4-wk administration of vitamin C or placebo. Blood was collected in plasma separator tubes with lithium heparin. Plasma was mixed with 90% methanol and 1 mM EDTA in HPLC-grade water (1:4, vol/vol) and centrifuged at 1,000  $g$  at 4°C for 10 min. The supernatant was stored at -70°C until the time of assay. For each sample, three aliquots were analyzed separately by HPLC with coulometric electrochemical detection as described previously (22, 43).

**FBF measurement.** On study days, brachial artery and antecubital vein catheters were placed in the forearm of each subject at ~7:30 AM after an overnight fast of at least 10 h. Twenty minutes after the placement of the catheters, baseline arterial and venous blood samples were obtained and FBF was measured by silastic strain-gauge venous-occlusion plethysmography, as described previously (15, 59). Endothelium-dependent and -independent vascular function were assessed by measuring FBF in response to graded intra-arterial infusions of ACh and sodium nitroprusside (SNP), respectively. Graded infusion rates were 7.5, 15, and 30  $\mu$ g/min for ACh, and 0.8, 1.6, and 3.2  $\mu$ g/min for SNP with a washout period of 20 min between the two drugs. Seven blood flow measurements conducted at 15-s intervals for each drug dose were determined, and the mean blood flow determination for each drug dose was used for further data analysis. The effect of insulin on blood flow was assessed by measuring FBF at baseline, 30, 60, 120, and 180 min after initiation of the insulin infusion during the glucose clamp procedure. Blood pressure was simultaneously determined by using a pressure transducer connected to the intra-arterial cannula. Forearm vascular resistance (FVR) was calculated by dividing mean arterial blood pressure by FBF values. Heart rate was recorded from an electrocardiographic lead. Blood flow data were measured and analyzed by using an EC6 Strain Gauge Plethysmograph with NIVP3 Arterial Inflow Software (Hokanson, Bellevue, WA).

**Hyperinsulinemic isoglycemic glucose clamp.** The glucose clamp study, conducted immediately after forearm plethysmography with ACh and SNP infusions, was completed according to methods previously described (33). An insulin solution (Humulin, Eli Lilly) was infused into a peripheral arm vein at a rate of 120  $\text{mU} \cdot \text{m}^{-2} \cdot \text{min}^{-1}$  for 3 h using a calibrated syringe pump (model A-99, Razel Industries, Stamford, CT). Potassium phosphate was infused at the same time (0.23  $\text{meq} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ) to prevent hypokalemia. Whole blood glucose concentrations were measured at the bedside every 5 min using a glucose analyzer (YSI 2700 Select, Yellow Springs Institute, Yellow Springs, OH), and an infusion of 20% dextrose was adjusted to maintain the blood glucose concentration at the fasting level. In addition, blood samples were collected at -10, -5, and 0 min (before starting insulin infusion) to obtain average baseline insulin levels. After the insulin infusion was started, blood samples were collected at 30, 60, 90, 110, 130, 150, and 180 min to measure plasma insulin concentrations (DPC Immulite 2000, Diagnostic Products, Los Angeles, CA). The steady-state period of the clamp was defined as a  $\geq 60$ -min period, where the coefficient of variation for blood glucose, plasma insulin, and glucose infusion rate was  $< 5\%$ . Mean values during the steady-state period were used to calculate  $\text{SI}_{\text{Clamp}}$ , a glucose clamp derived index of insulin sensitivity.  $\text{SI}_{\text{Clamp}}$  was defined as  $M/(G \times \Delta I)$  corrected for body weight, where  $M$  is steady-state glucose infusion rate (mg/min),  $G$  is steady-state blood glucose concentrations (mg/dl), and  $\Delta I$  is difference between basal and steady-state plasma insulin concentration ( $\mu\text{U}/\text{ml}$ ).

**Alternative indexes of insulin sensitivity.** In addition to  $\text{SI}_{\text{Clamp}}$ , we also calculated surrogate indexes, the quantitative insulin-sensitivity check index (QUICKI) (33), and the homeostasis model assessment (HOMA) (46). QUICKI was calculated as  $\text{QUICKI} = 1/[\log(I_0) + \log(G_0)]$ , where  $I_0$  is fasting insulin ( $\mu\text{U}/\text{ml}$ ) and  $G_0$  is fasting glucose (mg/dl). HOMA was calculated as  $G_0 \text{ (mmol/l)} \times I_0 \text{ (}\mu\text{U/ml)} / 22.5$ .

**Statistical analyses.** Changes in various parameters after treatment within a treatment group (placebo or vitamin C) were evaluated by using paired Student's  $t$ -tests where appropriate. Comparisons between parameters in the placebo and vitamin C group at baseline and after treatment were evaluated by using unpaired Student's  $t$ -tests. Changes in dose-response curves for FBF or FVR due to treatment with vitamin C or placebo and between placebo and vitamin C groups were analyzed by repeated-measures ANOVA. All data are means  $\pm$  SE. All  $P$  values reported represent two-tail analysis.  $P$  values of  $< 0.05$  were considered statistically significant. Study sample size was based on power calculations predicting that our sample size is ade-

Table 1. Study 1: clinical characteristics of subjects with Type 2 diabetes who were screened for inclusion in the protocol

Sex, M/W	Age, yr	Vitamin C, $\mu\text{M}$	BMI, $\text{kg}/\text{m}^2$	Random Glucose, $\text{mg}/\text{dl}$	HbA1c, %	Duration of Diabetes, yr	BP, mmHg	HR, beats/min	Cholesterol, $\text{mg}/\text{dl}$	LDL, $\text{mg}/\text{dl}$	HDL, $\text{mg}/\text{dl}$	TG, $\text{mg}/\text{dl}$
45/64	50 $\pm$ 1	36 $\pm$ 2	35 $\pm$ 1	165 $\pm$ 7	7.6 $\pm$ 0.2	5 $\pm$ 0.5	140 $\pm$ 2/77 $\pm$ 1	80 $\pm$ 1	196 $\pm$ 4	123 $\pm$ 4	49 $\pm$ 1	166 $\pm$ 12

Values are means  $\pm$  SE;  $n$  = 109 subjects. M and W, men and women, respectively; BMI, body mass index; HbA1c, hemoglobin-A1c; BP, blood pressure (values given are systolic and diastolic BP, respectively); HR, heart rate; TG, triglycerides.

quate to detect a 10% change in endothelial function and insulin sensitivity with more than 90% power.

## RESULTS

**Study subjects.** Clinical characteristics of 109 subjects with Type 2 diabetes screened for this study are reported in Table 1 (study 1). In this cohort, there were 48 Caucasians, 48 African Americans, 4 Hispanics, and 9 Asians. These subjects tended to be middle-aged and obese with elevated random glucose levels and hemoglobin-A1c (HbA1c). Notably, the average plasma vitamin C level in this diabetic cohort was strikingly low at 36  $\pm$  2  $\mu\text{M}$  [vitamin C levels in healthy individuals consuming at least the recommended dietary allowance of 75–90 mg/day are expected to be >50  $\mu\text{M}$  (43, 45)]. Thirty-seven of the 109 subjects whose plasma vitamin C levels were <40  $\mu\text{M}$  were enrolled for further study. Among these 37 subjects, 32 completed the entire study (study 2). Five subjects failed to complete the entire study due to personal choice, moving out of the area, or family emergencies. Seventeen subjects were randomized to the placebo treatment group, and 15 subjects were in the vitamin C treatment group. Baseline clinical characteristics of subjects in the placebo and vitamin C group were not statistically different (Table 2,  $P$  > 0.10 for comparison between placebo and vitamin C groups). Moreover, with the exception of vitamin C levels, the clinical characteristics of the 32 subjects who completed study 2 were similar to those of the 109 subjects who were initially screened in study 1 (Tables 1 and 2). Notably, the fasting vitamin C levels in the 32 subjects that completed study 2 were significantly lower than the already low levels observed in the 109 screened subjects from study 1 ( $P$  < 0.0001).

**Low vitamin C levels in diabetic subjects.** In the 109 diabetic subjects screened for study 1, the mean plasma vitamin C level was abnormally low (36  $\pm$  2  $\mu\text{M}$ ) when compared with levels in the general population (67, 79). When we analyzed our data for possible associations between low vitamin C levels and various clinical parameters, we did not find any significant associations. Plasma levels of vitamin C were similar between obese and nonobese (36  $\pm$  2 vs. 35  $\pm$  3  $\mu\text{M}$ ,  $P$  > 0.6), men and women (33  $\pm$  3 vs. 38  $\pm$  3  $\mu\text{M}$ ,  $P$  > 0.2), Caucasians and African Americans (35  $\pm$  3 vs. 38  $\pm$  3  $\mu\text{M}$ ,  $P$  > 0.6), subjects with and without hypertension (37  $\pm$  4 vs. 36  $\pm$  2  $\mu\text{M}$ ,  $P$  >

0.7), subjects with and without hypercholesterolemia (33  $\pm$  3 vs. 38  $\pm$  3  $\mu\text{M}$ ,  $P$  > 0.2), and subjects with >7% HbA1c compared with subjects with <7% (33  $\pm$  3 vs. 40  $\pm$  3  $\mu\text{M}$ ,  $P$  > 0.09). Furthermore, we did not observe significant correlations between plasma vitamin C levels and random blood glucose concentrations or the duration of diabetes. Likewise, we did not find significant correlations between vitamin C levels and body mass index (BMI), cholesterol, age, triglycerides, or HbA1c. Thus low vitamin C levels observed in our diabetic cohort could not be associated with or explained by any of the clinical parameters measured.

**Vitamin C levels before and after therapy with placebo or vitamin C.** In the 32 subjects who completed study 2, the baseline mean vitamin C level was exceptionally low (Table 3), but there were no significant differences between levels in subjects randomized to placebo or vitamin C treatment groups (21  $\pm$  3 vs. 23  $\pm$  2  $\mu\text{M}$ ,  $P$  > 0.8). As expected, in the placebo group, plasma vitamin C levels did not change significantly after 4-wk placebo treatment (Table 3,  $P$  > 0.7). In contrast, after subjects had taken daily oral doses of vitamin C for 4 wk, the plasma vitamin C level of subjects treated with vitamin C increased significantly from 23  $\pm$  2 to 48  $\pm$  6  $\mu\text{M}$  ( $P$  < 0.002). When changes in plasma vitamin C levels pre- and posttherapy ( $\Delta$  vitamin C) in the placebo group were compared with the changes in the vitamin C group, there was also a significant difference between  $\Delta$  vitamin C in the placebo group and the vitamin C group ( $P$  < 0.001). Interestingly, even though vitamin C levels substantially increased in our diabetic subjects with vitamin C treatment, the magnitude of this increase fell short of the full replenishment expected for healthy individuals. Indeed, healthy subjects with similarly low vitamin C levels would be expected to increase their fasting steady-state plasma vitamin C level to >80  $\mu\text{M}$  after 4 wk of vitamin C at 800 mg/day (47, 53). Taken together, these results indicate that oral supplementation with high-dose vitamin C for 4 wk only partially replenishes low vitamin C levels in subjects with Type 2 diabetes.

**Effects of placebo or vitamin C treatment on metabolic parameters.** To determine whether high-dose oral vitamin C treatment improves insulin sensitivity in diabetic subjects, we performed hyperinsulinemic isoglycemic glucose clamps on each subject before and after 4-wk administration of placebo or

Table 2. Study 2: baseline clinical characteristics of study subjects who completed full study

Treatment Group	$n$	Sex, M/W	Age, yr	Vitamin C, $\mu\text{M}$	BMI, $\text{kg}/\text{m}^2$	Fasting Glucose, $\text{mg}/\text{dl}$	Fasting, Insulin $\mu\text{U}/\text{ml}$	HbA1c, %	Duration of Diabetes, yr	BP, mmHg	HR, beats/min	Cholesterol, $\text{mg}/\text{dl}$	LDL, $\text{mg}/\text{dl}$	HDL, $\text{mg}/\text{dl}$	TG, $\text{mg}/\text{dl}$
Placebo	17	5/12	47 $\pm$ 3	22 $\pm$ 3	36 $\pm$ 2	159 $\pm$ 12	19 $\pm$ 7	8.3 $\pm$ 0.5	6.8 $\pm$ 1.2	138 $\pm$ 5/78 $\pm$ 2	77 $\pm$ 3	196 $\pm$ 9	129 $\pm$ 8	48 $\pm$ 3	163 $\pm$ 30
Vitamin C	15	8/7	49 $\pm$ 2	23 $\pm$ 2	34 $\pm$ 1	156 $\pm$ 11	14 $\pm$ 2	7.5 $\pm$ 0.3	4.3 $\pm$ 1.1	140 $\pm$ 4/77 $\pm$ 2	68 $\pm$ 5	202 $\pm$ 11	122 $\pm$ 11	45 $\pm$ 3	231 $\pm$ 44

Values are means  $\pm$  SE;  $n$ , number of subjects.



Table 3. Study 2: metabolic parameters of subjects before and after therapy with vitamin C or placebo

	Placebo (n = 17)				Vitamin C (n = 15)				P ( $\Delta$ Placebo vs. $\Delta$ Vitamin C)
	Pre-Tx	Post-Tx	$\Delta$	P	Pre-Tx	Post-Tx	$\Delta$	P	
Vitamin C, $\mu$ M	21 $\pm$ 3	19 $\pm$ 3	-1.6 $\pm$ 2.4	>0.7	23 $\pm$ 2	48 $\pm$ 6	24.9 $\pm$ 6.2	<0.002	<0.001
Fasting glucose, mg/dl	159 $\pm$ 12	147 $\pm$ 13	-12 $\pm$ 8	>0.1	156 $\pm$ 11	155 $\pm$ 9	-2 $\pm$ 11	>0.8	>0.4
Fasting insulin, $\mu$ U/ml	19 $\pm$ 6.8	16 $\pm$ 4.5	-2.5 $\pm$ 2.7	>0.3	14 $\pm$ 2.1	13 $\pm$ 1.4	-1.1 $\pm$ 1.3	>0.4	>0.6
SI <sub>Clamp</sub> , $10^{-4} \times \text{dl} \cdot \text{kg}^{-1} \cdot \text{min}^{-1} / (\mu\text{U/ml})$	2.06 $\pm$ 0.29	2.14 $\pm$ 0.31	0.09 $\pm$ 0.27	>0.7	2.71 $\pm$ 0.46	2.71 $\pm$ 0.37	0.00 $\pm$ 0.22	>0.9	>0.7
QUICKI	0.308 $\pm$ 0.008	0.315 $\pm$ 0.008	0.008 $\pm$ 0.005	>0.1	0.307 $\pm$ 0.006	0.308 $\pm$ 0.004	0.001 $\pm$ 0.005	>0.8	>0.3
HOMA	7.81 $\pm$ 3.02	5.28 $\pm$ 1.05	-2.53 $\pm$ 2.29	>0.2	5.48 $\pm$ 0.85	4.78 $\pm$ 0.49	-0.70 $\pm$ 0.73	>0.3	>0.4

Values are means  $\pm$  SE; n, number of subjects. Tx, therapy; SI<sub>Clamp</sub>, glucose clamp-derived index of insulin sensitivity; QUICKI, quantitative insulin-sensitivity check index; HOMA, homeostasis model assessment;  $\Delta$ , Post-Tx - Pre-Tx.

vitamin C. Values for fasting glucose, fasting insulin, SI<sub>Clamp</sub>, QUICKI, and HOMA before and after treatment with placebo or vitamin C are reported in Table 3. The mean glucose clamp index of insulin sensitivity (SI<sub>Clamp</sub>), QUICKI, and HOMA determined at baseline (before treatment) in both the placebo group and the vitamin C group were consistent with the insulin resistance characteristic of Type 2 diabetes (33). There was no significant change in SI<sub>Clamp</sub> when values for pre- and post-treatment were compared for either the placebo or vitamin C groups (Table 3). Moreover, when  $\Delta$ SI<sub>Clamp</sub> for placebo was compared with  $\Delta$ SI<sub>Clamp</sub> for vitamin C, there was no significant difference. Similarly, there was no significant change in fasting glucose, fasting insulin, QUICKI, or HOMA when pre- and posttreatment values were compared between groups given placebo or vitamin C. Finally, there was no significant difference between changes in the placebo and vitamin C groups for any of these metabolic parameters. These data suggest that partial replenishment of vitamin C levels achieved after high-dose oral vitamin C treatment is not sufficient to improve insulin resistance in subjects with Type 2 diabetes.

**Effects of placebo or vitamin C treatment on vascular function.** To determine the effects of placebo or vitamin C treatment on vascular function, we evaluated FBF in each subject in response to infusions of ACh, SNP, and insulin at baseline (before treatment) and after 4-wk treatment with either placebo or vitamin C. The ACh dose-response curves for the placebo group after treatment and the vitamin C group before and after treatment were similar [Fig. 1,  $P > 0.6$  by multiple ANOVA (MANOVA)] with an approximately twofold maximal increase in FBF. Despite random assignment to treatment groups, the ACh dose-response curves for the placebo and the vitamin C groups before treatment were significantly different (Fig. 1,  $P < 0.01$  by MANOVA). However, the maximal percentage changes in FBF were similar in both placebo and vitamin C groups both before and after treatment (Table 4). Thus endothelium-dependent increases in FBF in response to ACh in our diabetic subjects were unchanged by high-dose oral vitamin C treatment for 4 wk.

To assess endothelium-independent vasodilation, we measured FBF in response to SNP. As with ACh, the SNP dose-response curves for the placebo group after treatment and the vitamin C group before and after treatment were similar (Fig. 2,  $P > 0.9$  by MANOVA) with an  $\sim$ 1.6-fold maximal increase in FBF. The SNP dose-response curves for the placebo and the vitamin C groups before treatment were significantly different (Fig. 2,  $P < 0.01$  by MANOVA). Nevertheless, the maximal percentage changes in FBF in response to SNP were similar in

both the placebo and the vitamin C groups both before and after treatment (Table 4). Thus endothelial-independent increases in FBF in our diabetic subjects were unchanged by high-dose oral vitamin C treatment for 4 wk.

In healthy individuals, systemic intravenous infusion of insulin under glucose-clamp conditions results in significant twofold increases in limb blood flow (16, 71). In the present study, we observed that our diabetic subjects had very blunted vasodilator responses to insulin infusion of  $\sim$ 30% during the glucose clamp both before and after treatment with placebo or vitamin C for 4 wk (Fig. 3). Moreover, no significant differences in vasodilator responses to insulin were noted after treatment with vitamin C or placebo (Table 4). Thus our diabetic subjects had severe impairment in their endothelial response to insulin that was not improved by high-dose vitamin C treatment for 4 wk. In addition to measuring FBF, we also calculated FVR in response to ACh, SNP, and insulin in our




Fig. 1. Endothelium-dependent vasodilation in response to ACh does not significantly change after 4-wk treatment with either placebo or vitamin C. Forearm blood flow (FBF) in response to intra-arterial infusion of increasing doses of ACh was measured by strain-gauge plethysmography in study subjects before and after treatment with placebo or vitamin C for 4 wk as described in METHODS. Data are means  $\pm$  SE. ACh dose-response curves for subjects before vitamin C treatment, after vitamin C treatment, and after placebo treatment were similar [ $P > 0.6$  by multiple ANOVA (MANOVA)]. ACh dose-response curve for subjects before placebo treatment was significantly higher than that of subjects before vitamin C treatment ( $P < 0.01$  by MANOVA). Nevertheless, percent change in response to maximal ACh infusion was similar among all 4 groups ( $P > 0.6$ ).

Table 4. Study 2: maximal percent change over basal in forearm blood flow of subjects in response to ACh, SNP, and insulin before and after therapy with placebo or vitamin C

Drug	Placebo (n = 17)				Vitamin C (n = 15)				P, ( $\Delta$ Placebo vs. $\Delta$ Vitamin C)
	Pre-Tx	Post-Tx	$\Delta$	P	Pre-Tx	Post-Tx	$\Delta$	P	
ACh	271 $\pm$ 46	233 $\pm$ 56	-26 $\pm$ 57	>0.6	206 $\pm$ 31	210 $\pm$ 42	2 $\pm$ 34	>0.9	>0.6
SNP	180 $\pm$ 29	163 $\pm$ 23	-16 $\pm$ 37	>0.3	195 $\pm$ 17	163 $\pm$ 23	-34 $\pm$ 21	>0.1	>0.6
Insulin	36 $\pm$ 10	21 $\pm$ 10	-8 $\pm$ 14	>0.2	59 $\pm$ 25	31 $\pm$ 9	-29 $\pm$ 25	>0.2	0.02

Values are means  $\pm$  SE; n, number of subjects. SNP, sodium nitroprusside.

study subjects at baseline and after 4-wk administration of placebo or vitamin C. As with FBF, we did not observe significant changes in the FVR response to ACh, SNP, or insulin after subjects were treated with either placebo or vitamin C for 4 wk (data not shown). This was not surprising because mean arterial pressures for our study subjects during infusions of ACh, SNP, and insulin remained essentially constant (data not shown).

**Potential effects of variability in vitamin C levels.** In the vitamin C treatment group, there were four subjects whose plasma vitamin C levels did not change significantly after 4-wk treatment with vitamin C. This may be the result of physiological differences in these particular subjects or because these subjects failed to take their vitamin C. When queried carefully, all of these subjects claimed to have taken all of their doses of vitamin C. Reanalysis of our data with these four subjects excluded did not significantly change any of our results. Moreover, there was no significant correlation between the magnitude of increase in vitamin C levels and insulin sensitivity in the vitamin C group.

**Analysis of blood pressure as a covariate to effects of vitamin C.** To determine whether hypertension influences the effects of vitamin C on insulin sensitivity or endothelial func-

tion, we analyzed our data using blood pressure as a covariate. We did not observe any significant correlations between blood pressure (mean arterial pressure, systolic blood pressure, or diastolic blood pressure) and insulin sensitivity or endothelial function outcomes after treatment with either vitamin C or placebo (data not shown).

## DISCUSSION

Vitamin C is an antioxidant whose direct application to endothelium at pharmacological concentrations acutely improves endothelial dysfunction associated with Type 2 diabetes, acute hyperglycemia, or essential hypertension (9, 54, 75). In Type 2 diabetes, reduced serum antioxidant activity correlates with worsened glycemic control (47). Increased oxidative stress and low vitamin C levels are correlated with severity of diabetic neuropathy (83). In a recent review regarding vitamin C levels in Type 2 diabetes, Will and Byers (79) conclude that although people with diabetes may have lower vitamin C levels than normal, various methodological flaws in previous studies introduce uncertainty in vitamin C measurements. Because intra-arterial infusion of vitamin C is beneficial in diabetic subjects who may suffer from low vitamin C levels, full replenishment of vitamin C levels in diabetic subjects may improve their endothelial function. Pharmacological therapies that improve endothelial function also ameliorate insulin resis-




Fig. 2. Endothelium-independent vasodilation in response to sodium nitroprusside (SNP) does not significantly change after 4-wk treatment with either placebo or vitamin C. FBF in response to intra-arterial infusion of increasing doses of SNP before and after treatment with placebo or vitamin C for 4 wk as described in METHODS. Data are means  $\pm$  SE. SNP dose-response curves for subjects before vitamin C treatment, after vitamin C treatment, and after placebo treatment were similar ( $P > 0.9$  by MANOVA). SNP dose-response curve for subjects before placebo treatment was significantly higher than that of subjects before vitamin C treatment ( $P < 0.01$  by MANOVA). Nevertheless, percent change in response to maximal SNP infusion was similar among all 4 groups ( $P > 0.1$ ).




Fig. 3. Systemic intravenous insulin infusion during glucose clamp does not substantially increase FBF in diabetic subjects in placebo or vitamin C group before or after therapy. FBF was measured at various times over a 3-h period by strain-gauge plethysmography in study subjects undergoing a glucose clamp procedure with intravenous insulin infusion of 120  $\text{mU} \cdot \text{m}^{-2} \cdot \text{min}^{-1}$ . Data are means  $\pm$  SE. FBF after 180 min of insulin infusion was modestly higher than that at 0 min in all groups, but there was no statistical difference in time courses of subjects before or after treatment with vitamin C.

tance (36–39, 48). Therefore, we tested whether high dose oral vitamin C supplementation in Type 2 diabetes would improve endothelial function and insulin sensitivity.

Using a state-of-the-art coulometric electrochemical assay, we found that mean vitamin C levels in 109 subjects with Type 2 diabetes screened for our study were unequivocally low ( $36 \pm 2 \mu\text{M}$ ) consistent with previous studies (10, 67, 79). Vitamin C levels were not correlated with any of the other parameters we measured. Low vitamin C levels may be a consequence of diabetes because cellular uptake of vitamin C is regulated by glucose and insulin (11, 18, 69). Diabetic subjects may also have a higher turnover of vitamin C due to increased oxidative stress and oxidation of ascorbate to dehydroascorbic acid in mitochondria (66, 68). Patients with diabetic nephropathy have increased renal clearance of vitamin C (30). It would be of interest to know the oxidation status of our subjects before and after treatment with vitamin C. However, no reliable biomarker of *in vivo* oxidation has been validated for ascorbic acid or any other antioxidant (28, 57).  $\text{F}_2$ -isoprostanes are among the best potential biomarkers for lipid oxidation in humans (52). However, in diabetic subjects, ascorbic acid at higher doses than we used in our study has no effect on  $\text{F}_2$ -isoprostanes (19).

The 32 subjects who completed our intervention study had extremely low mean plasma vitamin C levels ( $22 \mu\text{M}$ ) and were chosen to maximize the likelihood that vitamin C supplementation would be beneficial. In contrast with the placebo group, we observed a substantial increase in mean plasma vitamin C levels to  $48 \mu\text{M}$  after 4-wk oral therapy (800 mg/day). Strikingly, this incomplete replenishment fell far short of the increase to  $80 \mu\text{M}$  expected for vitamin C-depleted healthy subjects on a similar dose of vitamin C (43–45). Although our subjects did not have overt nephropathy, it is possible that increased renal clearance of vitamin C may account for low vitamin C levels as well as for incomplete replenishment. Indeed, a previous study has documented low vitamin C levels in diabetic subjects despite questionnaire evidence that dietary vitamin C consumption was adequate (67). Thus low vitamin C levels in our subjects may be a consequence of diabetes itself rather than an inadequate intake of vitamin C.

We failed to find any beneficial effects of high-dose oral vitamin C therapy to improve endothelial function or insulin sensitivity in Type 2 diabetes. Our study is the first to report a lack of efficacy of oral vitamin C to improve insulin-mediated vasodilation in resistance arterioles. Our results are consistent with a previous study that did not observe significant differences in fasting glucose, HbA1c, cholesterol, or triglycerides in a double-blind crossover study with 50 diabetic patients taking vitamin C (500 mg/day) or placebo, each for 4 mo (12). Our results are discordant with another study by Paolisso et al. (60) that reported beneficial effects of oral vitamin C (1,000 mg/day for 4 mo) on glucose and lipid metabolism, and free radicals in subjects with Type 2 diabetes. Differences between that study and our study that may account for discordant outcomes include the mean age of the subjects (72 years old), as well as the longer duration and higher dose of vitamin C therapy. It is important to note that Paolisso et al. (60) conducted glucose clamps in only half of their subjects and that mean vitamin C levels after therapy increased to  $76 \mu\text{M}$ . In our study, we used an isoglycemic glucose clamp where blood glucose is main-

tained at the fasting steady state. This contrasts to euglycemic glucose clamps where blood glucose is maintained in the normal range. We chose the isoglycemic method to avoid confounding effects of acutely lowering blood glucose.

There are several potential reasons why our study was unable to document any beneficial effects of oral vitamin C therapy in Type 2 diabetes. One possibility is that our intervention duration and dose of vitamin C were inadequate. Steinberg (70) has hypothesized that antioxidant therapy may take more than 5 yr to demonstrate any benefit because the primary mechanism of these agents may be preventative. However, this reasoning may not apply to our study because direct administration of vitamin C to vascular endothelium is sufficient to acutely reverse endothelial dysfunction in diabetes (75), and other therapies that improve endothelial function improve insulin sensitivity within 1 mo (39, 48, 58). On the other hand, acute local improvement in forearm endothelial function in hypertensive subjects did not improve insulin-mediated glucose uptake in the treated forearm (54). However, mechanisms of endothelial dysfunction in hypertension and diabetes may differ.

It is unlikely that a higher dose of vitamin C would substantially increase levels of vitamin C in our subjects. The daily dose of 800 mg chosen in our study is a high and safe dose based on studies of depletion-replenishment of vitamin C in healthy subjects (44). In healthy subjects, vitamin C levels in plasma reach a maximum of  $\sim 80 \mu\text{M}$  at oral vitamin C doses of 400 mg/day and do not increase even at doses up to 2,500 mg/day (45). Thus the dose used in our study is twice that required to saturate plasma levels of vitamin C in healthy subjects and is much higher than the recommended dietary allowance (17, 45). It is also unlikely that our study is underpowered because our sample size is adequate to detect a 10% change in endothelial function and insulin sensitivity with more than 90% power. It seems more likely that a failure to achieve full replenishment of vitamin C levels accounts for our negative results. This may be due to a specific defect in vitamin C handling, because a recent study showed that oral antioxidant therapy with vitamin C and E combined was able to improve endothelial function only in Type 1 but not Type 2 diabetes (10). These results suggest that genetic contributions to insulin resistance play a role in the abnormal vitamin C handling observed in Type 2 diabetes. Another possible factor contributing to our negative results is that endothelial cells make oxygen radicals instead of eliminating them (27). Thus vitamin C studies may only work in the short term (62) because the ability to regenerate tetrahydrobiopterin may be time limited.

Improving bioavailability of NO is one mechanism that may improve endothelial function in Type 2 diabetes (56). SOD and vitamin C prevent the rapid inactivation of NO by superoxide anion (26). Interestingly, the concentration of vitamin C required to preserve NO-mediated arterial relaxation is higher than the concentration required to scavenge superoxide (32). However, high concentrations of vitamin C ( $>80 \mu\text{M}$ ) are required to preserve NO-dependent endothelial function because vitamin C only competes with NO for superoxide anion at these high concentrations (32, 65). Thus incomplete replenishment of vitamin C levels achieved in our subjects is the likely explanation for a lack of improvement in endothelial function and insulin sensitivity after high-dose oral vitamin C



therapy. Oxidized ascorbate (dehydroascorbic acid) may enter endothelial cells through glucose transporters. Thus, in diabetic subjects, dehydroascorbic acid transport may be inhibited by high glucose concentrations (25, 63), and intracellular ascorbate concentrations may have remained low despite supplementation. It remains possible that full replenishment of vitamin C levels in subjects with Type 2 diabetes may have beneficial effects to improve both endothelial dysfunction and insulin resistance. It is also possible that only high levels of vitamin C applied directly to the endothelium (not achievable by oral dosing) are sufficient to improve diabetic endothelial dysfunction. Understanding why high-dose oral vitamin C does not fully replenish patients with Type 2 diabetes is an important goal for future studies, as well as pharmacokinetic studies of vitamin C in patients with Type 2 diabetes to determine whether a renal leak may explain the incomplete replenishment of vitamin C that we observed.

## GRANTS

This research was supported by the Intramural Research Program of the National Institutes of Health, National Center for Complementary and Alternative Medicine.

## REFERENCES

1. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 27, Suppl 1: S5–S10, 2004.
2. Azen SP, Qian D, Mack WJ, Sevanian A, Selzer RH, Liu CR, Liu CH, and Hodis HN. Effect of supplementary antioxidant vitamin intake on carotid arterial wall intima-media thickness in a controlled clinical trial of cholesterol lowering. *Circulation* 94: 2369–2372, 1996.
3. Baron AD. Hemodynamic actions of insulin. *Am J Physiol Endocrinol Metab* 267: E187–E202, 1994.
4. Baron AD, Brechtel-Hook G, Johnson A, Cronin J, Leaming R, and Steinberg HO. Effect of perfusion rate on the time course of insulin-mediated skeletal muscle glucose uptake. *Am J Physiol Endocrinol Metab* 271: E1067–E1072, 1996.
5. Baron AD, Steinberg HO, Chaker H, Leaming R, Johnson A, and Brechtel G. Insulin-mediated skeletal muscle vasodilation contributes to both insulin sensitivity and responsiveness in lean humans. *J Clin Invest* 96: 786–792, 1995.
7. Baron AD, Tarshoby M, Hook G, Lazaridis EN, Cronin J, Johnson A, and Steinberg HO. Interaction between insulin sensitivity and muscle perfusion on glucose uptake in human skeletal muscle: evidence for capillary recruitment. *Diabetes* 49: 768–774, 2000.
8. Baynes JW. Role of oxidative stress in development of complications in diabetes. *Diabetes* 40: 405–412, 1991.
9. Beckman JA, Goldfine AB, Gordon MB, and Creager MA. Ascorbate restores endothelium-dependent vasodilation impaired by acute hyperglycemia in humans. *Circulation* 103: 1618–1623, 2001.
10. Beckman JA, Goldfine AB, Gordon MB, Garrett LA, Keaney JF Jr, and Creager MA. Oral antioxidant therapy improves endothelial function in Type 1 but not Type 2 diabetes mellitus. *Am J Physiol Heart Circ Physiol* 285: H2392–H2398, 2003.
11. Bigley R, Wirth M, Layman D, Riddle M, and Stankova L. Interaction between glucose and dehydroascorbate transport in human neutrophils and fibroblasts. *Diabetes* 32: 545–548, 1983.
12. Bishop N, Schorah CJ, and Wales JK. The effect of vitamin C supplementation on diabetic hyperlipidaemia: a double-blind, crossover study. *Diabet Med* 2: 121–124, 1985.
13. Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 414: 813–820, 2001.
14. Caballero AE. Endothelial dysfunction, inflammation, and insulin resistance: a focus on subjects at risk for type 2 diabetes. *Curr Diab Rep* 4: 237–246, 2004.
15. Cannon RO III, Schechter AN, Panza JA, Ognibene FP, Pease-Fye ME, Wacławski MA, Shelhamer JH, and Gladwin MT. Effects of inhaled nitric oxide on regional blood flow are consistent with intravascular nitric oxide delivery. *J Clin Invest* 108: 279–287, 2001.
16. Cardillo C, Kilcoyne CM, Nambi SS, Cannon RO III, Quon MJ, and Panza JA. Vasodilator response to systemic but not to local hyperinsulinemia in the human forearm. *Hypertension* 32: 740–745, 1998.
17. Carr AC and Frei B. Toward a new recommended dietary allowance for vitamin C based on antioxidant and health effects in humans. *Am J Clin Nutr* 69: 1086–1107, 1999.
18. Cunningham JJ. The glucose/insulin system and vitamin C: implications in insulin-dependent diabetes mellitus. *J Am Coll Nutr* 17: 105–108, 1998.
19. Darko D, Dornhorst A, Kelly FJ, Ritter JM, and Chowienzyk PJ. Lack of effect of oral vitamin C on blood pressure, oxidative stress and endothelial function in Type II diabetes. *Clin Sci (Lond)* 103: 339–344, 2002.
20. DeFronzo RA. Lilly lecture 1987. The triumvirate: beta-cell, muscle, liver. A collusion responsible for NIDDM. *Diabetes* 37: 667–687, 1988.
21. Delles C, Schneider MP, Oehmer S, Fleischmann I, Fleischmann EF, and Schmieder RE. Increased response of renal perfusion to the antioxidant vitamin C in type 2 diabetes. *Nephrol Dial Transplant* 19: 2513–2518, 2004.
22. Dhariwal KR, Hartzell WO, and Levine M. Ascorbic acid and dehydroascorbic acid measurements in human plasma and serum. *Am J Clin Nutr* 54: 712–716, 1991.
23. Engler MM, Engler MB, Malloy MJ, Chiu EY, Schloetter MC, Paul SM, Stuehlinger M, Lin KY, Cooke JP, Morrow JD, Ridker PM, Rifai N, Miller E, Witztum JL, and Mietus-Snyder M. Antioxidant vitamins C and E improve endothelial function in children with hyperlipidemia: Endothelial Assessment of Risk from Lipids in Youth (EARLY) Trial. *Circulation* 108: 1059–1063, 2003.
24. Ford ES, Mokdad AH, Giles WH, and Brown DW. The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. *Diabetes* 52: 2346–2352, 2003.
25. Goldenberg H and Schweinzer E. Transport of vitamin C in animal and human cells. *J Bioenerg Biomembr* 26: 359–367, 1994.
26. Gryglewski RJ, Palmer RM, and Moncada S. Superoxide anion is involved in the breakdown of endothelium-derived vascular relaxing factor. *Nature* 320: 454–456, 1986.
27. Guzik TJ, West NE, Black E, McDonald D, Ratnatunga C, Pillai R, and Channon KM. UltraRapid communications: vascular superoxide production by NAD(P)H Oxidase Association with endothelial dysfunction and clinical risk factors (Abstract). *Circ Res* 86: 1008, 2000.
28. Halliwell B and Whiteman M. Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? *Br J Pharmacol* 142: 231–255, 2004.
29. Hammes HP, Du X, Edelstein D, Taguchi T, Matsumura T, Ju Q, Lin J, Bierhaus A, Nawroth P, Hannak D, Neumaier M, Bergfeld R, Giordano I, and Brownlee M. Benfotiamine blocks three major pathways of hyperglycemic damage and prevents experimental diabetic retinopathy. *Nat Med* 9: 294–299, 2003.
- 29a. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of antioxidant vitamin supplementation in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 360: 23–33, 2002.
30. Hirsch IB, Atchley DH, Tsai E, Labbe RF, and Chait A. Ascorbic acid clearance in diabetic nephropathy. *J Diabetes Complications* 12: 259–263, 1998.
31. Hodis HN, Mack WJ, LaBree L, Cashin-Hemphill L, Sevanian A, Johnson R, and Azen SP. Serial coronary angiographic evidence that antioxidant vitamin intake reduces progression of coronary artery atherosclerosis. *JAMA* 273: 1849–1854, 1995.
32. Jackson TS, Xu A, Vita JA, and Keaney JF Jr. Ascorbate prevents the interaction of superoxide and nitric oxide only at very high physiological concentrations. *Circ Res* 83: 916–922, 1998.
33. Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, and Quon MJ. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 85: 2402–2410, 2000.
34. Kim JA, Koh KK, and Quon MJ. The union of vascular and metabolic actions of insulin in sickness and in health. *Arterioscler Thromb Vasc Biol* 25: 889–891, 2005.
35. Kim MK, Sasaki S, Sasazuki S, Okubo S, Hayashi M, and Tsugane S. Lack of long-term effect of vitamin C supplementation on blood pressure. *Hypertension* 40: 797–803, 2002.
36. Koh KK, Han SH, Quon MJ, Yeal Ahn J, and Shin EK. Beneficial effects of fenofibrate to improve endothelial dysfunction and raise adi-

- ponectin levels in patients with primary hypertriglyceridemia. *Diabetes Care* 28: 1419–1424, 2005.
37. Koh KK, Quon MJ, Han SH, Ahn JY, Jin DK, Kim HS, Kim DS, and Shin EK. Vascular and metabolic effects of combined therapy with ramipril and simvastatin in patients with type 2 diabetes. *Hypertension* 45: 1088–1093, 2005.
  38. Koh KK, Quon MJ, Han SH, Chung WJ, Ahn JY, Seo YH, Choi IS, and Shin EK. Additive beneficial effects of fenofibrate combined with atorvastatin in the treatment of combined hyperlipidemia. *J Am Coll Cardiol* 45: 1649–1653, 2005.
  39. Koh KK, Quon MJ, Han SH, Chung WJ, Ahn JY, Seo YH, Kang MH, Ahn TH, Choi IS, and Shin EK. Additive beneficial effects of losartan combined with simvastatin in the treatment of hypercholesterolemic, hypertensive patients. *Circulation* 110: 3687–3692, 2004.
  40. Laakso M, Edelman SV, Brechtel G, and Baron AD. Impaired insulin-mediated skeletal muscle blood flow in patients with NIDDM. *Diabetes* 41: 1076–1083, 1992.
  41. Lee DH, Folsom AR, Harnack L, Halliwell B, and Jacobs DR Jr. Does supplemental vitamin C increase cardiovascular disease risk in women with diabetes? *Am J Clin Nutr* 80: 1194–1200, 2004.
  42. Levine GN, Frei B, Koulouris SN, Gerhard MD, Keaney JF Jr, and Vita JA. Ascorbic acid reverses endothelial vasomotor dysfunction in patients with coronary artery disease. *Circulation* 93: 1107–1113, 1996.
  43. Levine M, Conry-Cantilena C, Wang Y, Welch RW, Washko PW, Dhariwal KR, Park JB, Lazarev A, Graumlich JF, King J, and Cantilena LR. Vitamin C pharmacokinetics in healthy volunteers: evidence for a recommended dietary allowance. *Proc Natl Acad Sci USA* 93: 3704–3709, 1996.
  44. Levine M, Rumsey SC, Daruwala R, Park JB, and Wang Y. Criteria and recommendations for vitamin C intake. *JAMA* 281: 1415–1423, 1999.
  45. Levine M, Wang Y, Padayatty SJ, and Morrow J. A new recommended dietary allowance of vitamin C for healthy young women. *Proc Natl Acad Sci USA* 98: 9842–9846, 2001.
  46. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, and Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28: 412–419, 1985.
  47. Maxwell SR, Thomason H, Sandler D, LeGuen C, Baxter MA, Thorpe GH, Jones AF, and Barnett AH. Poor glycaemic control is associated with reduced serum free radical scavenging (antioxidant) activity in noninsulin-dependent diabetes mellitus. *Ann Clin Biochem* 34: 638–644, 1997.
  48. McFarlane SI, Kumar A, and Sowers JR. Mechanisms by which angiotensin-converting enzyme inhibitors prevent diabetes and cardiovascular disease. *Am J Cardiol* 91: 30H–37H, 2003.
  49. Montagnani M and Quon MJ. Insulin action in vascular endothelium: potential mechanisms linking insulin resistance with hypertension. *Diabetes Obes Metab* 2: 285–292, 2000.
  50. Montagnani M, Chen H, Barr VA, and Quon MJ. Insulin-stimulated activation of eNOS is independent of  $Ca^{2+}$  but requires phosphorylation by Akt at Ser(1179). *J Biol Chem* 276: 30392–30398, 2001.
  51. Montagnani M, Ravichandran LV, Chen H, Esposito DL, and Quon MJ. Insulin receptor substrate-1 and phosphoinositide-dependent kinase-1 are required for insulin-stimulated production of nitric oxide in endothelial cells. *Mol Endocrinol* 16: 1931–1942, 2002.
  52. Morrow JD. Quantification of isoprostanes as indices of oxidant stress and the risk of atherosclerosis in humans. *Arterioscler Thromb Vasc Biol* 25: 279–286, 2005.
  53. Mullan BA, Young IS, Fee H, and McCance DR. Ascorbic acid reduces blood pressure and arterial stiffness in type 2 diabetes. *Hypertension* 40: 804–809, 2002.
  54. Natali A, Sironi AM, Toschi E, Camastra S, Sanna G, Perissinotto A, Taddei S, and Ferrannini E. Effect of vitamin C on forearm blood flow and glucose metabolism in essential hypertension. *Arterioscler Thromb Vasc Biol* 20: 2401–2406, 2000.
  55. Nishikawa T, Edelstein D, Du XL, Yamagishi S, Matsumura T, Kaneda Y, Yorek MA, Beebe D, Oates PJ, Hammes HP, Giordino I, and Brownlee M. Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature* 404: 787–790, 2000.
  56. Ogita H and Liao J. Endothelial function and oxidative stress. *Endothelium* 11: 123–132, 2004.
  57. Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O, Lee JH, Chen S, Corpe C, Dutta A, Dutta SK, and Levine M. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr* 22: 18–35, 2003.
  58. Paniagua JA, Lopez-Miranda J, Escribano A, Berral FJ, Marin C, Bravo D, Paz-Rojas E, Gomez P, Barcos M, Moreno JA, and Perez-Jimenez F. Cerivastatin improves insulin sensitivity and insulin secretion in early-state obese type 2 diabetes. *Diabetes* 51: 2596–2603, 2002.
  59. Panza JA, Casino PR, Kilcoyne CM, and Quyyumi AA. Role of endothelium-derived nitric oxide in the abnormal endothelium-dependent vascular relaxation of patients with essential hypertension. *Circulation* 87: 1468–1474, 1993.
  60. Paoilso G, Balbi V, Volpe C, Varricchio G, Gambardella A, Saccomanno F, Ammendola S, Varricchio M, and D'Onofrio F. Metabolic benefits deriving from chronic vitamin C supplementation in aged noninsulin dependent diabetics. *J Am Coll Nutr* 14: 387–392, 1995.
  61. Perticone F, Ceravolo R, Candigliota M, Ventura G, Iacopino S, Sinopoli F, and Mattioli PL. Obesity and body fat distribution induce endothelial dysfunction by oxidative stress: protective effect of vitamin C. *Diabetes* 50: 159–165, 2001.
  62. Regensteiner JG, Popylisen S, Bauer TA, Lindenfeld J, Gill E, Smith S, Oliver-Pickett CK, Reusch JE, and Weil JV. Oral L-arginine and vitamins E and C improve endothelial function in women with type 2 diabetes. *Vasc Med* 8: 169–175, 2003.
  63. Rumsey SC, Kwon O, Xu GW, Burant CF, Simpson I, and Levine M. Glucose transporter isoforms GLUT1 and GLUT3 transport dehydroascorbic acid. *J Biol Chem* 272: 18982–18989, 1997.
  64. Scherrer U, Randin U, Vollenweider P, Vollenweider L, and Nicod P. Nitric oxide release accounts for insulin's vascular effects in humans. *J Clin Invest* 94: 2511–2515, 1994.
  65. Sherman DL, Keaney JF Jr, Biegelsen ES, Duffy SJ, Coffman JD, and Vita JA. Pharmacological concentrations of ascorbic acid are required for the beneficial effect on endothelial vasomotor function in hypertension. *Hypertension* 35: 936–941, 2000.
  66. Sinclair AJ, Girling AJ, Gray L, Le Guen C, Lunec J, and Barnett AH. Disturbed handling of ascorbic acid in diabetic patients with and without microangiopathy during high dose ascorbate supplementation. *Diabetologia* 34: 171–175, 1991.
  67. Sinclair AJ, Taylor PB, Lunec J, Girling AJ, and Barnett AH. Low plasma ascorbate levels in patients with type 2 diabetes mellitus consuming adequate dietary vitamin C. *Diabet Med* 11: 893–898, 1994.
  68. Som S, Basu S, Mukherjee D, Deb S, Choudhury PR, Mukherjee S, Chatterjee SN, and Chatterjee IB. Ascorbic acid metabolism in diabetes mellitus. *Metabolism* 30: 572–577, 1981.
  69. Stankova L, Riddle M, Larned J, Burry K, Menashe D, Hart J, and Bigley R. Plasma ascorbate concentrations and blood cell dehydroascorbate transport in patients with diabetes mellitus. *Metabolism* 33: 347–353, 1984.
  70. Steinberg D. Clinical trials of antioxidants in atherosclerosis: are we doing the right thing? *Lancet* 346: 36–38, 1995.
  71. Steinberg HO, Brechtel G, Johnson A, Fineberg N, and Baron AD. Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent. A novel action of insulin to increase nitric oxide release. *J Clin Invest* 94: 1172–1179, 1994.
  72. Steinberg HO, Chaker H, Leaming R, Johnson A, Brechtel G, and Baron AD. Obesity/insulin resistance is associated with endothelial dysfunction. Implications for the syndrome of insulin resistance. *J Clin Invest* 97: 2601–2610, 1996.
  73. Taddei S, Virdis A, Ghiadoni L, Magagna A, and Salvetti A. Vitamin C improves endothelium-dependent vasodilation by restoring nitric oxide activity in essential hypertension. *Circulation* 97: 2222–2229, 1998.
  74. Timimi FK, Ting HH, Haley EA, Roddy MA, Ganz P, and Creager MA. Vitamin C improves endothelium-dependent vasodilation in patients with insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 31: 552–557, 1998.
  75. Ting HH, Timimi FK, Boles KS, Creager SJ, Ganz P, and Creager MA. Vitamin C improves endothelium-dependent vasodilation in patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 97: 22–28, 1996.
  76. Ting HH, Timimi FK, Haley EA, Roddy MA, Ganz P, and Creager MA. Vitamin C improves endothelium-dependent vasodilation in forearm resistance vessels of humans with hypercholesterolemia. *Circulation* 95: 2617–2622, 1997.



77. **Ulker S, McMaster D, McKeown PP, and Bayraktutan U.** Antioxidant vitamins C and E ameliorate hyperglycaemia-induced oxidative stress in coronary endothelial cells. *Diabetes Obes Metab* 6: 442–451, 2004.
78. **Vincent MA, Montagnani M, and Quon MJ.** Molecular and physiologic actions of insulin related to production of nitric oxide in vascular endothelium. *Curr Diab Rep* 3: 279–288, 2003.
79. **Will JC and Byers T.** Does diabetes mellitus increase the requirement for vitamin C? *Nutr Rev* 54: 193–202, 1996.
80. **Williams SB, Goldfine AB, Timimi FK, Ting HH, Roddy MA, Simonson DC, and Creager MA.** Acute hyperglycemia attenuates endothelium-dependent vasodilation in humans in vivo. *Circulation* 97: 1695–1701, 1998.
81. **Zeng G and Quon MJ.** Insulin-stimulated production of nitric oxide is inhibited by wortmannin. Direct measurement in vascular endothelial cells. *J Clin Invest* 98: 894–898, 1996.
82. **Zeng G, Nystrom FH, Ravichandran LV, Cong LN, Kirby M, Mostowski H, and Quon MJ.** Roles for insulin receptor, PI3-kinase, and Akt in insulin-signaling pathways related to production of nitric oxide in human vascular endothelial cells. *Circulation* 101: 1539–1545, 2000.
83. **Ziegler D, Sohr CG, and Nourooz-Zadeh J.** Oxidative stress and antioxidant defense in relation to the severity of diabetic polyneuropathy and cardiovascular autonomic neuropathy. *Diabetes Care* 27: 2178–2183, 2004.



Copyright of American Journal of Physiology: Heart & Circulatory Physiology is the property of American Physiological Society and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.