

Effects of Intravenous Cocaine and Cigarette Smoking on Luteinizing Hormone, Testosterone, and Prolactin in Men

JACK H. MENDELSON, MICHELLE B. SHOLAR, NICOLE H. MUTSCHLER, MARIA JASZYNA-GASIOR, NATHALIE V. GOLETIANI, ARTHUR J. SIEGEL, and NANCY K. MELLO

Alcohol and Drug Abuse Research Center, McLean Hospital/Harvard Medical School, Belmont, Massachusetts

Received April 10, 2003; accepted June 12, 2003

ABSTRACT

Cocaine and nicotine have a number of similar behavioral and neurobiological effects. This study compared the acute effects of cocaine and cigarette smoking on luteinizing hormone (LH), testosterone (T), and prolactin. Twenty-four men who met American Psychiatric Association Diagnostic and Statistical Manual criteria for cocaine abuse or nicotine dependence were given intravenous cocaine (0.4 mg/kg) or placebo-cocaine, or smoked a low or high nicotine cigarette under controlled conditions. Placebo-cocaine or low nicotine cigarette smoking did not change LH, T, or prolactin. Peak plasma levels of 254 ± 18 ng cocaine/ml and 22.6 ± 3.4 ng nicotine/ml were measured at 8 and 14 min, respectively. LH increased significantly after both i.v. cocaine and high nicotine cigarette smoking ($P < 0.01$). These LH increases were significantly correlated with increases

in cocaine and nicotine plasma levels ($P < 0.001$ – 0.003), and high nicotine cigarette smoking stimulated significantly greater increases in LH release than i.v. cocaine ($P < 0.05$). Testosterone levels did not change significantly after either cocaine or after high nicotine cigarette smoking. After i.v. cocaine, prolactin decreased significantly and remained below baseline levels throughout the sampling period ($P < 0.05$ – 0.01). After high nicotine cigarette smoking, prolactin increased to hyperprolactinemic levels within 6 min and remained significantly above baseline levels for 42 min ($P < 0.05$ – 0.03). The rapid increases in LH and reports of subjective “high” after both i.v. cocaine and high nicotine cigarette smoking illustrate the similarities between these drugs and suggest a possible contribution of LH to their abuse-related effects.

Cocaine abuse and cigarette smoking are major public health problems. Morbidity and mortality associated with cocaine abuse has been difficult to quantify (NIDA, 2002), but cigarette smoking is estimated to account for approximately 430,000 deaths each year from lung cancer, chronic obstructive pulmonary disease, and ischemic heart disease (CDC, 1993, 2001). There seem to be a number of parallels between the abuse-related effects of cocaine and nicotine. For example, recent clinical studies have shown that the positive subjective and physiological effects of i.v. nicotine are very similar to the effects of i.v. cocaine in cocaine abusers who smoke cigarettes (Jones et al., 1999). Nicotine and cocaine also have similar pharmacokinetic profiles after both inhalation and intravenous administration (Evans et al., 1996; Rose et al., 1999). The rapid distribution of both nicotine and cocaine, combined with the relatively brief duration of positive subjective effects, is characteristic of drugs with high abuse

liability (Balster and Schuster, 1973). Like cocaine abuse, cigarette smoking is very difficult to treat successfully, and relapse rates are high (Jaffe, 1990; Mendelson and Mello, 1996).

There is a general consensus that cigarette smoking, like cocaine abuse, is an addictive disorder (Jaffe, 1990; APA, 1994; CDC, 2001). Both clinical and preclinical studies have shown that i.v. nicotine is self-administered (Henningfield and Goldberg, 1983; Rose and Corrigan, 1997), and there is evidence that nicotine can induce tolerance and physical dependence (Jaffe, 1990; Henningfield et al., 1995). Nicotine, like cocaine, activates the mesolimbic dopamine system, and it has been postulated that the resulting increases in extracellular dopamine levels may mediate the abuse-related effects of both drugs (Kuhar et al., 1991; Corrigan et al., 1992; Di Chiara, 2000; Watkins et al., 2000). Interestingly, microdialysis studies suggest that administration of equipotent doses of nicotine and cocaine, given in combination, produce additive effects on nucleus accumbens dopamine release (Sziraki et al., 1999; Gerasimov et al., 2000). However, cocaine and nicotine seem to increase extracellular dopamine levels by different mechanisms. Nicotine stimulates dopamine release in the nucleus accumbens by stimulating nicotinic ace-

This research was supported in part by grants R01-DA15067, P01-DA14528, K05-DA00064, K05-DA00101, and T32-DA07252. Preliminary data were presented to the College on Problems of Drug Dependence in June 2002.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.

DOI: 10.1124/jpet.103.052928.

ABBREVIATIONS: LH, luteinizing hormone; T, testosterone; PRL, prolactin; DSM-IV, American Psychiatric Association Diagnostic and Statistical Manual-IV; VAS, visual analog scale; ANOVA, analysis of variance.

tylcholine receptors on the cell bodies of mesolimbic dopamine neurons (Watkins et al., 2000), whereas cocaine increases extracellular dopamine levels by blocking dopamine reuptake by the dopamine transporter (Kuhar et al., 1991).

In addition to dopaminergic effects, cocaine also stimulates release of some anterior pituitary, gonadal, and adrenal hormones (for review, see Mello and Mendelson, 2002). Preclinical studies suggest that these rapid hormonal changes may contribute to the abuse-related effects of cocaine under some conditions (Goeders, 1997; Mello and Mendelson, 2002). In comparison to cocaine, less is known about the acute effects of cigarette smoking on anterior pituitary and gonadal hormones, and the temporal relation of smoking-induced hormonal changes to reports of subjective "high" and "craving". Accordingly, the present study was designed to determine whether cigarette smoking and cocaine have similar or different effects on luteinizing hormone (LH), testosterone (T), and prolactin (PRL) and subjective effects measures. In recent clinical studies, cocaine (0.2 and 0.4 mg/kg i.v.) produced a rapid dose-dependent increase in LH levels (Mendelson et al., 2001). Because stimulation of LH release is a robust effect of cocaine administration in humans and in nonhuman primates (for review, see Mello and Mendelson, 2002), we compared the effects of i.v. cocaine versus placebo-cocaine and high versus low dose nicotine cigarette smoking on LH and T in men who met American Psychiatric Association Diagnostic and Statistical Manual (DSM-IV) criteria for cocaine abuse or nicotine dependence. Although gonadal steroid modulation of gonadotropin release through negative feedback mechanisms is well established (Yen et al., 1999), we are unaware of any studies of the interactions between LH and T after acute cocaine administration and cigarette smoking.

The acute effects of cocaine and cigarette smoking on prolactin release were also studied. Prolactin is under inhibitory dopaminergic control and administration of cocaine or exogenous dopamine significantly decreases prolactin in preclinical studies (Mello et al., 1994; for review, see Ben-Jonathan and Hnasko, 2001; Mello and Mendelson, 2002). To the extent that nicotine stimulates dopamine release, it might be predicted that cigarette smoking, like cocaine, would decrease prolactin levels after acute administration. Chronic smoking has been associated with low prolactin levels (Fuxe et al., 1989; CDC, 2001), but increases in prolactin levels have been reported after cigarette smoking (Wilkins et al., 1982; Seyler et al., 1986; Kirschbaum et al., 1994) and i.v. nicotine infusion (Newhouse et al., 1990). In rats, acute nicotine administration produces dose-dependent increases in prolactin (Sharp and Beyer, 1986; Fuxe et al., 1989; Hulihan-Giblin et al., 1990).

We now report that i.v. cocaine and high nicotine cigarette

smoking each stimulated significant increases in LH release that were temporally correlated with increases in plasma levels of cocaine and nicotine. Moreover, high nicotine cigarette smoking produced significantly greater stimulation of LH than i.v. cocaine. Testosterone levels did not change significantly after any treatment. Cocaine and high nicotine cigarette smoking had opposite effects on prolactin. After cocaine, prolactin decreased significantly, whereas after high nicotine cigarette smoking, rapid and sustained increases in prolactin were measured.

Materials and Methods

Subjects

Twenty-four men were recruited through newspaper advertisements and provided written informed consent for participation in this study. The study was approved by the Institutional Review Board of the McLean Hospital. Twelve men fulfilled DSM-IV criteria for a diagnosis of current cocaine abuse (305.6), and 12 men fulfilled DSM-IV criteria for current nicotine dependence (305.1). Volunteers with any lifetime DSM-IV Axis I disorder other than cocaine abuse and nicotine dependence were excluded. Men who were seeking treatment for cocaine abuse or nicotine dependence, or who were wearing a nicotine patch were excluded. All men selected for this study were in good physical health and had normal medical and laboratory screening examinations. The characteristics of subjects in the cocaine and nicotine groups are summarized in Table 1. These subjects did not differ significantly with respect to age and body mass index. As indicated in Table 1, 12 men were assigned to either cocaine or placebo-cocaine conditions, and 12 men were assigned to smoking either high nicotine cigarettes or low nicotine cigarettes.

Subjects were admitted to the clinical research ward on the morning of the study day. Subjects were asked to provide a urine specimen for drug testing, and levels of breath carbon monoxide and breath alcohol were measured. If subjects fulfilled the drug abstinence criteria described below, the study procedures were explained and any questions or concerns were discussed. After the study was completed, subjects remained on the clinical research ward for two or more hours. Lunch was provided and vital signs were measured at 30-min intervals. When subjects were medically stable and comfortable, they were discharged and transportation was provided. Subjects were paid for their participation in the study, consistent with National Institutes of Health regulations.

Drug Abstinence Requirements

It is important for subject safety, as well as to avoid confounding of the dependent variables, to ensure that subjects have not used any drugs before administration of intravenous cocaine. On the morning of each study day, urines were collected and analyzed with a Triage screen. The Triage Panel for Drugs of Abuse (Biosite Diagnostics, San Diego, CA) is a rapid multiple immunoassay system for the qualitative detection of the major metabolites of these drugs of abuse in urine at the following cut-off concentrations as recommended by the Substance Abuse and Mental Health Services Administration: phencyclidine, 25 ng/ml; benzodiazepines, 300 ng/ml; benzoyllecgo-

TABLE 1

Subject characteristics

Values are mean \pm S.E.M.

	Age	Education	BMI	Years Cocaine Use	Years Cigarette Smoking	Cigarettes Smoked Per Day
Cocaine ($n = 6$)	24.7 \pm 1.2	14.2 \pm 0.7	25.8 \pm 1.5	6.3 \pm 1.3	NA	NA
Placebo cocaine ($n = 6$)	27.0 \pm 1.3	15.0 \pm 0.6	25.2 \pm 1.5	7.2 \pm 1.5	NA	NA
Nicotine cigarettes ($n = 6$)	23.5 \pm 1.0	14.8 \pm 0.7	22.9 \pm 0.4	NA	5.0 \pm 0.5	15.7 \pm 1.3
Placebo cigarettes ($n = 6$)	23.8 \pm 0.6	15.5 \pm 0.3	24.4 \pm 0.5	NA	5.0 \pm 0.4	15.0 \pm 0.8

NA, not applicable.

nine, a metabolite of cocaine, 300 ng/ml; amphetamines, 1,000 ng/ml; tetrahydrocannabinol, 50 ng/ml; opiates, 300 ng/ml; and barbiturates, 300 ng/ml.

Smoking Abstinence Requirements. Cigarette smokers were asked to abstain from cigarettes and caffeinated beverages after midnight on the night before the study. Carbon monoxide (CO) levels were measured with a Vitalograph Breath CO Monitor (Vitalograph, Inc., Lenexa, KS) to assess compliance with smoking abstinence requirements. Cigarette smokers with a CO level above 10 ppm were not allowed to participate and were rescheduled. Cigarette smokers were also tested with the Triage screen on the study day. Cocaine abusers also were not allowed to smoke or drink caffeinated beverages before the study.

Cocaine and Nicotine Dose Selection

Cocaine. A 0.4 mg/kg i.v. dose of cocaine was selected on the basis of our previous clinical studies where it usually produced peak plasma cocaine levels over 200 ng/ml. This dose of cocaine has proved to be safe and induced significant changes in mood states and physiological responses in our previous clinical studies (Mendelson et al., 2001).

Nicotine. A commercially available, high-yield nicotine cigarette (2.3 mg) and a low nicotine cigarette were studied. The high-yield nicotine cigarette (Marlboro Red; Phillip Morris brand) contained 15.48 mg of nicotine and 16 mg of tar based on analysis by the Massachusetts Department of Public Health (MDPH, 1998). According to the Massachusetts Department of Public Health classification, cigarettes with a nicotine yield of 1.2 mg or higher are high nicotine cigarettes. In preliminary studies, smoking a Marlboro Red cigarette under the conditions described below produced peak plasma nicotine levels above 20 ng/ml. The low nicotine cigarettes contained 1.1 mg of nicotine and delivered 0.1 mg of nicotine and 2.8 mg of tar based on analyses provided by the manufacturer. Low nicotine cigarettes were acquired from Murty Pharmaceuticals Inc. (Lexington, KY).

Cocaine and Nicotine Administration Procedures. These studies were carried out on a clinical research ward. Cocaine or placebo-cocaine was administered intravenously into the antecubital vein of one arm over an interval of 1 min. Cigarettes were administered using a controlled smoking procedure designed to standardize puff volume and duration of inhalation (Griffiths et al., 1982). Every 30 s, subjects were asked to take a puff from a cigarette and hold the smoke for 5 s. At the end of 5 s, subjects were asked to exhale. Subjects took 24 puffs over a 12-min smoking period with an inter-puff interval of 25 s. This is equivalent to approximately two cigarettes. A new lighted cigarette was presented after every four puffs so that changes in cigarette length would not influence puff duration (Nemeth-Coslett and Griffiths, 1984a,b).

Subjects were studied in a semisupine position, and heart rate, blood pressure and EKGs were continuously monitored with an EKG monitor (model 78 352A; Hewlett Packard, Palo Alto, CA) for 10 min before intravenous cocaine administration or cigarette smoking and for 2 h after cocaine or cigarette administration. A physician certified in cardiopulmonary resuscitation was present during each study, and a cardiac defibrillator and appropriate emergency treatment medications were located in the study room.

Sample Collection Procedures. Baseline samples for analysis of LH, prolactin, testosterone, and plasma drug levels were collected 10 min before i.v. cocaine or placebo-cocaine injection or the onset of cigarette smoking. Samples for analysis of LH, prolactin, testosterone, and plasma cocaine or plasma nicotine were collected at 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 40, 50, 60, 80, and 120 min after completion of the cocaine injection or the onset of smoking a high nicotine cigarette. This rapid sampling frequency was based on previous observations that cocaine levels in plasma increase rapidly within 2 min after intravenous administration and reach peak levels within 4 to 5 min (Evans et al., 1996; Mendelson et al., 1999, 2001, 2002). Samples for analysis of plasma cocaine levels were collected at the same frequency after placebo-cocaine administration. Samples

were collected before and at 6, 12, 20, 30, 60, and 120 min after smoking a low nicotine placebo cigarette. All blood samples for hormone and cocaine analysis were collected from an intravenous catheter placed in the antecubital vein of the arm opposite the arm used for i.v. cocaine injection. The catheter site and collection tubes were covered during the smoking studies so that smoke in the air would not contaminate plasma samples.

Blood samples for hormone analysis and for nicotine analysis were collected in Vacutainer tubes without preservative. Blood samples for cocaine analysis were transferred to heparinized Vacutainer tubes containing sodium fluoride and acetic acid (to prevent the hydrolysis of cocaine). All samples were iced immediately, centrifuged, and plasma was removed and frozen at -70°C for cocaine analysis.

Subjective Effects Measures. Subjects were asked to rate how "high" they felt, how much they "craved" cigarettes or cocaine, and how "sick" they felt on a visual analog scale (VAS) that ranged from 0 to 100. Subjects were asked to provide ratings of high, craving, and sick before and every 2 min for the first 20 min after smoking or cocaine administration, then at less frequent intervals over the next 100 min.

Cocaine Hydrochloride Preparation. Cocaine hydrochloride was acquired from the National Institute of Drug Abuse in powder form and was dissolved in sterile water for intravenous injection by the McLean Hospital pharmacy. Sterility was ensured by passing the solution through a 0.22- μm Millipore filter and subjecting it to a limulus amoebocyte lysate test for detection of gram negative bacterial endotoxins. The test kit is manufactured by Cambrex Bio Science Inc. (Walkersville, MD). Placebo-cocaine was an equal volume vehicle control.

Assay Procedures

LH Assay. Serum LH was determined in duplicate by the ImmuChem hLH IRMA method, using kits (catalog no. 07-264-102) purchased from ICN Biomedicals, Inc. (Costa Mesa, CA). The assay sensitivity was 0.2 mIU/ml and the intra- and interassay CVs were 1.7 and 6.9%, respectively.

Prolactin Assay. Serum PRL was determined in duplicate by the ImmuChem hPRL IRMA method, using kits (catalog no. 07-274-102) purchased from ICN Biomedicals, Inc. The assay sensitivity was 0.1 ng/ml and the intra- and interassay CVs were 1.3 and 6.0%, respectively.

Testosterone Assay. Serum testosterone was determined in duplicate by the ImmuChem Testosterone IRMA method, using kits (catalog no. 07-289-102) purchased from ICN Biomedicals, Inc. The assay sensitivity was 2.8 ng/dl and the intra- and interassay CVs were 6.2 and 7.7%, respectively.

Plasma Cocaine Analysis. Plasma cocaine levels were measured in duplicate using a solid phase extraction method described by SPEC Instruction Manual by Ansys with a 5890 Series II gas chromatograph equipped with a capillary column and a 5971 Series mass selective detector (all from Hewlett Packard) (Abusada et al., 1993). The assay sensitivity was 10 ng/ml and the intra- and interassay coefficients of variation were 2.0 and 2.9%, respectively.

Plasma Nicotine Analysis. Plasma nicotine levels were measured in duplicate using a gas chromatography-mass spectrometry method described by Jacob and coworkers (Jacob et al., 2000). The nicotine assay sensitivity was 1.0 ng/ml and the intra- and interassay CVs were 4.7 and 11.5%, respectively. Nicotine assays were conducted in the laboratory of Dr. Peton Jacob III (Dept. of Medicine, University of California, San Francisco, CA).

Data Analysis

Plasma cocaine and nicotine levels, and hormone values were analyzed using a two-factor repeated measures ANOVA. If significant main effects were detected, one-way ANOVAs were performed to identify the time points that differed significantly from baseline

within each group. Comparisons between the effects of active and placebo-cocaine, and between the effects of high and low nicotine cigarettes, were also analyzed with ANOVA for repeated measures. Finally, the hormonal effects of acute cocaine administration and cigarette smoking were compared with ANOVA. The statistical significance of temporal covariance between LH, prolactin, and testosterone and plasma cocaine levels or plasma nicotine levels were determined by regression analyses. The relationship between peak ratings of sick and peak plasma prolactin levels after cigarette smoking was also evaluated with regression analyses.

Estimates of the primary kinetic parameters [i.e., time to peak plasma concentrations (T_{max}), peak plasma concentration (C_{max}), and half-life ($t_{1/2}$) of plasma nicotine, plasma cocaine, and LH] were obtained directly from a nonlinear regression-estimation software program based on the Manual of Pharmacologic Calculations with Computer Programs using PHARM/PCS version 4.2 (MicroComputer Specialist MCS, Philadelphia, PA). LH concentrations were fitted to a single dose, one-compartment model with bolus input, first order output, and elimination. Plasma concentrations were weighted by the reciprocal of the predicted concentrations. These pharmacokinetic parameters were analyzed with an ANOVA to determine whether there were any differences between the two drugs.

Results

Nicotine and Cocaine Plasma Levels (Fig. 1)

Baseline Nicotine and CO Levels. Plasma nicotine levels before and after low- and high-dose nicotine cigarette smoking are shown in Fig. 1 (top). Before cigarette smoking, low levels of nicotine that averaged 1.22 ± 0.27 and 1.33 ± 0.30 ng/ml were detected in the high and low nicotine cigarette groups, respectively. Baseline CO levels averaged 3.5 ± 1.4 ppm in the high nicotine cigarette group and 4.2 ± 1.1 ppm in the low nicotine cigarette group.

Nicotine Plasma Levels after Smoking. Nicotine plasma levels increased significantly within 2 min (or 4 puffs) ($P < 0.01$) and remained significantly above baseline throughout the 120-min sampling period. Peak plasma nicotine levels of 22.6 ± 3.4 ng/ml were detected within 14 min, and remained above 20 ng/ml between 12 and 20 min after cigarette smoking began. At the end of the 120-min sampling period, plasma nicotine levels averaged 8.22 ± 1.27 ng/ml. Plasma nicotine levels also increased significantly above presmoking baseline levels after smoking the low nicotine cigarette ($P < 0.01$). Peak plasma nicotine levels of 3.90 ± 0.77 ng/ml were detected at 12 min after low nicotine cigarette smoking began. At the end of the sampling period, plasma nicotine levels averaged 2.03 ± 0.49 ng/ml.

Cocaine Plasma Levels. Plasma cocaine levels after intravenous administration of 0.4 mg/kg cocaine are shown in Fig. 1 (bottom). Cocaine plasma levels were maximal at 8 min after i.v. cocaine administration and averaged 254 ± 18 ng/ml. Plasma cocaine levels gradually decreased over the remainder of the sampling period and averaged 58.8 ± 3.2 ng/ml at 120 min. The calculated $t_{1/2}$ of cocaine in plasma was 47.8 ± 0.5 min. No cocaine was detected in plasma after placebo-cocaine injection.

Luteinizing Hormone Levels after Cigarette Smoking and i.v. Cocaine (Fig. 2)

Baseline LH Levels in the Nicotine and Cocaine Groups. Before smoking, baseline LH levels averaged 5.14 ± 0.61 mIU/ml in the low nicotine cigarette group and $4.10 \pm$




Fig. 1. Plasma nicotine and plasma cocaine levels. Plasma drug levels (nanograms per milliliter) are shown on the left ordinates, and time (minutes) is shown on the abscissae. Points above baseline (BL) were collected 10 min before drug administration at time 0. Each data point is the average (\pm S.E.M.) of six subjects. Top, plasma nicotine levels after smoking a high-yield nicotine cigarette (filled circles) and a placebo cigarette (open circles). The 12-min cigarette smoking period is indicated by the gray rectangle. Bottom, plasma cocaine levels after administration of 0.4 mg/kg i.v. cocaine (closed circles). The vertical line indicates when cocaine was administered. Significant changes from the presmoking baseline are indicated by asterisks (*, $P < 0.05$; **, $P < 0.01$). Daggers indicate points at which plasma nicotine levels after smoking a low nicotine cigarette differed from plasma nicotine levels after smoking a high nicotine cigarette (\dagger , $P < 0.05$; \ddagger , $P < 0.01$).

0.44 mIU/ml in the high nicotine cigarette group. Baseline LH levels averaged 4.24 ± 0.64 mIU/ml in the placebo-cocaine group and 4.36 ± 0.87 mIU/ml in the active cocaine group. There were no significant differences in baseline LH levels before cocaine or placebo-cocaine administration or before low or high nicotine cigarette smoking.

Effects of Cigarette Smoking and Cocaine on LH. LH levels increased significantly within 14 min after high nicotine cigarette smoking compared with the presmoking base-




Fig. 2. Luteinizing hormone levels after cigarette smoking or i.v. cocaine administration. Plasma LH levels (milli-international units per milliliter) is shown on the abscissae. Points above baseline (BL) were collected 10 min before drug administration at time 0. Each data point is the average (\pm S.E.M.) of six subjects. Top, LH levels after smoking a high-yield nicotine cigarette (filled circles) and a placebo cigarette (open circles). The 12-min cigarette-smoking period is indicated by the gray rectangle. Bottom, LH levels after administration of 0.4 mg/kg i.v. cocaine (closed circles) or placebo-cocaine. The vertical line indicates when cocaine or placebo-cocaine was administered. Asterisks indicate points that were significantly different from baseline (*, $P < 0.05$; **, $P < 0.01$). Daggers indicate points at which LH was significantly different after high-yield nicotine cigarette smoking than after low nicotine cigarette smoking, or when LH was significantly different after active cocaine than after placebo-cocaine (\dagger , $P < 0.05$; $\dagger\dagger$, $P < 0.01$).

line ($P < 0.04$) (Fig. 2, top). LH reached peak levels of 8.15 ± 0.89 mIU/ml when plasma nicotine levels averaged 20.3 ± 3.1 ng/ml. LH remained significantly higher than baseline for 40 min (between 20 and 60 min) ($P < 0.01$ – 0.004). The increase in LH was significantly correlated with the increase in plasma nicotine levels shown in Fig. 1 ($r = 0.642$; $P = 0.003$). After low nicotine cigarette smoking, LH levels did not change significantly from baseline. LH levels were significantly higher after high nicotine cigarette smoking than

after low nicotine cigarette smoking at 18, 20, 25 and 30 min postsmoking ($P < 0.05$ or $P < 0.03$ – 0.02).

After administration of 0.4 mg/kg i.v. cocaine, LH increased significantly within 20 min compared with baseline ($P < 0.04$) (Fig. 2, bottom). LH reached peak levels of 5.8 ± 0.4 mIU/ml when plasma cocaine levels averaged 221.5 ± 16.9 ng/ml. LH remained significantly higher than baseline for 40 min (between 20 and 60 min) ($P < 0.04$). The increase in LH was significantly correlated with the increase in plasma cocaine levels shown in Fig. 1 ($r = 0.686$; $P = 0.0012$). After placebo-cocaine administration, LH levels did not change significantly from baseline. LH levels were significantly higher after cocaine than after placebo-cocaine at 18, 20, 25 and 30 min postinjection ($P < 0.05$).

Although high nicotine cigarette smoking and i.v. cocaine each produced significant increases in LH levels in men (Fig. 2), cigarette smoking produced higher levels of LH than i.v. cocaine over the same time period. LH levels after high nicotine cigarette smoking were significantly higher than LH levels after i.v. cocaine at 20 through 60 min after drug administration ($P < 0.02$ – 0.04). Pharmacokinetic analyses showed that the T_{\max} and C_{\max} for LH was significantly higher after cigarette smoking than after i.v. cocaine ($P < 0.02$ – 0.04). However, the $t_{1/2}$ of LH did not differ significantly after cigarette smoking and i.v. cocaine.

Testosterone Levels after Cigarette Smoking and i.v. Cocaine (Fig. 3)

Baseline Testosterone Levels in the Nicotine and Cocaine Groups. Baseline testosterone levels averaged 478.2 ± 38.45 ng/dl in the high nicotine cigarette group and 454.2 ± 37.3 in the low nicotine cigarette group, and these values were not significantly different. Baseline testosterone levels averaged 550.5 ± 30.1 ng/dl in the 0.4 mg/kg cocaine group and 541.17 ± 73.4 in the placebo-cocaine group, and these values were not significantly different. Also, there were no significant differences in baseline testosterone levels before active cocaine administration and high nicotine cigarette smoking.

Effects of Nicotine and Cocaine on Testosterone. After high nicotine cigarette smoking, testosterone levels tended to decrease, but these changes were not significantly different from baseline (Fig. 3). Testosterone averaged between 454 ± 28.8 and 434.7 ± 32.5 ng/ml at 20 to 60 min after smoking when LH levels were significantly higher than baseline (Fig. 2). At the end of the 120-min sampling period, testosterone averaged 460 ± 34.5 ng/ml (Fig. 3, top). There were no significant differences in testosterone between the high nicotine cigarette and the low nicotine cigarette groups.

After cocaine and placebo-cocaine administration, testosterone levels were quite variable. There were no significant changes in testosterone in comparison to baseline after either cocaine or placebo-cocaine administration. Moreover, testosterone levels did not differ significantly at any time point in the cocaine and placebo-cocaine groups.

Prolactin Levels after Cigarette Smoking and i.v. Cocaine (Fig. 4)

Baseline Prolactin Levels in the Nicotine and Cocaine Groups. Prolactin levels before low nicotine cigarette smoking averaged 12.00 ± 1.14 ng/ml, and these were not significantly different from baseline prolactin levels before




Fig. 3. Testosterone levels after cigarette smoking or i.v. cocaine administration. Plasma testosterone levels (nanograms per milliliter) are shown on the left ordinates, and time (minutes) is shown on the abscissae. Points above baseline (BL) were collected 10 min before drug administration at time 0. Each data point is the average (\pm S.E.M.) of six subjects. All other details are as described in the legend for Fig. 2.

high nicotine cigarette smoking (13.88 ± 1.1 ng/ml). Prolactin levels before placebo-cocaine administration averaged 10.82 ± 1.23 ng/ml and these were not significantly different from baseline prolactin levels before cocaine administration (10.29 ± 1.13 ng/ml). There were also no significant differences in baseline prolactin levels before high nicotine cigarette smoking and active cocaine administration.

Effects of Nicotine and Cocaine on Prolactin. Nicotine and cocaine produced opposite effects on prolactin in men (Fig. 4). After high nicotine cigarette smoking, prolactin increased rapidly to average 21.6 ± 5.21 ng/ml within 6 min after the smoking period ended. Prolactin remained significantly above baseline levels for 42 min ($P < 0.05$ – 0.03) and then gradually decreased to baseline levels within 120 min after cigarette smoking began. The increase in prolactin levels was not significantly correlated with the increase in plasma nicotine levels shown in Fig. 1 ($r = 0.433$; $P = 0.064$). Prolactin levels did not increase after low nicotine cigarette




Fig. 4. Prolactin levels after cigarette smoking or i.v. cocaine administration. Plasma prolactin levels (nanograms per milliliter) are shown on the left ordinates, and time (minutes) is shown on the abscissae. Points above baseline (BL) were collected 10 min before drug administration at time 0. Each data point is the average (\pm S.E.M.) of six subjects. All other details are as described in the legend for Fig. 2.

smoking and were significantly lower than after high nicotine cigarette smoking for 60 min (20–80 min) ($P < 0.05$).

After i.v. cocaine injection, prolactin levels decreased significantly from baseline at 40 min and remained below baseline throughout the sampling period ($P < 0.05$ – 0.01). The prolactin nadir averaged 5.65 ± 0.45 ng/ml, a 31% decrease from the precocaine baseline at 100 min postcocaine. At the end of the sampling period, prolactin levels after cocaine averaged 7.22 ± 1.67 ng/ml. The decrease in prolactin was significantly correlated with the increase in plasma cocaine levels shown in Fig. 1 ($r = 0.652$; $P = 0.003$). In contrast, after placebo-cocaine administration, prolactin levels remained relatively stable across the sampling period and ranged between 8.71 ± 1.6 and 11.31 ± 1.48 ng/ml. At the end of the sampling period after placebo-cocaine, prolactin

levels averaged 9.87 ± 1.41 ng/ml. Prolactin levels after cocaine and placebo-cocaine were not significantly different at any time point.

Subjective Effects of Cigarette Smoking and i.v. Cocaine (Fig. 5)

High Reports after Cigarette Smoking. Subjects reported feeling high immediately after smoking began, but the high intensity diminished during the 12-min smoking period (Fig. 5, top left). Subjects reported feeling high after both low and high nicotine cigarettes, but the high nicotine cigarette produced significantly greater subjective effects than the low nicotine cigarette ($P < 0.05$). High reports were not significantly correlated with plasma nicotine levels.

High Reports after Cocaine. Subjects reported feeling high immediately after i.v. cocaine administration, and high reports continued to increase for 8 min and then gradually declined (Fig. 5, lower left). High reports were significantly

correlated with plasma cocaine levels ($r = 0.812$; $P < 0.0003$). After placebo-cocaine, high reports did not differ significantly from the precocaine baseline and were significantly lower than after active cocaine.

Comparison of High Reports after Cigarette Smoking and Cocaine. The high feeling induced by cocaine was significantly greater than after smoking a high nicotine cigarette ($P < 0.05$). The high ratings after cocaine were significantly greater than after cigarette smoking between 8 and 40 min after drug administration began. Interestingly, the high ratings after smoking a low nicotine cigarette were significantly greater than high ratings measured after placebo cocaine administration.

Craving Reports after Cigarette Smoking. Baseline craving scores were elevated in both the low nicotine and high nicotine cigarette group, but these were not significantly different. When smoking began, reports of craving




Fig. 5. Reports of high and craving after high and low dose nicotine cigarette smoking and after cocaine and placebo-cocaine administration. Subjective ratings on a visual analog scale (0–100) are shown on the left ordinates, and time (minutes) is shown on the abscissae. Points above baseline (BL) were collected 10 min before drug administration at time 0. Each data point is the average (\pm S.E.M.) of five or six subjects. All other details are as described in the legend for Fig. 2.

decreased immediately in both groups, and continued to decrease during the smoking period. Craving scores remained low for 8 min after the smoking period ended and then began to increase in both groups. The low nicotine cigarette group reported greater craving than the high nicotine cigarette group until 48 min after the end of the smoking period.

Craving Reports after Cocaine. In contrast to the cigarette smokers, cocaine abusers reported no craving for cocaine during baseline. After placebo-cocaine, subjects continued to report minimal craving for cocaine throughout the sampling period. However, after active cocaine, subjects reported relatively low levels of cocaine craving for 16 min, but there was considerable variability between subjects.

Sick Reports after Cigarette Smoking. Only one subject reported feeling sick on one occasion at 2 min after smoking a low nicotine cigarette. Three subjects occasionally reported feeling sick after smoking a high nicotine cigarette. The maximal sick effect was reported at 18 min after smoking began and averaged 15.8 ± 9.7 on a VAS scale of 0 to 100. There was no significant correlation between peak reports of sick and peak levels of plasma prolactin.

Discussion

This is the first clinical study to compare the endocrine effects of i.v. cocaine and high nicotine cigarette smoking using a rapid (2-min) sampling procedure. Several similarities between the positive subjective and physiological effects of i.v. nicotine and i.v. cocaine were first reported by Jones et al. (1999), and our study confirms and extends those findings. We found that both i.v. cocaine and high nicotine cigarette smoking produced rapid increases in LH and reports of subjective high, whereas placebo-cocaine and low nicotine cigarettes had no effect on any endocrine or subjective report measures. Moreover, high nicotine cigarette smoking stimulated significantly greater increases in LH release than i.v. cocaine. In contrast, cocaine and high nicotine cigarette smoking had opposite effects on prolactin and reports of craving. The possible significance of these findings and some implications of the similarities and differences between cocaine abuse and cigarette smoking are discussed below.

Cocaine and Nicotine Plasma Levels. Intravenous cocaine and high nicotine cigarette smoking each produced rapid increases in drug plasma levels, although differences in the routes of administration produced differences in the time required to reach peak plasma levels and the half-life in plasma. Detection of peak cocaine levels within 8 min after i.v. administration is consistent with previous studies in men (Mendelson et al., 2001). Peak nicotine levels were detected 14 min after cigarette smoking began. Differences in smoking procedures limit comparisons with previous studies, but the peak plasma nicotine levels achieved are comparable with those reported by others (Benowitz et al., 1990). The low nicotine cigarette also produced measurable levels of nicotine in plasma but these had no significant behavioral or endocrine effects.

Comparison of Cocaine and Nicotine Effects on LH. Cocaine and high nicotine cigarette smoking each stimulated rapid increases in LH that were significantly correlated with increases in plasma levels of cocaine or nicotine. The time course and duration of LH increases after cocaine are consistent with previous clinical reports (Mendelson et al., 2001; for

review, see Mello and Mendelson, 2002). This is the first report that cigarette smoking, like cocaine, stimulates LH in men. Moreover, cigarette smoking produced significantly greater and more sustained increases in LH than i.v. cocaine. These findings are not consistent with previous clinical studies where no effect of cigarette smoking on LH was detected (Winternitz and Quillen, 1977; Seyler et al., 1986), and a number of procedural differences limit meaningful comparisons between those studies and the present study. However, differences in the frequency and duration of sample collection after cigarette smoking could account for these discrepant findings. For example, no changes in LH were detected when a single sample was collected after each cigarette (Winternitz and Quillen, 1977) or at 5-min intervals for 25 min after smoking (Seyler et al., 1986). In the present study, samples were collected every 2 min for 20 min after smoking began and then at 5- or 10-min intervals for 100 min. Significant increases in LH were not detected until 18 min after smoking began. Differences in LH assay sensitivity also may have contributed to detection of changes in LH in the present study; our LH assay sensitivity was 0.2 mIU/ml, but assay sensitivity was not reported in previous clinical studies (Winternitz and Quillen, 1977; Seyler et al., 1986). In rats, both stimulation and inhibition of LH after acute nicotine or cocaine administration have been reported (for review, see Fuxe et al., 1989; Mello and Mendelson, 2002).

Behavioral Correlates of Drug-Induced Increases in LH. The behavioral implications of rapid increases in LH release after i.v. cocaine and cigarette smoking are unclear. The significant correlations between increases in LH and plasma nicotine and plasma cocaine levels suggest that LH may contribute to the abuse-related effects of both drugs (for review, see Mello and Mendelson, 2002). In young men, LH increases were temporally related to behavioral and physiological measures of sexual arousal (La Ferla et al., 1978). LH is essential for normal reproductive function (Yen et al., 1999), and high LH levels during the periovulatory phase in females may also be associated with increased sexual receptivity (Mello and Mendelson, 2002).

Cocaine induced a greater increase in subjective high ratings than high nicotine cigarette smoking. This finding differs from a previous study in which the high ratings after the highest dose of i.v. nicotine (3 mg/70 kg) were greater than after i.v. cocaine (40 mg/70 kg or 0.571 mg/kg) (Jones et al., 1999). Moreover, the highest i.v. nicotine dose produced higher ratings on most of the subjective effects measures than i.v. cocaine (Jones et al., 1999). It is possible that i.v. nicotine has more potent subjective effects than smoked nicotine and is more comparable to i.v. cocaine. Drug plasma levels were not reported by Jones and coworkers, so nicotine dose levels cannot be directly compared with the present study.

The pattern of high ratings also differed after i.v. cocaine and cigarette smoking. High ratings were significantly correlated with increases in plasma cocaine levels, but not with increases in plasma nicotine levels. Rather, the peak high rating occurred within 2 min after smoking began and then decreased during the 12-min smoking period. In contrast, the cocaine-induced high continued to increase for 4-min postinjection and remained above a VAS rating of 50 for 14 min.

LH and Testosterone Interactions after Cocaine and Nicotine. The significant increases in LH after cocaine and

high nicotine cigarette smoking were not followed by changes in T levels. These findings are consistent with previous clinical reports where no changes in T were detected in men who smoked six cigarettes over 2 h (Winternitz and Quillen, 1977) or who were given intranasal cocaine (Heesch et al., 1996). However, stimulation of LH is usually followed by stimulation of T, which in turn inhibits LH release (Yen et al., 1999). For example, in rhesus monkeys, when LH was stimulated by an opioid antagonist, nalmefene, T increased significantly within 40 to 50 min after significant increases in LH (Mello et al., 2000). It may be that the magnitude of LH increases in the present study was not sufficient to stimulate T. LH increased by 80% from baseline after i.v. cocaine and by 107% after smoking a high nicotine cigarette. However, in rhesus monkey, nalmefene stimulated LH increases of 147 and 187% from baseline were followed by significant increases in T (Mello et al., 2000). Differences in LH assay procedures, as well as species differences limit these comparisons. Alternatively, the bioactivity of the LH isoform stimulated by cocaine and by high nicotine cigarette smoking might have been insufficient to alter T levels (cf. Bergendahl and Veldhuis, 2001). These data cannot directly address this possibility, because LH levels were analyzed with radioimmunoassay.

High LH levels and low T levels were also measured in dogs after a period of chronic smoking (Mittler et al., 1983). Based on analysis of liver 6- β -hydroxylase activity, these findings were interpreted to suggest that chronic cigarette smoking increased hepatic metabolism of testosterone (Mittler et al., 1983). In vitro studies indicate that both nicotine and cotinine inhibit LH-stimulated steroidogenesis in isolated mouse Leydig cells (Patterson et al., 1990).

Comparison of Cocaine and Nicotine Effects on Prolactin. Intravenous cocaine significantly decreased plasma prolactin levels within 50 to 120 min. These findings were anticipated because cocaine acts as an indirect dopamine agonist, and prolactin release is under inhibitory dopaminergic control (for review, see Ben-Jonathan and Hnasko, 2001; Mello and Mendelson, 2002). In contrast, high nicotine cigarette smoking produced significant and sustained elevations in prolactin. These findings were consistent with previous clinical reports that cigarette smoking can result in increased prolactin levels (Wilkins et al., 1982; Seyler et al., 1986; Kirschbaum et al., 1994). Intravenous infusion of nicotine (0.5 μ g/kg/min for 60 min) that yielded peak nicotine plasma levels of 13.72 ± 1.7 ng/ml, also increased prolactin in some normal men (Newhouse et al., 1990). Because prolactin is a stress labile hormone, malaise or nausea induced by smoking could increase prolactin release (Seyler et al., 1986). Although it is possible that discomfort associated with high nicotine cigarette smoking may have increased prolactin levels, three of the six subjects never reported feeling sick and peak prolactin levels were not significantly correlated with peak reports of sick. In rats, acute administration of nicotine by intravenous, intracerebroventricular, and intraperitoneal routes each consistently resulted in dose-dependent increases in prolactin (Sharp and Beyer, 1986; Hulihan-Giblin et al., 1990).

In the present study, prolactin increased to 50 to 78% above normal baseline levels, and average prolactin levels above 21 ng/ml were measured between 18 and 60 min after high nicotine cigarette smoking. In men, the normal range of prolactin is from 1 to 15 ng/ml and hyperprolactinemia is

defined as a persistent elevation of serum prolactin levels above 20 ng/ml (Yen et al., 1999). The regulation of prolactin is very complex, and abnormalities in prolactin can influence a variety of systems, including immune function and reproductive function. Chronic hyperprolactinemia can affect insulin and adrenal androgen regulation and can result in decreased bone density (Yen et al., 1999). The extent to which the repeated induction of transient hyperprolactinemia in chronic smokers may have adverse health consequences is unknown, but the incidence and severity of osteoporosis is greater in cigarette smokers (CDC, 2001).

Possible Mechanisms of Cocaine and Nicotine's Hormonal Effects. The contrasting effects of cocaine and nicotine on prolactin presumably reflect the fact that each drug increases extracellular dopamine levels by different mechanisms as noted in the Introduction (Kuhar et al., 1991; Watkins et al., 2000). Alternatively, high nicotine cigarette smoking may stimulate rapid release of prolactin by increasing endogenous opioids (for review, see Pomerleau, 1998), which in turn may inhibit dopamine release (Crowley, 1988). Yet, endogenous opioid peptides also inhibit release of hypothalamic luteinizing hormone releasing hormone (LHRH), which regulates LH release from the pituitary (Yen et al., 1999). A cigarette smoking-related increase in endogenous opioid peptides might predict a decrease, rather than an increase in LH.

A conceptual framework for reconciling concurrent stimulation of both LH and prolactin is provided by the pioneering studies of Yen et al. (1985). In clinical studies, infusion of the opioid antagonist naloxone led to an increase in release of both LH and prolactin in synchronous pulses. It was hypothesized that opioidergic control of both LH and prolactin may be mediated by LHRH. Consistent with this hypothesis, administration of LHRH to men stimulated prolactin release (Yen et al., 1985). A close temporal concordance of LH and prolactin pulsatile release in men has been consistently observed (Veldhuis and Johnson, 1988). In the present study, high nicotine cigarette smoking induced significant temporally concordant increases in both LH and prolactin ($r = 0.911$; $P < 0.0001$), and this could reflect nicotine's interactions with LHRH. Although the time course of LH increases after cocaine and cigarette smoking were similar, it is not clear whether these drugs stimulate LH release by similar mechanisms. Deconvolution analysis of cocaine's stimulation of LH suggested that the LH increase reflected a burst of LHRH (Mello and Mendelson, 2002). However, LHRH may be stimulated by norepinephrine, epinephrine, and neuropeptide Y, either stimulated or inhibited by dopamine and gonadal steroid hormones, and inhibited by endogenous opioid peptides (Yen et al., 1999). The relative contribution of these complex and inter-related systems to the acute effects of cocaine and nicotine on LH are unknown.

Acknowledgments

We are grateful to Murty Pharmaceuticals Inc. for generously supplying the denicotinized cigarettes used in this study. We thank Alicja Skupny and Gloria Cheng for excellent technical assistance in conducting the hormone and cocaine analyses. We also thank Dr. Peton Jacob III for conducting the nicotine assays.

References

- Abusada GM, Abukhalaf IK, Alford DD, Vinzon-Bautista I, Pramanik AK, Ansari NA, Manno JE, and Manno BR (1993) Solid-phase extraction and GC/MS quanti-

- tation of cocaine, ecgonine methyl ester, benzoylecgonine, and cocaethylene from meconium, whole blood and plasma. *J Anal Toxicol* **17**:353–358.
- APA (American Psychiatric Association) (1994) *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed., American Psychiatric Association, Washington, DC.
- Balster RL and Schuster CR (1973) Fixed-interval schedule of cocaine reinforcement: effect of dose and infusion duration. *J Exp Anal Behav* **20**:119–129.
- Ben-Jonathan N and Hnasko R (2001) Dopamine as a prolactin (PRL) inhibitor. *Endocrine Rev* **22**:724–763.
- Benowitz NL, Porchet H, and Jacob P, III (1990) Pharmacokinetics, metabolism and pharmacodynamics of nicotine, in *Nicotine Psychopharmacology: Molecular, Cellular and Behavioural Aspects* (Wonnacott S, Russell MAH, and Stolerman IP eds) pp 112–157, Oxford University Press, Oxford.
- Bergendahl M and Veldhuis JD (2001) Is there a physiological role for gonadotrophin oligosaccharide heterogeneity in humans? *Human Reprod* **16**:1058–1064.
- CDC (Centers for Disease Control) (1993) Cigarette smoking-attributable mortality and years of potential life lost - United States, 1990. *Morb Mortal Wkly Rep* **42**:645–649.
- CDC (Centers for Disease Control) (2001) Women and smoking: a report of the Surgeon General. *National Center for Chronic Disease Prevention and Prevention and Health Promotion*. Office on Smoking and Health, Washington, DC.
- Corrigall WA, Franklin KBI, Coen KM, and Clarke PBS (1992) The mesolimbic dopamine system is implicated in the reinforcing effects of nicotine. *Psychopharmacology* **107**:285–289.
- Crowley WF (1988) Role of endogenous opioid neuropeptides in the physiological regulation of luteinizing hormone and prolactin secretion, in *Peptide Hormones: Effects and Mechanisms of Action* (Negro-Vilar A and Conn PM eds) pp 79–116, CRC Press, Boca Raton, FL.
- Di Chiara G (2000) Role of dopamine in the behavioural actions of nicotine related to addiction. *Eur J Pharmacol* **393**:295–314.
- Evans SM, Cone EJ, and Henningfield JR (1996) Arterial and venous cocaine plasma concentrations in humans: relationship to route of administration, cardiovascular effects and subjective effects. *J Pharmacol Exp Ther* **279**:1345–1356.
- Fuxe K, Andersson K, Eneroth P, Harfstrand A, and Agnati LF (1989) Neuroendocrine actions of nicotine and of exposure to cigarette smoke: medical implications. *Psychoneuroendocrinology* **14**:19–41.
- Gerasimov MR, Franceschi M, Volkow ND, Rice O, Schiffer WK, and Dewey SL (2000) Synergistic interactions between nicotine and cocaine or methylphenidate depend on the dose of dopamine transporter inhibitor. *Synapse* **38**:432–437.
- Goeders NE (1997) A neuroendocrine role in cocaine reinforcement. *Psychoneuroendocrinology* **22**:237–259.
- Griffiths RR, Henningfield JE, and Bigelow GE (1982) Human cigarette smoking: manipulation of number of puffs per bout, interbout interval and nicotine dose. *J Pharmacol Exp Ther* **220**:256–265.
- Heesch CM, Negus BH, Bost JE, Keffer JH, Snyder RW, and Eichhorn EJ (1996) Effects of cocaine on anterior pituitary and gonadal hormones. *J Pharmacol Exp Ther* **278**:1195–1200.
- Henningfield JE and Goldberg SR (1983) Nicotine as a reinforcer in human subjects and laboratory animals. *Pharmacol Biochem Behav* **19**:989–992.
- Henningfield JE, Schuh LM, and Jarvik ME (1995) Pathophysiology of Tobacco Dependence, in *Psychopharmacology. The Fourth Generation of Progress* (Bloom FE and Kupfer DJ eds) pp 1715–1729, Raven Press, New York.
- Hulihan-Giblin BA, Lumpkin MD, and Kellar KJ (1990) Acute effects of nicotine on prolactin release in the rat: agonist and antagonist effects of a single injection of nicotine. *J Pharmacol Exp Ther* **252**:15–20.
- Jacob P, Wu S, Yu L, and Benowitz NL (2000) Simultaneous determination of mecamylamine, nicotine and cotinine in plasma by gas chromatography-mass spectrometry. *J Pharm Biomed Anal* **23**:653–661.
- Jaffe JH (1990) Tobacco smoking and nicotine dependence, in *Nicotine Psychopharmacology Molecular, Cellular and Behavioural Aspects* (Wonnacott S, Russell MAH, and Stolerman IP eds) pp 1–37, Oxford University Press, New York.
- Jones HE, Garrett BE, and Griffiths RR (1999) Subjective and physiological effects of intravenous nicotine and cocaine in cigarette smoking cocaine abusers. *J Pharmacol Exp Ther* **288**:188–197.
- Kirschbaum C, Scherer G, and Strasburger CJ (1994) Pituitary and adrenal hormone responses to pharmacological, physical and psychological stimulation in habitual smokers and nonsmokers. *Clin Invest* **72**:804–810.
- Kuhar MJ, Ritz MC, and Boja JW (1991) The dopamine hypothesis of the reinforcing properties of cocaine. *Trends Neurosci* **14**:299–302.
- La Ferla J, Anderson DL, and Schalch DS (1978) Psychoendocrine response to sexual arousal in human males. *Psychosomatic Med* **40**:166–172.
- MDPH (Massachusetts Department of Public Health) (1998) Massachusetts Department of Public Health 1997 Cigarette Nicotine Disclosure Report as required by Massachusetts General Laws Chapter 307B, *CMR 660,000*, January 16.
- Mello NK and Mendelson JH (2002) Cocaine, hormones and behavior: clinical and preclinical studies, in *Hormones, Brain and Behavior* (Pfaff DW, Arnold AP, Etgen AM, Fahrbach SE, and Rubin RT eds) pp 665–745, Academic Press, New York.
- Mello NK, Mendelson JH, Drieze JH, Teoh SK, Kelly ML, and Sholar JW (1994) Effects of dopamine on prolactin: interactions with cocaine self-administration by female rhesus monkeys. *J Pharmacol Exp Ther* **270**:1110–1120.
- Mello NK, Mendelson JH, and Kelly M (2000) Acute effects of nalmefene on LH, prolactin and testosterone in male rhesus monkeys. *Pharmacol Biochem Behav* **66**:275–284.
- Mendelson JH and Mello NK (1996) Management of cocaine abuse and dependence. *N Engl J Med* **334**:965–972.
- Mendelson JH, Mello NK, Sholar MB, Siegel AJ, Kaufman MJ, Levin JM, Renshaw PF, and Cohen BM (1999) Cocaine pharmacokinetics in men and in women during the follicular and luteal phase of the menstrual cycle. *Neuropsychopharmacology* **21**:294–303.
- Mendelson JH, Mello NK, Sholar MB, Siegel AJ, Mutschler N, and Halpern J (2002) Temporal concordance of cocaine effects on mood states and neuroendocrine hormones. *Psychoneuroendocrinology* **27**:71–82.
- Mendelson JH, Sholar M, Siegel AJ, and Mello NK (2001) Effects of cocaine on luteinizing hormone in women during the follicular and luteal phases of the menstrual cycle and in men. *J Pharmacol Exp Ther* **296**:972–979.
- Mittler JC, Pogach L, and Ertel NH (1983) Effects of chronic smoking on testosterone metabolism in dogs. *J Steroid Biochem* **18**:759–763.
- Nemeth-Coslett R and Griffiths RR (1984a) Determinants of puff duration in cigarette smokers: I. *Pharmacol Biochem Behav* **20**:965–971.
- Nemeth-Coslett R and Griffiths RR (1984b) Determinants of puff duration in cigarette smokers: II. *Pharmacol Biochem Behav* **21**:903–912.
- Newhouse PA, Sunderland T, Narang PK, Mellow AM, Fertig JB, Lawlor BA, and Murphy DL (1990) Neuroendocrine, physiologic and behavioral responses following intravenous nicotine in nonsmoking healthy volunteers and in patients with Alzheimer's Disease. *Psychoneuroendocrinology* **15**:471–484.
- NIDA (National Institute on Drug Abuse) (2002) *Epidemiologic Trends in Drug Abuse*. National Institutes of Health Publication No. 03-5109A.
- Patterson TR, Stringham JD, and Meikle AW (1990) Nicotine and cotinine inhibit steroidogenesis in mouse Leydig cells. *Life Sci* **46**:265–272.
- Pomerleau OF (1998) Endogenous opioids and smoking: a review of progress and problems. *Psychoneuroendocrinology* **23**:115–130.
- Rose JE, Behm FM, Westman EC, and Coleman RE (1999) Arterial nicotine kinetics during cigarette smoking and intravenous nicotine administration: implications for addiction. *Drug Alcohol Depend* **56**:99–107.
- Rose JE and Corrigall WA (1997) Nicotine self-administration in animals and humans: similarities and differences. *Psychopharmacology* **130**:28–40.
- Seyler LE Jr, Pomerleau OF, Fertig JB, Hunt D, and Parker K (1986) Pituitary hormone response to cigarette smoking. *Pharmacol Biochem Behav* **24**:159–162.
- Sharp BM and Beyer HS (1986) Rapid desensitization of the acute stimulatory effects of nicotine on rat plasma adrenocorticotropin and prolactin. *J Pharmacol Exp Ther* **238**:486–491.
- Sziraki I, Sershen H, Benuck M, Hashim A, and Laitha A (1999) Differences in receptor system participation between nicotine- and cocaine-induced dopamine overflow in nucleus accumbens. *NY Acad Sci* **877**:800–802.
- Veldhuis JD and Johnson ML (1988) Operating characteristics of the hypothalamo-pituitary-gonadal axis in men: circadian, ultradian and pulsatile release of prolactin and its temporal coupling with luteinizing hormone. *J Clin Endocrinol Metab* **67**:116–123.
- Watkins SS, Koob GF, and Markou A (2000) Neural mechanisms underlying nicotine addiction: acute positive reinforcement and withdrawal. *Nicotine Tob Res* **2**:19–37.
- Wilkins JN, Carlson HE, Van Vunakis H, Hill MA, Gritz E, and Jarvik ME (1982) Nicotine from cigarette smoking increases circulating levels of cortisol, growth hormone and prolactin in male chronic smokers. *Psychopharmacology* **78**:305–308.
- Winternitz WW and Quillen D (1977) Acute hormonal response to cigarette smoking. *J Clin Pharmacol* **17**:389–397.
- Yen SSC, Jaffe RB, and Barbieri RL (1999) *Reproductive Endocrinology: Physiology, Pathophysiology and Clinical Management*, 4th ed., W. B. Saunders Company, Philadelphia.
- Yen SSC, Quigley ME, Reid RL, Ropert JF, and Cetel NS (1985) Neuroendocrinology of opioid peptides and their role in the control of gonadotropin and prolactin secretion. *Am J Obstet Gynecol* **152**:485–493.

Address correspondence to: Dr. Jack H. Mendelson, Alcohol and Drug Abuse Research Center, Harvard Medical School-McLean Hospital, 115 Mill St., Belmont, MA 02478. E-mail: jmendel@mclean.harvard.edu