

Inter-sexual selection and choice

12 February 2019

Why are individuals of the resource-limited sex choosy?



- ▶ Males hold territories, but territory quality and size are unrelated to feather length¹
- ▶ Red-collard widowbird body condition is positivey correlated with tail length¹
- ▶ Male body condition declines more slowly when tails are shortened
- ▶ More females nested in the territories of males with long tails ('control')
- ▶ Females prefer long-tailed males

¹Pryke, S R, & S Andersson (2005). Biol. J. Linnean. Soc. 86:35-43.

²Image: *Euplectes ardens* Nigel Voaden. CC 2.0

Why are individuals of the resource-limited sex choosy?

- ▶ **Mate-limited:** For the sex limited by access to mates, sexual selection expected for traits increasing competitive ability



¹Pryke, S R, & S Andersson (2005). Biol. J. Linnean. Soc. 86:35-43.

²Image: *Euplectes ardens* Nigel Voaden. CC 2.0

Why are individuals of the resource-limited sex choosy?



- ▶ **Mate-limited:** For the sex limited by access to mates, sexual selection expected for traits increasing competitive ability
- ▶ **Resource-limited:** For the sex limited by resources available to rear offspring (e.g., energy investment in large gametes, parental care, etc.), sexual selection expected for choosiness to increase offspring quality

¹Pryke, S R, & S Andersson (2005). Biol. J. Linnean. Soc. 86:35-43.

²Image: *Euplectes ardens* Nigel Voaden. CC 2.0

Why are individuals of the resource-limited sex choosy?



- ▶ **Mate-limited:** For the sex limited by access to mates, sexual selection expected for traits increasing competitive ability
- ▶ **Resource-limited:** For the sex limited by resources available to rear offspring (e.g., energy investment in large gametes, parental care, etc.), sexual selection expected for choosiness to increase offspring quality
- ▶ In most species, females are resource-limited, & males are mate-limited

¹Pryke, S R, & S Andersson (2005). Biol. J. Linnean. Soc. 86:35-43.

²Image: *Euplectes ardens* Nigel Voaden. CC 2.0

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else
- ▶ **Good genes hypothesis:** Choose mates with high quality genes for viability to increase the fitness of offspring

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else
- ▶ **Good genes hypothesis:** Choose mates with high quality genes for viability to increase the fitness of offspring
- ▶ **Runaway sexual selection:** Assortative mating between preference and preferred trait (e.g., females choose attractive males to produce attractive male offspring)

Direct benefits: resource provisioning




Figure 1: Copulating hangingflies

¹**Image:** Byers, G W. (2002). Kansas School Naturalist. 48:0022-877X

Direct benefits: resource provisioning

- ▶ Thornhill¹ studied *Bittacus apicalis* in Michigan, USA
- ▶ Males catch prey, then use pheromones to attract females
- ▶ If females accept prey, copulation ensues while female eats
- ▶ Female gets food from the male, and does not need to risk predation hunting for herself

¹Thornhill, R (1976). Am. Nat. 110:529-548

Direct benefits: resource provisioning




FIG. 3.—The relationship between the duration of copulation and nuptial prey size and palatability in *Bittacus apicalis* in nature. The numbers above the dots indicate more than one observation for a prey size. $N = 118$.

¹Thornhill, R (1976). Am. Nat. 110:529-548

Direct benefits: resource provisioning




FIG. 4.—The number of sperm transferred as a function of the duration of copulation in *Bittacus apicalis*. $N = 71$.

¹Thornhill, R (1976). Am. Nat. 110:529-548

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else
- ▶ **Good genes hypothesis:** Choose mates with high quality genes for viability to increase the fitness of offspring
- ▶ **Runaway sexual selection:** Assortative mating between preference and preferred trait (e.g., females choose attractive males to produce attractive male offspring)

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ Proctor¹ studied the water mite *Neumania papillator* in Ontario, Canada

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ Proctor¹ studied the water mite *Neumania papillator* in Ontario, Canada
- ▶ Mites live around aquatic plants, and are ambush predators on copepods

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ Proctor¹ studied the water mite *Neumania papillator* in Ontario, Canada
- ▶ Mites live around aquatic plants, and are ambush predators on copepods
- ▶ Very limited eyesight – use smell and touch to hunt
 - ▶ Stands on four hind legs, spreads out four front legs
 - ▶ Waits for water vibrations, then ambushes the source

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ Proctor¹ studied the water mite *Neumania papillator* in Ontario, Canada
- ▶ Mites live around aquatic plants, and are ambush predators on copepods
- ▶ Very limited eyesight – use smell and touch to hunt
 - ▶ Stands on four hind legs, spreads out four front legs
 - ▶ Waits for water vibrations, then ambushes the source
- ▶ Males attract females by trembling their legs, then transfer spermatophores

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ Proctor¹ studied the water mite *Neumania papillator* in Ontario, Canada
- ▶ Mites live around aquatic plants, and are ambush predators on copepods
- ▶ Very limited eyesight – use smell and touch to hunt
 - ▶ Stands on four hind legs, spreads out four front legs
 - ▶ Waits for water vibrations, then ambushes the source
- ▶ Males attract females by trembling their legs, then transfer spermatophores
- ▶ **Proctor hypothesised that female choice for trembling behaviour evolved because it mimics vibrations of prey**

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait




Figure 1. Courtship in *N. papillator*: (a) the male vibrates his foreleg in front of the female, which is in net-stance; (b) the female has turned slightly in response to the trembling, and the male fans his fourth legs over the spermatophores he has just deposited. Sausage-shaped objects on top of spermatophore stalks are sperm packets.

¹Proctor, H. (1991). Anim. Behav. 42:589-598

Pleiotropy: Choosiness as a side-effect of a trait

Key results

- ▶ Male water mites tremble at a frequency that is within the frequency range of copepod prey

Pleiotropy: Choosiness as a side-effect of a trait

Key results

- ▶ Male water mites tremble at a frequency that is within the frequency range of copepod prey
- ▶ Female behaviour for clutching spermatophore similar to that of ambushing prey

Pleiotropy: Choosiness as a side-effect of a trait

Key results

- ▶ Male water mites tremble at a frequency that is within the frequency range of copepod prey
- ▶ Female behaviour for clutching spermatophore similar to that of ambushing prey
- ▶ Less well fed females responded to males more readily than well fed females

Pleiotropy: Choosiness as a side-effect of a trait

Key results

- ▶ Male water mites tremble at a frequency that is within the frequency range of copepod prey
- ▶ Female behaviour for clutching spermatophore similar to that of ambushing prey
- ▶ Less well fed females responded to males more readily than well fed females
- ▶ Males tremble to determine female direction, and deposit spermatophores in front of them more often than expected by chance

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ **But did female predatory behaviour evolve before male mating behaviour?**

¹Proctor, H. (1992). Anim. Behav. 44:745-752

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ **But did female predatory behaviour evolve before male mating behaviour?**
- ▶ Proctor¹ investigated mite phylogeny to see which behaviour evolved first

¹Proctor, H. (1992). Anim. Behav. 44:745-752

Pleiotropy: Choosiness as a side-effect of a trait

- ▶ But did female predatory behaviour evolve before male mating behaviour?
- ▶ Proctor¹ investigated mite phylogeny to see which behaviour evolved first
- ▶ Two evolutionary histories appear most likely (left)
 - ▶ Insufficient evidence for predatory behaviour evolving first (top)
 - ▶ Predatory behaviour evolved first, male mating behaviour independently evolved twice (bottom)




Figure 4. Behavioural character states overlaid on the cladogram, showing two equally parsimonious evolutionary scenarios (T = male courtship trembling; N = net-snare). Presence of the behaviour in a species is indicated by a filled box above the species abbreviation and absence by an empty box. Blank lines indicate unknown character states.

¹Proctor, H. (1992). Anim. Behav. 44:745-752

Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else
- ▶ **Good genes hypothesis:** Choose mates with high quality genes for viability to increase the fitness of offspring^{1,2}
- ▶ **Runaway sexual selection:** Assortative mating between preference and preferred trait (e.g., females choose attractive males to produce attractive male offspring)

¹Zahavi, A. (1975). J. Theor. Biol. 53:205-214

²Grafen, A. (1990). J. Theor. Biol. 144:517–546.

Good genes: Choosiness to increase offspring fitness

- ▶ Individuals of the mate-limited sex (usually males) are chosen by members of the resource limited sex (usually females) based on traits that are indicators of high fitness
- ▶ Chosen traits are often elaborate ornaments or behaviours



¹Zahavi, A. (1975). J. Theor. Biol. 53:205-214

²Grafen, A. (1990). J. Theor. Biol. 144:517–546.

³Image: *Fregata magnificens* Julian Hammer. CC 3.0

Good genes: Female choosiness in grey tree frogs




- ▶ Grey tree frogs (*Hyla chrysoscelis* & *H. versicolor*) endemic to North American woodlands
- ▶ During breeding season, males make pulsed calls to females ('trills')
- ▶ Males vary in **call rate (CR)** & **pulse number (PN)**
- ▶ Gerhardt et al.¹ hypothesised that females prefer higher CR & PR

¹Gerhardt, H C, et al. (1996). Behav. Ecol. 7:7-18

²Image: *Hyla versicolor* Trisha Shears. CC 3.0

Good genes: Female choosiness in grey tree frogs



¹Gerhardt, H C, et al. (1996). Behav. Ecol. 7:7-18

Good genes: Female choosiness in grey tree frogs




Figure 5

Percentage of female choices (one per female; HV = *H. versicolor*, HC = *H. chrysoscelis*) for high-PN (H-PN) or fast-CR (F-CR) calls over low-PN (L-PN) or slow-CR (S-CR) calls. In Experiments 1 and 2, a difference in SPL of 6 dB at the release point in favor of low-PN or slow-CR calls relative to high-PN and fast-CR calls was created by differential amplification; females were released midway between the speakers as in Figure 2b. In Experiments 3 and 4, the 6-dB difference was created by distance as shown in Figure 2c. Error bars represent 95% confidence limits on the proportions of females that responded to the preferred stimulus; raw data are also presented in each bar as the number of females choosing the preferred stimulus divided by the number of responding females. *p*-values for two-tailed binomial tests of the null hypothesis of no preference: Experiment 1—HC (<.001); HV (.099); Experiment 2—HC (.016); HV (.07); Experiment 3—HV (<.01); Experiment 4—HV (.12).


Gerhardt et al.'s¹ hypothesis supported; in experiments, females prefer:

- ▶ Higher call rates (CR)
- ▶ Higher pulse number (PN)

Preferences significant despite higher CR & PN speakers played more softly, or placed farther away from choosing female

¹Gerhardt, H C, et al. (1996). Behav. Ecol. 7:7-18

Good genes: Female choosiness in grey tree frogs



Further, when a loudspeaker playing a call with a relatively high pulse number (PN) was placed in the same direction but farther away from a loudspeaker with a short PN call, females tended to bypass the latter entirely and go to the farther high PN call.

¹Gerhardt, H C, et al. (1996). Behav. Ecol. 7:7-18

Good genes: Female choosiness in grey tree frogs




Figure 5. Phonotaxis scores of *H. versicolor* females to song models varying in pulse number. Each point represents the mean (\pm SE) of 10 females. A logarithmic curve is fitted to the data. The standard song model used in the control trials had 18 pulses/call.

- ▶ Bush et al.¹ further quantified *H. versicolor* response to PN.
- ▶ Calculated a phonotaxis score: ratio of time to reach loudspeaker during control trial to test trial ($t_{control}/t_{test}$)
- ▶ Control trial set to be as attractive as typically observed calls

¹Bush, S L et al. (2002). Anim. Behav. 63:7-14

Good genes: Female choosiness in grey tree frogs




Figure 5. Phonotaxis scores of *H. versicolor* females to song models varying in pulse number. Each point represents the mean (\pm SE) of 10 females. A logarithmic curve is fitted to the data. The standard song model used in the control trials had 18 pulses/call.

- ▶ Bush et al.¹ further quantified *H. versicolor* response to PN.
- ▶ Calculated a phonotaxis score: ratio of time to reach loudspeaker during control trial to test trial ($t_{control}/t_{test}$)
- ▶ Control trial set to be as attractive as typically observed calls
- ▶ **Do males with higher call rates and pulse numbers have better genes?**

¹Bush, S L et al. (2002). Anim. Behav. 63:7-14

Good genes: Female choosiness in grey tree frogs



- ▶ Welch et al.¹ collected unfertilised eggs from wild female tree frogs
- ▶ Fertilised some of a female's eggs with sperm of males with high pulse number, and some with sperm of males with low pulse number
- ▶ Measured fitness related traits of offspring from each type of male on a low and high food diet

¹Welch, A M, et al. (1998). Science 280:1928-1930

Tree frog table

Welch et al.¹ found that offspring of males with higher pulse numbers had higher fitness, or no difference in fitness, across all measures of fitness

Fitness measure	High food diet	Low food diet
Larval growth	No difference	Long-call better
Time to metamorphosis	Long-call better	No difference
Mass at metamorphosis	No difference	Long-call better
Larval survival	Long-call better	No difference

¹ Welch, A M, et al. (1998). Science 280:1928-1930

Tree frog table

Welch et al.¹ found that offspring of males with higher pulse numbers had higher fitness, or no difference in fitness, across all measures of fitness

Fitness measure	High food diet	Low food diet
Larval growth	No difference	Long-call better
Time to metamorphosis	Long-call better	No difference
Mass at metamorphosis	No difference	Long-call better
Larval survival	Long-call better	No difference

If the choosing sex is always selecting for ornamentation, why is there still variation in fitness?

¹ Welch, A M, et al. (1998). Science 280:1928-1930

The lek paradox, and one potential resolution

The lek paradox: Given many generations of sustained selection for ornamentation (i.e., choice), genetic variation for ornamentation should erode, followed by variation for choosiness¹

¹Borgia, G. (1979). Sexual selection and the evolution of mating systems. In Sexual selection and reproductive competition in insects (eds M. S. Blum & N. A. Blum), pp. 19–80. New York, NY: Academic Press.

²Hamilton, W D & Zuk, M (1982). Science 281:384–387

The lek paradox, and one potential resolution

The lek paradox: Given many generations of sustained selection for ornamentation (i.e., choice), genetic variation for ornamentation should erode, followed by variation for choosiness¹

Marlene Zuk and Bill Hamilton² suggest that female choice for ornamentation is related to signalling for parasite resistance by males

- ▶ Males that are genetically resistant to common parasites will have *good genes*
- ▶ Males that are not genetically resistant will have *bad genes*
- ▶ Selection should favour females who choose males with good genes for passing parasite resistance to offspring

¹Borgia, G. (1979). Sexual selection and the evolution of mating systems. In Sexual selection and reproductive competition in insects (eds M. S. Blum & N. A. Blum), pp. 19–80. New York, NY: Academic Press.

²Hamilton, W D & Zuk, M (1982). Science 281:384–387

The lek paradox, and one potential resolution

Bright ornaments (e.g., bird plumage), might serve as a reliable indicator of good health due to parasite resistance.

Males ornamentation	Susceptible to parasites	Parasite resistant
None	Appears healthy	Appears healthy
Costly ornament	Appears unhealthy	Appears healthy

Host-parasite coevolution will cause alleles underlying parasite resistance to change over time, maintaining genetic variation²

¹Borgia, G. (1979). Sexual selection and the evolution of mating systems. In Sexual selection and reproductive competition in insects (eds M. S. Blum & N. A. Blum), pp. 19–80. New York, NY: Academic Press.

²Hamilton, W D & Zuk, M (1982). Science 281:384-387


Why are individuals of the resource-limited sex choosy?

Direct benefits: Food, parental care, or some other resource is provided

Indirect benefits: Better genes transmitted to offspring

- ▶ **Pleiotropy:** Choosiness a side-effect of a trait that evolved by natural selection for something else
- ▶ **Good genes hypothesis:** Choose mates with high quality genes for viability to increase the fitness of offspring
- ▶ **Runaway sexual selection:** Assortative mating between preference and preferred trait (e.g., females choose attractive males to produce attractive male offspring)

Runaway sexual selection: a positive feedback loop



- ▶ The ‘runaway’ of selection is a positive feedback loop between ornamentation and preference
- ▶ Feedback loop generates linkage disequilibrium between alleles for ornamentation and preference for ornamentation

Runaway sexual selection: stalk-eyed flies




- ▶ Multiple species across the globe
- ▶ Roost at dawn and dusk on root hairs, and mate
- ▶ Males have longer eye stalks than females
- ▶ Some evidence male-male combat selects for longer eye stalks¹

¹Panhuis, T M & Wilkinson, G S. (1999). Behav. Ecol. Sociobiol. 46:221-227


²Image: Diopsidae Guido Bohne. CC 2.0

Runaway sexual selection: stalk-eyed flies


(a) Variation in eyestalks and preferences should lead to assortative mating:



(b) Assortative mating should produce genetic correlations between sons and daughters within families:




(c) Selection on male eyestalks should produce a response in female preference:



¹Freeman, S., & Herron, J. C. (2007). Evolutionary analysis. Upper Saddle River, NJ: Pearson Prentice Hall.


Runaway sexual selection: stalk-eyed flies

- ▶ Wilkinson & Reillo¹ tested whether or not selection on male eye stalk length leads to a correlated response in female preference



¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6


Runaway sexual selection: stalk-eyed flies



- ▶ Wilkinson & Reillo¹ tested whether or not selection on male eye stalk length leads to a correlated response in female preference
- ▶ Established 3 laboratory populations
 - ▶ Control line with females and males picked at random
 - ▶ Long-selected line with females picked at random, but males with longest eye stalks
 - ▶ Short-selected line with females picked at random, but males with shortest eye stalks

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

Runaway sexual selection: stalk-eyed flies



- ▶ Wilkinson & Reillo¹ tested whether or not selection on male eye stalk length leads to a correlated response in female preference
- ▶ Established 3 laboratory populations
 - ▶ Control line with females and males picked at random
 - ▶ Long-selected line with females picked at random, but males with longest eye stalks
 - ▶ Short-selected line with females picked at random, but males with shortest eye stalks
- ▶ After 13 generations, male eye stalk length evolved

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

Runaway sexual selection: stalk-eyed flies

- ▶ Females from each line placed in a cage between a long-stalk and short-stalk¹ (separated by a clear barrier with a hole small enough for females to pass)

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

Runaway sexual selection: stalk-eyed flies

- ▶ Females from each line placed in a cage between a long-stalk and short-stalk¹ (separated by a clear barrier with a hole small enough for females to pass)
- ▶ Observed preferences of females from each line of male eye stalk selection




Figure 4. Difference between the number of females (mean \pm s.e.) roosting either with an L or s male in the four mate choice experiments referred to in the text. Unfilled bar indicates experiment 1 where males could interact. Filled bars indicate, from left to right, experiments 2–4 in which a transparent perforated partition separated L and s males in each cage.

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

Runaway sexual selection: stalk-eyed flies

- ▶ Female stalk-eyed flies are choosy

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

²Freeman, S., & Herron, J C. (2007). Evolutionary analysis. Upper Saddle River, NJ: Pearson Prentice Hall.

Runaway sexual selection: stalk-eyed flies

- ▶ Female stalk-eyed flies are choosy
- ▶ Both male eye stalk length and *female preference* for eye stalk length are heritable

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

²Freeman, S., & Herron, J C. (2007). Evolutionary analysis. Upper Saddle River, NJ: Pearson Prentice Hall.

Runaway sexual selection: stalk-eyed flies

- ▶ Female stalk-eyed flies are choosy
- ▶ Both male eye stalk length and *female preference* for eye stalk length are heritable
- ▶ Sexual selection on the trait of one sex can cause an evolutionary response in the trait of the other sex

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

²Freeman, S., & Herron, J C. (2007). Evolutionary analysis. Upper Saddle River, NJ: Pearson Prentice Hall.

Runaway sexual selection: stalk-eyed flies

- ▶ Female stalk-eyed flies are choosy
- ▶ Both male eye stalk length and *female preference* for eye stalk length are heritable
- ▶ Sexual selection on the trait of one sex can cause an evolutionary response in the trait of the other sex
- ▶ Genetic correlation between male eye stalk length and female preference for eye stalk length^{1,2}

¹Wilkinson, G S & Reillo, P R. (1994). Proc. B. Soc. B. 255:1-6

²Freeman, S., & Herron, J C. (2007). Evolutionary analysis. Upper Saddle River, NJ: Pearson Prentice Hall.