

# Foundations of ecological and evolutionary change

A. Bradley Duthie<sup>1,a,\*</sup> and Victor J. Luque<sup>2,a</sup>

[1] Department of Biological and Environmental Sciences, University of Stirling, Stirling,  
Scotland [2] Department of Philosophy, University of Valencia, Valencia, Spain [\*]  
Corresponding author: [alexander.duthie@stir.ac.uk](mailto:alexander.duthie@stir.ac.uk) [a] Equal contribution

## Abstract

Biological evolution is realised through the same mechanisms of birth and death that underlie change in population density. The deep interdependence between ecology and evolution is well-established, but much theory in each discipline has been developed in isolation. While recent work has accomplished eco-evolutionary synthesis, a gap remains between the logical foundations of ecology and evolution. We bridge this gap with a new equation that defines a summed value for a characteristic across individuals in a population, from which the fundamental equations of population ecology and evolutionary biology (the Price equation) are derived. We thereby unify the fundamental equations of population ecology and biological evolution under a general framework. Our unification further demonstrates the equivalence between mean population growth rate and evolutionary fitness and links this change to ecosystem function. Finally, we outline how our proposed framework can be used to unify social evolution and density-dependent population growth.

**Key words:** Ecology, Evolution, Eco-Evolutionary Theory, Fundamental Theorem, Price Equation, Population Growth

## Introduction

Theoretical unification of phenomena is a powerful tool for scientific advancement. Such unification has been a major goal in scientific research throughout history ([Kitcher 1993](#)), and its value is perhaps most evident in reconciling unconnected models and revealing new and unexpected empirical predictions. In evolutionary

biology, the Price equation (Box 1) provides a unifying framework for evolutionary theory by exhaustively  
 24 and exactly describing evolutionary change for any closed evolving population (Price 1970; Luque 2017;  
 Lehtonen et al. 2020). The Price equation is therefore fundamental, in the sense that it binds together  
 26 all of evolutionary theory by formally defining what evolutionary change is and is not (Price 1970; Rice  
 2004; Gardner 2008; Frank 2017; Luque 2017; Luque and Baravalle 2021). Using this formal definition,  
 28 the scope of, and relationships among, sub-disciplines within evolutionary theory can be clarified. For  
 example, fundamental equations of both population and quantitative genetics can be derived from the Price  
 30 equation (Queller 2017). This provides conceptual clarity by demonstrating the logical consistency of different  
 theoretical frameworks within evolutionary biology. Our aim here is to propose an equation that extends this  
 32 conceptual clarity to include population change and thereby provide a formal and exact definition of joint  
 ecological and evolutionary change.

34 In biological populations, ecological change is caused by the same processes of individual birth and death  
 that cause changes in allele frequencies and phenotypes (Turchin 2001; Connor and Hartl 2004; Barfield et al.  
 36 2011). As with evolution, a fundamental equation can exhaustively and exactly define population change.  
 Unlike the Price equation, this fundamental equation is perhaps self-evident; population change is simply the  
 38 addition of births and subtraction of deaths from current population size ( $N_t$ ) to recover the new population  
 size ( $N_{t+1}$ ; Box 2). By definition, this relationship of  $N_{t+1} = N_t + Births - Deaths$  applies to any closed  
 40 population. Turchin (2001) argues that general principles are needed to establish a logical foundation for  
 population ecology, and this simple birth and death model and the consequences that logically follow from it  
 42 (e.g., exponential population growth) is fundamental to population ecology. Any unifying definition of joint  
 ecological and evolutionary change must be able to derive both the Price equation and this birth and death  
 44 model.

The interdependence of ecological and evolutionary processes has long been recognised (e.g., Darwin 1859;  
 46 Fisher 1958; Pelletier et al. 2009), but the rise of eco-evolutionary models, which incorporate both, is relatively  
 recent following a widespread recognition that ecology and evolution can happen on similar timescales (Govaert  
 48 et al. 2019; Yamamichi et al. 2023). Currently, a universally recognised formal definition of eco-evolutionary  
 change is lacking, with some theoreticians broadly interpreting “eco-evolutionary dynamics” to allow for a

separation of ecological and evolutionary timescales (Lion et al. 2023) and others advocating for a more narrow interpretation in which no such separation is permitted (Bassar et al. 2021). Bassar et al. (2021) identify two types of eco-evolutionary models that follow from these interpretations. The first type uses separate equations to model population change versus evolutionary change, thereby allowing for any number of ecological, evolutionary, or environmental feedbacks (e.g., Lion 2018; Patel et al. 2018; Lion et al. 2023). The second type models population demographics as functions of quantitative traits, with ecological and evolutionary change following from demographic processes and trait distributions operating on the same timescale (e.g., Barfield et al. 2011; Simmonds et al. 2020; Jaggi et al. 2024). Both model types can be very general, but like all predictive models, they rely on simplifying assumptions for tractability (Levins 1966; Luque 2017). These simplifying assumptions are often grounded in the Price equation to demonstrate accuracy and logical consistency when modelling evolutionary change (e.g., Coulson and Tuljapurkar 2008; Barfield et al. 2011; Rees and Ellner 2016; Lion 2018). For example, Barfield et al. (2011) link their model back to Price (1970), which they consider to be a “universal law of evolution”, to place their conclusions concerning stage structured evolution in the broader context of evolutionary theory. The role of fundamental equations is therefore important for unifying theory (Luque and Baravalle 2021), and we believe that a fundamental equation of eco-evolutionary change is needed.

We present an equation from which the fundamental equations of ecology and evolutionary biology can be derived. Derivation follows by adding assumptions that are specific to population ecology or evolution in the same way that key equations of population genetics (e.g., average excess) or quantitative genetics (e.g., Breeder’s equation) can be derived from the Price equation (Queller 2017). We propose our equation as a formal definition of eco-evolutionary change.

---

**Box 1:** The Price equation is an abstract formula to represent evolutionary change. Formulated originally in the early 1970s by George Price (Price 1970, 1972), it postulates some basic properties that all evolutionary systems must satisfy: change over time, ancestor and descendant relations, and a character or phenotype (Rice 2004). Using simple algebraic language, the Price equation

represents evolutionary change with the predominant notation,

$$\bar{w}\Delta\bar{z} = \text{Cov}(w, z) + E(w\Delta z).$$

In the above equation,  $\Delta\bar{z}$  is the change in the average character value  $z$  over a time step of arbitrary length,  $w$  is an individual's fitness, and  $\bar{w}$  average population fitness. On the right-hand side of the equation, the first term is the covariance between a character value  $z$  and fitness  $w$ , which reflects  $\bar{z}$  change attributable to differential survival and reproduction. The second term is the expected value of  $\Delta z$ , which reflects the extent to which offspring deviate from parents in  $z$  (Rice 2004; Okasha 2006; Frank 2012). A more specific version of the covariance term was already known within the quantitative and population genetics tradition (Robertson 1966), usually representing the action of natural selection. The Price equation adds an expectation term and abstracts away from any specific mechanisms of replication or reproduction, or mechanisms of inheritance. Its definitional nature and lack of substantive biological assumptions has been portrayed both as a strength (Rice 2004; Frank 2012; Luque 2017; Baravalle and Luque 2022), and its greatest weakness. The abstract nature of the Price equation places it at the top of the hierarchy of fundamental theorems of evolution from which the rest (Robertson's theorem, Fisher's fundamental theorem, breeder's equation, Hamilton's rule, adaptive dynamics, etc.) can be easily derived (Lehtonen 2016, 2018; Queller 2017). This abstractness is also key to developing a more general view of evolution (Rice 2020; Luque and Baravalle 2021; Edelaar et al. 2023). In contrast, some researchers consider the Price equation just a triviality (even tautological), and useless without further modelling assumptions (van Veelen 2005; van Veelen et al. 2012). The debate remains open (van Veelen 2020; Baravalle et al. 2025).

---

---

**Box 2:** The number of individuals in any closed population ( $N$ ) at any given time ( $t + 1$ ) is determined by the existing number of individuals ( $N_t$ ), plus the number of births (*Births*) minus

the number of deaths (*Deaths*),

$$N_{t+1} = N_t + Births - Deaths.$$

This equation is necessarily true for any closed population. Despite its simplicity, it is a general equation for defining population change and a starting point for understanding population ecology. Turchin (2001) notes that a consequence of this fundamental equation is the tendency for populations to grow exponentially (technically geometrically in the above case where time is discrete). This inherent underlying tendency towards exponential growth persists even as the complexities of real populations, such as structure, stochasticity, or density-dependent effects are added to population models (Turchin 2001). Given the assumption that all individuals in the population are identical, a per capita rate of birth,  $Births = bN_t$  and death,  $Deaths = dN_t$  can be defined. Rearranging and defining  $\lambda = 1 + b - d$  gives,  $N_{t+1} = N_t\lambda$ . Here  $\lambda$  is the finite rate of increase (Gotelli 2001), and note that because  $0 \leq d \leq 1$ ,  $\lambda \geq 0$ . Verbally, the change in size of any closed population equals its existing size times its finite rate of increase.

---

## A foundation for biological evolution and population ecology

To fully unify biological evolution and population ecology, we must reconcile the the Price equation (Box 1) with the general equation for population change (Box 2). The Price equation is critical for partitioning different components of biological change (Price 1970; Frank 1997; Gardner 2008; Luque 2017; Queller 2017; Lehtonen 2018). It has also been highly useful for integrating evolutionary theory across disciplines (Fox 2006; Brantingham 2007; MacCallum et al. 2012; Frank 2015; Godsoe et al. 2021; Ulrich et al. 2024). These properties would seem to make it an intuitive starting point for a logical foundation of ecology and evolution, perhaps through some kind of mathematical equivalence (Page and Nowak 2002) or addition of terms (Collins and Gardner 2009), or through the use of its recursive structure (Kerr and Godfrey-Smith 2009; Frank 2012). But despite its flexibility, the Price equation still relies on relative frequencies, which must by definition sum

to one (Frank 2015). This is because the Price equation describes the average change in a population; the frequency of entities is scaled thereby conserving total probability (Frank 2015, 2016). But to recover the fundamental principle of exponential population growth (Turchin 2001), this scaling must be avoided in a fundamental equation of ecology and evolution.

We therefore begin with the most fundamental axioms underlying the ecology and evolution of living systems (Rice 2004; Rice and Papadopoulos 2009). In such systems, diversity is discontinuous and can be defined in terms of discrete entities (Dobzhansky 1970). Our framework is general enough that entities can be anything discrete, but we will focus on each entity  $i$  as an individual organism. Individuals give rise to new individuals through birth such that  $\beta_i$  is the number of births attributable to  $i$ . Individuals are removed from the population through death such that  $\delta_i$  is an indicator variable that takes a value of 1 (death of  $i$ ) or 0 (persistence of  $i$ ). All individuals are defined by some characteristic  $z_i$ , and  $\Delta z_i$  defines any change in  $z_i$  from one time step  $t$  to the next  $t + 1$ . The total number of individuals in the population is  $N$ . From this foundation, we can define  $\Omega$  to be a summed characteristic across  $N$  entities,

$$\Omega = \sum_{i=1}^N (\beta_i - \delta_i + 1) (z_i + \Delta z_i). \quad (1)$$

The foundation of eco-evolutionary change is an interaction between the demographic processes of birth and death ( $\beta_i - \delta_i + 1$ ) and some characteristic of an individual ( $z_i + \Delta z_i$ ). The value of  $\Omega$  is a summation of individual characteristics, which takes the same units as  $z$ . From eqn 1, we can derive the most fundamental equations of population ecology (Box 2) and evolutionary biology (Box 1) through an appropriate interpretation of  $z$ . Under some limited interpretations of  $z$ , we can also interpret  $\Omega$  as a metric of ecosystem function.

## Population ecology

To recover the general equation for population ecology (Box 2), we define  $z_i$  as the identity of  $i$  belonging to the population. In other words, we set  $z_i = 1$  to simply indicate that  $i$  is a member of the population. In this case,  $z$  is count, which takes the unit 1 [note that ‘individual’ is a label, not a unit; Newell and Tiesinga

(2019)]. Further, we assume that individuals do not change species by setting  $\Delta z_i = 0$ . In this case,

$$\Omega = \sum_{i=1}^N (\beta_i - \delta_i + 1).$$

146 We can now interpret  $\Omega$  as the population size at  $t + 1$ ,  $N_{t+1}$ ,

$$N_{t+1} = N_t + \text{Births} - \text{Deaths}. \quad (2)$$

If we assume that individuals are identical, then we can drop the subscript  $i$  such that  $\beta_i = \beta$  and  $\delta_i = \delta$  and  
 148 note that  $N\beta$ ,  $N\delta$ , and  $N$  are the total births, total deaths, and size of the population at  $t$ , respectively,

Summing from 1 to  $N$ , we can rewrite the above,

$$\Omega = N(1 + \beta - \delta).$$

150 If we define  $\lambda = 1 + \beta - \delta$  (Box 2), then we can rewrite,

$$N_{t+1} = N_t \lambda. \quad (3)$$

We therefore recover the general equation for population ecology (eqn 2) and the fundamental property of  
 152 exponential growth in populations (Turchin 2001) (eqn 3).

## Evolutionary biology

154 Recovering the Price equation requires a few more steps. We start by defining individual fitness,

$$w_i = \beta_i - \delta_i + 1. \quad (4)$$

In this definition, the longevity of the individual matters. An individual that survives from  $t$  to  $t + 1$  has a higher fitness than one that dies, even if both have the same reproductive output. With this definition of fitness (eqn 4), we substitute,

$$\Omega = \sum_{i=1}^N (w_i z_i + w_i \Delta z_i). \quad (5)$$

We can break eqn 5 down further and multiply each side by  $1/N$ ,

$$\frac{1}{N}\Omega = \frac{1}{N} \sum_{i=1}^N (w_i z_i) + \frac{1}{N} \sum_{i=1}^N (w_i \Delta z_i). \quad (6)$$

We can rewrite the terms on the right-hand side of eqn 6 as expected values and remove the subscripts,

$$\frac{1}{N}\Omega = E(wz) + E(w\Delta z). \quad (7)$$

Now we must consider the total conservation of probability (Frank 2015, 2016). In eqn 7,  $\Omega$  is the total sum trait values ( $z_i$ ) across the entire population at  $t + 1$  divided by the number of individuals ( $N$ ) in the population at  $t$ . But the size of the population can change from  $t$  to  $t + 1$ . To recover mean trait change for the Price equation (and therefore conserve total probability), we need to account for this change in population size. The left-hand side of eqn 7 describes contributions to the sum trait value from the new population at  $t + 1$ . But we cannot treat  $\Omega/N$  as the mean of  $z$  at  $t + 1$  ( $\bar{z}'$ ) because we need to weigh  $N$  by the mean fitness of the population at  $t$  to account for any change in population size from  $t$  to  $t + 1$ .

For example, if mean fitness at  $t$  is 2 (i.e.,  $\bar{w} = 2$ ), then half as many individuals will have contributed to  $\Omega$  in  $t + 1$  than would have if  $\bar{w} = 1$  (i.e., there are  $N$  individuals at  $t$  and  $2N$  individuals at  $t + 1$ ). We therefore need to multiply the mean trait value  $\bar{z}'$  (at  $t + 1$ ) by the mean fitness  $\bar{w}$  (at  $t$ ) to recover the mean contribution of the  $N$  individuals at  $t$  to the total  $\Omega$ . Consequently,

$$\Omega = N\bar{w}\bar{z}' \quad (8)$$



Equation 8 conserves the total probability and now clarifies  $\Omega$  as a summed trait value, which has the same  
 172 units as  $z$  and equals expected population growth at  $t$  times mean trait value at  $t + 1$ . This is consistent with  
 the population ecology derivation from the previous section where  $z_i = 1$  by definition, and  $\Omega = N_{t+1}$  (note  
 174  $\lambda = \bar{w}$ ). We can therefore rewrite eqn 7,

$$\bar{w}\bar{z}' = E(wz) + E(w\Delta z). \quad (9)$$

We can rearrange eqn 9 to derive the Price equation by expressing covariance as,  $\text{Cov}(X, Y) = E(XY) -$   
 176  $E(X)E(Y)$ , and therefore  $E(XY) = \text{Cov}(X, Y) + E(X)E(Y)$ . Substituting into eqn 9,

$$\bar{w}\bar{z}' = \text{Cov}(w, z) + \bar{w}\bar{z} + E(w\Delta z).$$

From here,

$$\bar{w}(\bar{z}' - \bar{z}) = \text{Cov}(w, z) + E(w\Delta z).$$

178 Since  $\Delta\bar{z} = (\bar{z}' - \bar{z})$ ,

$$\bar{w}\Delta\bar{z} = \text{Cov}(w, z) + E(w\Delta z). \quad (10)$$

From eqn 1, which describes fundamental birth and death processes in a population, we can derive both the  
 180 most fundamental model of population ecology (eqn 2; Box 2) and the fundamental equation of evolution  
 (eqn 10; Box 1).

## 182 Ecosystem function

In some cases,  $\Omega$  could also be interpreted as the total contribution of a population to ecosystem function. This  
 184 will be restricted to cases in which  $z$  is an individual characteristic defining an absolute quantity measured

at the whole organism level, such as biomass, seed production, carbon capture, flower visits, or nutrient  
 186 consumption. In these cases, the sum across individuals gives a meaningful total quantity for the population.  
 When  $z$  is instead defined by relative organism-level measurements such wing loading, nutrient ratio, or diet  
 188 composition, or when  $z$  is measured at a level of biological organisation below the organism (e.g., average cell  
 volume or leaf surface area),  $\Omega$  does not have a clear population-level interpretation.

190 Box 3 provides a toy example of three plants with different fitnesses and carbon captures. We use this toy  
 example to demonstrate a concrete example of our proposed framework.

---

**Box 3:** As a toy example of our framework, consider a population of  $N_t = 3$  annual plants in  
 194 which individual fruit mass (kg) is measured, and change is observed over a year. For all plants,  
 $\delta_i = 1$ , and let plant fecundities be  $\beta_1 = 1$ ,  $\beta_2 = 1$ , and  $\beta_3 = 2$ . Applying eqn 1 to population  
 196 change such that  $\Omega = N_{t+1}$ ,  $z_i = 1$  and  $\Delta z_i = 0$ ,  $N_{t+1} = (1 - 1 + 1)(1 + 0) + (1 - 1 + 1)(1 +$   
 $0) + (2 - 1 + 1)(1 + 0) = 4$  (note  $\bar{w} = 4/3$ , so  $N_t \bar{w} = 4$ ). Focusing next on the characteristic  
 198 of total fruit mass, let  $z_1 = 0.8$ ,  $z_2 = 1.0$ , and  $z_3 = 1.5$ . Also let  $\Delta z_i = 0.1$  for all plants to  
 reflect a change in soil environment from  $t$  to  $t + 1$ . In this case, total fruit yield at  $t + 1$  is  
 200  $\Omega = (1 - 1 + 1)(0.8 + 0.1) + (1 - 1 + 1)(1.0 + 0.1) + (2 - 1 + 1)(1.5 + 0.1) = 5.2$  kg. At  $t$ , mean  
 fruit yield per plant was 1.1 kg, but at  $t + 1$ , mean fruit yield per plant is 1.3 kg. Note that  
 202  $\text{Cov}(w, z) = 2/15$  and  $E(w\Delta z) = 2/15$ , so applying the Price equation,  $\bar{w}\Delta z = 2/15 + 2/15 = 4/15$ .  
 Since  $\bar{w} = 4/3$ , multiplying both sides of the equation by  $3/4$  returns  $\Delta z = 0.2$ , which is the mean  
 204 difference in fruit yield between  $t + 1$  and  $t$ . The framework expressed in eqn 1 thereby links  
 population change, evolutionary change, and ecosystem function.

---

## Discussion

208 A classical sign of scientific progress is the ability to connect disparate theories and models to show how  
 empirical and theoretical models are logical (mathematical) consequences of more fundamental ones ([Nagel](#)

1961; Morrison 2000). Rather than making simplifying assumptions, as is the approach for most ecological and evolutionary models, we focus on fundamental axioms that are universal to closed biological systems: discrete individuals, birth, death, and change over time. We define an abstract sum ( $\Omega$ ), to which all individuals within the population contribute. From the simple assumptions of population identity ( $z_i = 1$ ) and invariability ( $\Delta z_i = 0$ ), we recover the most general equation of population ecology (Box 2) and principle of exponential growth ( $N_{t+1} = N_t \lambda$ ). By defining individual fitness ( $w_i$ ) and applying the total conservation of probability to individual frequencies (Frank 2015, 2016), we recover the most fundamental equation of evolution (Box 1). Our eqn 1 thereby provides a foundation for defining eco-evolutionary change in any population.

The Price equation provides a complete and exact description of evolution in any closed evolving system (Price 1970; Frank 2012). It is derived by rearranging the mathematical notation defining changes in the frequencies and characteristics of any type of entity (Price 1970; Gardner 2008; Luque 2017) (e.g., individuals, alleles). This derivation partitions total characteristic change into different components, making it possible to isolate evolutionary mechanisms (e.g., selection) and levels of biological organisation (e.g., group, individual) (Frank 1995, 2012; Kerr and Godfrey-Smith 2009; Luque 2017; Okasha and Otsuka 2020). Because of its abstract nature and lack of any system-specific assumptions, the Price equation is not dynamically sufficient and makes no predictions about what will happen in any particular system (Gardner 2020). Its role is not to predict, but to formally and completely define and separate components of evolutionary change. The same is true of the general equation for population change (eqns 2 and 3), at least as we have used it here where it serves to define what population growth means in ecology. This equation formally and completely describes population change in terms of births and deaths. In eqn 1, we therefore have a fundamental equation from which we can derive complete ecological and evolutionary change in any closed biological population, but one that is revealed to be necessarily abstract and not dynamically sufficient. We anticipate that this will be useful for eco-evolutionary theory in the same way that the Price equation is useful for evolutionary theory: facilitating specific model development and identifying new conceptual insights, unresolved errors, and sources of model disagreements.

Our unification recovers the equivalence between the finite rate of increase  $\lambda$  (Box 2) (Gotelli 2001) and population mean evolutionary fitness  $\bar{w}$  (Box 1). The population growth equation  $N_{t+1} = N_t \lambda$  can therefore

be rewritten as  $N_{t+1} = N_t \bar{w}$ . This specific equivalence has been proposed before (Lande 1976), as has the  
 238 broader relationship between population growth rate and evolutionary fitness (Fisher 1930; Charlesworth  
 1980; Lande 1982; Roff 2008; Lion 2018). We show this from first principles and clarify the relationship  
 240 between fitness and population growth. Over an arbitrary length of time, fitness is properly defined as  
 $w_i = \beta_i - \delta_i + 1$ . Over an individual’s lifetime (which, by definition, includes death), fitness is therefore  $\beta_i$ .  
 242 The rate of change in ecology and evolution are reflected in the first and second statistical moments of fitness,  
 respectively. Population growth rate reflects mean fitness  $\bar{w}$ , while the rate of evolutionary change reflects  
 244 the variance in fitness  $\text{Var}(w)/\bar{w}$  (i.e., Fisher’s fundamental theorem) (Frank 1997; Rice 2004; Queller 2017).

Our unification may also help explain, at least partially, some of the success of classical population genetic  
 246 models. For decades, population genetics (and to some extent quantitative genetics) has been accused of  
 being a reductionist view of evolution, reducing everything to changes in allele frequencies and abstracting  
 248 away from individuals and their environments (the ecological interactions) (MacColl 2011). This has been a  
 line of argumentation by some defenders of the so-called Extended Synthesis (Pigliucci 2009), especially in  
 250 relation to niche construction (Odling-Smee et al. 2003). Famously, Mayr (1959) characterized population  
 genetics as a simple input and output of genes, analogous to “the adding of certain beans to a beanbag and  
 252 the withdrawing of others” (also called “beanbag genetics”). Historical critics of population genetics could  
 not articulate a clear explanation for why it works so well despite all of its idealisations and simplifications.  
 254 From the Price equation, we are able to recover classical population and quantitative genetic models (Queller  
 2017) and develop new ones (Rice 2004, 2020; Luque 2017; Lion 2018). Our eqn 1 contains ecology at its core,  
 256 and we show how the Price equation logically follows from it after accounting for absolute population growth  
 (eqn 8). We therefore conclude that population and quantitative genetic equations contain ecology (no matter  
 258 how hidden), and the ecological nature of evolution is implicit in population and quantitative genetic models.

We have focused on the dynamics of a closed population, and in doing so leave ecological and evolutionary  
 260 change attributable to migration for future work. In population ecology, immigration and emigration can  
 be incorporated by adding a term for each to the right-hand side of the equation in Box 2 (Gotelli 2001).  
 262 In evolution, because the Price equation relies on mapping ancestor-descendant relationships, accounting  
 for migration is more challenging. Kerr and Godfrey-Smith (2009) demonstrate how the Price equation can

be extended to allow for arbitrary links between ancestors and descendants, thereby extending the Price equation to allow for immigration and emigration. Frank (2012) presents a simplified version of Kerr and Godfrey-Smith (2009) that allows some fraction of descendants to be unconnected to ancestors. In both ecology and evolution, accounting for migration is done through the use of additional terms on the right-hand side of the equations.

Our fundamental equation is complete and exact. It therefore implicitly includes any effects of density dependence on population growth (see Box 2), or any social effects on evolutionary change (see Box 1). Both of these effects can be made explicit by specifying how other individuals in a population affect birth and death of a focal individual. Assuming that effects of individuals are additive, independent, and identical, we can derive well-established logistic growth equations and Hamilton’s rule from eqn 1 (Supporting Information S1).

We have shown that we can derive the fundamental equations of population ecology and biological evolution from a single unifying equation. Lastly, we propose our eqn 1 as a potential starting point for defining ecosystem function and further conceptual unification between ecology, evolution, and ecosystem function. The Price equation has previously been used to investigate ecosystem function (Loreau and Hector 2001; Fox 2006), but not with any attempt towards conceptual unification with evolutionary biology. For example, Fox (2006) applied the abstract properties of the Price equation to partition total change in ecosystem function into separate components attributable to species richness, species composition, and context dependent effects. This approach provides a framework for comparing the effects of biodiversity on ecosystem function in empirical systems (Fox 2006; Winfree et al. 2015; Mateo-Tomás et al. 2017). Instead, our eqn 1 defines  $\Omega$  as total ecosystem function contributed by a focal population. It is therefore possible to investigate ecological, evolutionary, and ecosystem function change from the same shared framework.

## Acknowledgements

This manuscript was supported by joint funding between the French Foundation for Research on Biodiversity (FRB) Centre for the Synthesis and Analysis of Biodiversity (CESAB) and the German Centre for Integrative

Biodiversity Research (sDiv). It was written as part of the Unification of Modern Coexistence Theory and Price Equation (UNICOP) project. Victor J. Luque was also supported by the Spanish Ministry of Science and Innovation (Project: PID2021-128835NB-I00), and the Conselleria d’Innovació, Universitats, Ciència i Societat Digital – Generalitat Valenciana (Project: CIGE/2023/16). We are grateful for many conversations with Sébastien Lion, Kelsey Lyberger, Swati Patel, and especially Lynn Govaert, whose questions helped us clarify the relationship between population growth and fitness. Brad Duthie would also like to thank Brent Danielson and Stan Harpole. Victor J. Luque would also like to thank Lorenzo Baravalle, Pau Carazo, Santiago Ginnobili, Manuel Serra, and Ariel Roffé.

## Author Contributions

Both authors came up with the question idea. Following many discussions between the authors, ABD proposed the initial equation with subsequent exploration and development from both authors. Both authors contributed to the writing.

## Competing Interests

The authors declare no competing interests.

## Data Availability

This work does not include any data.

## References

- Baravalle, L., and V. J. Luque. 2022. [Towards a pricean foundation for cultural evolutionary theory](#). *Theoria* 37:209–231.
- Baravalle, L., A. Roffé, V. J. Luque, and S. Ginnobili. 2025. [The value of price](#). Pages 12–24 *in* *Biological Theory* (Vol. 20).

- Barfield, M., R. D. Holt, and R. Gomulkiewicz. 2011. [Evolution in stage-structured populations](#). *American Naturalist* 177:397–409.
- Bassar, R. D., T. Coulson, J. Travis, and D. N. Reznick. 2021. [Towards a more precise – and accurate – view of eco-evolution](#). *Ecology Letters* 24:623–625.
- Brantingham, P. J. 2007. [A unified evolutionary model of archaeological style and function based on the Price equation](#). *American Antiquity* 72:395–416.
- Charlesworth, B. 1980. *Evolution in age-structured populations*. Cambridge studies in mathematical biology. Cambridge University Press, Cambridge.
- Collins, S., and A. Gardner. 2009. [Integrating physiological, ecological and evolutionary change: A Price equation approach](#). *Ecology Letters* 12:744–757.
- Connor, J., and D. L. Hartl. 2004. *A premier of ecological genetics*. Sinauer Associates Incorporated.
- Coulson, T., and S. Tuljapurkar. 2008. [The dynamics of a quantitative trait in an age-structured population living in a variable environment](#). *American Naturalist* 172:599–612.
- Darwin, C. 1859. *The origin of species*. Penguin.
- Dobzhansky, T. 1970. *Genetics of the evolutionary process* (Vol. 139). Columbia University Press.
- Edelaar, Pim, J. Otsuka, and V. J. Luque. 2023. [A generalised approach to the study and understanding of adaptive evolution](#). *Biological Reviews* 98:352–375.
- Fisher, R. A. 1930. *The genetical theory of natural selection*. Oxford University Press, Oxford, UK.
- Fisher, R. A. 1958. *The genetical theory of natural selection* (2nd ed.). Dover.
- Fox, J. W. 2006. [Using the price equation to partition the effects of biodiversity loss on ecosystem function](#). *Ecology* 87:2687–2696.
- Frank, S. A. 1995. [George Price’s contributions to evolutionary genetics](#). *Journal of Theoretical Biology* 175:373–388.
- . 1997. [The Price equation, Fisher’s fundamental theorem, kin selection, and causal analysis](#). *Evolution* 51:1712–1729.
- . 2012. [Natural selection. IV. The Price equation](#). *Journal of Evolutionary Biology* 25:1002–1019.
- . 2015. [D’Alembert’s direct and inertial forces acting on populations: The price equation and the](#)

fundamental theorem of natural selection. *Entropy* 17:7087–7100.

———. 2016. [Common probability patterns arise from simple invariances](#). *Entropy* 18:1–22.

———. 2017. [Universal expressions of population change by the Price equation: natural selection, information, and maximum entropy production](#). *Ecology and Evolution* 1–16.

Gardner, A. 2008. [The Price equation](#). *Current Biology* 18:198–202.

———. 2020. [Price’s equation made clear](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 375:20190361.

Godsoe, W., K. E. Eisen, D. Stanton, and K. M. Sirianni. 2021. [Selection and biodiversity change](#). *Theoretical Ecology* 14:367–379.

Gotelli, N. J. 2001. *A primer of ecology*. Sinauer associate. Inc. Sunderland, MA.

Govaert, L., E. A. Fronhofer, S. Lion, C. Eizaguirre, D. Bonte, M. Egas, A. P. Hendry, et al. 2019.

[Eco-evolutionary feedbacks—Theoretical models and perspectives](#). *Functional Ecology* 33:13–30.

Jaggi, H., W. Zuo, R. Kentie, J. M. Gaillard, T. Coulson, and S. Tuljapurkar. 2024. [Density dependence](#)

[shapes life-history trade-offs in a food-limited population](#). *Ecology letters* 27:e14551.

Kerr, B., and P. Godfrey-Smith. 2009. [Generalization of the price equation for evolutionary change](#). *Evolution* 63:531–536.

Kitcher, P. 1993. *The advancement of science*. Oxford University Press, New York.

Lande, R. 1976. Natural selection and random genetic drift in phenotypic evolution. *Evolution* 30:314–334.

———. 1982. [A quantitative genetic theory of life history evolution](#). *Ecology* 63:607–615.

Lehtonen, J. 2016. [Multilevel selection in kin selection language](#). *Trends in Ecology and Evolution* xx:1–11.

———. 2018. [The Price equation, gradient dynamics, and continuous trait game theory](#). *American Naturalist* 191:146–153.

Lehtonen, J., S. Okasha, and H. Helanterä. 2020. [Fifty years of the Price equation](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 375:20190350.

Levins, R. 1966. [The strategy of model building in population biology](#). *American Naturalist*.

Lion, S. 2018. [Theoretical approaches in evolutionary ecology: environmental feedback as a unifying perspective](#). *American Naturalist* 191.



- Lion, S., A. Sasaki, and M. Boots. 2023. [Extending eco-evolutionary theory with oligomorphic dynamics](#). Ecology Letters 26:S22–S46.
- Loreau, M., and A. Hector. 2001. [Partitioning selection and complementarity in biodiversity experiments](#). Nature 412:72–76.
- Luque, V. J. 2017. [One equation to rule them all: a philosophical analysis of the Price equation](#). Biology and Philosophy 32:1–29.
- Luque, V. J., and L. Baravalle. 2021. [The mirror of physics: on how the price equation can unify evolutionary biology](#). Synthese 199:12439–12462.
- MacCallum, R. M., M. Mauch, A. Burt, and A. M. Leroi. 2012. [Evolution of music by public choice](#). Proceedings of the National Academy of Sciences 109:12081–12086.
- MacColl, A. D. C. 2011. [The ecological causes of evolution](#). Trends in Ecology and Evolution 26:514–522.
- Mateo-Tomás, P., P. P. Olea, M. Moleón, N. Selva, and J. A. Sánchez-Zapata. 2017. [Both rare and common species support ecosystem services in scavenger communities](#). Global Ecology and Biogeography 26:1459–1470.
- Mayr, E. 1959. [Where are we? Genetics and twentieth century darwinism](#). Pages 1–14 *in* Cold spring harbor symposia on quantitative biology (Vol. 24).
- Morrison, M. 2000. Unifying scientific theories: Physical concepts and mathematical structures. Cambridge University Press.
- Nagel, E. 1961. The structure of science: Problems in the logic of scientific explanation. Harcourt, Brace & World, New York, NY, USA.
- Newell, D. B., and E. Tiesinga. 2019. [The international system of units \(SI\)](#).
- Odling-Smee, F. J., K. N. Laland, and M. W. Feldman. 2003. Niche construction: The neglected process in evolution. Princeton University Press.
- Okasha, S. 2006. [Evolution and the levels of selection](#). Oxford University Press.
- Okasha, S., and J. Otsuka. 2020. [The Price equation and the causal analysis of evolutionary change](#). Philosophical Transactions of the Royal Society B: Biological Sciences 375:20190365.
- Page, K. M., and M. A. Nowak. 2002. [Unifying evolutionary dynamics](#). Journal of theoretical biology 219:93–98.

- Patel, S., M. H. Cortez, and S. J. Schreiber. 2018. [Partitioning the effects of eco-evolutionary feedbacks on community stability](#). *American Naturalist* 191:1–29.
- Pelletier, F., D. Garant, and A. P. Hendry. 2009. [Eco-evolutionary dynamics](#). *Philosophical Transactions of the Royal Society London B* 364:1483–1489.
- Pigliucci, M. 2009. [An extended synthesis for evolutionary biology](#). *Annals of the New York Academy of Sciences* 1168:218–228.
- Price, G. R. 1970. [Selection and covariance](#).
- . 1972. [Extension of covariance selection mathematics](#). *Annals of Human Genetics* 35:485–490.
- Queller, D. C. 2017. [Fundamental theorems of evolution](#). *American Naturalist* 189:000–000.
- Rees, M., and S. P. Ellner. 2016. [Evolving integral projection models: Evolutionary demography meets eco-evolutionary dynamics](#). *Methods in Ecology and Evolution* 7:157–170.
- Rice, S. H. 2004. *Evolutionary theory: mathematical and conceptual foundations*. Sinauer Associates Incorporated.
- . 2020. [Universal rules for the interaction of selection and transmission in evolution](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 375.
- Rice, S. H., and A. Papadopoulos. 2009. [Evolution with stochastic fitness and stochastic migration](#). *PLoS One* 4.
- Robertson, A. 1966. [A mathematical model of the culling process in dairy cattle](#). *Animal Science* 8:95–108.
- Roff, D. A. 2008. [Defining fitness in evolutionary models](#). *Journal of Genetics* 87:339–348.
- Simmonds, E. G., E. F. Cole, B. C. Sheldon, and T. Coulson. 2020. [Phenological asynchrony: A ticking time-bomb for seemingly stable populations?](#) *Ecology Letters*. Blackwell Publishing Ltd.
- Turchin, P. 2001. [Does population ecology have general laws?](#) *Oikos* 94:17–26.
- Ulrich, W., N. J. Gotelli, G. Strona, and W. Godsoe. 2024. [Reconsidering the price equation: Benchmarking the analytical power of additive partitioning in ecology](#). *Ecological Modelling* 491:110695.
- van Veelen, M. 2005. [On the use of the Price equation](#). *Journal of Theoretical Biology* 237:412–426.
- . 2020. [The problem with the Price equation](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 375:20190355.

- 418 van Veelen, M., J. García, M. W. Sabelis, and M. Egas. 2012. [Group selection and inclusive fitness are not  
equivalent; the Price equation vs. models and statistics](#). *Journal of Theoretical Biology* 299:64–80.
- 420 Winfree, R., J. W. Fox, N. M. Williams, J. R. Reilly, and D. P. Cariveau. 2015. [Abundance of common  
species, not species richness, drives delivery of a real-world ecosystem service](#). *Ecology Letters* 18:626–635.
- 422 Yamamichi, M., S. P. Ellner, and N. G. Hairston. 2023. [Beyond simple adaptation: Incorporating other  
evolutionary processes and concepts into eco-evolutionary dynamics](#). *Ecology Letters* 26:S16–S21.