

3 Apportionment

3.2 Jefferson's Method

Definition 3.2.1 Jefferson's Method:

1. Find the standard divisor
2. Find each states lower quota
3. If the lower quotas do not add up to the number of seats available, then adjust the divisor, and go back to step 2.

Example 3.2.2 Let's redo the first example from last time, this time with Jefferson's method.

County	Population
Kent	162,310
New Castle	538,479
Sussex	197,145
Total	1,052,567

Example 3.2.3 Use Jefferson’s method to apportion the 75 seats of Rhode Island’s House of Representatives among its five counties.

County	Population
Bristol	49,875
Kent	166,158
Newport	82,888
Providence	626,667
Washington	126,979
Total	1,052,567

Example 3.2.4 Consider a city with four districts, named A , B , C , and D . The local city council has a total of 12 seats, which are to be divided up between the four districts according to population. Using the table below, determine the number of seats each district gets using Jefferson's method.

District	Population
A	435
B	324
C	134
D	612
Total	1505

Example 3.2.5 Use **Hamilton's** method to apportion the 10 seats of the city council between the three districts A , B , C according to the following table:

District	Population
A	6
B	6
C	2

What happens if the number of seats is increased to 11?

Example 3.2.6 Three people invest in a treasure dive, each investing the amount listed below. The dive results in 36 gold coins. Apportion those coins to the investors using Jefferson's method.

Alice \$7,600

Ben: \$5,900

Calos: \$1,400