CHRISTEYE / SCIENTIFIC DEVELOPMENT DOSSIER

SYLPH

Adaptive Cancer Surveillance, Neural Training & Controlled Cell Therapy

Integrated research hypotheses • Living Periodic Table • Preclinical studies • Conditional clinical-trial pathway

VERSION 1.0 | 2 NOVEMBER 2026 | WE DEVELOP SOFTWARE LLC

RESEARCH STATUS

A comprehensive exploratory program. Not a validated product, medical device, cancer-screening method, prevention method, or treatment. No human administration, oral living-neuron vial, or in-body autonomous neural network is authorized or proposed for immediate testing.

Prepared as a development and collaboration brief for oncology researchers, cell biologists, bioelectronic engineers, clinical-trial designers, and regulatory advisors.




Figure 1. Recommended modular architecture; not an established physiological pathway. The biological controller is optional until it proves incremental value over electronic control.

01 Executive summary and claim boundaries

ChristEye Sylph is a proposed closed-loop research platform that combines longitudinal biomarker interpretation, optional cultured-neuron computation, and separately controlled therapeutic cells. The immediate objective is not universal cancer prevention. It is to determine whether an adaptive controller can correctly distinguish normal, injured, and malignant biological states, and whether linking that decision to an independently validated treatment provides a measurable advantage without added risk.

What the project contributes

- A common Living Table vocabulary to connect molecular mechanisms, neural signaling, support cells, therapeutic effectors, safety feedback, and learning history.
- A falsifiable neural-training program with separate measures for network learning, axonal propagation, endurance, viability, and whole-system energy consumption.
- A staged oncology pathway: monitor a defined biomarker pattern; verify signal interpretation; test selective tumor response; assess safety and only then consider a clinical program.
- A longitudinal patient-baseline hypothesis with strict safeguards against false positives, false negatives, drift, tumor promotion, and uncontrolled cellular persistence.

What Sylph does not yet establish

It has not been demonstrated that neurons independently identify cancer, circulating axons form a stable network, a daily oral vial creates a surveillance system, stem cells automatically heal tumors, or any version prevents or cures cancer. “Bad vibrations” is used here only as a metaphor for repeated biological injury and abnormal signaling. mRNA is normal and essential to cellular protein synthesis; a pathological RNA profile is not equivalent to all RNA being dangerous.

DECISION PRINCIPLE

The default comparator is an ordinary electronic controller and standard oncology practice. A living controller must earn its place with repeatable incremental benefit, not merely successful biological activity.

02 Target use cases and primary hypotheses

Track	Near-term target	Why it is separate
A. Baseline and early warning	Research-grade prediction of a narrowly defined cancer-related or tissue-injury state using longitudinal measurements.	Prediction is not diagnosis; screening usefulness requires clinical-outcome evidence.
B. Biohybrid neural control	Contained cultured-neuron system trained on sensor-derived surrogate patterns.	It may offer no advantage over software.
C. Localized therapy	One tumor indication; approved or investigational effector evaluated independently.	Tumor suppression and tissue regeneration require distinct endpoints.
D. Long-horizon prevention	Future concept only after benefit, harms, and clinical validity are proven.	Healthy populations face especially high safety and false-alarm burdens.

Pre-registered hypotheses

ID	Proposed research hypothesis	Null / failure condition
H1	A patient-specific longitudinal model adds diagnostic discrimination beyond an accepted non-personalized model.	No reproducible improvement on held-out patient cohorts.
H2	Feedback-trained neural cultures improve task performance over input-only, untrained, and replay controls.	No above-control effect across independent cultures.
H3	Axonal transmission remains reliable under task demands, and adaptation is quantifiable without compromising viability.	Activity rises without task gains, or reliability/viability worsens.
H4	A selected therapy reduces malignant-model survival while preserving healthy counterpart tissue.	No selectivity or increased tumor-supportive growth.
H5	Independent stop/deactivation controls reliably limit therapeutic cell activity and persistence.	Unacceptable escape, persistence, migration, or failure to deactivate.
H6	The integrated module improves a predefined clinical-relevant outcome over the conventional comparator.	No net advantage after harms, costs, or delay are included.

03 ChristEye ontology: framework mapped to measurable systems

Framework term	Proposed role in Sylph	Operational observable / boundary
Soul	Integration and continuity of the entire monitored system.	System availability, coordinated operation, patient safety; philosophical term, not a detected biological entity.
Lotus	Processing, adaptive decision-making, training and coordinated response.	Classifier performance, action accuracy, model drift, neural readout.
Marrow	DNA, inherited cell identity, structural integrity and expression competence.	Genotype identity, transcript/protein markers, lineage stability, mutation monitoring.
Roots	Retained development/history and learned response patterns.	Longitudinal baseline, trained parameters, patient history and calibration.
Winds / Spirit	In this model, signal changes, biochemical and ionic communication.	Signals quantified as molecules, currents, concentrations and time; spiritual meaning not a measured force.
Heart	Priorities, needs, homeostasis and protective control.	Explicit clinical objectives, allowable operating range, escalation and stop rules.
Knowledge	Observed or learned relationships linking signals and outcomes.	Traceable datasets, prediction uncertainty, reproducible parameter estimates.

Framework term	Proposed role in Sylph	Operational observable / boundary
Understanding	Robust use of information to support the specified task.	Generalization, calibration, durability, physiological benefit and no net harm.

The Living Table is a functional engineering taxonomy across levels of biology. It is not a substitute for the chemical periodic table and does not claim the elements listed are interchangeable.

04 The Living Periodic Table: exhaustive candidate inventory

Layer 1 — atoms, ions, and essential chemical ingredients

Type	Candidate and known role	Sylph-relevant question
Chemical elements	C, H, O, N, P, S are structural building blocks of cells, proteins, lipids and nucleic acids.	Do available nutrients and matter meet system demands?
Electrolytes	Na+, K+, Ca2+, Cl- and Mg2+ participate in gradients, transmission and regulation.	Are ionic conditions compatible with reliable neural operation?
Trace metals	Fe, Zn and Cu support enzymes and metabolism but can be harmful at excess levels.	Do assays detect toxic imbalance and oxidative stress?
Subatomic context	Electrons and protons govern chemistry; neutrons are nuclear constituents.	Avoid confusing electrochemical charge transport with neuron firing or fusion energy.

Layer 2 — biochemical information and cellular machinery

Code / group	Established role	Key readout or guardrail
DNA / Marrow	Cellular genetic information and regulatory architecture.	Identity, instability, tumor-associated alterations.
RNA / mRNA	Transmits instructions for protein synthesis; abnormal expression can be disease-related.	Select validated transcripts; do not treat mRNA itself as toxic.
Rb — ribosome	Translates mRNA into proteins.	Protein identity, fidelity and bounded expression.
Mt — mitochondrion	ATP production and local metabolic regulation.	Respiration, ATP availability, stress signals.
ATP / metabolites	Chemical energy handling and turnover.	Energy budget, viable-cell demand, waste products.
Ion channels / pumps	Electrical excitability and restoration of Na/K gradients.	Action-potential fidelity, recovery and metabolic burden.
Cytoskeleton / tau	Structural support and intracellular transport.	Morphological integrity; tau is protein, not a new cell type.
Vesicles / receptors	Package and detect cellular signals.	Release identity, thresholds, off-target activation.

Layer 3 — neuronal structures and support cells

Code	Function in the Living Table	Sylph role / caution
Nr — neurons	Electrical integration and network adaptation.	Optional isolated processing substrate; not automatically tumor-killing.
Ax — axons	Propagate spikes and transport cellular material.	Measure signal delivery, fatigue and cost; not bloodstream vehicles.
Sy — synapses	Regulate effective neural connectivity.	Feedback-dependent plasticity and retained response.
As — astrocytes	Maintain neural milieu, metabolism and synaptic support.	Support research culture; assess tumor-supportive signaling risk.
Ol — oligodendrocytes	Central nervous system myelination and metabolic support.	Optional mechanism to test conduction; not a general medicine.
Sc — Schwann cells	Peripheral nerve support and myelination.	Peripheral-nerve model only where justified.
Mi — microglia	Resident CNS immune surveillance and inflammatory response.	Examine neuroimmune cross-talk; not assumed safe cancer killers.

Layer 4 — therapeutic/experimental modules

Code	Module	Purpose and boundary
Ns / Nc	Neural stem cells / neural crest progenitors	Developmental candidate models; neural crest is embryonic and may migrate unpredictably.
Ef	Specified effector cells	Target-defined immune or engineered therapeutic function; validate independently.
St	Storage, scaffold, or support substrate	Supports regulated local function; does not guarantee oral survival or safe engraftment.
Se	Validated sensor / reporter	Connect a defined observable to a harmless output in early testing.
Ct	Independent safety controller	Fail-closed gating, logging, external stop and clinical escalation.

05 Stem-cell and cellular-team selection

The candidate list is deliberately broad. It is NOT a proposed mixture for injection, ingestion, or implantation. Each cell class has a different lineage, potency, persistence and risk profile. Inclusion in the Living Table does not imply use in patients.

Candidate	Possible research function	Principal safety limitation / design decision
Neural stem/progenitor cells (NSCs)	Study migration and localized delivery concepts in selected brain-tumor models; generate neural lineages in vitro.	Risk of unexpected differentiation, persistence and tumor-supportive effects. Qualified specialist use.
Neural crest progenitors (NCCs)	Study migratory development and peripheral-neural support mechanisms.	Highly variable lineage outcomes; not a default therapeutic or circulating controller.
Induced pluripotent stem cells (iPSCs)	Source well-characterized laboratory neural and non-neural cell models.	Residual pluripotency and genetic instability require careful control; not administered uncharacterized.
Embryonic pluripotent stem cells (ESCs)	Research comparator for differentiation and developmental biology.	Ethical provenance, lineage fidelity, tumorigenicity and regulation.
Mesenchymal stromal/stem cells (MSCs)	Study tissue signals, inflammation and engineered secretion in vitro.	Can exert immunosuppressive or tumor-supportive effects; not presumed cancer-preventive.
Hematopoietic stem cells (HSCs)	Contextual reference for established transplant medicine and blood lineage generation.	Not equivalent to a generalized DNA-repair or cancer-detection system.
Tissue-specific epithelial progenitors	Study normal and wounded organoid repair states.	Proliferation signals can overlap with malignant states.
Cancer stem-like cell populations	Model therapy resistance and tumor recurrence.	Research targets, not beneficial treatment ingredients.
Astrocytes / oligodendrocyte lineage	Stabilize and characterize specific neural cultures.	Context-dependent effects; non-neural cancers may respond unpredictably.
Schwann cells / peripheral neural support	Mechanistic comparison for peripheral axonal models.	Potential unwanted innervation and growth signaling.
Microglia / macrophage models	Examine inflammatory discrimination and clearance pathways.	Activation can harm normal tissue; model-specific interpretation.
Engineered T cells and NK cells	Oncology effector comparator: specific tumor recognition and killing.	Immune toxicity and off-target effects; must be indication-specific.
Dendritic/antigen-presenting cells	Potential research comparator for immune recognition.	Requires antigen validation; not a direct neural relay.
Endothelial cells / pericytes	Model vascular barriers, tissue perfusion and biodistribution.	Vessel growth can support tumors; control angiogenic effects.
Fibroblasts and cancer-associated fibroblasts	Model wound-healing vs tumor-microenvironment differences.	Some fibroblast phenotypes promote malignancy.
Patient-derived tumor organoids and healthy organoids	Paired research models to compare specificity and harm.	Model heterogeneity and limited immune/vascular representation.

SELECTED FIRST-GENERATION TEAM

Prioritize patient-derived tumor + matched healthy tissue models, a standard software controller, and a small isolated neuronal culture with astrocyte support for learning research. Do NOT combine all stem-cell populations into a single patient product.

06 Neural Lotus: axon and neuron training program

6.1 Closed-loop learning architecture

Defined sensor proxy input → encoder → cultured neuronal network → fixed or tracked decoder → harmless reporter / simulated action → outcome-contingent feedback → repeated assessment. This is a research system external to the patient. Closed-loop neuronal adaptation is supported in culture; oncology-specific performance is not yet established [R1, R2].

6.2 Distinguish five mechanisms

Process	Candidate observable	What it does NOT prove
Synaptic plasticity	Change in evoked connection strength and task accuracy.	Does not prove selective cancer recognition.
Network adaptation	Change in responses under contingent feedback.	Does not imply consciousness or unlimited learning.
Homeostatic regulation	Return of firing and viability toward acceptable ranges.	More spikes are not necessarily better.
Axonal propagation	Delivered/initiated spikes, transmission delay, failure rate.	Does not imply axons move bulk energy or patrol blood.
Support-cell / myelin adaptation	Conduction and metabolic changes in defined co-cultures.	Not evidence that making extra nerves near tumors is safe.

6.3 Preclinical training comparisons

1. Predefine a narrow simulated classification task and hold out entire donors/batches; define success before viewing outcomes.
2. Compare feedback-trained cultures with untrained cultures, stimulus-only cultures, replayed or non-contingent feedback, and non-living electronic controllers.
3. Keep the decoder fixed in the primary attribution analysis; report any software-decoder learning separately from cellular adaptation.
4. Measure response accuracy, class sensitivity, false alarms, drift, retention after rest, generalization and robustness to noise.
5. Measure axonal signal fidelity and recovery separately from overall task performance. Track morphology, viability, metabolic cost, and support-cell needs.
6. Repeat across independent biological preparations and blinded analysis; avoid treating each spike as an independent biological replicate.

6.4 Axonal operating-envelope metrics from the Living Table

Symbol	Operational measurement	Interpretation
A_{task}	Correct outputs / tested outputs, with class balance and chance level declared.	Task success, not cell potency.
R_{axon}	Distally delivered spikes / proximally initiated spikes on defined pathway.	Transmission fidelity.
$\tau_{response}$	Input-to-decoded-action delay, with an explicit event definition.	Control latency.

Symbol	Operational measurement	Interpretation
T_endurance	Period during which accuracy, fidelity and viability stay within predeclared limits.	Sustainable operation.
T_recovery	Time to recover a defined reference state following controlled workload.	Resilience.
E_correct	Whole-system energy consumed per correctly completed task.	Energy efficiency including support apparatus.
V_viable	Fraction of cells meeting a specified viability assessment.	Cell safety / culture health.
GE	1 − (energy per correct task after / energy per correct task before).	Illustrative efficiency gain; both terms measured under comparable conditions.

Trainability is distinct from cell proliferation. An increase in axon number, length, myelin or activity is not itself a therapeutic benefit. Extra innervation can be a cancer risk in some settings [R6].

07 Sensing, baseline learning and the repeated-use hypothesis

7.1 Translating “repeated damaging vibrations” into biology

The user’s poetic term describes repeated exposures or processes that injure tissue: chronic inflammation, oxidative stress, certain infections, carcinogenic exposures and harmful dysregulated signaling. Not all tissue damage becomes cancer. Each proposed biomarker must be tied to a defined clinical phenomenon and validated for that intended use. Normal mRNA is essential to life; transcript changes may serve as one possible class of measurement [R7].

7.2 Candidate inputs, not diagnoses

Signal family	Possible research proxy	Major confounder
Inflammation	Longitudinal cytokine or inflammatory marker patterns.	Infections, injuries and many benign conditions.
Tissue damage	Validated cell-injury biomarkers for a specific organ.	Exercise, procedures and normal turnover.
Tumor-derived material	Indication-specific ctDNA / cell-free DNA or molecular signature.	Very low disease prevalence, shedding variability and false alarms.
RNA / proteins	Expression patterns selected for a particular tumor setting.	Normal cellular responses and sample variability.
Imaging / pathology	Imaging or biopsy review as independent reference when clinically appropriate.	Reader variability, invasive sampling and sampling error.
Time / medication history	Treatment, infections, exposures and laboratory timing.	Confounding by interventions, missed follow-up.

7.3 How personalization would accumulate

The near-term baseline resides in auditable software, not an assumption that daily ingestion adds neurons. The model maintains a time-stamped patient history, quantifies personal variation, identifies drift and requests independent confirmation of suspicious findings. A future living, resident or repeated-dose version is a distinct product hypothesis requiring its own delivery, persistence and safety studies.

7.4 Separate questions for prevention and therapy

- Prevention/early detection: Does a screening or monitoring strategy reduce clinically meaningful cancer harm, not merely increase the number of positive tests?
- Localized therapy: Does a defined agent safely improve tumor control in an identified cancer and patient population?
- Tissue healing: Does healthy tissue recover without accelerating malignant growth or masking residual disease?

08 Proposed system architecture and safety boundaries

Module	Design responsibility	Independent safety gate
Sensor / data collector	Measure preselected, traceable laboratory or clinical inputs.	Data-quality controls and clinical reference standards.
Software reference model	Primary baseline comparator; personalization and decision calibration.	Versioning, audit logs, clinician review.
Optional neural Lotus	Research classification/control using contained neural cells.	No direct authority to deliver medicine.
Cytotoxic / secretion effector	Only a specific product already justified for the chosen indication.	Separate release authorization, bounded potency, external stop.
Support / scaffold	Localize cells or support the ex vivo circuit where appropriate.	No uncontrolled persistence or tissue invasion.
Safety controller / Heart	Monitor outputs, enforce allowed states and fail closed.	Independent from the learning model.
Clinical investigator	Interpret results and decide treatment eligibility.	Regulated consent, clinical supervision, rescue procedures.

Safety-by-design requirements

- Treat the cancer-targeting and regenerative actions as separate indications; never deploy growth-promoting effectors solely because a classifier sees damage.
- Use defined molecular targets and confirm tumor selectivity against matched normal tissue.
- Investigate independent deactivation/elimination controls, restricted persistence and traceable cell identity; never assume complete elimination.
- Watch for neural-tumor interactions, unwanted innervation, immunogenicity, off-target secretion, migration, tumorigenicity and treatment resistance [R6, R9].
- Keep any living neural controller contained outside the body until its safety and incremental utility are established. Do not use oral or daily self-administered neurons in a person.

09 Preclinical test plan: laboratory stages

WP0 — Intended use, standards and study design

Choose a single indication with an oncology collaborator (example research candidate: recurrent high-grade glioma, given existing neural stem-cell delivery investigations [R3]). Build a biomarker-to-decision causal model and a formal target product profile. Prespecify negative controls, biological replicates, safety signals and go/no-go criteria.

Inputs	Deliverables	Advance only if
Indication, clinical comparator, target tissue, approved/experimental treatment context.	Target product profile, risk register, statistical analysis plan and independent advisory review.	Use case is narrow, identifiable and medically relevant.

WP1 — Retrospective baseline and diagnostic discrimination

Use de-identified, appropriately licensed longitudinal datasets with independent ground-truth labels. Compare patient-specific software models against published clinical or simple statistical baselines, including temporal and external holdouts. Analyze calibration, per-patient false alarms, positive predictive value at realistic disease prevalence and subgroup stability. No treatment decisions are generated.

WP2 — Contained cultured-neuron training

In a qualified laboratory, couple a small neural culture to a non-harmful sensor surrogate and recording interface. Test whether feedback improves classification or switching of a harmless reporter. Keep identical input data for software and neural arms, and compare trained versus untrained and noncontingent-feedback controls. Measure the full axonal operating envelope and culture safety.

WP3 — Specificity in paired tissue models

Use matched healthy, inflamed/damaged and malignant cell or organoid models to assess whether the intended effector treats malignant states without suppressing necessary healthy repair. Patient-derived organoids and other models are available through NCI resources [R10]. Do not use animal or human interventions as substitutes for missing cell-culture specificity data.

WP4 — Localized therapeutic-cell concept and containment

If a cell therapy is scientifically justified, separately evaluate identity, localized action, biodistribution, unintended differentiation, abnormal proliferation, persistence, immunogenicity and controlled termination in specialist-led nonclinical models. The effectors are tested independently of the ChristEye learning component first. Investigations using animals, when justified, require oversight and humane endpoints.

WP5 — Integrated simulation and independent replication

Integrate only the validated sensing, controller and effector interfaces. Test failure modes in simulation or contained ex vivo settings; stress-test missing data, wrong classifications, drift, incomplete shutdown and noisy measurements. Seek replication by a different laboratory, with locked methods and independent analysis.

10 Experiment matrix and decisive comparisons

Study comparison	Primary question	Key endpoint / failure signal

Personalized vs non-personalized software	Does within-person history improve detection?	External sensitivity/specificity, calibration, alarms per patient-time; no hidden-label leakage.
Trained neurons vs untrained / replay controls	Is improvement attributable to meaningful biological feedback?	Held-out accuracy and retained performance across independent cultures.
Neural controller vs electronic controller	Does biology add practical value?	Net correct decisions, latency, durability, energy cost and operating complexity.
Malignant vs damaged vs healthy tissue	Is the intervention selective?	Cancer-cell response without undue damage to healthy/repairing tissue.
Effector alone vs integrated controller	Does connection to Sylph help?	Added benefit, no uncontrolled therapy activation.
Shutdown triggered vs normal operation	Can activation and persistence be restrained?	Residual active cells, escape, unwanted survival and delayed harm.
Cross-lab replication	Does the result generalize beyond one setting?	Prespecified effects reproduced with independent materials and analysts.

Statistical credibility and reproducibility

- Pre-register the hypothesis, independent experimental unit, primary endpoint, analysis plan, missing-data approach and stopping rules before the decisive study.
- Determine sample sizes using pilot variance and a clinically meaningful effect; do not invent participant counts or assume a high success probability.
- Blind outcome assessors where feasible. Randomize allocations and independently repeat key results across days, culture batches and donors.
- Report confidence intervals, calibration, sensitivity, specificity, precision/positive predictive value, false-negative cost and subgroup results.
- For neural systems report biological replicate count separately from neuron count, spike count and observation count.
- Publish null and adverse results; establish independent external data and assay verification.

11 Decision gates and program stopping rules

Gate	Evidence required to advance	Stop / redesign condition
G0: fit-for-purpose	Indication, biomarker and comparator clearly identified.	No reliable disease-relevant signal or wrong target indication.
G1: valid prediction	Independent external performance better than comparator at an acceptable false-alarm burden.	Overfitting, hidden confounding or unacceptable missed disease.
G2: neural added value	Causal improvement from actual neuronal training, not just software adaptation.	No advantage, instability or excess metabolic cost.
G3: tumor selectivity	Favorable malignant vs healthy/damaged tissue response.	Tumor growth acceleration, toxicity or poor specificity.

Gate	Evidence required to advance	Stop / redesign condition
G4: controllability	Potency, identity, persistence and independent stop mechanism validated.	Uncontrolled proliferation, migration, unintended effects or unreliable shutdown.
G5: translation	Independent reproducibility, favorable risk-benefit, regulatory readiness.	No clinically meaningful incremental value or unresolved safety hazards.
G6: clinical gate	Regulatory authorization, IRB approval, safety monitoring and appropriate consent.	No IND clearance/activation or clinical oversight.

12 Living-cell product feasibility, storage and dosing boundaries

An eventual therapeutic product may contain well-defined cells, a validated carrier and controlled-release components. The phrase “daily refrigerated vial” is a delivery hypothesis, not a selected formulation. Refrigeration and an expiration label alone do not demonstrate stability or safety.

Product attribute	Required research and manufacturing demonstration
Identity / purity	Precisely defined cell population; absence of unintended lineages or contamination.
Potency	Release assay causally linked to intended therapeutic function and safety, with lot-to-lot control [R11].
Viability	Storage stability and post-administration function within a validated range.
Sterility / adventitious agents	Validated controls and release criteria under applicable manufacturing standards.
Genetic / phenotypic stability	No concerning drift, transformation or unanticipated therapeutic product expression.
Dose / repeat exposure	Justified from pharmacology/toxicology; daily dosing is not assumed beneficial.
Route / biodistribution	Clinically defensible delivery and proof of target localization; ingestion of neurons not established.
Persistence / reversibility	Validated clearance/stop strategy and long-term monitoring appropriate to cell type.
Packaging / cold chain	Supported by measured shelf life, excursion testing and traceable handling.

ADMINISTRATION BOUNDARY

No patient should ingest, inject, implant or otherwise self-administer neurons, stem cells or experimental living-cell mixtures. A future clinical route and dose can only be selected after appropriate nonclinical evidence and regulatory review.

13 Clinical-trial development pathway — conditional

This section is an outline for scientific planning, not a trial protocol or permission to begin patient studies. FDA regulates many proposed cell and gene therapies through an Investigational New Drug (IND) application; ethics review and qualified investigators are essential [R4, R8].

Milestone	Illustrative trial/research question	Primary evidence and safeguards
Pre-IND meeting and CMC planning	What is the exact product, indication, manufacturing method, potency test and main risk?	Formal FDA interaction, product characterization, tox package, clinical protocol concept.
First-in-human / Phase I (only if justified)	Is a narrowly targeted therapy tolerable in patients with the selected disease?	Safety, dose-limiting toxicities, biodistribution/persistence, stopping rules, pharmacodynamic signals.
Phase II	Does the therapeutic approach show preliminary clinically meaningful activity beyond expected background?	Prespecified response/progression outcome, healthy-tissue harm, immune events, comparator adequacy.
Phase III, if supported	Does the intervention improve patient outcomes versus accepted care?	Randomized adequate trial, survival/function where suitable, adverse events, patient-reported outcomes.
Long-term follow-up	Are delayed growth, recurrence, immune, genetic or off-target effects evident?	Risk-based longitudinal follow-up; FDA guidance for relevant gene therapies [R12].
Future prevention studies — separate indication	Does screening or prevention benefit otherwise healthy people?	Only after separate evidence of effectiveness and exceptionally favorable safety; no substitute for standard screening.

Clinical governance and oversight requirements

- Disease indication chosen with oncology investigators; inclusion/exclusion criteria justified by existing cancer care and anticipated risks.
- Independent institutional review board, patient-centered informed consent, adverse-event reporting, and data monitoring/safety committee when appropriate.
- Independent confirmation of cancer diagnosis; no treatment solely on an experimental neural classifier output.
- Standard-of-care treatment protected: control groups and rescue plans must not deny effective established treatment.
- Patient privacy, biomarker-data governance, biospecimen consent, monitoring for late effects and transparent registration of applicable trials.

14 Principal risks and mitigation requirements

Hazard	Why it matters	Mandatory control before advancement
Tumor promotion by nerves or stromal signals	Cancer can exploit nerve growth and local trophic pathways [R6].	Tumor co-culture risk assessment, no default pro-growth intervention.
Unwanted stem-cell growth / residual iPSC	Persistent dividing cells can form abnormal tissue.	Lineage identity, unwanted-growth studies, persistence and removal plan.

Hazard	Why it matters	Mandatory control before advancement
False cancer alarm	Triggers unnecessary biopsy, treatment and anxiety.	External validation, confirmation and clinician decision authority.
False negative	May delay necessary care.	Coverage limits, uncertainty outputs and continued standard screening.
Immune toxicity or infection	Living-cell products can provoke serious harm [R9].	Validated microbiological controls, toxicity monitoring and escalation.
Cell migration / unanticipated engraftment	Cells may persist in non-target tissue.	Biodistribution testing, local delivery, safe clearance and surveillance.
Neural controller drift or failure	Biological networks adapt, age and change.	Drift limits, periodic recalibration, safe default and independent off switch.
Refrigeration misuse / unstable batch	Cold storage alone says little about viability or sterility.	Release specification, stability data, traceability and recall procedures.
Overinterpreting spiritual language	Philosophical labels can obscure experimentally measurable claims.	Declare scientific observable for every proposed mechanism.

15 Program sequence and resourcing

Workstream	Earliest sensible sequence	Lead capabilities
A. Clinical problem and study design	Start immediately; narrow to one cancer indication.	Medical oncologist, translational scientist, biostatistician, patient advisor.
B. Data baseline model	After data access/ethics approvals; first quantitative test.	Data scientist, bioinformatician, clinical lab specialist.
C. Contained neural training	Parallel research, entirely external to patients.	Neuroscientist, electrophysiologist, cell-culture specialist, engineer.
D. Cellular disease models	After target/assay selection; work with tumor and normal organoids.	Oncology cell biologist, pathologist, organoid specialist.
E. Therapeutic effector selection	Only after mechanism and target are defined.	Cell-therapy specialist, toxicologist, manufacturing scientist.
F. Clinical translation	Only after G0–G5 and FDA/IRB readiness.	Clinical PI, GMP/CMC, regulatory affairs, medical monitor.

Suggested research collaborations

- Cancer center / organoid core for an indication-specific tumor/healthy-tissue model.
- Electrophysiology or biohybrid computing group for blinded neuron-vs-software comparison.
- Cell-therapy translational group for therapeutic effector identity, containment and nonclinical safety.
- Regulatory and clinical-trial experts for intended-use definition, FDA pre-IND discussions, ethics and long-term follow-up.

Planning packages for funding proposals

7. Package 1 — Sylph data demonstrator: de-identified biomarkers, calibration, negative controls, locked evaluation.

8. Package 2 — Living Table neural demonstrator: defined culture and neural readout, feedback controls, axon reliability and energy accounting.
9. Package 3 — Tumor-selectivity dossier: independently verified malignant vs healthy tissue outcomes and safety findings.
10. Package 4 — Translational feasibility assessment: manufacturing, regulatory pathway, clinical need and rejection criteria.

16 Data management, governance and reproducibility

- Maintain immutable records of study version, protocol changes, data provenance, instrument calibration, trained models, neural-culture lineage and cell-product lots.
- Use secure, access-controlled clinical data; minimize patient-identifiable information and define retention and consent boundaries.
- Separate hypothesis-generation datasets from locked confirmatory studies and prevent the same donors from leaking between partitions.
- Report incidents, failed controls, contamination, attrition, adverse findings and negative results in addition to promising results.
- An independent scientific advisory group should review go/no-go decisions, especially where therapeutic cells or human testing are considered.

17 What would count as a breakthrough — and what would falsify the program?

Finding	Meaning
Reproducible biomarker discrimination better than ordinary baselines	Supports the baseline-learning concept, not a cancer cure.
Neural-culture advantage over software on a narrow clinical surrogate	Supports a potential biohybrid component; still requires reliability/cost justification.
Therapeutic selectivity in matched cancer and healthy organoids	Supports the selected effector under test, not universal prevention.
Independent fail-closed control of therapeutic-cell activity	Necessary safety evidence, not proof of complete safety.
No incremental neural benefit; software is superior	Preserve the patient baseline and abandon the living-controller component for that task.
Tumor promotion, unsafe persistence, or no tumor selectivity	Stop the therapeutic-cell branch; reassess the target and mechanism.
Consistent poor predictive value in the intended population	Do not advance a cancer-screening indication.
Any serious uncontained growth or unexpected pathology	Suspend progression pending independent safety review.

18 Selected scientific and regulatory references

R1 Kagan et al. 2022, Neuron. In vitro neuronal learning using closed-loop feedback. https://www.sciencedirect.com/science/article/pii/S0896627322008066

R2 Khajehnejad et al. 2025. Dynamic network plasticity and sample efficiency in biological neural cultures. https://pmc.ncbi.nlm.nih.gov/articles/PMC12320521/

R3 NCI study NCT05139056: neural stem cell therapy for recurrent high-grade gliomas.
https://www.cancer.gov/research/participate/clinical-trials-search/v?id=NCI-2022-10170

R4 FDA: Investigational New Drug application requirements.
https://www.fda.gov/drugs/types-applications/investigational-new-drug-application-ind

R5 FDA: Oncology Cell and Gene Therapy program.
https://www.fda.gov/about-fda/oncology-center-excellence/oncology-cell-and-gene-therapy

R6 Yaniv et al. 2024. Targeting the peripheral neural-tumour microenvironment for cancer therapy.
https://www.nature.com/articles/s41573-024-01017-z

R7 NCI: cancer screening tests and limitations of multi-cancer detection.
https://www.cancer.gov/about-cancer/screening/screening-tests

R8 FDA: Frequently Asked Questions — Developing Potential Cellular and Gene Therapy Products (Aug. 2026).
https://www.fda.gov/regulatory-information/search-fda-guidance-documents/frequently-asked-questions-developing-potential-cellular-and-gene-therapy-products

R9 FDA: risks from unapproved regenerative medicine therapies.
https://www.fda.gov/vaccines-blood-biologics/consumers-biologics/important-patient-and-consumer-information-about-regenerative-medicine-therapies

R10 NCI Patient-Derived Models Repository: organoids, matched model resources.
https://dctd.cancer.gov/drug-discovery-development/reagents-materials/pdmr

R11 FDA: Potency Assurance for Cellular and Gene Therapy Products (Nov. 2026).
https://www.fda.gov/regulatory-information/search-fda-guidance-documents/guidance-industry-potency-assurance-cellular-and-gene-therapy-products

R12 FDA: Long-Term Follow-up after administration of human gene therapy products.
https://www.fda.gov/regulatory-information/search-fda-guidance-documents/long-term-follow-after-administration-human-gene-therapy-products

R13 NCI: cancer screening overview — harms and evidence limitations of multi-cancer detection.
https://www.cancer.gov/about-cancer/screening/hp-screening-overview-pdq

R14 NIH/NHLBI: clinical-trial purposes and phase framework.
https://www.nhlbi.nih.gov/research/clinical-trials/how-studies-work

R15 ChristEye Living Table & Energy Framework v0.5 (internal project source), Nov. 2, 2026.
Internal research draft; not peer-reviewed.

R16 ChristEye Sylph Cancer Surveillance Concept v0.2 (internal project source), Nov. 2, 2026.
Internal research draft; not peer-reviewed.

Study interpretation

External references establish components of neuroscience, oncology, biological safety and regulatory procedure. None independently validates the ChristEye Sylph integrated concept. Research plans, names, predicted benefits and commercialization claims remain hypotheses until confirmed by suitably controlled studies.

19 Glossary and definitions

Term	Project meaning
Biohybrid	Device or research system combining living biological components with external electronic sensing or computation.
Ex vivo / in vitro	Investigations conducted outside a patient in a controlled experimental setting.
Patient-specific baseline	Longitudinal reference model of a person’s measured biological

Term	Project meaning
	variability.
Effector cell	Cell with a defined therapeutic or reporter function; not necessarily a neuron or stem cell.
Stem cell	A cell population with specific self-renewal/differentiation properties; different stem cells are not interchangeable.
Axon / synapse	Neural signal-transmission process and cellular junction involved in effective communication.
Potency assay	Test relating cell-product performance to its intended biological function.
Selectivity	Differential effect on malignant versus healthy or injured tissue.
IND / IRB	U.S. investigational new drug submission and independent human-subjects ethics oversight.
Go/no-go gate	Evidence requirement used to authorize or halt a research phase, not a prediction of success.

END OF RESEARCH DEVELOPMENT SPECIFICATION

Document control: Version 1.0 | 2 November 2026 | Working proposal for scientific review | Not a human-use product specification or investigational trial protocol.