

SEED STENCILS: HIGH-ORDER FINITE DIFFERENCES AND RBF-FD THROUGH MATERIAL EDGES TOO STEEP FOR THE GRID

BRADLEY P. MARTIN

ABSTRACT. Between a material whose parameters vary smoothly on the scale of the grid and one with a jump lies the case of an edge that is smooth but changes over a distance δ far smaller than the grid spacing h . Standard high-order finite differences sample the coefficients at the nodes and lose their order there until $h \lesssim \delta$, and the interface-aware stencils built for a jump do not apply, since there is no jump. We build stencils on the same equispaced (1-D) or scattered (2-D) nodes whose basis functions are not monomials but *seeds*: the $t = 0$ profiles of solutions of the wave equation that are polynomial in time, continued through the edge by ordinary differential equations in the coordinate normal to it. The seeds form an extended complete Chebyshev system for every δ , so the stencil solve is nonsingular, and as $\delta \rightarrow 0$ they reduce to the translated polynomial basis of the interface-aware stencils, so one construction covers the jump, the sub-grid edge and the resolved edge. In one dimension, standard fourth-order finite differences have a knee at $h = \delta$ while the seed stencils are fourth order at every resolution with the errors of the jump case. In two dimensions, for the elastic wave equation discretised with radial basis function-generated finite differences (RBF-FD) on scattered nodes, the seeds of a straight edge reduce to triangular systems of ordinary differential equations in the normal coordinate; the seed-augmented stencils are stable at the standard hyperviscosity, cut the error through an unresolved edge by up to an order of magnitude or more on the node sets tested, and lead to the rule: seed when $\delta \leq h$. The same seeds marched along the true normal handle a curved edge at the same accuracy on those node sets.

1. INTRODUCTION

[Stub, #55: the sub-grid smooth edge between “smooth” and “jump”; what standard high-order FD does there (the knee at $h = \delta$); the thesis: one construction from the jump (algebra) through the ODE march to the resolved edge (Fornberg); contributions; relation to prior work from LITERATURE.md (#53); outline.]

1.1. Relation to prior work. *[Stub, #53 supplies this paragraph; #55 places it.]*

Date: September 20, 2026.

2. SETTING

[Stub, #55: the 1-D first-order system $\rho u_t = f_x$, $f_t = Ku_x$; the 2-D elastic system in five fields; equispaced and scattered node sets; RBF-FD with polynomial augmentation, hyperviscosity, RK4; the interface-aware stencils of the jump as background (translated basis, continuity matrices, the two straddling rows, curvature by local expansion); what the implementation leaves out (triple stencils, variable Lamé parameters, the FD/RBF hybrid), stated as scope.]

3. SEEDS IN ONE DIMENSION

3.1. Seeds as the profiles of solutions polynomial in time. [Stub, #56: $u_{tt} = L_u u$ with $L_u = \rho^{-1} \partial_x K \partial_x$; the chain $\phi_0 = 1$, $K\phi'_1 = K(x_e)$, $L_u \phi_k = k(k-1)c_e^2 \phi_{k-2}$ with $\phi_k(x_e) = \phi'_k(x_e) = 0$; the even/odd (u/f) alternation; first-order form so that K is never differentiated.]

3.2. Nested kernels and the jump as the limit. [Stub, #56: $\{1\} \subset \ker L_u \subset \{L_u g = \text{const}\} \subset \ker L_u^2 \subset \dots$; the seed weights converge to the interface-aware weights at first order in δ .]

3.3. Well-posedness. [Stub, #56: the seeds are an extended complete Chebyshev system (Pólya, Karlin–Studden), so the per-stencil solve is non-singular on any distinct nodes for every δ .]

3.4. Why the Taylor route does not reach a sub-grid edge. [Stub, #56: notes §1.5; the quasi-Trefftz comparison.]

3.5. Stencil assembly and cost. [Stub, #56: the Fornberg-style solve on the seeds; one ODE march per stencil.]

4. ONE-DIMENSIONAL RESULTS

[Stub, #57: test problem (medium, contrast, pulse, $\delta \in \{0, 0.0025, 0.01\}$); the Fourier pseudo-spectral reference; the knee of standard FD4 at $h = \delta$; seeds fourth order at every n with the jump case's errors; condition numbers 20–60 from $\delta = 10^{-5}$ to 1.]

5. SEEDS IN TWO DIMENSIONS

5.1. Design: a straight edge. [Stub, #58: the x' -polynomial ansatz turns the 2-D chain into triangular systems of 1-D ODEs in the normal coordinate; tangential dependence stays polynomial; tractions as flux variables.]

5.2. The elastic seeds. [Stub, #58: all 30 seeds of one stencil marched as one 600-state system anchored at the evaluation node; constant-material, jump-limit, residual and conditioning checks.]

5.3. Stencil assembly. [Stub, #58: 19-node seed rows for the elastic operator; Δ^3 rows on the naive 30-node footprint with the seeds annihilated; the seeded region (19δ) and its trim.]

5.4. Curved edges. *[Stub, #58: route (a), the seeds marched along the true normal through the foot point, zeroth order in curvature, the same approximation as the jump stencils; what route (b) would be.]*

6. TWO-DIMENSIONAL RESULTS

6.1. Test problems and references. *[Stub, #59: 1-D spectral at normal incidence; Fourier-in- x for oblique; product-grid for the curved $\delta > 0$; the 122,500-node jump-aware run for the curved $\delta = 0$.]*

6.2. Naive baseline. *[Stub, #59: the resolution floor hides the edge error in v ; the spurious u separates unresolved from resolved edges $10\times$ more sharply.]*

6.3. Stability. *[Stub, #59: spectra at $n = 900$ and 2500 , standard γ , every δ , flat and curved.]*

6.4. The flat sweep and the rule. *[Stub, #59: seeds fourth order through $\delta = 0.0025$ at every n , $20\times$ below naive in v and $67\times$ in spurious u at 19,600 nodes; crossover at $h \approx \delta$; a resolved edge $1.4\text{--}2\times$ worse seeded; the rule: seed when $\delta \leq h$.]*

6.5. Oblique incidence. *[Stub, #59: seeds $1.4\text{--}1.9\times$ over naive at 26.6° , the ablation worse than naive, every scheme about 2.5 from the converted- S floor, reported as it is.]*

6.6. The curved edge. *[Stub, #59: jump stencils at the resolution floor at every n ; seeds equal the flat seeds at their own floor, $2.2\text{--}7\times$ ($\delta = 0.005$) and $3\text{--}12\times$ ($\delta = 0.0025$) below naive; where route (a)'s geometry first shows; the fine-end rate and the trim; costs.]*

7. DISCUSSION

7.1. Limitations. *[Stub, #60: FD4 / degree 3 only, one contrast, at most 19,600 nodes, the flat translation symmetry unexploited, one march per stencil, the port's exclusions.]*

7.2. Future work. *[Stub, #60: curvilinear evaluation of the marched seeds; seeded-width tuning; the seeds' true Δ^3 as hyperviscosity right-hand sides; Mühlbach's recurrence in place of the dense solve; higher order; other contrasts; thin bands; variable Lamé parameters; 3-D; the oblique train on the curved geometry.]*

7.3. Conclusions. *[Stub, #60: the rule and the one-construction claim, in #53's wording.]*

ACKNOWLEDGMENTS

[Stub, #62: Bengt Fornberg, for the 2016-era work this construction extends (the dissertation and the interface treatment of [MF17a]); the AI-assistance statement crediting Claude (Fable 5.1) for the re-derivation, the code and the drafting.]

REFERENCES

- [AS55] D. N. de G. Allen and R. V. Southwell, *Relaxation methods applied to determine the motion, in two dimensions, of a viscous fluid past a fixed cylinder*, The Quarterly Journal of Mechanics and Applied Mathematics **8** (1955), no. 2, 129–145.
- [Bac62] George E. Backus, *Long-wave elastic anisotropy produced by horizontal layering*, Journal of Geophysical Research **67** (1962), no. 11, 4427–4440.
- [BCO94] Ivo Babuška, Gabriel Caloz, and John E. Osborn, *Special finite element methods for a class of second order elliptic problems with rough coefficients*, SIAM Journal on Numerical Analysis **31** (1994), no. 4, 945–981.
- [BFFB17] Victor Bayona, Natasha Flyer, Bengt Fornberg, and Gregory A. Barnett, *On the role of polynomials in RBF-FD approximations: II. Numerical solution of elliptic PDEs*, Journal of Computational Physics **332** (2017), 257–273.
- [BGL17] Lehel Banjai, Emmanuil H. Georgoulis, and Oluwaseun Lijoka, *A Trefftz polynomial space-time discontinuous Galerkin method for the second order wave equation*, SIAM Journal on Numerical Analysis **55** (2017), no. 1, 63–86.
- [BMSK21] Jure Močnik Berljavac, Pankaj K. Mishra, Jure Slak, and Gregor Kosec, *RBF-FD analysis of 2D time-domain acoustic wave propagation in heterogeneous media*, Computers & Geosciences **153** (2021), 104796.
- [BO83] I. Babuška and J. E. Osborn, *Generalized finite element methods: their performance and their relation to mixed methods*, SIAM Journal on Numerical Analysis **20** (1983), no. 3, 510–536.
- [Bro84] David L. Brown, *A note on the numerical solution of the wave equation with piecewise smooth coefficients*, Mathematics of Computation **42** (1984), no. 166, 369–391.
- [CGM10a] Yann Capdeville, Laurent Guillot, and Jean-Jacques Marigo, *1-D non-periodic homogenization for the seismic wave equation*, Geophysical Journal International **181** (2010), no. 2, 897–910.
- [CGM10b] ———, *2-D non-periodic homogenization to upscale elastic media for P–SV waves*, Geophysical Journal International **182** (2010), no. 2, 903–922.
- [CJZ26] Tao Cheng, Lili Ju, and Xiaoping Zhang, *A high-order radial basis function finite difference method for linear anisotropic elliptic interface problems*, Journal of Scientific Computing **108** (2026), no. 1, 8.
- [Cop71] W. A. Coppel, *Disconjugacy*, Lecture Notes in Mathematics, Springer, Berlin, Heidelberg, 1971.
- [DV14] Kenneth Duru and Kristoffer Virta, *Stable and high order accurate difference methods for the elastic wave equation in discontinuous media*, Journal of Computational Physics **279** (2014), 37–62.
- [FF15a] Bengt Fornberg and Natasha Flyer, *A primer on radial basis functions with applications to the geosciences*, Society for Industrial and Applied Mathematics, Philadelphia, 2015.
- [FF15b] ———, *Solving PDEs with radial basis functions*, Acta Numerica **24** (2015), 215–258.

- [FFBB16] Natasha Flyer, Bengt Fornberg, Victor Bayona, and Gregory A. Barnett, *On the role of polynomials in RBF-FD approximations: I. Interpolation and accuracy*, Journal of Computational Physics **321** (2016), 21–38.
- [FL11] Bengt Fornberg and Erik Lehto, *Stabilization of RBF-generated finite difference methods for convective PDEs*, Journal of Computational Physics **230** (2011), no. 6, 2270–2285.
- [For88a] Bengt Fornberg, *Generation of finite difference formulas on arbitrarily spaced grids*, Mathematics of Computation **51** (1988), no. 184, 699–706.
- [For88b] ———, *The pseudospectral method: accurate representation of interfaces in elastic wave calculations*, Geophysics **53** (1988), no. 5, 625–637.
- [FRR⁺06] A. Farjadpour, David Roundy, Alejandro Rodriguez, M. Ibanescu, Peter Bermel, J. D. Joannopoulos, Steven G. Johnson, and G. W. Burr, *Improving accuracy by subpixel smoothing in the finite-difference time domain*, Optics Letters **31** (2006), no. 20, 2972.
- [Ger12] G. A. Gerolymos, *A general recurrence relation for the weight-functions in Mühlbach–Neville–Aitken representations with application to WENO interpolation and differentiation*, Applied Mathematics and Computation **219** (2012), no. 8, 4133–4142.
- [GW04] Bertil Gustafsson and Per Wahlund, *Time compact difference methods for wave propagation in discontinuous media*, SIAM Journal on Scientific Computing **26** (2004), no. 1, 272–293.
- [GW25] Andreas Granath and Siyang Wang, *A hybrid numerical method for elastic wave propagation in discontinuous media with complex geometry*, Journal of Computational Physics **524** (2025), 113745.
- [HW97] Thomas Y. Hou and Xiao-Hui Wu, *A multiscale finite element method for elliptic problems in composite materials and porous media*, Journal of Computational Physics **134** (1997), no. 1, 169–189.
- [IG25] Lise-Marie Imbert-Gérard, *Local Taylor-based polynomial quasi-Trefftz spaces for scalar linear equations*, 2025, arXiv:2505.18480.
- [IGMPS25] Lise-Marie Imbert-Gérard, Andrea Moiola, Chiara Perinati, and Paul Stocker, *Polynomial quasi-Trefftz DG for PDEs with smooth coefficients: elliptic problems*, IMA Journal of Numerical Analysis **45** (2025), no. 6, 3473–3506, arXiv:2408.00392.
- [IGMS23] Lise-Marie Imbert-Gérard, Andrea Moiola, and Paul Stocker, *A space-time quasi-Trefftz DG method for the wave equation with piecewise-smooth coefficients*, Mathematics of Computation **92** (2023), no. 341, 1211–1249, arXiv:2011.04617.
- [Il’69] A. M. Il’in, *Differencing scheme for a differential equation with a small parameter affecting the highest derivative*, Mathematical Notes of the Academy of Sciences of the USSR **6** (1969), no. 2, 596–602.
- [JHY21] Jaeyong Jeong, Sanghyun Ha, and Donghyun You, *An immersed interface method for acoustic wave equations with discontinuous coefficients in complex geometries*, Journal of Computational Physics **426** (2021), 109932.
- [JZ24] Luqian Jiang and Wei Zhang, *Efficient implementation of equivalent medium parametrization in finite-difference seismic wave simulation methods*, Geophysical Journal International **239** (2024), no. 1, 675–693.
- [KMCK17] Jozef Kristek, Peter Moczo, Emmanuel Chaljub, and Miriam Kristekova, *An orthorhombic representation of a heterogeneous medium for the finite-difference modelling of seismic wave propagation*, Geophysical Journal International **208** (2017), no. 2, 1250–1264.
- [KMPS16] Fritz Kretzschmar, Andrea Moiola, Ilaria Perugia, and Sascha M. Schnepp, *A priori error analysis of space-time Trefftz discontinuous Galerkin methods for*

- wave problems*, IMA Journal of Numerical Analysis **36** (2016), no. 4, 1599–1635.
- [KS66] Samuel Karlin and William J. Studden, *Tchebycheff systems: With applications in analysis and statistics*, Pure and Applied Mathematics, vol. 15, Interscience Publishers, New York, 1966.
- [KWR22] Erik F. M. Koene, Jens Wittsten, and Johan O. A. Robertsson, *Finite-difference modelling of 2-D wave propagation in the vicinity of dipping interfaces: a comparison of anti-aliasing and equivalent medium approaches*, Geophysical Journal International **229** (2022), no. 1, 70–96.
- [LL94] Randall J. LeVeque and Zhilin Li, *The immersed interface method for elliptic equations with discontinuous coefficients and singular sources*, SIAM Journal on Numerical Analysis **31** (1994), no. 4, 1019–1044.
- [LP04] Bruno Lombard and Joël Piraux, *Numerical treatment of two-dimensional interfaces for acoustic and elastic waves*, Journal of Computational Physics **195** (2004), no. 1, 90–116.
- [M73] G. Mühlbach, *A recurrence formula for generalized divided differences and some applications*, Journal of Approximation Theory **9** (1973), no. 2, 165–172.
- [Mar16] Bradley P. Martin, *Application of RBF-FD to wave and heat transport problems in domains with interfaces*, Ph.D. thesis, University of Colorado Boulder, 2016, ProQuest 10151046.
- [MDEN92] Francis Muir, Joe Dellinger, John Etgen, and Dave Nichols, *Modeling elastic fields across irregular boundaries*, Geophysics **57** (1992), no. 9, 1189–1193.
- [MF17a] Bradley Martin and Bengt Fornberg, *Seismic modeling with radial basis function-generated finite differences (RBF-FD) – a simplified treatment of interfaces*, Journal of Computational Physics **335** (2017), 828–845.
- [MF17b] ———, *Using radial basis function-generated finite differences (RBF-FD) to solve heat transfer equilibrium problems in domains with interfaces*, Engineering Analysis with Boundary Elements **79** (2017), 38–48.
- [MFSC15] Bradley Martin, Bengt Fornberg, and Amik St-Cyr, *Seismic modeling with radial-basis-function-generated finite differences*, Geophysics **80** (2015), no. 4, T137–T146.
- [MHI08] Ken Mattsson, Frank Ham, and Gianluca Iaccarino, *Stable and accurate wave-propagation in discontinuous media*, Journal of Computational Physics **227** (2008), no. 19, 8753–8767.
- [Mic93] Ronald E. Mickens, *Nonstandard finite difference models of differential equations*, World Scientific, 1993.
- [MKG14] Peter Moczo, Jozef Kristek, and Martin Gális, *The finite-difference modelling of earthquake motions*, Cambridge University Press, 2014.
- [MKV⁺02] Peter Moczo, Jozef Kristek, Václav Vavryčuk, Ralph J. Archuleta, and Ladislav Halada, *3D heterogeneous staggered-grid finite-difference modeling of seismic motion with volume harmonic and arithmetic averaging of elastic moduli and densities*, Bulletin of the Seismological Society of America **92** (2002), no. 8, 3042–3066.
- [MN06] Ken Mattsson and Jan Nordström, *High order finite difference methods for wave propagation in discontinuous media*, Journal of Computational Physics **220** (2006), no. 1, 249–269.
- [MP18] Andrea Moiola and Ilaria Perugia, *A space-time Trefftz discontinuous Galerkin method for the acoustic wave equation in first-order formulation*, Numerische Mathematik **138** (2018), no. 2, 389–435.
- [OZ07] Houman Owhadi and Lei Zhang, *Metric-based upscaling*, Communications on Pure and Applied Mathematics **60** (2007), no. 5, 675–723, arXiv:math/0505223.

- [OZ08] ———, *Numerical homogenization of the acoustic wave equations with a continuum of scales*, Computer Methods in Applied Mechanics and Engineering **198** (2008), no. 3–4, 397–406, arXiv:math/0604380.
- [P22] G. Pólya, *On the mean-value theorem corresponding to a given linear homogeneous differential equation*, Transactions of the American Mathematical Society **24** (1922), no. 4, 312–324.
- [PL01] Joël Piraux and Bruno Lombard, *A new interface method for hyperbolic problems with discontinuous coefficients: one-dimensional acoustic example*, Journal of Computational Physics **168** (2001), no. 1, 227–248.
- [RST08] Hans-Görg Roos, Martin Stynes, and Lutz Tobiska, *Robust numerical methods for singularly perturbed differential equations: Convection-diffusion-reaction and flow problems*, Springer Series in Computational Mathematics, Springer, Berlin, Heidelberg, 2008.
- [Sam01] A. A. Samarskii, *The theory of difference schemes*, Pure and Applied Mathematics, Marcel Dekker, 2001.
- [SG69] D. L. Scharfetter and H. K. Gummel, *Large-signal analysis of a silicon Read diode oscillator*, IEEE Transactions on Electron Devices **16** (1969), no. 1, 64–77.
- [SMP⁺23] Roberto Sabatini, Alessandro Monti, Yan Pailhas, Angeliki Xenaki, and Paul Cristini, *An arbitrary-order immersed interface method for the two-dimensional propagation of acoustic and elastic waves*, Physics of Fluids **35** (2023), no. 10, 107106.
- [ST09] William W. Symes and Igor S. Terentyev, *Subgrid modeling via mass lumping in constant density acoustics*, SEG Technical Program Expanded Abstracts 2009, 2009, pp. 2572–2576.
- [SV67] M. H. Schultz and R. S. Varga, *L-splines*, Numerische Mathematik **10** (1967), no. 4, 345–369.
- [SV09] William W. Symes and Tetyana Vdovina, *Interface error analysis for numerical wave propagation*, Computational Geosciences **13** (2009), no. 3, 363–371.
- [TE03] Anna-Karin Tornberg and Björn Engquist, *Regularization techniques for numerical approximation of PDEs with singularities*, Journal of Scientific Computing **19** (2003), no. 1–3, 527–552.
- [TE06] ———, *Regularization for accurate numerical wave propagation in discontinuous media*, Methods and Applications of Analysis **13** (2006), no. 3, 247–274.
- [TE08] ———, *Consistent boundary conditions for the Yee scheme*, Journal of Computational Physics **227** (2008), no. 14, 6922–6943.
- [TS62] A. N. Tikhonov and A. A. Samarskii, *Homogeneous difference schemes*, USSR Computational Mathematics and Mathematical Physics **1** (1962), no. 1, 5–67.
- [VLTR14] Dmitry Vishnevsky, Vadim Lisitsa, Vladimir Tcheverda, and Galina Reshetova, *Numerical study of the interface errors of finite-difference simulations of seismic waves*, Geophysics **79** (2014), no. 4, T219–T232.
- [VSKO20] Zoltan Vizvari, Zoltan Sari, Mihaly Klincsik, and Peter Odry, *Exact schemes for second-order linear differential equations in self-adjoint cases*, Advances in Difference Equations **2020** (2020), 497.
- [ZL97] Chaoming Zhang and Randall J. LeVeque, *The immersed interface method for acoustic wave equations with discontinuous coefficients*, Wave Motion **25** (1997), no. 3, 237–263.

Email address: 1bradley.martin1@gmail.com