

Isochrone overturning in steady glacier flow

Bradley Paul LIPOVSKY

Department of Earth and Space Sciences, University of Washington, Seattle, USA

Correspondence: Bradley Paul Lipovsky, bp17@uw.edu

ABSTRACT. Ancient ice recovered at Allan Hills extends into the Miocene, but steep layers, age reversals and small-scale folds complicate its interpretation. Existing steady-flow theory shows that isochrones cannot overturn where horizontal speed increases upward. Full-Stokes flow over rough beds can contain local extrusion, so this pointwise condition is stronger than necessary. Here, we derive a signed order Jacobian that measures whether particles deposited in sequence retain their horizontal order along an isochrone. Stratigraphic order depends on the cumulative competition between normal and reverse vertical shear, weighted most strongly where horizontal motion is slow. Reverse shear must erase both the ordering imposed at deposition and the accumulated normal shear before a layer becomes vertical. Full-Stokes mechanics provides no universal sign guarantee, although shallow-ice flow preserves order. We derive a full-Stokes threshold, bounds for non-stagnant localized obstacles, a first-order cancellation result for periodic roughness about plug flow and a trajectory diagnostic. Idealized full-Stokes calculations then produce recumbent folds directly from an ordered surface-deposition map, without an inherited layer perturbation. These results distinguish topographic draping from overturning at Allan Hills and other old-ice sites.

Keywords: glacier flow; ice stratigraphy; recumbent folds; blue ice; full-Stokes modelling

1. INTRODUCTION

Recovering old ice is useful only where its age and stratigraphic context can be established. The European Beyond EPICA–Oldest Ice project used radar stratigraphy and ice-flow modelling at Little Dome C to target a continuous record extending beyond 1 Ma (Lilien and others, 2021; Chung and others, 2025). Antarctic blue-ice areas provide a complementary archive: surface mass loss and upward ice motion expose discontinuous climatic intervals that can be much older than conventional vertical cores (Bintanja, 1999; Spaulding and others, 2012). At Allan Hills, shallow cores first yielded a disturbed section near 1 Ma (Higgins and others, 2015), then atmospheric snapshots older than 2 Ma (Yan and others, 2019), and most recently Pliocene and Miocene ice and air with reported ages approaching 6 Ma (Shackleton and others, 2025). These cores are a focus of NSF COLDEX, which seeks to reconstruct atmospheric composition and Antarctic climate beyond the continuous 800 ka ice-core record.

These records are stratigraphically difficult to interpret. Allan Hills ice is strongly thinned and locally out of order. The ALHIC1901 age–depth relation includes multiple reversals, while the deepest ice can represent very large time intervals per unit core length (Shackleton and others, 2025). Three-dimensional electrical conductivity measurements resolve mean layer dips of approximately 29° and 69° in two recent Allan Hills cores (Kirkpatrick and others, 2025). Microstructural observations from ALHIC1901 indicate pervasive simple shear, strongly elongated bubbles, crystal-orientation stripes, and centimetre-scale kink and z-folds (Stoll and others, 2026). Radar and flow modelling independently show that the duration and continuity of the archive vary sharply across the blue-ice area and with basal geometry (Kehrl and others, 2018). A recent three-dimensional full-Stokes model predicts near-stagnant basal ice upstream of the local nunatak and faster flow into the Cul-de-sac from a distinct accumulation environment (Manos and others, 2026). Deformation and transport history therefore control which ages are present, how they are ordered and how much time a sample averages.

A basic unresolved question is whether a fixed bed obstacle in an otherwise steady glacier can create the finite-amplitude disturbance required for a recumbent fold. Flow over topography certainly bends internal layers. Full mechanical calculations predict draping over long-wavelength topography and overriding of shorter-wavelength relief (Hindmarsh and others, 2006). Rough-bed Stokes solutions can also contain local extrusion, meaning that horizontal speed increases downward over part of the ice column (Gudmundsson, 1997a; Waddington, 2010). At larger bed slopes, separated secondary circulation is possible even in creeping

53 flow (Gudmundsson, 1997b). These results appear, at first sight, to undermine the common intuition that
 54 a steady obstacle can produce only a standing layer disturbance rather than a recumbent fold.

55 These statements concern different parts of the layer geometry. A material tangent is an infinitesimal
 56 line segment joining neighboring particles within a layer. Flow around an obstacle can bend the layer and
 57 rotate this segment locally. The layer remains ordered, however, if particles deposited at successively larger
 58 upstream positions also occur at successively larger downstream positions everywhere along an equal-age
 59 surface. A recumbent fold begins only where that horizontal ordering vanishes and then reverses. Curvature
 60 and local tangent rotation can therefore occur without an order reversal.

61 The local theory of Waddington and others (2001), hereafter WBA, formalized how a pre-existing wrinkle
 62 is flattened by longitudinal extension or overturned by shear. WBA also stated the key qualitative principle
 63 that layers spending their entire history in a steady-state flow field are not passively folded; a transient
 64 process must first create a disturbance relative to the steady stratigraphy. Jacobson and Waddington (2004)
 65 replaced the local criterion with finite-strain integration along trajectories, and Jacobson and Waddington
 66 (2005) showed that divide migration can place an inherited Raymond arch in a shear regime that turns
 67 it into a recumbent fold. For steady stratigraphy, Parrenin and others (2006) introduced logarithmic flux
 68 coordinates for analytical age solutions, and Parrenin and Hindmarsh (2007) generalized the construction
 69 to horizontally non-uniform velocity profiles. Their Appendix A showed that an isochrone cannot pass
 70 through vertical when horizontal velocity increases monotonically upward. Corrections for flow-induced
 71 layer thinning date at least to Nye (1963) and the early depth–age model of Dansgaard and Johnsen (1969).
 72 The same characteristic framework subsequently underpinned numerical age solvers (Hindmarsh and others,
 73 2009; Parrenin and others, 2025) and an explicit vertical thinning function (Parrenin, 2013; Parrenin and
 74 others, 2025).

75 Monotonic upward speed is sufficient but not necessary for preserving order because full-Stokes glacier
 76 flow does not impose $u_z \geq 0$ pointwise. The relevant quantity is cumulative: how much reverse shear
 77 can a particle experience before the depositional ordering of the isochrone is lost? Here, we derive a
 78 physical-coordinate representation of the signed inverse vertical-thinning factor and interpret it as horizontal
 79 order along an isochrone. The resulting budget separates surface-fast and reverse vertical shear. We then
 80 decompose the reverse-shear contribution using the full-Stokes stress balance, identify regimes in which
 81 glacier mechanics preserves order, and obtain a sufficient no-overturning bound for localized obstacles and
 82 a first-order cancellation result for periodic small-amplitude roughness. Finally, we formulate a trajectory

83 diagnostic and construct steady full-Stokes examples in which surface-generated isochrones overturn without
 84 an inherited perturbation.

85 The analysis builds directly on established results. The monotonic-shear no-overturning condition is due
 86 to Parrenin and Hindmarsh (2007); the order Jacobian derived below is the reciprocal of the vertical
 87 thinning function developed by Parrenin (2013) and implemented by Parrenin and others (2025); the local
 88 wrinkle criterion and finite-strain interpretation are due to WBA and Jacobson and Waddington (2004);
 89 and rough-bed extrusion is established (Gudmundsson, 1997a). The paper first relates these descriptions of
 90 folding, then derives the exact order budget, examines its mechanical constraints, and tests its two predicted
 91 overturning routes in idealized calculations. The distinct step is to write the sign of inverse thinning as an
 92 exact positive–negative u_z/u^3 shear budget. This form extends steady-isochrone theory without introducing
 93 a new state variable or claiming a universal no-folding theorem.

94 **2. EXISTING CRITERIA FOR STRATIGRAPHIC FOLDING**

95 **2.1. Passive and active folding**

96 Glacier stratification is a material structure produced at the surface and subsequently deformed by flow.
 97 Structural glaciology distinguishes passive deformation, in which layers are markers that do not alter the
 98 velocity field, from active folding, in which rheological or anisotropic contrasts feed back on stress and
 99 motion (Hudleston, 2015; Jennings and Hambrey, 2021). The distinction matters because a prescribed
 100 smooth velocity field can amplify or overturn an existing perturbation without selecting its wavelength,
 101 whereas active deformation can generate disturbances and select spatial scales.

102 Microstructural observations show where stratigraphy and rheology may be mechanically coupled. At
 103 GISP2, crystal-orientation stripes coincide with localized shear and grain-scale folding, and spatial variation
 104 in stripe density may produce larger viscosity contrasts (Alley and others, 1997). Numerical models predict
 105 that anisotropy and convergent flow can produce folds spanning scales from cloudy bands to radar reflectors
 106 (Bons and others, 2016, 2025; Zhang and others, 2024). The analysis below instead treats age as a passive
 107 scalar in a prescribed velocity field. It gives a necessary kinematic test for folding, but it does not describe
 108 the growth of an active instability.

109 **2.2. The WBA local wrinkle criterion**

Let $m = \delta z / \delta x$ be the slope of an infinitesimal material-line segment $\delta \mathbf{x}$ in a two-dimensional velocity field
 (u, w). The continuum-kinematic relation $D\delta \mathbf{x} / Dt = (\nabla \mathbf{u})\delta \mathbf{x}$ describes the deformation of that segment

(Gurtin and others, 2010). Taking the ratio of its vertical and horizontal components gives

$$\frac{Dm}{Dt} = w_x + (w_z - u_x)m - u_z m^2. \quad (1)$$

Equation (1), the Riccati form of the material-line equation used by WBA, is therefore standard continuum kinematics. WBA's distinctive step was to compare the rotation of a disturbed tangent with the rotation of the local steady isochrone tangent m_0 . A second equal-rotation direction separates disturbances that initially flatten toward m_0 from those that rotate away from it. Their shear number expresses the competition between shear-driven rotation and longitudinal extension.

Equation (1) is local. Applied pointwise along a complete curve, it contains the exact tangent kinematics of that curve. The published WBA stability diagnostic, however, does not by itself determine whether an entire equal-age surface loses its horizontal ordering. WBA emphasized this limitation over rough topography: a disturbed tangent can open relative to the local steady isochrone on the lee side of a bump while merely recovering deformation accumulated on the stoss side (Waddington and others, 2001).

2.3. Finite strain and steady isochrone geometry

Jacobson and Waddington (2004) integrated finite deformation along particle paths rather than relying on a homogeneous instantaneous velocity gradient. This is the appropriate framework for deciding whether an inherited open fold becomes recumbent after a long and spatially variable strain history. Their later divide-migration calculation supplied a concrete source of the inherited perturbation (Jacobson and Waddington, 2005).

A separate line of analysis concerns the geometry of isochrones generated continuously in a steady flow. Parrenin and others (2006) introduced logarithmic total-flux and normalized-streamfunction coordinates in which particle trajectories are straight characteristics when the vertical velocity shape is horizontally uniform. Hindmarsh and others (2006) compared streamlines and isochrones over basal topography and showed that they are close when

$$\frac{\dot{a}}{U} \frac{L}{H} \ll 1, \quad (2)$$

where $\dot{a}$ is accumulation, U a horizontal speed, L the forcing wavelength and H the ice thickness. Parrenin and Hindmarsh (2007) generalized the characteristic analysis to horizontally non-uniform velocity profiles and wrote isochrone slope as an instantaneous normalized-streamfunction contribution plus a path-history term. In their notation, the sign of $1 - \alpha$ determines whether neighboring particles preserve their order, and $\alpha > 1$ corresponds to an isochrone that has passed through vertical. Section 3.5 relates $1 - \alpha$ directly

Table 1. Relationship between established folding analyses and the present formulation.

Study	Primary question	Diagnostic and scope
Waddington and others (2001)	Does a pre-existing local wrinkle flatten or rotate toward overturning?	Instantaneous tangent slope relative to a steady reference isochrone.
Jacobson and Waddington (2004)	Does a finite-strain history overturn an injected disturbance?	Deformation gradient integrated along trajectories.
Parrenin and others (2006); Parrenin and Hindmarsh (2007)	How does steady flow shape isochrones and preserve particle order?	Logarithmic flux coordinates, characteristic age solutions and a path factor; monotonic upward speed precludes vertical isochrones.
Hindmarsh and others (2006)	Do layers drape or override bed and basal-boundary perturbations?	Perturbation and full-mechanical models; wavelength controls architecture.
Hindmarsh and others (2009); Parrenin and others (2025)	How can age and thinning be computed accurately in steady flow tubes?	Eulerian and characteristic solvers for age, origin and vertical thinning.
Gudmundsson (1997a)	Can rough-bed Stokes flow contain local extrusion?	Analytical velocity extrema and reverse vertical shear near sinusoidal relief.
This study	Has cumulative reverse shear erased depositional ordering?	Signed inverse-thinning factor, positive–negative shear budget, full-Stokes threshold, obstacle bounds and trajectory diagnostic.

131 to the order Jacobian derived below. Parrenin and Hindmarsh (2007) proved that $\alpha > 1$ cannot occur if
132 horizontal velocity increases upward. Hindmarsh and others (2009) used these solutions to verify a large-
133 scale numerical age model, and Parrenin and others (2025) recently implemented the same flux-coordinate
134 framework for age, particle origin and vertical thinning.

135 Table 1 locates the present analysis within this literature.

3. STEADY AGE TRANSPORT AND THE ORDER JACOBIAN

3.1. Setting and assumptions

Consider a steady two-dimensional glacier velocity

$$\mathbf{u}(x, z) = \begin{bmatrix} u(x, z) \\ w(x, z) \end{bmatrix} \quad (3)$$

where u and w are the horizontal and vertical velocity components, respectively, and x and z are horizontal and vertical coordinates. The domain is bounded above by the steady surface $z = s(x)$. The kinematic result requires only the following assumptions on the region and trajectories of interest:

(A1) The velocity is continuously differentiable and incompressible,

$$u_x + w_z = 0. \quad (4)$$

(A2) Horizontal motion is one-way,

$$u(x, z) > 0, \quad (5)$$

after orienting x downstream.

(A3) Every trajectory considered originates in one upstream surface accumulation zone. The accumulation rate, expressed in vertical ice-equivalent units, satisfies $\dot{a}(x) > 0$ where the trajectory enters the ice.

(A4) Stratigraphic age is a passive, non-diffusive scalar. Its steady value $\tau(x, z)$ satisfies

$$\frac{D\tau}{Dt} = \underbrace{\tau_t}_{=0} + u\tau_x + w\tau_z = u\tau_x + w\tau_z = 1, \quad \tau = 0 \quad \text{at deposition.} \quad (6)$$

Equation (6) distinguishes a steady age field from the age of a moving parcel. The local derivative τ_t vanishes at fixed position, while the material age increases at one unit of time per unit time along a trajectory.

An impermeable bed is not required for the local result. Basal melting may terminate some trajectories without altering the derivation before termination. Basal freezing, in contrast, introduces a second material source and requires separate origin labels. Assumption (A2) excludes horizontal stagnation and closed circulation. Assumption (A3) requires special care in a blue-ice area because ice leaves through the local surface ablation zone. Its material origin must instead be traced to an upstream surface accumulation zone or to basal freeze-on. Appendix B gives the corresponding mass-flux construction when density varies.




Figure 1. Material map for steady plane flow. Neighboring particles enter through the surface accumulation zone at x_0 and $x_0 + \delta x_0$, follow distinct streamlines and reach positions separated by δx at the same age τ . The order Jacobian is the limit of $\delta x / \delta x_0$. The curves enforce the kinematic requirements used in the derivation—surface origin, non-intersecting streamlines and transverse intersections with the isochrone—but are not a calculated Stokes solution.

In a two-dimensional incompressible Stokes solution, mass conservation permits the velocity to be represented by a scalar streamfunction ψ . Curves of constant ψ are streamlines because $\mathbf{u} \cdot \nabla \psi = 0$, and the difference in ψ between two streamlines is their volume flux per unit width. The streamfunction is defined up to an arbitrary additive constant and follows from incompressibility alone; no additional rheological assumption is needed. We choose the orientation

$$u = \psi_z, \quad w = -\psi_x. \quad (7)$$

Combining Equations (6) and (7) gives the exact Jacobian identity

$$\begin{vmatrix} \frac{\partial \tau}{\partial x} & \frac{\partial \tau}{\partial z} \\ \frac{\partial \psi}{\partial x} & \frac{\partial \psi}{\partial z} \end{vmatrix} = \tau_x \psi_z - \tau_z \psi_x = u \tau_x + w \tau_z = 1. \quad (8)$$

153 Thus age and streamfunction form local area-preserving coordinates. Isochrones and streamlines intersect
 154 transversely wherever the assumptions hold. This is a classical flow-coordinate construction; it is useful here
 155 because its inverse directly measures stratigraphic order. Appendix A gives the inverse-coordinate identities
 156 used below. Figure 1 shows the deposition coordinate, two neighboring trajectories and their positions on
 157 an equal-age surface.

158 3.2. Surface origin and travel time

The streamfunction restricted to the surface is

$$Q(x) = \psi(x, s(x)). \quad (9)$$

The steady surface kinematic condition is

$$u_s s_x - w \Big|_{z=s(x)} = \dot{a}, \quad (10)$$

where $u_s = u(x, s(x))$. Differentiating Q with the chain rule, substituting Equation (7), and then applying Equation (10) gives

$$Q'(x) = \psi_x \Big|_{z=s(x)} + \psi_z \Big|_{z=s(x)} s_x = -w \Big|_{z=s(x)} + u_s s_x = \dot{a}(x) > 0. \quad (11)$$

The increment $d\psi$ is the ice flux per unit width between neighboring streamlines. At the surface, this flux is $\dot{a}_0 dx_0$, so larger accumulation places the same flux interval within a smaller deposition interval. Every streamline label ψ therefore has a unique deposition position $x_0(\psi)$ satisfying

$$Q(x_0) = \psi, \quad \frac{dx_0}{d\psi} = \frac{1}{\dot{a}_0}, \quad (12)$$

159 where $\dot{a}_0 \equiv \dot{a}(x)|_{x=x_0} = \dot{a}(x_0)$ is the accumulation rate at deposition.

At fixed ψ , Equation (6) gives

$$\frac{\partial \tau}{\partial x} \Big|_{\psi} = \frac{1}{u}. \quad (13)$$

Consequently the age at (x, ψ) is the travel time from deposition. Here, ξ is the horizontal coordinate of integration and $z(\xi, \psi)$ is the elevation of the streamline labelled by ψ :

$$\tau(x, \psi) = \int_{x_0(\psi)}^x \frac{d\xi}{u[\xi, z(\xi, \psi)]}. \quad (14)$$

At fixed x , Equation (7) implies

$$z_{\psi} = \frac{1}{u}. \quad (15)$$

Stratigraphic order compares travel times on neighboring streamlines, which are neighboring values of ψ . We therefore differentiate Equation (14) with respect to ψ at fixed downstream position. Using Equations (12) and (15) gives

$$\tau_{\psi} = -\frac{1}{\dot{a}_0 u_{s0}} - \int_{x_0}^x \frac{u_z}{u^3} d\xi, \quad (16)$$

where $u_{s0} = u[x_0, s(x_0)]$ is the horizontal speed at deposition. The first term records the separation imposed at deposition; the integral records the subsequent differential travel time between neighboring streamlines.

3.3. Exact order theorem

Parameterize an isochrone of fixed age by the surface deposition coordinate x_0 . The relevant quantity is not its curvature but the horizontal projection of its tangent.

Definition 1 (Isochrone-order Jacobian). The isochrone-order Jacobian is

$$\mathcal{J}(x, x_0) = \left. \frac{\partial x}{\partial x_0} \right|_{\tau}. \quad (17)$$

It measures whether particles deposited in increasing order of x_0 remain in increasing downstream order on an equal-age surface.

Theorem 1 (Cumulative order criterion). *Under assumptions (A1)–(A4), the isochrone-order Jacobian is*

$$\mathcal{J} = \frac{u(x, z)}{u_{s0}} \left(1 + \dot{a}_0 u_{s0} \int_{x_0}^x \frac{u_z}{u^3} d\xi \right). \quad (18)$$

It is also related exactly to the vertical age gradient by

$$\mathcal{J} = -\dot{a}_0 \tau_z. \quad (19)$$

Hence $\mathcal{J} > 0$ denotes ordered stratigraphy, $\mathcal{J} = 0$ a vertical tangent, and $\mathcal{J} < 0$ an overturned limb.

Proof. At fixed age, differentiation of $\tau(x, \psi)$ gives

$$\left. \frac{\partial x}{\partial \psi} \right|_{\tau} = -\frac{\tau_{\psi}}{\tau_x|_{\psi}} = -u\tau_{\psi}. \quad (20)$$

Multiplication by $d\psi/dx_0 = \dot{a}_0$ and substitution of Equation (16) gives Equation (18). At fixed x , $\tau_z = \tau_{\psi}\psi_z = u\tau_{\psi}$, which gives Equation (19). $\square$

The correction of annual-layer thickness for cumulative ice deformation goes back at least to Nye (1963) and Dansgaard and Johnsen (1969). In the present notation, the vertical thinning function λ_z is the current vertical thickness of an infinitesimal age layer divided by its ice-equivalent thickness at deposition. For an ordered layer, a positive age increment $d\tau$ corresponds to the present vertical thickness $-dz = -d\tau/\tau_z$ and the initial thickness $\dot{a}_0 d\tau$. Hence

$$\lambda_z = -\frac{1}{\dot{a}_0 \tau_z} = \mathcal{J}^{-1}, \quad \mathcal{J} > 0. \quad (21)$$

Thus $\mathcal{J}$ is not a new independent state variable: in the ordered regime it is the inverse of the established vertical thinning function (Parrenin, 2013; Parrenin and others, 2025). Equation (19) provides its signed continuation through a vertical tangent. The standard positive thinning function diverges as $\tau_z \rightarrow 0$, whereas $\mathcal{J}$ crosses zero regularly and records orientation reversal.

At deposition, $x = x_0$, $u = u_{s0}$, and $\mathcal{J} = 1$. At the first vertical tangent, $\tau_z = 0$, but the material curve remains regular. Inverting Equation (8) shows that

$$\left. \frac{\partial z}{\partial x_0} \right|_{\tau} = \dot{a}_0 \tau_x. \quad (22)$$

When $\mathcal{J} = 0$, Equation (6) gives $\tau_x = 1/u$, and therefore

$$\left. \frac{\partial z}{\partial x_0} \right|_{\tau, \mathcal{J}=0} = \frac{\dot{a}_0}{u} > 0. \quad (23)$$

The divergence of the conventional slope $m = dz/dx$ at overturning is therefore a coordinate singularity, not a singularity of the layer itself.

3.4. Positive and reverse shear budgets

Define

$$u_z^+ = \max(u_z, 0), \quad u_z^- = \max(-u_z, 0), \quad (24)$$

and the dimensionless cumulative contributions

$$\mathcal{P}(x, x_0) = \dot{a}_0 u_{s0} \int_{x_0}^x \frac{u_z^+}{u^3} d\xi, \quad (25)$$

$$\mathcal{R}(x, x_0) = \dot{a}_0 u_{s0} \int_{x_0}^x \frac{u_z^-}{u^3} d\xi. \quad (26)$$

Equation (18) becomes

$$\boxed{\mathcal{J} = \frac{u}{u_{s0}} \mathcal{B}, \quad \mathcal{B} \equiv 1 + \mathcal{P} - \mathcal{R}.} \quad (27)$$

The dimensionless factor $\mathcal{B}$ is an order budget: deposition contributes the initial value one, normal shear adds $\mathcal{P}$, and reverse shear subtracts $\mathcal{R}$. Because $u/u_{s0} > 0$, the necessary-and-sufficient condition for preservation of horizontal order is

$$\boxed{\mathcal{R} < 1 + \mathcal{P}.} \quad (28)$$

The first vertical tangent occurs at

$$\boxed{\mathcal{R} = 1 + \mathcal{P},} \quad (29)$$




Figure 2. Conceptual order budget. The sign of $\mathcal{B} = 1 + \mathcal{P} - \mathcal{R}$ is the sign of the order Jacobian. Local reverse shear need not cause overturning; a vertical tangent appears only after cumulative reverse shear removes the initial depositional contribution and the accumulated normal-shear contribution.

177 and overturning occurs when $\mathcal{R} > 1 + \mathcal{P}$.

178 The unit term in Equations (27)–(29) represents the initial ordering imposed at deposition. Positive u_z ,
 179 the usual surface-fast glacier profile, adds to the budget. Reverse shear must remove both contributions.
 180 Thus a local region with $u_z < 0$ is not itself a fold criterion.

In travel-time coordinates, $dx = u \, dt$, so

$$\mathcal{R} = \dot{a}_0 u_{s0} \int_0^T \frac{u_z^-}{u^2} \, dt. \quad (30)$$

181 where T is the travel time from deposition to the point of interest. Reverse shear is amplified by residence
 182 time and by slow horizontal motion. This weighting explains why local extrusion in fast throughflow may
 183 leave stratigraphic order unchanged, while modest reverse shear near stagnation can dominate the order
 184 budget.

185 **Corollary 1** (Monotonic-shear result). *If $u_z \geq 0$ everywhere along a trajectory, then $\mathcal{R} = 0$, $\mathcal{J} > 0$, and*
 186 *the steady isochrone cannot overturn.*

187 This corollary is the physical-coordinate statement of the no-overturning result in Appendix A of Parrenin
 188 and Hindmarsh (2007). It is sufficient, not necessary. Equation (28) permits finite intervals of reverse shear
 189 provided their cumulative weighted contribution remains below the order budget. Figure 2 summarizes the
 190 three contributions and the point at which the budget reaches zero.

191 3.5. Relation to the Parrenin–Hindmarsh path factor

Parrenin and Hindmarsh (2007) used a normalized streamfunction and logarithmic flux coordinates. For two neighboring particles, their Appendix A gives a mapping of the form

$$\delta\pi = (1 - \alpha) \delta\pi_0, \quad (31)$$

where π_0 and π are their logarithmic flux coordinates at deposition and at the point of interest, respectively, and α is their path-history factor. An isochrone exceeds vertical when $\alpha > 1$. Their present and surface-origin coordinates are monotone functions of x and x_0 wherever assumptions (A2)–(A3) hold. Consequently,

$$\text{sgn}(1 - \alpha) = \text{sgn}\left(\frac{\partial x}{\partial x_0}\bigg|_{\tau}\right) = \text{sgn}(\mathcal{J}) = \text{sgn}(-\tau_z). \quad (32)$$

192 The present result is therefore not an independent no-overturning theorem. Nor is $\mathcal{J}$ independent of the
 193 established thinning variable: Equation (21) identifies it as the signed inverse vertical-thinning factor. The
 194 contribution of the physical-space representation is that it exposes the positive and negative parts of vertical
 195 shear, remains regular at the first vertical tangent and can be evaluated directly from a numerical velocity
 196 field. The `age_flow_line` numerical age model of Parrenin and others (2025) assumes strictly positive
 197 accumulation for its horizontal flux coordinate and no reverse horizontal flow for its vertical flux coordinate;
 198 it computes age and positive vertical thinning within that admissible regime.

199 4. FULL-STOKES CONSTRAINTS ON ISOCHRONE ORDER

200 4.1. Exact stress decomposition of vertical shear

The order theorem is kinematic. To determine whether glacier mechanics forces Equation (28) to hold, consider steady, isothermal or thermomechanically prescribed, isotropic generalized-Newtonian Stokes flow,

$$\nabla \cdot \mathbf{u} = 0, \quad (33)$$

$$-\nabla p + \nabla \cdot \boldsymbol{\tau} + \rho \mathbf{g} = \mathbf{0}, \quad (34)$$

$$\boldsymbol{\tau} = 2\eta \mathbf{D}, \quad \mathbf{D} = \frac{1}{2} \left(\nabla \mathbf{u} + \nabla \mathbf{u}^T \right), \quad \eta > 0. \quad (35)$$

201 Here, p is pressure, ρ is ice density, $\mathbf{g}$ is gravitational acceleration, $\boldsymbol{\tau}$ is deviatoric stress, $\mathbf{D}$ is the strain-rate
 202 tensor and $\eta > 0$ is the scalar effective viscosity. The viscosity may depend on temperature and effective
 203 strain rate. Anisotropic rheology is discussed separately below.

With z vertical and gravity having no horizontal component, the horizontal momentum equation is

$$-p_x + \tau_{xx,x} + \tau_{xz,z} = 0. \quad (36)$$

Write

$$p = p_{\text{atm}} + \rho g(s - z) + p_{\text{nh}}, \quad (37)$$

where $g = |\mathbf{g}|$, p_{atm} is atmospheric pressure and p_{nh} contains all nonhydrostatic pressure, including the finite-slope surface correction. Integrating Equation (36) downward from the free surface, with ζ as the vertical coordinate of integration, gives

$$\tau_{xz}(x, z) = \tau_{xz}^s - \rho g s_x(s - z) + \int_z^s (\tau_{xx,x} - p_{\text{nh},x}) \, d\zeta. \quad (38)$$

Because $\tau_{xz} = \eta(u_z + w_x)$,

$$u_z = \underbrace{\frac{\rho g(s - z)(-s_x)}{\eta}}_G + \underbrace{\left[-w_x + \frac{\tau_{xz}^s}{\eta} + \frac{1}{\eta} \int_z^s (\tau_{xx,x} - p_{\text{nh},x}) \, d\zeta \right]}_{R_{\text{FS}}}. \quad (39)$$

where the superscript s denotes surface stress. For a surface descending downstream, $s_x \leq 0$, the gravity-driven term G is nonnegative and favors order preservation. The residual R_{FS} contains the full-Stokes effects omitted from the shallow-ice shear balance: gradients of vertical velocity, longitudinal stress, nonhydrostatic pressure and the Cartesian surface shear-stress component.

Substitution into Equation (18) gives

$$\mathcal{J} = \frac{u}{u_{s0}} (1 + \mathcal{G} + \mathcal{F}), \quad (40)$$

with

$$\mathcal{G} = \dot{a}_0 u_{s0} \int_{x_0}^x \frac{G}{u^3} \, d\xi \geq 0, \quad (41)$$

$$\mathcal{F} = \dot{a}_0 u_{s0} \int_{x_0}^x \frac{R_{\text{FS}}}{u^3} \, d\xi. \quad (42)$$

The exact full-Stokes condition for a vertical or overturned isochrone is

$$\boxed{\mathcal{F} \leq -(1 + \mathcal{G})}. \quad (43)$$

A conservative sufficient condition for no overturning is

$$\boxed{\dot{a}_0 u_{s0} \int_{x_0}^x \frac{[R_{\text{FS}}]_-}{u^3} d\xi < 1.} \quad (44)$$

This inequality discards all positive contributions, including gravity-driven shear, and is therefore stronger than necessary but easy to test.

4.2. There is no universal full-Stokes sign theorem

Neither incompressibility nor the Stokes equations fixes the sign of u_z . The constitutive relation constrains $u_z + w_x$, while longitudinal stress and nonhydrostatic pressure can alter the Cartesian vertical gradient of horizontal speed. Analytical rough-bed solutions predict local velocity maxima and minima, and therefore regions with $u_z < 0$, even for gravity-driven linear viscous flow (Gudmundsson, 1997a). The historical extrusion-flow literature likewise shows that surface-fast velocity profiles are not a pointwise theorem of full-Stokes mechanics (Waddington, 2010).

At stronger roughness, the one-way-flow assumption itself can fail. Gudmundsson (1997b) found separated secondary circulation in troughs of sufficiently steep sinusoidal beds. Ideal no-slip Stokes flow near a sharp corner can contain nested Moffatt eddies (Moffatt, 1964); Meyer and Creyts (2017) extended this mechanism to Glen-law ice in subglacial mountain valleys and showed that oblique flow can produce three-dimensional eddies. These examples are not direct models of Allan Hills: glacier beds may slide, Glen flow is nonlinear, and real corners are rounded. Their theoretical significance is that steady creeping-flow equations do not forbid recirculation. When $u = 0$ or trajectories close, the downstream travel-time coordinate and unique surface-origin map cease to exist, so Equations (18)–(28) no longer apply globally.

The distinction between local extrusion and cumulative order reversal is central. Published rough-bed solutions establish that u_z can be negative. They do not, by that fact alone, establish that $\mathcal{R}$ reaches $1 + \mathcal{P}$. The latter is an order-one integrated threshold.

5. REGIMES THAT PRESERVE ORDER

5.1. Shallow-ice flow

Under the conventional shallow-ice approximation,

$$\tau_{xz} \simeq -\rho g(s - z)s_x. \quad (45)$$

For downstream flow with $s_x < 0$, $\tau_{xz} > 0$. An isotropic Glen-type rheology gives u_z with the same sign as τ_{xz} , while leading-order basal sliding contributes a depth-independent horizontal speed (Cuffey and Paterson, 2010). Therefore

$$u_z \geq 0, \quad \mathcal{R} = 0, \quad \mathcal{J} > 0. \quad (46)$$

230 A fixed bed obstacle can produce arches, troughs and finite curvature in shallow-ice stratigraphy, but it
231 cannot overturn a steady isochrone under these assumptions.

232 5.2. Regular thin-film full-Stokes flow

Let

$$\varepsilon = \frac{H}{L} \ll 1 \quad (47)$$

be the aspect ratio of a smooth glacier. Higher-order glacier models reproduce the Stokes solution to second order in ε under their regular asymptotic assumptions, including a range of basal slip regimes (Blatter, 1995; Schoof and Hindmarsh, 2010). Suppose the non-shallow contribution scales as

$$R_{\text{FS}} = O\left(\varepsilon^2 \frac{U}{H}\right) \quad (48)$$

through a region of length $L = H/\varepsilon$, and suppose u remains comparable to U . Then

$$\mathcal{F} = O\left(\frac{\dot{a}}{U}\varepsilon\right). \quad (49)$$

Thus small regular full-Stokes corrections cannot erase the initial depositional contribution when

$$\frac{\dot{a}}{U}\varepsilon \ll 1. \quad (50)$$

233 This is an asymptotic sufficient regime, not a uniform theorem. The approximation can break down near
234 divides, abrupt sliding transitions, contact zones, steep obstacles, very slow flow or other regions where the
235 thin-film expansion is singular or nonuniform.

236 5.3. A sufficient bound for a localized obstacle

237 The cumulative criterion yields a direct obstacle bound without requiring pointwise monotonic shear.

Proposition 1 (Localized reverse-shear bound). *Suppose reverse shear is confined to a trajectory segment of horizontal length ℓ . On that segment, assume*

$$u \geq U_{\min} > 0, \quad u_{s0} \leq U_{\text{dep}}, \quad u_z^- \leq \dot{\gamma}_{\text{rev}}, \quad (51)$$

where $U_{\min}$ is the minimum horizontal speed on the segment, U_{dep} bounds the horizontal speed at deposition, and $\dot{\gamma}_{\text{rev}}$ bounds the reverse vertical shear rate. Then

$$\mathcal{R} \leq \frac{\dot{a}_0 U_{\text{dep}} \dot{\gamma}_{\text{rev}} \ell}{U_{\min}^3}. \quad (52)$$

A sufficient condition for no overturning is that the right-hand side be less than one.

Proof. Substitute Equation (51) into Equation (26) and bound the integral by its maximum integrand multiplied by ℓ . $\square$

For an isolated obstacle whose horizontal and vertical influence scales are both $O(H)$, with non-stagnant throughflow,

$$\mathcal{R} = O\left(\frac{\dot{a}}{U}\right). \quad (53)$$

In a typical interior or flank setting where $U \gg \dot{a}$, an isolated fixed bump can create local extrusion and visible draping while remaining far from the reverse-shear value required for a vertical tangent. More generally, setting the right-hand side of Equation (52) equal to one and solving for length gives

$$\ell \gtrsim \frac{U_{\min}^3}{\dot{a}_0 U_{\text{dep}} \dot{\gamma}_{\text{rev}}}. \quad (54)$$

Slow throughflow reduces this required length through the $U_{\min}^3$ numerator. For scale, take $\dot{a}_0 = 0.05 \text{ m yr}^{-1}$, $U_{\text{dep}} = 10 \text{ m yr}^{-1}$, $\dot{\gamma}_{\text{rev}} = 0.01 \text{ yr}^{-1}$ and $\ell = 1 \text{ km}$. Equation (52) gives $\mathcal{R} \leq 0.005$ when $U_{\min} = 10 \text{ m yr}^{-1}$, which rules out overturning under these assumed bounds. With $U_{\min} = 1 \text{ m yr}^{-1}$, the bound is $\mathcal{R} \leq 5$ and no longer rules it out. These illustrative values are not estimates for Allan Hills; they show that a tenfold reduction in minimum speed raises the bound by a factor of one thousand.

Equation (52) makes precise why “flow over a bump cannot fold an isochrone” is often a good glacier approximation but not a universal Stokes theorem. The approximation depends on non-stagnant flow, a localized deformation zone and a small accumulation-to-speed ratio.

5.4. First-order cancellation over periodic roughness about plug flow

A second reason that fixed roughness may have weak cumulative effect is cancellation, but the result requires a plug-flow background. Let

$$u(x, z; \epsilon_b) = U_b + \epsilon_b u_1(x, z) + O(\epsilon_b^2), \quad U_b > 0, \quad (55)$$

where ϵ_b is a small amplitude parameter, $U_b > 0$ is spatially uniform, u_1 is periodic in x , and k is its wavenumber. The perturbation has wavelength $2\pi/k$ and zero horizontal mean at each elevation. Along the unperturbed trajectory $z = z_0$,

$$\frac{u_z}{u^3} = \epsilon_b \frac{u_{1,z}(x, z_0)}{U_b^3} + O(\epsilon_b^2). \quad (56)$$

Therefore, over one complete wavelength,

$$\int_x^{x+2\pi/k} \frac{u_z}{u^3} d\xi = O(\epsilon_b^2). \quad (57)$$

Appendix C gives the expansion leading to Equation (57). The perturbative frictionless rough-bed solution of Gudmundsson (1997a) has this plug-flow structure: local extrusion occurs, but its first-order stoss- and lee-side contributions cancel over a complete symmetric wavelength.

Cancellation is not guaranteed for a sheared background, a solitary asymmetric obstacle, a finite path ending mid-cycle, strongly nonlinear roughness or near-stagnant trajectories. Equation (57) is therefore a restricted asymptotic result, not a general symmetry theorem.

6. STEADY OVERTURNING AND LIMITS OF THE ASSUMPTIONS

The theorem does not fail when assumptions (A1)–(A4) hold. Instead, it predicts whether the order budget remains positive or reaches the vertical-tangent threshold. This section first describes overturning within the theorem and then identifies settings in which one or more assumptions are not satisfied, so that the two-dimensional steady coordinate construction does not apply.

6.1. Overturning within the theorem: cumulative extrusion

The direct steady-flow mechanism is cumulative reverse shear. A glacier with $u > 0$ can overturn a steady isochrone if reverse shear is sufficiently strong, sufficiently persistent or sufficiently concentrated where u is small. Equation (29) gives the exact threshold:

$$\dot{a}_0 u_{s0} \int_{x_0}^x \frac{u_z^-}{u^3} d\xi = 1 + \dot{a}_0 u_{s0} \int_{x_0}^x \frac{u_z^+}{u^3} d\xi. \quad (58)$$

A change in boundary conditions is not mechanically necessary if a steady full-Stokes velocity field satisfies this equality. This outcome is predicted by the theorem rather than representing a failure of it. The extrusion-flow literature establishes the necessary local reverse shear but does not evaluate this cumulative isochrone-order threshold.

6.2. Stagnation, separation and closed circulation

If u reaches zero, the x -travel-time representation becomes singular. If a secondary circulation forms, a streamline may not have a unique surface origin and may intersect an equal-age surface in a topology not represented by a single downstream coordinate. Strong roughness, cavities and sharp corners are therefore not merely large values of $\mathcal{R}$; they can invalidate the coordinate system on which the theorem rests. Numerical calculations should treat $u = 0$ as an event and switch to a more general Lagrangian deformation analysis.

6.3. Surface ablation and sublimation

Newly deposited material enters the glacier through a surface accumulation zone with $\tau = 0$. Existing isochrones terminate where ice emerges in a surface ablation zone. In a blue-ice area, the surface streamfunction need not be globally monotone because

$$Q'(x) = \dot{a}(x) \tag{59}$$

changes sign between accumulation and ablation zones. A streamline may enter through an upstream accumulation zone and emerge through a downstream ablation zone. The theorem can still be applied along an interior trajectory if that trajectory has a unique origin in the upstream accumulation zone, but Q^{-1} must be constructed only over that part of the surface. Multiple surface accumulation zones or basal freeze-on require a more general material-source label.

Steady ablation most naturally produces upward-bent and truncated layers, not an intact fold. However, migration, onset or cessation of an ablation zone changes the age field relative to the new steady solution and supplies the inherited disturbance considered by WBA. Surface sublimation followed by burial can also preserve an unconformity with steep or truncated limbs. At Allan Hills, present ablation terminates trajectories, whereas changes in the position of the ablation zone introduce time dependence that lies outside the steady theorem.

6.4. Basal melting and freeze-on

Basal melting removes trajectories and truncates layers. Basal freeze-on adds ice at the bed, so one surface deposition coordinate cannot label the entire stratigraphy. Numerical models show that freeze-on can generate plume-like structures and complex internal layering extending through a large fraction of the ice thickness (Leysinger Vieli and others, 2018). A moving melt/freeze boundary additionally makes the velocity and age problem time dependent.

6.5. Time-dependent geometry and basal traction

If the bed, free surface, divide position or basal traction changes, the inherited age field is not the steady age solution of the new velocity field. The perturbation then obeys a homogeneous advection equation only after the change; it need not vanish initially. Divide migration converts a formerly steady Raymond arch into a fold seed (Jacobson and Waddington, 2005). Traveling slippery or sticky patches produce moving-frame critical layers and thickness-scale overturning (Wolovick and others, 2014; Wolovick and Creyts, 2016). Bedrock uplift, rotation or erosion similarly changes the reference stratigraphy and can inject a finite-amplitude disturbance even when subsequent flow is steady.

6.6. Active rheology, anisotropy and three-dimensional flow

Equation (39) assumes isotropic scalar viscosity. Crystal fabric, impurities, grain size and damage can alter both the stress–strain relation and the velocity field in response to stratigraphy. Full-Stokes anisotropic divide models already demonstrate substantial changes in isochrone shape and age–depth relations (Martín and others, 2009a). More strongly coupled heterogeneity can generate folds internally rather than merely advecting an imposed perturbation (Alley and others, 1997; Thorsteinsson and Waddington, 2002; Bons and others, 2016, 2025). The Allan Hills microstructure indicates that anisotropy and localized shear cannot be ignored (Stoll and others, 2026).

The order theorem is also intrinsically two-dimensional. In three dimensions, incompressibility provides a vector potential rather than a scalar streamfunction, and neighboring surface origins can exchange order in one horizontal projection while remaining separated in the other. Divide models show that along-ridge flow and divide migration alter isochrone architecture even when cross-flow is modest (Martín and others, 2009b). A three-dimensional generalization should track the two-dimensional Jacobian mapping a surface deposition patch to an equal-age surface. The scalar $\mathcal{J}$ is recovered only when cross-flow is negligible and a coherent flow tube exists.

7. IMPLICATIONS FOR OLD-ICE SITES

Allan Hills lies close to several limits in which a two-dimensional fixed-obstacle approximation becomes unreliable. Present-day ice rises into a surface ablation zone above steep nunatak topography (Spaulding and others, 2012; Kehrl and others, 2018; Stoll and others, 2026). The three-dimensional full-Stokes model of Manos and others (2026) predicts nearly stagnant basal ice upstream of the nunatak, where the bottom 20 m averages approximately $0.4\text{--}0.5\text{ mm yr}^{-1}$. Modeled speed remains nonzero but locally falls to order

micrometres per year. In contrast, modeled basal speeds in the Cul-de-sac are approximately 46–49 mm yr⁻¹.

These modeled scales distinguish local preservation in slow ice from transport of old ice into the Cul-de-sac.

Near-stagnant flow has two consequences for the present calculation. First, the u^{-3} weighting in Equation (26) becomes large and numerically sensitive even though assumption (A2) may remain technically satisfied. Second, the modeled flow is fundamentally three-dimensional. It contains important local bridging stresses associated with horizontal gradients of vertical velocity; the corresponding $-w_x$ term appears explicitly in the full-Stokes shear decomposition, Equation (39). A two-dimensional order budget is therefore meaningful only within a coherent flow tube for which cross-flow is small. The Cul-de-sac presents a separate provenance problem: its geochemistry and modeled transport indicate a distinct accumulation environment (Manos and others, 2026). The assumption that each trajectory can be represented by one surface accumulation zone is therefore unlikely to describe all Cul-de-sac ice, and a three-dimensional, multi-source material map is needed there.

The structural observations constrain, but do not determine, the deformation history. The 69° mean layer dip in ALHIC2302 is steep but remains below vertical (Kirkpatrick and others, 2025). Centimetre-scale z-folds in ALHIC1901 record local folding, but they do not by themselves explain every core-scale age reversal (Stoll and others, 2026). The inverted age–depth relation in ALHIC1901 more directly indicates that τ_z changes sign over some sampled intervals (Shackleton and others, 2025). Equation (19) then implies $\mathcal{J} < 0$ if the samples lie on one continuous family of surface-deposited layers. Multiple source regions, unconformities or faults could instead produce an apparent age reversal.

The same calculation is relevant to deep old-ice drilling. At the Beyond EPICA site, radar and flow modelling indicate strong basal thinning and possible stagnant or disturbed ice near the bed (Lilien and others, 2021; Chung and others, 2025). Evaluating $\mathcal{J}$ along model trajectories would test whether the steady velocity field preserves order above this basal unit. The boundary conditions differ from those at Allan Hills, but both settings require a mechanical link between modelled flow and the stratigraphic age interpretation.

For old-ice stratigraphic interpretation, the equations define four tests of whether steady local topography can produce an observed fold or age reversal.

- (1) **Steady fixed geometry.** Compute the present or reconstructed steady full-Stokes field and evaluate $\mathcal{P}$, $\mathcal{R}$ and $\mathcal{J}$ along source-to-core trajectories. If the order budget $1 + \mathcal{P} - \mathcal{R}$ remains well above zero, fixed present-day geometry cannot produce the observed recumbent folding.

(2) **Steady but singular or multi-source flow.** Test for stagnation, recirculation, basal freeze-on and surface origin multiplicity. If these occur, the simple order theorem is inapplicable and the full material map must be used.

(3) **Inherited unsteady stratigraphy.** If the steady model preserves order, evaluate changes in ablation pattern, ice thickness, flow direction, divide position, bed geometry and basal traction. These mechanisms inject finite disturbances that can later be amplified by shear.

(4) **Active deformation.** If kinematic models underpredict observed folds, couple fabric and rheological heterogeneity to flow. The microstructural observations make this mechanism plausible at centimetre scale.

This hierarchy distinguishes regimes in which changing boundary conditions are mechanically required. In regular, shallow, non-stagnant flow, Equation (52) keeps the order budget well above zero, so a time-dependent or rheological mechanism is needed. In more general full-Stokes flow, sufficiently cumulative extrusion or loss of one-way flow can overturn steady isochrones without a boundary change.

8. TRAJECTORY DIAGNOSTIC FOR FULL-STOKES MODELS

The order criterion can be evaluated without explicitly solving the age equation or advecting a dense family of isochrones. Start trajectories in a surface accumulation zone and record x_0 , $\dot{a}_0$ and u_{s0} . With t denoting time since deposition, integrate along each trajectory

$$\frac{dx}{dt} = u, \quad \frac{dz}{dt} = w, \quad (60)$$

$$\frac{d\mathcal{P}}{dt} = \dot{a}_0 u_{s0} \frac{u_z^+}{u^2}, \quad \frac{d\mathcal{R}}{dt} = \dot{a}_0 u_{s0} \frac{u_z^-}{u^2}, \quad (61)$$

with $\mathcal{P}(0) = \mathcal{R}(0) = 0$. Then

$$\boxed{\mathcal{J}(t) = \frac{u(t)}{u_{s0}} [1 + \mathcal{P}(t) - \mathcal{R}(t)]}. \quad (62)$$

The diagnostic is

$$\min_t [1 + \mathcal{P}(t) - \mathcal{R}(t)] \begin{cases} > 0, & \text{order preserved,} \\ = 0, & \text{first vertical tangent,} \\ < 0, & \text{overturned steady isochrone.} \end{cases} \quad (63)$$

A robust implementation should include four checks. First, compare Equation (62) with $-\dot{a}_0 \tau_z$ from an independently solved steady age equation. Second, advect at least one pair of neighboring particles to

365 verify the sign of $\partial x/\partial x_0|_\tau$. Third, stop the downstream-coordinate integration when u approaches zero
 366 and diagnose recirculation with the full trajectory topology. Fourth, compute u_z using the finite-element
 367 gradient rather than differencing noisy interpolated velocities. Separate storage of $\mathcal{P}$ and $\mathcal{R}$ is preferable
 368 to integrating their difference because large compensating contributions can otherwise obscure numerical
 369 error.

370 For transient flow, Equation (62) is not valid because the steady travel-time representation fails. The
 371 corresponding calculation should integrate the deformation of the two-parameter deposition map through
 372 time. Nevertheless, the steady order budget remains useful as a baseline and as a diagnostic applied to
 373 quasi-steady snapshots.

374 9. FULL-STOKES NUMERICAL EXAMPLES

375 The calculations test whether a steady full-Stokes velocity field can generate a recumbent fold from ordered
 376 surface deposition, without first imposing a finite-amplitude layer perturbation. They isolate this question
 377 in controlled two-dimensional geometries and are not reconstructions of Allan Hills or another old-ice site.
 378 The domain is periodic, 800 m long and 160 m thick, with a flat traction-free surface and a smooth raised-
 379 cosine bed obstacle. Ice is incompressible and linear viscous, with $\eta = 2.5 \times 10^8$ Pa yr and $\rho = 917$ kg m⁻³.
 380 Gravity acts at a mean surface slope of 0.03 rad. We solve for velocity and pressure with Taylor–Hood
 381 elements, using quadratic velocity and linear pressure.

The basal boundary is a sliding end member. The bed is fixed, the horizontal basal ice speed is
 $U_b = 0.20$ m yr⁻¹, and no penetration gives

$$\mathbf{u}_b = U_b \begin{bmatrix} 1 \\ b_x \end{bmatrix}. \quad (64)$$

382 This condition removes the large surface-fast shear generated by a no-slip bed and allows form drag to control
 383 the deformation around the obstacle. It is a specified basal-velocity model, not a sliding-law prediction. The
 384 flat surface is held steady by the diagnosed accumulation rate $\dot{a} = -w|_{z=H}$. We use only surface intervals
 385 where $\dot{a} > 0$ and trajectories on which $u > 0$.

386 The numerical isochrones are generated from the deposition boundary. At age zero, a particle labelled
 387 by x_0 is placed at $[x_0, H]^\top$. Integrating the steady velocity for a common age τ gives the material curve
 388 $[x(x_0, \tau), z(x_0, \tau)]^\top$. We do not initialize an internal horizontal layer, wrinkle or fold. Therefore, a negative
 389 value of $\partial x/\partial x_0|_\tau$ measures order reversal created between the surface and the plotted isochrone.

390 The first example isolates persistent extrusion. The obstacle is 150 m high, with horizontal half-widths of
 391 300 m and 80 m. Surface-generated isochrones remain ordered at 1 kyr, first contain an overturned interval
 392 by 2 kyr, and have a clear recumbent limb at 2.5 kyr (Figures 3a–b). The minimum material-map Jacobian
 393 at 2.5 kyr is -0.455 on the finest mesh. Along the corresponding trajectory, horizontal speed does not fall
 394 below its value at deposition. Reverse shear nevertheless removes the initial order contribution because it
 395 persists along the path.

396 The second example isolates slow-flow amplification. The 140 m-high obstacle has half-widths of 45 m
 397 and 260 m. A surface-origin trajectory slows substantially upstream of the steep face while retaining $u > 0$.
 398 The corresponding isochrone reaches vertical by approximately 0.7 kyr and has a minimum material-map
 399 Jacobian of -0.270 at 0.8 kyr (Figures 3c–d). This calculation realizes the residence-time mechanism in
 400 Equation (30): slow horizontal motion gives reverse shear greater weight even though the velocity field
 401 remains steady.

402 The split shear budget gives the same mechanism distinction (Figure 4). In the persistent-extrusion case,
 403 $\mathcal{P}$ remains negligible and $\mathcal{R}$ crosses one while $u/u_{s0} \geq 1$. In the slow-flow case, the minimum value of u/u_{s0}
 404 is 0.080, so horizontal speed decreases by more than an order of magnitude before $\mathcal{R}$ removes the order
 405 budget. The first calculation therefore crosses the threshold through sustained reverse shear; the second
 406 combines reverse shear with the slow-speed weighting in Equation (30).

407 We repeat the calculation on meshes with 64 by 14, 96 by 21 and 128 by 28 divisions. The minimum
 408 neighboring-trajectory estimates of $\partial x / \partial x_0|_\tau$ are -0.452 , -0.454 and -0.455 for persistent extrusion and
 409 -0.282 , -0.279 and -0.270 for slow-flow amplification (Figure 5). This calculation is independent of the
 410 vertical-shear quadrature in Equation (62). The slow-flow budget is more sensitive in magnitude because
 411 of the u^{-2} weight, but the material-map calculation retains the sign and the overturned interval under
 412 refinement.




Figure 3. Recumbent isochrones generated by steady full-Stokes flow. The curves are equal-age positions of particles deposited over an ordered surface interval; no internal layer perturbation is prescribed. (a) Persistent extrusion over an asymmetric sliding-bed obstacle. (b) Detail of the 2.5 kyr isochrone. Color gives the surface deposition coordinate x_0 , and open circles mark the backward-running interval where $\partial x / \partial x_0|_{\tau} < 0$. (c) Isochrones upstream of a steeper obstacle, where horizontal motion becomes slow. (d) Detail of the 0.8 kyr overturned limb. Gray regions show the fixed bed; horizontal and vertical coordinates are in m.




Figure 4. Two routes to order reversal in the steady calculations. The left panels show the cumulative positive shear $\mathcal{P}$, reverse shear $\mathcal{R}$ and order budget $1 + \mathcal{P} - \mathcal{R}$ along the trajectory with the most negative material-map Jacobian. The right panels show horizontal speed relative to its value at deposition along the same trajectory. The dotted line marks the first zero of the integrated order budget. Persistent extrusion crosses the threshold without slow flow, whereas the second example is amplified by a large decrease in horizontal speed.




Figure 5. Mesh check for the boundary-generated folds. The ordinate is the minimum value of $\partial x / \partial x_0|_\tau$ evaluated from neighboring surface-origin trajectories at 2.5 kyr and 0.8 kyr for the persistent-extrusion and slow-flow examples, respectively. Both calculations retain a negative interval on the three meshes.

These examples are constructive rather than site specific. Their topographic relief is large, the basal speed is prescribed and the rheology is linear viscous. They show that the steady age field itself can fold when cumulative reverse shear crosses the threshold; they do not show that this mechanism produced the observed Allan Hills stratigraphy. Applying the calculation at Allan Hills requires measured three-dimensional geometry, a basal traction model and source-to-core trajectories consistent with the observed surface mass balance.

10. DISCUSSION

10.1. A theorem about deposition order, not about curvature

The curvature equation for an advected material line contains first and second spatial derivatives of velocity. It correctly shows that an obstacle can create curvature from an initially straight layer. The present order criterion answers a different question. A recumbent fold requires the horizontal projection of an equal-age surface to reverse, or equivalently τ_z to change sign. An obstacle may generate large curvature while $\mathcal{J}$ remains positive everywhere. This is the rigorous sense in which a steady bump can disturb isochrones without supplying the order-reversing seed required for a recumbent fold.

WBA's statement that steady-state layers are not passively folded can now be made conditional. If a unique steady age field exists and Equation (28) holds, the steady surface-deposition map remains ordered. If cumulative reverse shear crosses the threshold, the steady age field itself contains a fold; no perturbation relative to that solution is required. This boundary-generated problem is distinct from placing an arbitrary material line in the velocity field and asking whether an inherited wrinkle grows.

10.2. Why local extrusion is usually insufficient

Full-Stokes solutions contain local extrusion, but local extrusion does not imply order reversal. Equation (27) can be read as an order budget. Deposition supplies an initial contribution of one, surface-fast shear adds $\mathcal{P}$, and reverse shear subtracts $\mathcal{R}$. Reverse shear around a localized or periodic obstacle may also be short-lived or cancel between stoss and lee sides. Equation (52) bounds this subtraction, while Equation (57) gives the perturbative cancellation.

Near stagnation, the reverse-shear subtraction becomes much more sensitive to speed. The u^{-3} spatial weighting and u^{-2} temporal weighting imply that a slowly moving basal zone can accumulate a large order change even when the pointwise shear is moderate. This is also the regime in which age becomes large, numerical conditioning deteriorates and recirculation becomes possible. A steady-obstacle explanation for

442 Allan Hills folding must therefore identify a slow reverse-shear trajectory and show that the order budget
 443 reaches zero in Equation (29); showing $u_z < 0$ is insufficient.

444 10.3. Boundary changes as fold seeds

445 Time-dependent boundary processes provide several mechanisms for creating finite inherited disturbances.
 446 Surface sublimation can truncate and steepen layers; migration of an ablation zone can leave that geometry
 447 to be buried or sheared. Basal melt and freeze-on alter material sources and sinks. Divide migration changes
 448 the reference velocity profile. Traveling traction anomalies create a moving-frame critical layer and material
 449 trapping. Bedrock uplift or rotation changes both geometry and stress. Each mechanism breaks at least one
 450 assumption behind the steady order theorem and can therefore create a nonzero perturbation relative to
 451 the steady stratigraphy.

452 The order framework does not identify which mechanism operated at Allan Hills. It supplies a rejection
 453 test. If a high-resolution, history-consistent full-Stokes model of fixed geometry keeps the order budget well
 454 above zero on all relevant trajectories, then steady topographic draping is excluded as the cause of recumbent
 455 age reversals. The remaining explanation must invoke time dependence, multiple material sources, active
 456 rheology, three-dimensional exchange or structural discontinuity.

457 10.4. Novelty and relationship to established theory

458 Several components of this manuscript are established. Equation (8) is a standard consequence of steady
 459 incompressible transport. Logarithmic flux coordinates and characteristic age solutions were developed by
 460 Parrenin and others (2006) and Parrenin and Hindmarsh (2007), and the latter paper contains both the
 461 monotonic-shear corollary and the path dependence of isochrone slope. The vertical thinning function was
 462 derived in the same framework by Parrenin (2013) and is now computed operationally by Parrenin and
 463 others (2025); Equation (21) shows that the present order factor is its inverse in the ordered regime. The
 464 need for a transient wrinkle and the competition between shear and extension are due to Waddington and
 465 others (2001). Finite-strain amplification is treated by Jacobson and Waddington (2004). Local extrusion
 466 and rough-bed recirculation are known full-Stokes phenomena (Gudmundsson, 1997a,b).

467 The analysis therefore reformulates established age–streamfunction and thinning theory for use with
 468 numerical velocity fields. Its distinct result is the physical-space identity that writes signed inverse thinning
 469 as a cumulative u_z/u^3 integral. The positive–negative decomposition in Equations (25)–(28) then gives the
 470 full-Stokes residual threshold in Equation (43), the localized-obstacle bound in Equation (52), the plug-flow
 471 cancellation result in Equation (57) and the split trajectory diagnostic in Equation (62). These results are

a cautious extension of Parrenin–Hindmarsh theory rather than a universal theorem about incompressible Stokes flow.

The idealized calculations in Section 9 provide constructive examples of steady order reversal, but they do not test Allan Hills geometry. The next step is to apply the same material-map and order-budget calculations to reconstructed geometry and flow history. An order budget that remains well above zero would exclude fixed roughness as the cause of overturning, whereas a trajectory crossing $\mathcal{J} = 0$ would identify where steady deformation is sufficient. Layer attitudes, age reversals and fabric provide independent observations against which to evaluate either result.

10.5. Limitations

The analysis neglects diffusion of age markers, fracture and discontinuous displacement. It assumes a smooth two-dimensional velocity field and that each trajectory originates in one surface accumulation zone. The full-Stokes decomposition assumes isotropic scalar viscosity. It does not establish a universal mechanical bound on R_{FS} , because none follows from the Stokes equations alone. The localized-obstacle estimate depends on externally supplied bounds on minimum speed, deposition speed and reverse shear. The periodic cancellation result is first order and can be destroyed by asymmetry or nonlinear geometry.

Most importantly, the theorem diagnoses overturning of a continuous material isochrone. An observed age reversal can also result from faulting, unconformity, multiple source regions, freeze-on ice, sampling across a sharply dipping but discontinuous structure, or unresolved three-dimensional geometry. The order diagnostic should therefore be used with structural observations rather than as a stand-alone interpretation.

11. CONCLUSIONS

A steady two-dimensional isochrone preserves depositional order if and only if its signed order Jacobian remains positive (cf. Equation (18)). The equivalent observable condition is $\tau_z < 0$ (Equation (19)), and in the ordered regime $\mathcal{J}$ is the reciprocal of the established vertical thinning function (Equation (21)). Pointwise surface-fast shear, $u_z \geq 0$, is sufficient but not necessary. The exact condition is the cumulative order budget $\mathcal{R} < 1 + \mathcal{P}$ (Equation (28)).

Full-Stokes mechanics does not guarantee this condition for arbitrary glacier geometry. Gravity-driven shear favors order, but longitudinal stress, nonhydrostatic pressure and horizontal gradients of vertical velocity can generate reverse shear (cf. Equation (39)). The shallow-ice regime preserves order (Equation (46)); non-stagnant localized obstacles and small-amplitude periodic roughness obey the more

restricted results in Equations (52) and (57). Stagnation and recirculation invalidate the one-way coordinate construction.

The idealized full-Stokes examples generate recumbent isochrones directly from ordered surface-deposition maps (Figures 3–5). One crosses the threshold through persistent reverse shear while horizontal speed remains comparable to its deposition value. The other combines reverse shear with slow-flow amplification. Both retain $u > 0$, and both retain a negative neighboring-trajectory Jacobian under mesh refinement. The intuition that fixed obstacles produce draping rather than folds is therefore conditional: outside ordinary fast, shallow throughflow, a steady velocity field can generate a fold without an inherited perturbation if the diagnostic in Equation (62) reaches zero.

For NSF COLDEX at Allan Hills, the next calculation is to integrate $\mathcal{P}$, $\mathcal{R}$ and $\mathcal{J}$ along reconstructed source-to-core trajectories using Equations (60)–(62). A three-dimensional material map is required where cross-flow or multiple source regions are important. The same diagnostic can test stratigraphic order at continuous old-ice sites such as Beyond EPICA.

DATA AVAILABILITY STATEMENT

No new observational data are presented. The analytical results are fully specified by the equations in this manuscript. Three executed notebooks contain the full-Stokes solves, trajectory calculations, numerical checks and figure generation and are available at github.com/bradlipovsky/allan-hills-stratigraphy.

AUTHOR CONTRIBUTIONS

Bradley Paul Lipovsky developed the analysis and wrote the manuscript draft.

COMPETING INTERESTS

To be completed by the author before submission.

ACKNOWLEDGEMENTS

To be completed before submission. Support and contributions from NSF COLDEX and collaborating investigators should be documented here.

REFERENCES

- Alley RB, Gow AJ, Meese DA, Fitzpatrick JJ, Waddington ED and Bolzan JF (1997) Grain-scale processes, folding, and stratigraphic disturbance in the GISP2 ice core. *Journal of Geophysical Research: Oceans*, **102**(C12), 26819–26830 (doi: 10.1029/96JC03836)
- Bintanja R (1999) On the glaciological, meteorological, and climatological significance of antarctic blue ice areas. *Reviews of Geophysics*, **37**(3), 337–359 (doi: 10.1029/1999RG900007)
- Blatter H (1995) Velocity and stress fields in grounded glaciers: a simple algorithm for including deviatoric stress gradients. *Journal of Glaciology*, **41**(138), 333–344 (doi: 10.3189/S002214300001621X)
- Bons PD, Jansen D, Mundel F, Bauer CC, Binder T, Eisen O, Jessell MW, Llorens MG, Steinbach F, Steinhage D and Weikusat I (2016) Converging flow and anisotropy cause large-scale folding in greenland’s ice sheet. *Nature Communications*, **7**, 11427 (doi: 10.1038/ncomms11427)
- Bons PD, Hu Y, Llorens MG, Franke S, Stoll N, Weikusat I, Westhoff J and Zhang Y (2025) Folding due to anisotropy in ice, from drill-core-scale cloudy bands to km-scale internal reflection horizons. *The Cryosphere*, **19**, 5095–5109 (doi: 10.5194/tc-19-5095-2025)
- Chung A, Parrenin F, Mulvaney R, Vittuari L, Frezzotti M, Zanutta A, Lilien DA, Cavitte MGP and Eisen O (2025) Age, thinning and spatial origin of the Beyond EPICA ice from a 2.5d ice flow model. *The Cryosphere*, **19**(10), 4125–4140 (doi: 10.5194/tc-19-4125-2025)
- Cuffey KM and Paterson WSB (2010) *The Physics of Glaciers*. Elsevier, Amsterdam, 4 edition
- Dansgaard W and Johnsen SJ (1969) A flow model and a time scale for the ice core from Camp Century, Greenland. *Journal of Glaciology*, **8**(53), 215–223 (doi: 10.3189/S0022143000031208)
- Gudmundsson GH (1997a) Basal-flow characteristics of a linear medium sliding frictionless over small bedrock undulations. *Journal of Glaciology*, **43**(143), 71–79 (doi: 10.3189/S0022143000002823)
- Gudmundsson GH (1997b) Basal-flow characteristics of a non-linear flow sliding frictionless over strongly undulating bedrock. *Journal of Glaciology*, **43**(143), 80–89 (doi: 10.3189/S0022143000002835)
- Gurtin ME, Fried E and Anand L (2010) *The Mechanics and Thermodynamics of Continua*. Cambridge University Press, Cambridge, UK, ISBN 9780521405980
- Higgins JA, Kurbatov AV, Spaulding NE, Brook EJ, Introne DS, Chimiak LM, Yan Y, Mayewski PA and Bender ML (2015) Atmospheric composition 1 million years ago from blue ice in the allan hills, antarctica. *Proceedings of the National Academy of Sciences*, **112**(22), 6887–6891 (doi: 10.1073/pnas.1420232112)
- Hindmarsh RCA, Leysinger Vieli GJMC, Raymond MJ and Gudmundsson GH (2006) Draping or overriding: The effect of horizontal stress gradients on internal layer architecture in ice sheets. *Journal of Geophysical Research: Earth Surface*, **111**(F2), F02018 (doi: 10.1029/2005JF000309)

- Hindmarsh RCA, Leysinger Vieli GJMC and Parrenin F (2009) A large-scale numerical model for computing isochrone geometry. *Annals of Glaciology*, **50**(51), 130–140 (doi: 10.3189/172756409789097450)
- Hudleston PJ (2015) Structures and fabrics in glacial ice: A review. *Journal of Structural Geology*, **81**, 1–27 (doi: 10.1016/j.jsg.2015.09.003)
- Jacobson HP and Waddington ED (2004) Recumbent folding in ice sheets: a core-referential study. *Journal of Glaciology*, **50**(168), 3–16 (doi: 10.3189/172756504781830204)
- Jacobson HP and Waddington ED (2005) Recumbent folding of divide arches in response to unsteady ice-divide migration. *Journal of Glaciology*, **51**(173), 201–209 (doi: 10.3189/172756505781829412)
- Jennings SJA and Hambrey MJ (2021) Structures and deformation in glaciers and ice sheets. *Reviews of Geophysics*, **59**(3), e2021RG000743 (doi: 10.1029/2021RG000743)
- Kehrl L, Conway H, Holschuh N, Campbell S, Kurbatov AV and Spaulding NE (2018) Evaluating the duration and continuity of potential climate records from the allan hills blue ice area, east antarctica. *Geophysical Research Letters*, **45**(9), 4096–4104 (doi: 10.1029/2018GL077511)
- Kirkpatrick LR, Carter AJ, Marks-Peterson J, Shackleton S and Fudge TJ (2025) Three-dimensional multitrack electrical conductivity method for interpretation of complex ice core stratigraphy. *Journal of Glaciology*, **71**, e105 (doi: 10.1017/jog.2025.10081)
- Leysinger Vieli GJMC, Martín C, Hindmarsh RCA and Lüthi MP (2018) Basal freeze-on generates complex ice-sheet stratigraphy. *Nature Communications*, **9**, 4669 (doi: 10.1038/s41467-018-07083-3)
- Lilien DA, Steinhage D, Taylor D, Parrenin F, Ritz C, Mulvaney R, Martín C, Yan JB, O’Neill C, Frezzotti M, Miller H, Gogineni P, Dahl-Jensen D and Eisen O (2021) Brief communication: New radar constraints support presence of ice older than 1.5 Myr at little dome c. *The Cryosphere*, **15**(4), 1881–1888 (doi: 10.5194/tc-15-1881-2021)
- Manos JM, Shaya M, Kirkpatrick L, Stoll N, Banerjee A, Hishamunda V, Horlings AN, Hudak A, Shapero DR, Fudge TJ, Higgins JA, Steig EJ, Brook EJ and Lipovsky BP (2026) Ice flow controls on old ice preservation in the Allan Hills, Antarctica, preprint
- Martín C, Gudmundsson GH, Pritchard HD and Gagliardini O (2009a) On the effects of anisotropic rheology on ice flow, internal structure, and the age–depth relationship at ice divides. *Journal of Geophysical Research: Earth Surface*, **114**(F4), F04001 (doi: 10.1029/2008JF001204)
- Martín C, Hindmarsh RCA and Navarro FJ (2009b) On the effects of divide migration, along-ridge flow, and basal sliding on isochrones near an ice divide. *Journal of Geophysical Research: Earth Surface*, **114**(F2), F02006 (doi: 10.1029/2008JF001025)
- Meyer CR and Creyts TT (2017) Formation of ice eddies in subglacial mountain valleys. *Journal of Geophysical Research: Earth Surface*, **122**(9), 1574–1588 (doi: 10.1002/2017JF004329)

- 589 Moffatt HK (1964) Viscous and resistive eddies near a sharp corner. *Journal of Fluid Mechanics*, **18**(1), 1–18 (doi:
590 10.1017/S0022112064000015)
- 591 Nye JF (1963) Correction factor for accumulation measured by the thickness of the annual layers in an ice sheet.
592 *Journal of Glaciology*, **4**(36), 785–788 (doi: 10.3189/S0022143000028367)
- 593 Parrenin F (2013) *Sur l'âge de la glace et des bulles d'air dans les calottes polaires*. Ph.D. thesis, Université Joseph
594 Fourier, Grenoble, France
- 595 Parrenin F and Hindmarsh RCA (2007) Influence of a non-uniform velocity field on isochrone geometry along a
596 steady flowline of an ice sheet. *Journal of Glaciology*, **53**(183), 612–622 (doi: 10.3189/002214307784409298)
- 597 Parrenin F, Hindmarsh RCA and Rémy F (2006) Analytical solutions for the effect of topography, accumulation
598 rate and lateral flow divergence on isochrone layer geometry. *Journal of Glaciology*, **52**(177), 191–202 (doi:
599 10.3189/172756506781828728)
- 600 Parrenin F, Chung A and Martín C (2025) age_flow_line-1.0: a fast and accurate numerical age model for a pseudo-
601 steady flow tube of an ice sheet. *Geoscientific Model Development*, **18**, 8203–8216 (doi: 10.5194/gmd-18-8203-2025)
- 602 Schoof C and Hindmarsh RCA (2010) Thin-film flows with wall slip: An asymptotic analysis of higher order
603 glacier flow models. *The Quarterly Journal of Mechanics and Applied Mathematics*, **63**(1), 73–114 (doi:
604 10.1093/qjmam/hbp025)
- 605 Shackleton S, Hishamunda V, Davidge L, Brook EJ, Marks-Peterson J, Carter A, Aarons S, Kurbatov AV, Introne DS,
606 Yan Y, Nesbitt IM, Buizert C, Steig EJ, Schauer AJ, Morgan J, Neff PD, Epifanio JA, Severinghaus JP, Bender
607 ML and Higgins JA (2025) Miocene and pliocene ice and air from the allan hills blue ice area, east antarctica.
608 *Proceedings of the National Academy of Sciences*, **122**(44), e2502681122 (doi: 10.1073/pnas.2502681122)
- 609 Spaulding NE, Spikes VB, Hamilton GS, Mayewski PA, Dunbar NW, Harvey RP, Schutt J and Kurbatov AV
610 (2012) Ice motion and mass balance at the allan hills blue-ice area, antarctica, with implications for paleoclimate
611 reconstructions. *Journal of Glaciology*, **58**(208), 399–406 (doi: 10.3189/2012JoG11J176)
- 612 Stoll N, Shaya M, Kirkpatrick L, Freitag J, Hishamunda V, Manos JM, Jansen D, Weikusat I, Fegyveresi J, Lipovsky B,
613 Brook EJ, Shackleton S, Higgins JA and Fudge TJ (2026) Fabric, texture, and bubble characteristics of the million-
614 year old allan hills blue ice core ALHIC1901. *Journal of Geophysical Research: Solid Earth*, **131**(5), e2026JB033890
615 (doi: 10.1029/2026JB033890)
- 616 Thorsteinsson T and Waddington ED (2002) Folding in strongly anisotropic layers near ice-sheet centers. *Annals of*
617 *Glaciology*, **35**, 480–486 (doi: 10.3189/172756402781816708)
- 618 Waddington ED (2010) Life, death and afterlife of the extrusion flow theory. *Journal of Glaciology*, **56**(200), 973–996
619 (doi: 10.3189/002214311796406022)
- 620 Waddington ED, Bolzan JF and Alley RB (2001) Potential for stratigraphic folding near ice-sheet centers. *Journal*
621 *of Glaciology*, **47**(159), 639–648 (doi: 10.3189/172756501781831756)

- 622 Wolovick MJ and Creyts TT (2016) Overturned folds in ice sheets: Insights from a kinematic model of traveling sticky
 623 patches and comparisons with observations. *Journal of Geophysical Research: Earth Surface*, **121**(5), 1065–1083
 624 (doi: 10.1002/2015JF003698)
- 625 Wolovick MJ, Creyts TT, Buck WR and Bell RE (2014) Traveling slippery patches produce thickness-scale folds in
 626 ice sheets. *Geophysical Research Letters*, **41**(24), 8895–8901 (doi: 10.1002/2014GL062248)
- 627 Yan Y, Bender ML, Brook EJ, Clifford HM, Kemeny PC, Kurbatov AV, Mackay S, Mayewski PA, Ng J, Severinghaus
 628 JP and Higgins JA (2019) Two-million-year-old snapshots of atmospheric gases from antarctic ice. *Nature*, **574**,
 629 663–666 (doi: 10.1038/s41586-019-1692-3)
- 630 Zhang Y, Sachau T, Franke S, Yang H, Li D, Weikusat I and Bons PD (2024) Formation mechanisms
 631 of large-scale folding in greenland’s ice sheet. *Geophysical Research Letters*, **51**(16), e2024GL109492 (doi:
 632 10.1029/2024GL109492)

633 Appendix A. INVERSE-COORDINATE IDENTITIES

Equation (8) can be written

$$\begin{bmatrix} \tau_x & \tau_z \\ \psi_x & \psi_z \end{bmatrix} \begin{bmatrix} dx \\ dz \end{bmatrix} = \begin{bmatrix} d\tau \\ d\psi \end{bmatrix}, \quad \det = 1. \quad (\text{A1})$$

The inverse transformation is

$$\begin{bmatrix} dx \\ dz \end{bmatrix} = \begin{bmatrix} \psi_z & -\tau_z \\ -\psi_x & \tau_x \end{bmatrix} \begin{bmatrix} d\tau \\ d\psi \end{bmatrix} = \begin{bmatrix} u & -\tau_z \\ w & \tau_x \end{bmatrix} \begin{bmatrix} d\tau \\ d\psi \end{bmatrix}. \quad (\text{A2})$$

Therefore

$$x_\tau|_\psi = u, \quad z_\tau|_\psi = w, \quad x_\psi|_\tau = -\tau_z, \quad z_\psi|_\tau = \tau_x. \quad (\text{A3})$$

- 634 The first two identities state that advancing one unit in age at fixed streamline follows the velocity for
 635 one unit of time. The latter two specify the equal-age tangent. Multiplication by $d\psi/dx_0 = \dot{a}_0$ yields
 636 Equation (19) and the corresponding vertical tangent component.

637 Appendix B. EXTENSION TO STEADY VARIABLE-DENSITY FLOW

Below the firn, glacier ice is accurately treated as incompressible with nearly constant density. The coordinate construction extends to a steady variable-density flow satisfying

$$\nabla \cdot (\rho \mathbf{u}) = 0. \quad (\text{B4})$$

Define the mass-flux streamfunction by

$$\rho u = \psi_z, \quad \rho w = -\psi_x. \quad (\text{B5})$$

Then

$$\begin{vmatrix} \tau_x & \tau_z \\ \psi_x & \psi_z \end{vmatrix} = \rho(u\tau_x + w\tau_z) = \rho. \quad (\text{B6})$$

638 The mapping remains nonsingular for $\rho > 0$, and the same order interpretation follows after replacing
 639 volume flux by mass flux. The constant-density formula used in the main text is therefore not conceptually
 640 tied to strict incompressibility, although the weights in the cumulative integral acquire density factors.

641 Appendix C. FIRST-ORDER PERIODIC CANCELLATION ABOUT PLUG FLOW

Let

$$u(x, z; \epsilon_b) = U_b + \epsilon_b u_1(x, z) + O(\epsilon_b^2), \quad U_b > 0, \quad (\text{C7})$$

where U_b is spatially uniform and u_1 is periodic in x with wavelength $2\pi/k$ and zero horizontal mean at each elevation. Then

$$u_z = \epsilon_b u_{1,z} + O(\epsilon_b^2), \quad u^{-3} = U_b^{-3} + O(\epsilon_b). \quad (\text{C8})$$

Because the background integrand is identically zero, the $O(\epsilon_b)$ displacement of the trajectory affects the integral only at $O(\epsilon_b^2)$. Evaluating the first-order term along the unperturbed streamline $z = z_0$ gives

$$\int_x^{x+2\pi/k} \frac{u_z}{u^3} d\xi = \frac{\epsilon_b}{U_b^3} \int_x^{x+2\pi/k} u_{1,z}(\xi, z_0) d\xi + O(\epsilon_b^2) = O(\epsilon_b^2), \quad (\text{C9})$$

642 because differentiation with respect to z preserves the zero horizontal mean of u_1 . This establishes
 643 Equation (57). For a vertically sheared background, trajectory displacement can act on the nonzero
 644 background integrand at first order, so the cancellation does not follow without additional assumptions.