What controls the variability of CO_2 fluxes in eastern boundary upwelling systems?

Riley X. Brady¹, Nicole S. Lovenduski¹, Michael A. Alexander², Michael Jacox³, and Nicolas Gruber⁴

¹University of Colorado Boulder and Institute for Arctic and Alpine Research

²NOAA Boulder

³NOAA SWFSC

⁴ETH Zurich

Correspondence to: Riley X. Brady (riley.brady@colorado.edu)

Abstract. TEXT

Copyright statement. TEXT

- 1 Introduction
- 2 Methods

5 2.1 Model Configuration and Upwelling Regions

We utilize monthly output from 34 members of the Community Earth System Model Large Ensemble Project (CESM-LENS), which is derived from a fully coupled Atmosphere-Ocean General Circulation Model (AOGCM) with ocean biogeochemistry (Kay et al., 2015). Round-off level perturbations are made to the atmospheric temperature in 1920, leading to an ensemble of simulations that diverge solely due to the influence of internally generated variability. This provides us with a set of 34 independent representations of climate variability, with which we can robustly assess the controls on air-sea CO₂ flux variability in EBUS. The ensemble is forced with historical radiative forcing from 1920–2005 and RCP8.5 radiative forcing from 2006–2100. Output is made available at approximately 1°x1° global resolution.

2.2 Statistical Analysis

Air-sea CO₂ fluxes in CESM are computed following the parameterization of Wanninkhof (2014):

15
$$F = k \cdot K_0 \cdot (pCO_2^o - pCO_2^a)$$
 (1)

where k represents the gas transfer velocity (dependent on the wind speed squared), K_0 the solubility of CO_2 in seawater, and pCO_2^o and pCO_2^a the partial pressures of CO_2 in the surface ocean and atmosphere, respectively. Surface ocean pCO_2 tends to be the largest term in this equation, since the variability of ocean pCO_2 is much larger than that of the atmosphere (Takahashi

et al., 2002; Gruber et al., 2002). However, the influence of wind variability on k cannot be ignored in EBUS, since these systems are aligned with the eastern flank of subtropical gyres that exhibit interannual to decadal variability (Hasanean, 2004; Schroeder et al., 2013). Further, both upwelling and biological productivity in EBUS are highly sensitive to the structure, intensity, and persistence of alongshore winds (Enriquez and Friehe, 1995; Capet et al., 2004; Botsford et al., 2006). This sensitivity feeds back onto the CO₂ fluxes, since these processes in turn influence DIC, Alk, and SST.

We use a linear Taylor expansion to quantify the relative contribution of each parameter to internally generated variability in air-sea CO_2 flux following Lovenduski et al. (2007) and Turi et al. (2014).

$$\Delta F = \frac{\partial F}{\partial U} \Delta U + \frac{\partial F}{\partial p C O_2^{oc}} \Delta p C O_2^{oc}$$
(2)

where $\frac{\partial F}{\partial U}$ and $\frac{\partial F}{\partial pCO_2^{oc}}$ are determined from the model equations and mean values in each EBUS, and the influence of sea ice was assumed to be negligible. Δ 's represent the linear regression of the given variable onto a climate index. The contributions from ΔpCO_2^{oc} is further decomposed into DIC, Alk, SST, and salinity terms.

$$\Delta pCO_2^{oc} = \frac{\partial pCO_2^{oc}}{\partial DIC} \Delta DIC + \frac{\partial pCO_2^{oc}}{\partial Alk} \Delta Alk + \frac{\partial pCO_2^{oc}}{\partial T} \Delta T + \frac{\partial pCO_2^{oc}}{\partial S} \Delta S \tag{3}$$

Because DIC and Alk can be diluted by freshwater fluxes, as well as altered by physical circulation and biology, we further partitioned the linear expansion. We introduce salinity-normalized DIC (sDIC) and Alk (sAlk), in addition to a separate freshwater term.

$$\Delta F = \frac{\partial F}{\partial U} \Delta U + \frac{S}{S_0} \frac{\partial F}{\partial DIC} \Delta sDIC + \frac{S}{S_0} \frac{\partial F}{\partial Alk} \Delta sAlk + \frac{\partial F}{\partial fw} \Delta fw + \frac{\partial F}{\partial T} \Delta T + \frac{\partial F}{\partial S} \Delta S \tag{4}$$

To compensate for autocorrelation that is characteristic of climate indices and also introduced from smoothing, we replace the t statistic sample size N with an effective sample size, N_{eff} :

$$N_{eff} = N\left(\frac{1 - r_1 r_2}{1 + r_1 r_2}\right) \tag{5}$$

where r_1 and r_2 are the lag-1 autocorrelation coefficients of the two time series being correlated (Bretherton et al., 1999; Lovenduski and Gruber, 2005). N_{eff} represents the number of statistically independent measurements, thus requiring a higher r-value for a significant correlation with high autocorrelation in one or both time series.

2.3 Model Evaluation

CESM-LENS air-sea CO_2 fluxes were compared to the SOM-FFN (Self-Organizing Map-Feed Forward Network) neural network product (Landschützer et al., 2017) along the four major EBUS outlined by Chavez and Messié (2009). The SOM-FFN was generated by a two step process. First, the global oceans were grouped in 16 biogeochemical provinces, based on common relationships between sea surface temperature, sea surface salinity, mixed layer depth, and pCO_2 climatology from Takahashi et al. (2009). Secondly, nonlinear relationships were determined between an expanded set of predictor variables and the Surface Ocean Carbon Atlas version 4 (Bakker et al., 2016) database of surface ocean CO_2 measurements to interpolate pCO_2 to monthly resolution spanning 1982-2015 at $1^o x 1^o$ global resolution. Extensive details on and validation of the procedure can be found in Landschützer et al. (2013) and Landschützer et al. (2016).

3 Results **Internal Variability in Upwelling Systems** California Current 3.2 We find that natural CO2 flux anomalies in the CalCS are most strongly correlated with the North Pacific Gyre Oscillation (NPGO). On the other hand, correlations between the CalCS and El Nino Southern Oscillation (ENSO) are modest. However, recent literature shows that the CalCS responds uniquely to any given ENSO event (Fiedler and Mantua, 2017; Turi et al., 2017; Frischknecht et al., 2017). 3.3 **Humboldt Current** 3.4 **Canary Current** 3.5 Benguela Current 10 **Conclusions TEXT** Code availability. TEXT Data availability. TEXT Code and data availability. TEXT 15

Appendix A

Author contributions. TEXT

Competing interests. TEXT

 $\mathbf{A1}$

Disclaimer: TEXT

Acknowledgements. TEXT

References

5

20

30

- J. E. Kay, C. Deser, A. Phillips, A. Mai, C. Hannay, G. Strand, J. M. Arblaster, S. C. Bates, G. Danabasoglu, J. Edwards, M. Holland, P. Kushner, J.-F. Lamarque, D. Lawrence, K. Lindsay, A. Middleton, E. Munoz, R. Neale, K. Oleson, L. Polvani, and M. Vertenstein. The community earth system model (cesm) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8):1333–1349, 2015. https://doi.org/10.1175/BAMS-D-13-00255.1. URL https://doi.org/10.1175/BAMS-D-13-00255.1.
- Rik Wanninkhof. Relationship between wind speed and gas exchange over the ocean revisited: Gas exchange and wind speed over the ocean. *Limnology and Oceanography: Methods*, 12(6):351–362, June 2014. ISSN 15415856. https://doi.org/10.4319/lom.2014.12.351. URL http://doi.wiley.com/10.4319/lom.2014.12.351.
- Taro Takahashi, Stewart C. Sutherland, Colm Sweeney, Alain Poisson, Nicolas Metzl, Bronte Tilbrook, Nicolas Bates, Rik Wanninkhof, Richard A. Feely, Christopher Sabine, Jon Olafsson, and Yukihiro Nojiri. Global sea–air CO2 flux based on climatological surface ocean pCO2, and seasonal biological and temperature effects. *Deep Sea Research Part II: Topical Studies in Oceanography*, 49(9):1601–1622, January 2002. ISSN 0967-0645. https://doi.org/10.1016/S0967-0645(02)00003-6. URL http://www.sciencedirect.com/science/article/pii/S0967064502000036.
- Nicolas Gruber, Charles D. Keeling, and Nicholas R. Bates. Interannual Variability in the North Atlantic Ocean Carbon Sink. *Science*, 298(5602):2374–2378, December 2002. ISSN 0036-8075, 1095-9203. https://doi.org/10.1126/science.1077077. URL http://science.sciencemag.org/content/298/5602/2374.
 - H. M. Hasanean. Variability of the north atlantic subtropical high and associations with tropical sea-surface temperature. *International Journal of Climatology*, 24(8):945–957, 2004. ISSN 1097-0088. https://doi.org/10.1002/joc.1042. URL http://dx.doi.org/10.1002/joc.1042.
 - Isaac D. Schroeder, Bryan A. Black, William J. Sydeman, Steven J. Bograd, Elliott L. Hazen, Jarrod A. Santora, and Brian K. Wells. The north pacific high and wintertime pre-conditioning of california current productivity. *Geophysical Research Letters*, 40(3):541–546, 2013. ISSN 1944-8007. https://doi.org/10.1002/grl.50100. URL http://dx.doi.org/10.1002/grl.50100.
- A. G. Enriquez and C. A. Friehe. Effects of Wind Stress and Wind Stress Curl Variability on Coastal Upwelling. *Jour-* nal of Physical Oceanography, 25(7):1651–1671, July 1995. ISSN 0022-3670, 1520-0485. https://doi.org/10.1175/1520-0485(1995)025<1651:EOWSAW>2.0.CO;2. URL http://journals.ametsoc.org/doi/abs/10.1175/1520-0485%281995%29025%3C1651%3AEOWSAW%3E2.0.CO%3B2.
 - X. J. Capet, P. Marchesiello, and J. C. McWilliams. Upwelling response to coastal wind profiles. Geophys. Res. Lett., 31(13):L13311, July 2004. ISSN 1944-8007. https://doi.org/10.1029/2004GL020123. URL http://onlinelibrary.wiley.com/doi/10.1029/2004GL020123/abstract.
 - Louis W. Botsford, Cathryn A. Lawrence, Edward P. Dever, Alan Hastings, and John Largier. Effects of variable winds on biological productivity on continental shelves in coastal upwelling systems. *Deep Sea Research Part II: Topical Studies in Oceanography*, 53(25): 3116 3140, 2006. ISSN 0967-0645. https://doi.org/https://doi.org/10.1016/j.dsr2.2006.07.011. URL http://www.sciencedirect.com/science/article/pii/S0967064506002281. The Role of Wind-Driven Flow in Shelf Productivity.
- Nicole S. Lovenduski, Nicolas Gruber, Scott C. Doney, and Ivan D. Lima. Enhanced CO2 outgassing in the Southern Ocean from a positive phase of the Southern Annular Mode. *Global Biogeochem. Cycles*, 21(2):GB2026, June 2007. ISSN 1944-9224. https://doi.org/10.1029/2006GB002900. URL http://onlinelibrary.wiley.com/doi/10.1029/2006GB002900/abstract.

- G. Turi, Z. Lachkar, and N. Gruber. Spatiotemporal variability and drivers of pCO2 and air–sea CO2 fluxes in the California Current System: an eddy-resolving modeling study. *Biogeosciences*, 11(3):671–690, February 2014. ISSN 1726-4189. https://doi.org/10.5194/bg-11-671-2014. URL http://www.biogeosciences.net/11/671/2014/.
- Christopher S. Bretherton, Martin Widmann, Valentin P. Dymnikov, John M. Wallace, and Ileana Bladé. The Effective Number of Spatial Degrees of Freedom of a Time-Varying Field. *Journal of Climate*, 12(7):1990–2009, July 1999. ISSN 0894-8755, 1520-0442. https://doi.org/10.1175/1520-0442(1999)012<1990:TENOSD>2.0.CO;2. URL http://journals.ametsoc.org/doi/abs/10.1175/1520-0442% 281999%29012%3C1990%3ATENOSD%3E2.0.CO%3B2.

5

10

15

20

- Nicole S. Lovenduski and Nicolas Gruber. Impact of the Southern Annular Mode on Southern Ocean circulation and biology. *Geophys. Res. Lett.*, 32(11):L11603, June 2005. ISSN 1944-8007. https://doi.org/10.1029/2005GL022727. URL http://onlinelibrary.wiley.com/doi/10.1029/2005GL022727/abstract.
- Peter Landschützer, Nicolas Gruber, and Dorothee C. E. Bakker. An updated observation-based global monthly gridded sea surface pco2 and air-sea co2 flux product from 1982 through 2015 and its monthly climatology (ncei accession 0160558). Version 2.2. NOAA National Centers for Environmental Information. Dataset., July 2017.
- Francisco P. Chavez and Monique Messié. A comparison of eastern boundary upwelling ecosystems. *Progress in Oceanography*, 83(1):80 96, 2009. ISSN 0079-6611. https://doi.org/10.1016/j.pocean.2009.07.032. URL http://www.sciencedirect.com/science/article/pii/S0079661109000998. Eastern Boundary Upwelling Ecosystems: Integrative and Comparative Approaches.
 - Taro Takahashi, Stewart C. Sutherland, Rik Wanninkhof, Colm Sweeney, Richard A. Feely, David W. Chipman, Burke Hales, Gernot Friederich, Francisco Chavez, Christopher Sabine, Andrew Watson, Dorothee C. E. Bakker, Ute Schuster, Nicolas Metzl, Hisayuki Yoshikawa-Inoue, Masao Ishii, Takashi Midorikawa, Yukihiro Nojiri, Arne Körtzinger, Tobias Steinhoff, Mario Hoppema, Jon Olafsson, Thorarinn S. Arnarson, Bronte Tilbrook, Truls Johannessen, Are Olsen, Richard Bellerby, C. S. Wong, Bruno Delille, N. R. Bates, and Hein J. W. de Baar. Climatological mean and decadal change in surface ocean pCO2, and net sea–air CO2 flux over the global oceans. *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(8):554–577, April 2009. ISSN 0967-0645. https://doi.org/10.1016/j.dsr2.2008.12.009. URL http://www.sciencedirect.com/science/article/pii/S0967064508004311.
- Dorothee C E Bakker, Benjamin Pfeil, Kevin M O'Brien, Kim I Currie, Stephen D Jones, Camilla S Landa, Siv K Lauvset, Nicolas Metzl, 25 David R Munro, Shin-Ichiro Nakaoka, Are Olsen, Denis Pierrot, Shu Saito, Karl Smith, Colm Sweeney, Taro Takahashi, Chisato Wada, Rik Wanninkhof, Simone R Alin, Meike Becker, Richard G J Bellerby, Alberto Vieira Borges, Jacqueline Boutin, Yann Bozec, Eugene Burger, Wei-Jun Cai, Robert D Castle, Catherine E Cosca, Michael D DeGrandpre, Matthew Donnelly, Greg Eischeid, Richard A Feely, Thanos Gkritzalis, Melchor González-Dávila, Catherine Goyet, Antoine Guillot, Nicolas J Hardman-Mountford, Judith Hauck, Mario Hoppema, Matthew P Humphreys, Christopher W Hunt, J Severino P Ibánhez, Tadafumi Ichikawa, Masao Ishii, Lauren W Juranek, Vas-30 silis Kitidis, Arne Körtzinger, Urbain K Koffi, Alexander Kozyr, Akira Kuwata, Nathalie Lefèvre, Claire Lo Monaco, Ansley Manke, Pierre Marrec, Jeremy T Mathis, Frank J Millero, Natalie Monacci, Pedro M S Monteiro, Akihiko Murata, Timothy Newberger, Yukihiro Nojiri, Isao Nonaka, Abdirahman M Omar, Tsuneo Ono, Xose Antonio Padín, Gregor Rehder, Anna Rutgersson, Christopher L Sabine, Joe Salisbury, Juana Magdalena Santana-Casiano, Daisuke Sasano, Ute Schuster, Rainer Sieger, Ingunn Skjelvan, Tobias Steinhoff, Kevin Sullivan, Stewart C Sutherland, Adrienne Sutton, Kazuaki Tadokoro, Maciej Telszewski, Helmuth Thomas, Bronte Tilbrook, 35 Steven Van Heuven, Doug Vandemark, Douglas WR Wallace, and Ryan Woosley. Surface Ocean CO2 Atlas (SOCAT) V4, 2016. DOI: 10.1594/PANGAEA.866856.

- P. Landschützer, N. Gruber, D. C. E. Bakker, U. Schuster, S. Nakaoka, M. R. Payne, T. P. Sasse, and J. Zeng. A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink. Biogeosciences, 10(11):7793–7815, November 2013. ISSN 1726-4189. https://doi.org/10.5194/bg-10-7793-2013. URL https://www.biogeosciences.net/10/7793/2013/.
- Peter Landschützer, Nicolas Gruber, and Dorothee C. E. Bakker. Decadal variations and trends of the global ocean carbon sink. Global Biogeochemical Cycles, 30(10):1396–1417, 2016. ISSN 1944-9224. https://doi.org/10.1002/2015GB005359. URL http://dx.doi.org/10. 1002/2015GB005359. 2015GB005359.
- Paul C. Fiedler and Nathan J. Mantua. How are warm and cool years in the California Current related to ENSO? Journal of Geophysical Research: Oceans, 122(7):5936-5951, July 2017. ISSN 21699275. https://doi.org/10.1002/2017JC013094. URL http://doi.wiley.com/ 10.1002/2017JC013094.
- G. Turi, Michael A. Alexander, Nicole S. Lovenduski, Antonietta Capotondi, James D. Scott, Charles A. Stock, John P. Dunne, Jasmin John, and Michael G. Jacox. Response of O2 and pH to ENSO in the California Current System in a high resolution global climate model. Ocean Sciences Discussion, August 2017.
 - Martin Frischknecht, Matthias Münnich, and Nicolas Gruber. Local atmospheric forcing driving an unexpected California Current System response during the 2015-2016 El Niño. Geophys. Res. Lett., 44(1):2016GL071316, January 2017. ISSN 1944-8007.
- 15 https://doi.org/10.1002/2016GL071316. URL http://onlinelibrary.wiley.com/doi/10.1002/2016GL071316/abstract.

5