


Guide-support robustness of the actual constant W/HW blend


Own-baseline gains — I: primary +0.001954, identity +0.003575; R: primary +0.001489, identity +0.002917.

21–51 electrodes/session; original reference margins 0.010 primary / 0.005 identity. Changed-support sensitivities cannot reset the original-support incumbent.









Mihili draw exceptions (IDs 0–7): nonpositive means, primary [4], identity [none]; nonpositive omission in primary [2,4], identity [4].

Chewie draw exceptions (IDs 0–7): nonpositive means, primary [none], identity [none]; nonpositive omission in primary [none], identity [none].

◆ Mihili mean ◆ Chewie mean ■ Equal-animal mean ○ Individual draw mean (D)

A/B small dots: all 16 signed session effects; solid bars: fixed-prediction session-omission ranges. Both panels share one complete linear scale.
D uses its own full signed scale: open circles/dotted spans show all 8 draw means/their range; filled markers show the mean-R contrast.
These ranges are not confidence intervals or p-values. Overlapping support views are not biological replicates: 2 animals, with 6 and 10 sessions.
I uses recorded guide membership; R uses the exact eight ordinary subsets of equal size. True trial pairing remains intact; these are not pairing-null models.
Full blend and prior I/R-HW keep their original selections. New-λ own-HW comparisons use I-blend's selected penalty; both versions remain separately displayed.
Only reduced current-W states are newly fit; saved HW predictions stay fixed. Mixtures retain error cross-terms, and R averages SSEs rather than predictions.
The compact source retains full-native-error cross-products and separate W/HW trace/df: neither is an isolated neural mechanism. Equal counts or λ do not match capacity.
Guide membership does not identify spike contamination. Exposed development data; no promotion. CSV includes every draw's sessions/omissions and selected α/λ values.