A time-varying mixture does not improve the constant blend's primary score

Primary 0.002647 < 0.010; identity 0.004602 < 0.005. Primary change versus blend -0.0000322.

All λ=1 · nine folds reuse the constant half-blend exactly · seven choose HW weight 0.25 → 0.75

A Selected coefficients and fixed reversal



B Other-15-session selections: 4 Mihili + 3 Chewie active

C Direct primary prediction differences



D Direct trial-pairing differences



Every active Chewie session loses primary accuracy against both constant and reversed schedules.

Chewie identity change versus constant is only +0.0000064; omitting 2015-06-29 makes it negative.

◆ Mihili mean ◆ Chewie mean ■ Equal-animal mean

C/D small dots: all 16 signed session effects. Solid bars: fixed-prediction session-omission ranges; not confidence intervals or p-values.
Both effect panels share one full signed linear scale. Two animals (6 and 10 sessions), equally weighted after within-animal averaging.
The reversal is an unretuned descriptive contrast, not an independent control or extra trial. At zero slope it is literally the same prediction.
Identity is affine in prediction: selected-minus-reversed is exactly twice selected-minus-half-blend here, not independent corroboration.
All original W/HW predictions, trials, native XY baseline and denominator remain fixed. α was selected using only the other fifteen sessions.
A fixed mean mixing coefficient does not fix mean prediction contribution or capacity. These schedules do not identify physiological timing or causal lag effects.
Exposed development data · both practical margins still fail · no incumbent reset · CSV preserves all selected coefficients, session effects and omissions.