

Lecture 10: LU decomposition Revisited

Lecturer: Prof. Yue Qiu & Prof. Ziping Zhao

Scribe: Zhihang Xu

Disclaimer: These notes have not been subjected to the usual scrutiny reserved for formal publications. They may be distributed outside this class only with the permission of the Instructor.

The intention of this notes is to further explain the definition of LU decomposition and to modify some ambiguity in [Homework 2](#).

Definition 1 (General definition of LU decomposition¹). A matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is said to have an LU decomposition/factorization if it can be factored as

$$\mathbf{A} = \mathbf{L}\mathbf{U},$$

where $\mathbf{L} \in \mathbb{R}^{n \times n}$ is unit lower triangular (lower triangular with unit diagonal elements, i.e., $\ell_{ii} = 1$ for all i), and $\mathbf{U} \in \mathbb{R}^{n \times n}$ is upper triangular.

In this case, a natural question is, does every matrix have an LU decomposition? The answer is no.

Theorem 1. If $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $\det(\mathbf{A}_{\{1, \dots, k\}}) \neq 0$ for $k = 1, \dots, n-1$, then \mathbf{A} has an LU decomposition. If this is the case and \mathbf{A} is nonsingular, then the factorization is unique².

So far, we have shown the sufficient condition for the LU decomposition. However, what if \mathbf{A} is singular, does \mathbf{A} still admit an LU decomposition?

From the general definition of LU decomposition, as long as we can find matrices \mathbf{L} and \mathbf{U} which are unit lower triangular and upper triangular such that $\mathbf{A} = \mathbf{L}\mathbf{U}$, we can claim that \mathbf{A} has LU decomposition. For example, recall [Problem 2 of Homework 2](#), consider the 4×4 matrix

$$\begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix}$$

we can find its LU decomposition via

$$\begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & 0 & d-c \end{bmatrix}. \quad (1)$$

When $a = 0$, or $b = a$, or $c = b$, \mathbf{U} is singular and consequently \mathbf{A} is singular. Therefore, even singular matrix can have an LU decomposition in the general sense. And in this case, is the LU

¹This definition is consistent with the definition given in Slides.

²The proof of this theorem is omitted here, you can find the proof in Section 3.2.5 of [2].

decomposition unique? The answer is no. For example, when $a = 0$ and b, c, d are randomly given, then \mathbf{A} can be factored as

$$\begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ \ell_{21} & 1 & 0 & 0 \\ \ell_{31} & 1 & 1 & 0 \\ \ell_{41} & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & 0 & d-c \end{bmatrix},$$

where $\ell_{21}, \ell_{31}, \ell_{41}$ can take any values. Therefore, such LU decomposition is not unique.

Definition 2 (General definition of LDM/LDU decomposition). *A matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is said to have an LDM/LDU decomposition/factorization if it can be factored as*

$$\mathbf{A} = \mathbf{LDM}^T,$$

where $\mathbf{L} \in \mathbb{R}^{n \times n}$ and \mathbf{M} are unit lower triangular (lower triangular with unit diagonal elements), and $\mathbf{D} = \text{diag}(d_1, \dots, d_n)$ is a diagonal matrix.

If \mathbf{A} is nonsingular and has LU decomposition $\mathbf{A} = \mathbf{LU}$ (in this case, the diagonal elements of \mathbf{U} are all nonzero), then the LDM decomposition can be given as

$$\mathbf{A} = \mathbf{LDM}^T,$$

where

$$\mathbf{D} = \text{diag}(u_{11}, \dots, u_{nn}), \quad \mathbf{M} = \mathbf{U}^T \mathbf{D}^{-1},$$

therefore, in the case of nonsingular \mathbf{A} , the existence of LDM decomposition follows the existence of LU decomposition.

In the case of singular \mathbf{A} , take (2) for example, the LDM decomposition can be given by

$$\begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & 0 & 0 & 0 \\ 0 & b-a & 0 & 0 \\ 0 & 0 & c-b & 0 \\ 0 & 0 & 0 & d-c \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}. \quad (2)$$

But if the existence of LDM decomposition follows the existence of LU decomposition in the singular case? Let the LU decomposition for singular \mathbf{A} be given by (denote $\mathbf{L} = (\ell_{i,j})_{n \times n}$, $\mathbf{U} = (u_{i,j})_{n \times n}$).

$$\mathbf{A} = (a_{i,j})_{n \times n} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ \ell_{31} & \ell_{32} & 1 & \\ \vdots & \vdots & & \ddots \\ \ell_{n1} & \ell_{n2} & \cdots & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ & u_{22} & \cdots & u_{2n} \\ & & \ddots & \vdots \\ & & & u_{nn} \end{bmatrix},$$

suppose that $u_{ii} = 0$ for some i , but $u_{i,j} \neq 0$ for some $j \in \{i+1, \dots, n\}$, then consider the LDM

decomposition (denote $\mathbf{M} = (m_{i,j})_{n \times n}$)

$$\mathbf{A} = (a_{i,j})_{n \times n} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ \ell_{31} & \ell_{32} & 1 & \\ \vdots & \vdots & & \ddots \\ \ell_{n1} & \ell_{n2} & \cdots & 1 \end{bmatrix} \begin{bmatrix} u_{11} & & & \\ & u_{22} & & \\ & & \ddots & \\ & & & u_{ii} = 0 \\ & & & \ddots & \\ & & & & u_{nn} \end{bmatrix}$$

$$\begin{bmatrix} 1 & m_{21} & \cdots & m_{n1} \\ & 1 & \cdots & m_{n2} \\ & & \ddots & \vdots \\ & & & 1 \end{bmatrix},$$

which indicates that $u_{ij} = \mathbf{D}(i,:) \mathbf{M}(:,j) = 0$ and it contradicts the assumption. Therefore, in this case, \mathbf{A} which has an LU decomposition does not have an LDM decomposition. Whereas, if the LU decomposition for \mathbf{A} satisfies $u_{ii} = 0$ for some i , and $u_{i,j} = 0$ for $j = i+1, \dots, n$, then the LDM decomposition for \mathbf{A} exists,

$$\mathbf{A} = (a_{i,j})_{n \times n} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ \ell_{31} & \ell_{32} & 1 & \\ \vdots & \vdots & & \ddots \\ \ell_{n1} & \ell_{n2} & \cdots & 1 \end{bmatrix} \begin{bmatrix} u_{11} & & & \\ & u_{22} & & \\ & & \ddots & \\ & & & u_{ii} = 0 \\ & & & \ddots & \\ & & & & u_{nn} \end{bmatrix}$$

$$\begin{bmatrix} 1 & m_{21} = \frac{u_{12}}{u_{11}} & \cdots & m_{n1} = \frac{u_{1n}}{u_{11}} \\ & 1 & \cdots & m_{n2} = \frac{u_{2n}}{u_{22}} \\ & & \ddots & \vdots \\ & & & 1 & m_{i+1,i} & \cdots & m_{n,i} \\ & & & & \ddots & & \vdots \\ & & & & & & 1 \end{bmatrix}$$

where $m_{j,i}$ ($j = i+1, \dots, n$) can take any value, and therefore is not unique in general. To sum up, the problem description in [problem 4 of Homework 2](#) is ambiguous. The modified version can be found in [1](#) (The modified part is marked with red).

However, when discussing LU decomposition, we almost always consider the problem of solving linear equations,

$$\mathbf{Ax} = \mathbf{b},$$

and \mathbf{A} is nonsingular by default. And the singular \mathbf{A} case may be trivial. In other contexts, the definitions of LU decomposition may vary. For example, in section 3.10 of [1], the LU factorization this textbook discussed is restricted to nonsingular \mathbf{A} case. In section 3.2.10 of [2], LU decomposition of a rectangular matrix is defined. You can read section 3.2 of [2] for more details.

1 Problem 4 of Homework 2 modified

Given a matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$, suppose that the LDM (LDU) decomposition of \mathbf{A} exists, prove that

1. suppose \mathbf{A} is nonsingular, then the LDM (LDU) decomposition of \mathbf{A} is *uniquely* determined;
2. if \mathbf{A} is a nonsingular symmetric matrix, then its LDM (LDU) decomposition must be $\mathbf{A} = \mathbf{L}\mathbf{D}\mathbf{L}^T$, which is called LDL decomposition in this case;
3. \mathbf{A} is a symmetric and positive definite matrix if and only if its Cholesky decomposition exists (i.e., there exists a matrix $\mathbf{G} \in \mathbb{R}^{n \times n}$ such that $\mathbf{A} = \mathbf{G}\mathbf{G}^T$, where \mathbf{G} is lower triangular with *positive* diagonal entries);
4. if \mathbf{A} is a symmetric and positive definite matrix, then its Cholesky decomposition is *uniquely* determined.

Hint: You can directly utilize the following lemmas,

- the inverse (if it exists) of a lower (resp. upper) triangular matrix is also lower (resp. upper) triangular;
- the product of two lower (resp. upper) triangular matrices is lower (resp. upper) triangular;
- also, if such two lower (resp. upper) triangular matrices have unit diagonal entries, then their product also has unit diagonal entries.

Solution:

1. Assume that \mathbf{A} has two LDM decompositions as $\mathbf{A} = \mathbf{L}_1\mathbf{D}_1\mathbf{M}_1^T = \mathbf{L}_2\mathbf{D}_2\mathbf{M}_2^T$, we expect to prove that $\mathbf{L}_1 = \mathbf{L}_2$, $\mathbf{D}_1 = \mathbf{D}_2$, and $\mathbf{M}_1 = \mathbf{M}_2$. For nonsingular matrix \mathbf{A} , ~~First, note that the existence of the LDM decomposition implies that \mathbf{A} is nonsingular. Hence,~~ the determinant of \mathbf{A} satisfies $|\mathbf{A}| \neq 0$. Besides, since $|\mathbf{A}| = |\mathbf{L}_1| \times |\mathbf{D}_1| \times |\mathbf{M}_1| = |\mathbf{L}_2| \times |\mathbf{D}_2| \times |\mathbf{M}_2|$, we have that \mathbf{L}_1 , \mathbf{D}_1 , \mathbf{M}_1 , \mathbf{L}_2 , \mathbf{D}_2 , and \mathbf{M}_2 are all nonsingular (i.e., invertible).

Second, since $\mathbf{L}_1\mathbf{D}_1\mathbf{M}_1^T = \mathbf{L}_2\mathbf{D}_2\mathbf{M}_2^T$, we have

$$\mathbf{L}_2^{-1}\mathbf{L}_1 = \mathbf{D}_2\mathbf{M}_2^T(\mathbf{M}_1^T)^{-1}\mathbf{D}_1^{-1}.$$

We can check that the left hand side $\mathbf{L}_2^{-1}\mathbf{L}_1$ is lower triangular, while the right hand side $\mathbf{D}_2\mathbf{M}_2^T(\mathbf{M}_1^T)^{-1}\mathbf{D}_1^{-1}$ is upper triangular. Hence, the left hand side and the right hand side are both diagonal matrices.

Third, the diagonal entries of the left hand side $\mathbf{L}_2^{-1}\mathbf{L}_1$ must be one, which implies $\mathbf{L}_2^{-1}\mathbf{L}_1 = \mathbf{I}$ and accordingly $\mathbf{L}_1 = \mathbf{L}_2$. Similarly, we can derive that $\mathbf{M}_2^T(\mathbf{M}_1^T)^{-1} = \mathbf{D}_2^{-1}\mathbf{L}_2^{-1}\mathbf{L}_1\mathbf{D}_1$, which can further deduce that $\mathbf{M}_1 = \mathbf{M}_2$. Finally, we have $\mathbf{D}_1 = \mathbf{L}_1^{-1}\mathbf{A}(\mathbf{M}_1^T)^{-1} = \mathbf{L}_2^{-1}\mathbf{A}(\mathbf{M}_2^T)^{-1} = \mathbf{D}_2$, which concludes this proof.

2. For nonsingular \mathbf{A} , \mathbf{A} must have LDM decomposition and it is unique. Let $\mathbf{A} = \mathbf{L}\mathbf{D}\mathbf{M}^T$, and since \mathbf{A} is symmetric, we have $\mathbf{L}\mathbf{D}\mathbf{M}^T = \mathbf{A} = \mathbf{A}^T = \mathbf{M}\mathbf{D}\mathbf{L}^T$. From the proof in 1), we learn that \mathbf{L} and \mathbf{M}^T are both invertible. Then we can derive that

$$\mathbf{D}\mathbf{M}^T(\mathbf{L}^T)^{-1} = \mathbf{L}^{-1}\mathbf{M}\mathbf{D}.$$

Note that $(\mathbf{L}^T)^{-1}$ and \mathbf{M}^T are both upper triangular, while \mathbf{L}^{-1} and \mathbf{M} are both lower triangular. We can check that the left hand side $\mathbf{D}\mathbf{M}^T(\mathbf{L}^T)^{-1}$ is upper triangular and the right hand side $\mathbf{L}^{-1}\mathbf{M}\mathbf{D}$ is lower triangular. Hence, the left hand side and the right hand side are both diagonal matrices. Since \mathbf{D} is a diagonal matrix, $\mathbf{L}^{-1}\mathbf{M}$ is also a diagonal matrix. Moreover, the diagonal entries of $\mathbf{L}^{-1}\mathbf{M}$ must be one, which implies $\mathbf{L}^{-1}\mathbf{M} = \mathbf{I}$ and accordingly $\mathbf{L} = \mathbf{M}$.

3. \mathbf{A} is a symmetric and positive definite matrix \Rightarrow its Cholesky decomposition exists)
 Note that a positive definite matrix must be nonsingular. According to the conclusion in 2), we have $\mathbf{A} = \mathbf{L}\mathbf{D}\mathbf{L}^T$. For any vector $\mathbf{x} \in \mathbb{R}^n$, there exists a vector $\mathbf{y} \in \mathbb{R}^n$ such that $\mathbf{y} = \mathbf{L}^T\mathbf{x}$. Since \mathbf{A} is a positive definite matrix, we can derive that

$$\mathbf{y}^T\mathbf{D}\mathbf{y} = \mathbf{x}^T\mathbf{L}\mathbf{D}\mathbf{L}^T\mathbf{x} = \mathbf{x}^T\mathbf{A}\mathbf{x} > 0.$$

Hence, the diagonal entries of \mathbf{D} are all positive. Let \mathbf{D}' denote a diagonal matrix, where $\mathbf{D}'_{ii} = \sqrt{\mathbf{D}_{ii}}$ for $i = 1, 2, 3, \dots, n$. Then we have

$$\mathbf{A} = \mathbf{L}\mathbf{D}\mathbf{L}^T = \mathbf{L}\mathbf{D}'(\mathbf{D}')^T\mathbf{L}^T = \mathbf{L}\mathbf{D}'(\mathbf{L}\mathbf{D}')^T.$$

Let $\mathbf{G} = \mathbf{L}\mathbf{D}'$, then $\mathbf{A} = \mathbf{G}\mathbf{G}^T$, where \mathbf{G} is lower triangular with positive diagonal elements.

(\mathbf{A} is a symmetric and positive definite matrix \Leftarrow its Cholesky decomposition exists)
First, since $\mathbf{A}^T = (\mathbf{G}\mathbf{G}^T)^T = \mathbf{G}\mathbf{G}^T = \mathbf{A}$, we can learn that \mathbf{A} is a symmetric matrix.
Second, for any non-zero vector $\mathbf{x} \in \mathbb{R}^n$, we have

$$\mathbf{x}^T\mathbf{A}\mathbf{x} = \mathbf{x}^T\mathbf{G}\mathbf{G}^T\mathbf{x} = (\mathbf{G}^T\mathbf{x})^T(\mathbf{G}^T\mathbf{x}) > 0,$$

which implies that \mathbf{A} is a positive definite matrix.

4. **First we have that \mathbf{A} is nonsingular since \mathbf{A} is PD.** Assume that \mathbf{A} has two Cholesky decompositions as $\mathbf{A} = \mathbf{G}_1\mathbf{G}_1^T = \mathbf{G}_2\mathbf{G}_2^T$, we expect to prove that $\mathbf{G}_1 = \mathbf{G}_2$.
First, since \mathbf{G}_1 and \mathbf{G}_2 are both lower triangular with positive diagonal entries, \mathbf{G}_1 and \mathbf{G}_2 must be invertible. *Second*, since $\mathbf{A} = \mathbf{G}_1\mathbf{G}_1^T = \mathbf{G}_2\mathbf{G}_2^T$, we have

$$\mathbf{G}_1^{-1}\mathbf{G}_2 = (\mathbf{G}_1^{-1}\mathbf{G}_2)^T.$$

We can check that the left hand side $\mathbf{G}_1^{-1}\mathbf{G}_2$ is lower triangular, while the right hand side $(\mathbf{G}_1^{-1}\mathbf{G}_2)^T$ is upper triangular. Hence, the left hand side and the right hand side are both diagonal matrices. Let $\mathbf{G}_0 = \mathbf{G}_1^{-1}\mathbf{G}_2$, then \mathbf{G}_0 is a diagonal matrix.

Third, by $\mathbf{G}_1\mathbf{G}_1^T = \mathbf{G}_2\mathbf{G}_2^T$, we can also derive that

$$\mathbf{I} = \mathbf{G}_1^{-1}\mathbf{G}_2\mathbf{G}_2^T(\mathbf{G}_1^T)^{-1} = \mathbf{G}_0\mathbf{G}_0^T.$$

Hence, the diagonal entries of \mathbf{G}_0 must be 1 or -1 . Note that $\mathbf{G}_2 = \mathbf{G}_1\mathbf{G}_0$. Considering the diagonal entries of \mathbf{G}_2 and \mathbf{G}_1 , we have $(\mathbf{G}_2)_{ii} = (\mathbf{G}_1)_{ii} \times (\mathbf{G}_0)_{ii}$ for $i = 1, 2, \dots, n$. Since the diagonal entries of \mathbf{G}_2 and \mathbf{G}_1 are required to be positive, $(\mathbf{G}_0)_{ii}$ can only be 1 for $i = 1, 2, \dots, n$, i.e., $\mathbf{G}_0 = \mathbf{I}$. Accordingly, $\mathbf{G}_1 = \mathbf{G}_2$, which concludes this proof.

References

- [1] Meyer, Carl D. Matrix analysis and applied linear algebra. Vol. 71. Siam, 2000.
- [2] Gene H. Golub, Chales F. Van Loan. Matrix Computations. The John Hopkins University Press.