

SI231b Review: SVD related topic

Mao Song

ShanghaiTech University

maosong@shanghaitech.edu.cn

December 10, 2020

Overview

- 1 Main Results
- 2 Computation
- 3 Relation to Linear System
- 4 Relation to Matrix Norm
- 5 Relation to Subspace
- 6 Application

Definition

Given a matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$, the Singular Value Decomposition (SVD) of \mathbf{A} is given by

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T \quad (1)$$

where $\mathbf{U} \in \mathbb{R}^{m \times m}$ and $\mathbf{V} \in \mathbb{R}^{n \times n}$ are orthonormal matrices. $\mathbf{\Sigma} \in \mathbb{R}^{m \times n}$ is a diagonal matrix, and $\Sigma_{ii} = \sigma_i(\mathbf{A})$ with $\sigma_1(\mathbf{A}) \geq \cdots \geq \sigma_{\min\{m,n\}}(\mathbf{A}) \geq 0$. This type of SVD is also called Full SVD.

Relatively, we have Thin SVD or Truncated SVD of \mathbf{A} (Suppose that $\text{rank}(\mathbf{A}) = r$), which is given by (more economical!)

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T \quad (2)$$

where $\mathbf{U} \in \mathbb{R}^{m \times r}$ and $\mathbf{V} \in \mathbb{R}^{n \times r}$ are semi-orthogonal matrices ($\mathbf{U}^T \mathbf{U} = \mathbf{I}_r$, $\mathbf{V}^T \mathbf{V} = \mathbf{I}_r$). $\mathbf{\Sigma} \in \mathbb{R}^{r \times r}$ is a diagonal matrix, and $\Sigma_{ii} = \sigma_i(\mathbf{A})$ with $\sigma_1(\mathbf{A}) \geq \cdots \geq \sigma_r(\mathbf{A}) > 0$.

Basic Properties

- The columns of \mathbf{U} and the columns of \mathbf{V} are called the **left-singular vectors** and **right-singular vectors** of \mathbf{A} , respectively.
- From the SVD point of view, $x \rightarrow \mathbf{A}x$ can be seen as : rotating x (\mathbf{V}), scaling($\mathbf{\Sigma}$), and rotating again(\mathbf{U}). *Thus, the singular values can be interpreted as the magnitude of the semiaxes of an ellipse.*
- SVD of a matrix always exists.
- The outer product form: $\mathbf{A} = \sum_{i=1}^r \sigma_i(\mathbf{A}) \mathbf{u}_i \mathbf{v}_i^T$.
- The SVD of a matrix is not unique. ($\mathbf{\Sigma}$ is unique if Σ_{ii} sorted in descending order, but \mathbf{U} and \mathbf{V} are not unique).
- When we talk about SVD, we often assume its singular values are non-negative and sorted in descending order!

Basic Idea

Suppose $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\text{rank}(\mathbf{A}) = r$.

The eigen-decompositions of $\mathbf{A}\mathbf{A}^T$ and $\mathbf{A}^T\mathbf{A}$ are given by

$$\mathbf{U}^T \mathbf{A} \mathbf{A}^T \mathbf{U} = \text{diag}(\sigma_1^2(\mathbf{A}), \dots, \sigma_r^2(\mathbf{A}), \underbrace{0, \dots, 0}_{m-r})$$

$$\mathbf{V}^T \mathbf{A}^T \mathbf{A} \mathbf{V} = \text{diag}(\sigma_1^2(\mathbf{A}), \dots, \sigma_r^2(\mathbf{A}), \underbrace{0, \dots, 0}_{n-r})$$

where \mathbf{U} and \mathbf{V} are eigenvectors of $\mathbf{A}\mathbf{A}^T$ and $\mathbf{A}^T\mathbf{A}$ respectively.

How to compute SVD by hand

To compute the SVD of $\mathbf{A} \in \mathbb{R}^{m \times n}$ by hand, execute the following steps:

- 1 Calculate $\mathbf{A}\mathbf{A}^T$.
- 2 Compute the eigen-decomposition of $\mathbf{A}\mathbf{A}^T$

$$\begin{bmatrix} \mathbf{U}_1^T \\ \mathbf{U}_2^T \end{bmatrix} \mathbf{A}\mathbf{A}^T \begin{bmatrix} \mathbf{U}_1 & \mathbf{U}_2 \end{bmatrix} = \begin{bmatrix} \Delta_r^2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad (3)$$

where $\mathbf{U}_1 \in \mathbb{R}^{m \times r}$ is the matrix of eigenvectors corresponding to Δ_r^2 .

- 3 Let $\mathbf{V}_1 = \mathbf{A}^T \mathbf{U}_1 \Delta_r^{-1}$, and find $\mathbf{V}_2 \in \mathbb{R}^{n \times (n-r)}$ such that $[\mathbf{V}_1, \mathbf{V}_2]$ is an orthonormal matrix. (Eigenvectors of $\mathbf{A}^T \mathbf{A}$ corresponding to eigenvalue 0)
- 4 Extend Δ_r to $\Sigma \in \mathbb{R}^{m \times n}$. Finally, the Full SVD of \mathbf{A} is given by $\mathbf{A} = \mathbf{U}\Sigma\mathbf{V}^T$, where

$$\mathbf{U} = \begin{bmatrix} \mathbf{U}_1 & \mathbf{U}_2 \end{bmatrix} \in \mathbb{R}^{m \times m}, \mathbf{V} = \begin{bmatrix} \mathbf{V}_1 & \mathbf{V}_2 \end{bmatrix} \in \mathbb{R}^{n \times n}, \Sigma = \begin{bmatrix} \Delta_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{m \times n} \quad (4)$$

Example

Find the SVD of \mathbf{A} , where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \\ 1 & 1 \end{bmatrix} \quad (5)$$

Solution Step 1

$\mathbf{A}\mathbf{A}^T$ is given by

$$\mathbf{A}\mathbf{A}^T = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 0 & 2 \end{bmatrix} \quad (6)$$

Example

Solution Step 2

$\det(\lambda \mathbf{I} - \mathbf{A}\mathbf{A}^T) = \lambda^2(\lambda - 4)$. thus the eigenvalues of $\mathbf{A}\mathbf{A}^T$ are $\lambda_1 = 4$, $\lambda_2 = \lambda_3 = 0$, and $\sigma_1(\mathbf{A}) = \sqrt{4} = 2$, $\mathbf{\Delta}_1 = [2]$ ($r = 1$).

The unit eigenvector of $\mathbf{A}\mathbf{A}^T$ corresponding to $\lambda_1 = 4$ is given by $\mathbf{u}_1 = [1/\sqrt{2}, 0, 1/\sqrt{2}]^T$, $\mathbf{U}_1 = \mathbf{u}_1 = [1/\sqrt{2}, 0, 1/\sqrt{2}]^T$.

Solution Step 3

$$\mathbf{V}_1 = \mathbf{A}^T \mathbf{U}_1 \mathbf{\Delta}_1^{-1} = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}.$$

Thus the Thin SVD of \mathbf{A} is given by $\mathbf{A} = \mathbf{U}_1 \mathbf{\Delta}_1 \mathbf{V}_1^T$

Example

Solution step 3

Solve the system $\mathbf{A}\mathbf{A}^T\mathbf{v} = 0\mathbf{v}$ to obtain the eigenvectors $\mathbf{U}_2 \in \mathbb{R}^{3 \times 2}$ corresponding to 0. Then

- ① use the Gram-Schmidt process to orthogonalize \mathbf{U}_2 .
- ② normalize the column vectors of \mathbf{U}_2 .

we have

$$\mathbf{U}_2 = \begin{bmatrix} -1/\sqrt{2} & 0 \\ 0 & 1 \\ 1/\sqrt{2} & 0 \end{bmatrix} \quad (7)$$

then we solve the system $\mathbf{A}^T\mathbf{A}\mathbf{v} = 0\mathbf{v}$ to obtain the eigenvectors $\mathbf{V}_2 \in \mathbb{R}^{2 \times 1}$ corresponding to 0. we have $\mathbf{V}_2 = [-1/\sqrt{2}, 1/\sqrt{2}]^T$

Example

Solution Step 4

Thus, the Full SVD of \mathbf{A} is given by

$$\mathbf{A} = [\mathbf{U}_1 \quad \mathbf{U}_2] \mathbf{\Sigma} [\mathbf{V}_1 \quad \mathbf{V}_2]^T \quad (8)$$

$$= \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & 1 \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}^T \quad (9)$$

Note

- 1 Do not forget the normalization.
- 2 Corresponds the eigenvector and the eigenvalue.

Pseudo inverse

Definition

Given a matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$, the **pseudo inverse** of \mathbf{A} , $\mathbf{A}^\dagger \in \mathbb{R}^{n \times m}$ is a matrix satisfying the *Penrose-Moore equations*:

$$\begin{aligned}\mathbf{A}\mathbf{A}^\dagger\mathbf{A} &= \mathbf{A}, & \mathbf{A}^\dagger\mathbf{A}\mathbf{A}^\dagger &= \mathbf{A}^\dagger \\ (\mathbf{A}\mathbf{A}^\dagger)^T &= \mathbf{A}\mathbf{A}^\dagger, & (\mathbf{A}^\dagger\mathbf{A})^T &= \mathbf{A}^\dagger\mathbf{A}\end{aligned}$$

Pseudo inverse & SVD

If SVD of \mathbf{A} is given by $\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$, then

$$\mathbf{A}^\dagger = \mathbf{V}\mathbf{\Sigma}^\dagger\mathbf{U}^T \quad (10)$$

where $\mathbf{\Sigma}^\dagger \in \mathbb{R}^{n \times m}$ is diagonal matrix such that

$$\Sigma_{ii}^\dagger = \begin{cases} \frac{1}{\Sigma_{ii}}, & \text{if } \Sigma_{ii} > 0 \\ 0, & \text{else} \end{cases} \quad (11)$$

Pseudo inverse & linear system

Solution via pseudo inverse

For a linear system $\mathbf{Ax} = \mathbf{b}$, $\mathbf{x} = \mathbf{A}^\dagger \mathbf{b}$ is always a solution regardless of \mathbf{A} is full rank or not.

When \mathbf{A} is invertible, we have $\mathbf{A}^\dagger = \mathbf{A}^{-1}$.

Theorem

If $\text{rank}[\mathbf{A}, \mathbf{b}] \neq \text{rank}(\mathbf{A})$, that is, $\mathbf{b} \notin \mathcal{R}(\mathbf{A})$, then $\mathbf{x}^ = \mathbf{A}^\dagger \mathbf{b}$ is the solution to the least square problem*

$$\min_{\mathbf{x}} \|\mathbf{Ax} - \mathbf{b}\|_2 \quad (12)$$

moreover, \mathbf{x}^ is of minimum 2-norm among all solutions, i.e., if \mathbf{u} is also a solution to problem (12), then*

$$\|\mathbf{u}\|_2 \geq \|\mathbf{x}^*\|_2 \quad (13)$$

See HW3, Problem 6 for proof.

Frobenius norm

Frobenius norm is an extension of vector 2-norm. If we vectorize a matrix into a vector (along column or row), the result will become clear. Thus Frobenius norm is often used to measure the “distance” between two matrices.

$$\|\mathbf{A}\|_F = \sum_{i=1}^n \sum_{j=1}^m A_{ij}^2 = \sqrt{\text{Trace}(\mathbf{A}^T \mathbf{A})} \quad (14)$$

Property

$$\|\mathbf{A}\|_F = \sqrt{\sum_i \sigma_i^2(\mathbf{A})} \quad (15)$$

The proof is clear if we use the SVD of \mathbf{A} , and note that \mathbf{U} and \mathbf{V} are both orthonormal.

Matrix 2 norm

The matrix 2-norm of $\mathbf{A} \in \mathbb{R}^{m \times n}$ is given by

$$\|\mathbf{A}\|_2 = \max_{\mathbf{x} \neq 0} \frac{\|\mathbf{Ax}\|_2}{\|\mathbf{x}\|_2} = \max_{\|\mathbf{x}\|_2=1} \|\mathbf{Ax}\|_2 \quad (16)$$

Block 3

Suppose SVD of \mathbf{A} is given by $\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$, then $\|\mathbf{A}\|_2 = \sigma_1(\mathbf{A})$

Proof.

$$\max_{\|\mathbf{x}\|_2=1} \|\mathbf{Ax}\|_2^2 = \mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{x}^T \mathbf{Q}^T \mathbf{D} \mathbf{Q} \mathbf{x} \quad (17)$$

$$= \max_{\|\mathbf{y}\|_2=1} \|\mathbf{Dy}\|_2^2 \leq \lambda_{\max}(\mathbf{A}^T \mathbf{A}) \quad (18)$$

where $\mathbf{Q}^T \mathbf{D} \mathbf{Q}$ is the eigen-decomposition of $\mathbf{A}^T \mathbf{A}$ with \mathbf{Q} orthonormal. By SVD of \mathbf{A} , we have $\lambda_{\max}(\mathbf{A}^T \mathbf{A}) = \sigma_1^2(\mathbf{A})$. On the other hand, take $\mathbf{x} = \mathbf{v}_1$, we have $\max_{\|\mathbf{x}\|_2=1} \|\mathbf{Ax}\|_2 \geq \sigma_1(\mathbf{A})$. □

Condition number

Recall the definition of condition number:

$$\kappa = \|\mathbf{A}\| \|\mathbf{A}^\dagger\| \quad (19)$$

If we use matrix 2-norm, then the condition number is given by

$$\kappa_2 = \frac{\sigma_1(\mathbf{A})}{\sigma_r(\mathbf{A})} \quad (20)$$

where $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\text{rank}(\mathbf{A}) = r$.

Projection onto the four subspaces

Suppose that SVD of \mathbf{A} is given by

$$\mathbf{A} = \begin{bmatrix} \mathbf{U}_1 & \mathbf{U}_2 \end{bmatrix} \begin{bmatrix} \boldsymbol{\Sigma} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{V}_1^T \\ \mathbf{V}_2^T \end{bmatrix} \quad (21)$$

then we have

- $\mathbf{V}_1 \mathbf{V}_1^T$ is the orthogonal projection onto $\mathcal{R}(\mathbf{A}^T)$.
- $\mathbf{V}_2 \mathbf{V}_2^T$ is the orthogonal projection onto $\mathcal{N}(\mathbf{A})$.
- $\mathbf{U}_1 \mathbf{U}_1^T$ is the orthogonal projection onto $\mathcal{R}(\mathbf{A})$.
- $\mathbf{U}_2 \mathbf{U}_2^T$ is the orthogonal projection onto $\mathcal{N}(\mathbf{A}^T)$.

Remark

- 1 Do not misunderstand the relation of four subspaces:
 $\mathcal{R}(\mathbf{A}^T) \perp \mathcal{N}(\mathbf{A})$ and $\mathcal{R}(\mathbf{A}) \perp \mathcal{N}(\mathbf{A}^T)$.
- 2 We can also write the above result in the form of pseudo inverse, see [Pseudo Projection](#).

Numerical rank

Though the rank function has a closed form, it is not easy to calculate in the presence of errors in the matrix elements.

Rounding errors and fuzzy data make the rank determination a nontrivial exercise. That is why numerical rank comes out. The numerical rank is useful, especially when the matrix is ill-conditioned. See [Example](#).

The numerical rank of a matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$ is defined as:

$$\text{numerical rank}(\mathbf{A}) = \#\{\sigma_i(A) \mid \sigma_i(A) \geq \epsilon\} \quad (22)$$

where ϵ often depends on the machine precision.

Low Rank Matrix Completion

Low Rank Matrix Completion is a popular topic in recent years, aiming to find a lower rank estimation of a given matrix with partially observed entries. Mathematically, given $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\Omega \subseteq [m] \times [n]$, we aim to solve the problem

$$\begin{aligned} \min_{\mathbf{X} \in \mathbb{R}^{m \times n}} \quad & \frac{1}{2} \|\mathcal{P}_{\Omega}(\mathbf{X}) - \mathbf{A}\|_F^2 \\ \text{s.t.} \quad & \text{rank}(\mathbf{X}) \leq d \end{aligned}$$

where \mathcal{P}_{Ω} is the projection such that $\mathcal{P}_{\Omega}(\mathbf{X}_{ij}) = \mathbf{X}_{ij}$ if $(i, j) \in \Omega$ and $\mathcal{P}_{\Omega}(\mathbf{X}_{ij}) = 0$ if $(i, j) \notin \Omega$.

A popular method to solve the LRMC problem is to use the Nuclear norm to replace the lower rank constraints, the nuclear norm is defined as

$\|\mathbf{X}\|_* = \sum_{i=1}^{\min\{m,n\}} \sigma_i(\mathbf{X})$. You may refer to the famous paper [2] written by Candes for more details.

Spectral functions*

The spectral function reveals the relation of a special class of functions $f : \mathbb{R}^{m \times n} \rightarrow (-\infty, \infty]$ to the class of functions $g : \mathbb{R}^{\min\{m,n\}} \rightarrow (-\infty, \infty]$ via singular values. For example,

$$\|\mathbf{A}\|_F = \|\sigma(\mathbf{A})\|_2, \|\mathbf{A}\|_* = \|\sigma(\mathbf{A})\|_1, \|\mathbf{A}\|_2 = \|\sigma(\mathbf{A})\|_\infty \quad (23)$$

Thus we can reduce the complexity of the problem.

If you are working with optimization problem on matrices, you may refer to [1] Chapter 7 for more details.

Principal Component Analysis

SVD is also a popular tool for PCA, which is powerful for Dimension Reduction in machine learning and De-noising in signal processing, many packages in machine learning have implemented the PCA model via SVD. You may refer to [3] for more details or select the course SI232 Subspace Learning to investigate the theory behind the SVD.

Reference



Amir Beck.

First Order Methods in Optimization.
2010.



Emmanuel J Candes and Benjamin Recht.

Exact low-rank matrix completion via convex optimization.
In 2008 46th Annual Allerton Conference on Communication, Control, and Computing, pages 806–812. IEEE, 2008.



René Vidal and Yi Ma.

Generalized Principal Component Analysis.
2019.

The End