



Multi-probe scanning tunneling spectroscopy for in-plane electronic transport: theory and experiment

Pedro Brandimarte¹, Marek Kolmer^{2,3}, Hiroyo Kawai⁴, Thomas Frederiksen^{1,5},
Aran Garcia-Lekue^{1,5}, Nicolas Lorente⁶, Jakub Lis², Rafal Zuzak²,
Szymon Godlewski², Christian Joachim⁷, Marek Szymonski², Daniel Sánchez-Portal^{1,6}

1 Donostia International Physics Center, Spain

2 NANOSAM - Jagiellonian University, Poland

3 CNMS - Oak Ridge National Laboratory, USA

4 IMRE - National University of Singapore, Singapore

5 IKERBASQUE, Basque Foundation for Science, Spain

6 Centro de Física de Materiales CSIC-UPV/EHU, Spain

7 CEMES-CNRS, France

October 3, 2019



Multi-probe scanning tunneling spectroscopy for in-plane electronic transport: **theory** and experiment

Pedro Brandimarte¹, Marek Kolmer^{2,3}, Hiroyo Kawai⁴, Thomas Frederiksen^{1,5},
Aran Garcia-Lekue^{1,5}, Nicolas Lorente⁶, Jakub Lis², Rafal Zuzak²,
Szymon Godlewski², Christian Joachim⁷, Marek Szymonski², Daniel Sánchez-Portal^{1,6}

1 Donostia International Physics Center, Spain

2 NANOSAM - Jagiellonian University, Poland

3 CNMS - Oak Ridge National Laboratory, USA

4 IMRE - National University of Singapore, Singapore

5 IKERBASQUE, Basque Foundation for Science, Spain

6 Centro de Física de Materiales CSIC-UPV/EHU, Spain

7 CEMES-CNRS, France

October 3, 2019

Multi-probe scanning tunneling spectroscopy for in-plane electronic transport: theory and experiment

Pedro Brandimarte¹, Marek Kolmer^{2,3}, Hiroyo Kawai⁴, Thomas Frederiksen^{1,5},
Aran Garcia-Lekue^{1,5}, Nicolas Lorente⁶, Jakub Lis², Rafal Zuzak²,
Szymon Godlewski², Christian Joachim⁷, Marek Szymonski², Daniel Sánchez-Portal^{1,6}

1 Donostia International Physics Center, Spain

2 NANOSAM - Jagiellonian University, Poland

3 CNMS - Oak Ridge National Laboratory, USA

4 IMRE - National University of Singapore, Singapore


5 IKERBASQUE, Basque Foundation for Science, Spain

6 Centro de Física de Materiales CSIC-UPV/EHU, Spain


7 CEMES-CNRS, France

October 3, 2019


Two-probe STM at the atomic level




UNIVERSITAS JAGELLONICA
CRACOVIENSIS




Two-probe STM at the atomic level





UNIVERSITAS JAGELLONICA
CRACOVENSIS




Two-probe STM at the atomic level




UNIVERSITAS JAGELLONICA
CRACOVIENSIS




Two-probe STM at the atomic level



Two-probe STM at the atomic level



Two-probe STM at the atomic level



Methods

Density-Functional Theory (DFT)

SIESTA

E. Artacho *et al.* *Phys. Stat. Sol. (b)* **215**, 809 (1999).
J. M. Soler *et al.* *J. Phys. Condens. Matter.* **14**, 2745 (2002).

Methods

Density-Functional Theory (DFT)

+

Non-Equilibrium Green's Function (NEGF)

TranSIESTA

E. Artacho *et al.* *Phys. Stat. Sol. (b)* **215**, 809 (1999).

J. M. Soler *et al.* *J. Phys. Condens. Matter.* **14**, 2745 (2002).

M. Brandbyge *et al.* *Phys. Rev. B* **65**, 165401 (2002).

N. Papior *et al.* *Comp. Phys. Comm.* **212**, 8 (2017).

Methods

Density-Functional Theory (DFT)

+

Non-Equilibrium Green's Function (NEGF)

TranSIESTA

Multi-terminal!!!



E. Artacho *et al.* *Phys. Stat. Sol. (b)* **215**, 809 (1999).

J. M. Soler *et al.* *J. Phys. Condens. Matter.* **14**, 2745 (2002).





M. Brandbyge *et al.* *Phys. Rev. B* **65**, 165401 (2002).

N. Papior *et al.* *Comp. Phys. Comm.* **212**, 8 (2017).





Ge(001)x(4x2) surface






Ge(001)x(4x2) surface





Ge(001)x(4x2) surface



Ge(001)x(4x2) surface




Ge(001) surface: 2-terminal setup




of atoms/orbitals: 2240/16000


Ge(001) surface: 2-terminal setup



Ge(001) surface: 2-terminal setup





Standard (one probe) Scanning Tunneling Spectroscopy (STS)




of atoms/orbitals: 2462/18221

Standard (one probe) Scanning Tunneling Spectroscopy (STS)





Proof of ballistic transport




Proof of ballistic transport


dI/dV (arb. units) low high




Proof of ballistic transport




Proof of ballistic transport




Proof of ballistic transport



Proof of ballistic transport




Proof of ballistic transport




Coherence length
up to **50 nm!!!**

Ge(001) surface addressed by two tips: 4-terminal setup





of atoms/orbitals: 4924/36442



Ge(001) surface addressed by two tips: 4-terminal setup






Ge(001) surface addressed by two tips: 4-terminal setup





Ge(001) surface addressed by two tips: 4-terminal setup




Ge(001) surface addressed by two tips: 4-terminal setup



Ge(001) surface addressed by two tips: 4-terminal setup




Ge(001) surface addressed by two tips: 4-terminal setup



Conclusions

- Identification and characterization of the **quasi 1D transport** surface channels **along a single dimer row** on bare Ge(001)-c(4×2) surface;
- SP-STM/STS and calculated eigenchannels on a step-edge confirms a **coherent transport length up to 50nm**;
- **TP-STs planar transconductance** resonances measured with a tip-to-tip distance down to **30nm** and **confirmed by multi-terminal DFT-NEGF simulations**.



M. Kolmer, P. Brandimarte* et al. *Nature Communications* **10**, 1573 (2019)

