

Aspects of graphene nanoribbon device simulations

Pedro Brandimarte^{1,2}, Nick R. Papior^{3,4}, Mads Englund^{1,2}, Aran Garcia-Lekue^{2,5}, Thomas Frederiksen^{2,5}, Eduard Carbonell-Sanromà⁶, Martina Corso^{1,5,6}, Richard Balog⁶, Shigeki Kawai^{7,8}, Shohei Saito^{8,9}, Shinichiro Osumi⁹, Shigehiro Yamaguchi^{9,10,11}, Jose I. Pascual^{1,5,6}, and Daniel Sánchez-Portal^{1,2}

¹Centro de Física de Materiales (CFM), Spain

²Donostia International Physics Center (DIPC), Spain

³Catalan Institute of Nanoscience Nanotechnology - Campus UAB, Spain

⁴Center for Nanostructured Graphene (CNG) - DTU Nanotech, Denmark

⁵IKERBASQUE, Basque Foundation for Science, Spain

⁶CIC nanoGUNE, Spain

⁷Department of Physics - University of Basel, Switzerland

⁸PRESTO - Japan Science and Technology Agency, Japan

⁹Department of Chemistry from Graduate School of Science - Nagoya University, Japan

¹⁰Institute of Transformative Bio-molecules - Nagoya University, Japan

¹¹CREST - Japan Science and Technology Agency, Japan

Graphene nanoribbons (GNRs) have been considered as strong candidates for electronics, since they incorporate some of the remarkable properties from graphene while presenting a band gap [1]. These structures can be fabricated nowadays with a high control on the edge structure with the so-called bottom-up approaches [2]. Together with these experimental breakthrough a major effort has been devoted to the theoretical and computational methods developments. In this talk I will show some challenging features found when simulating GNRs based devices, such as: 1. The effects on the transport properties of 4-terminal crossed GNRs devices when varying the structural parameters determining the crossing; 2. The effect of substitutional chemical defects on the transmission channels and the importance of a proper description of the electrostatics in quasi 1-dimension systems.




Figure 1: Direct (left) and inter-ribbon (right) transmissions at $V=0$ in a crossbar system with 60° rotation angle.




Figure 2: Project density of states from a substitutional boron-doped armchair graphene nanoribbon.

[1] L. Yang *et al.*, Phys. Rev. Lett. **99**, 186801 (2007).

[2] J. Cai *et al.*, Nature **466**, 470-473 (2010).