
Realistic multi-terminal fist-principles transport simulations of two-probe STM measurements on Ge(001) surface: demonstration of quasi-ballistic transport through dangling-bond dimer wires

<u>Pedro Brandimarte</u>¹, Marek Kolmer^{2,3}, Hiroyo Kawai⁴, Thomas Frederiksen^{1,5}, brandimarte@pm.me

Aran Garcia-Lekue^{1,5}, Nicolas Lorente⁶, Jakub Lis², Rafal Zuzak², Szymon Godlewski², Christian Joachim⁷, Marek Szymonski², Daniel Sánchez-Portal^{1,6}

1 Donostia International Physics Center, Spain 2 NANOSAM – Jagiellonian University, Poland 3 CNMS – Oak Ridge National Laboratory, USA 4 IMRE – National University of Singapore, Singapore 5 IKERBASQUE, Basque Foundation for Science, Spain 6 Centro de Física de Materiales CSIC-UPV/EHU, Spain 7 CEMES-CNRS, France



M. Kolmer, P. Brandimarte* et al. Nature Communications 10, 1573 (2019)


















Two-probe STM at the atomic level








M. Kolmer et al. J. Phys: Cond. Mat. 29, 444004 (2017).

Ge(001)x(4x2) surface



tip2

 $-\mu_0$

Methodology

Density-Functional Theory (DFT)


+

Non-Equilibrium Green's Function (NEGF)


TranSIESTA

E. Artacho et al. Phys. Stat. Sol. (b) 215, 809 (1999).
J. M. Soler et al. J. Phys. Condens. Matter. 14, 2745 (2002).
M. Brandbyge et al. Phys. Rev. B 65, 165401 (2002).
N. Papior et al. Comp. Phys. Comm. 212, 8 (2017).


Ge(001) step edge: coherence length








Ge(001) surface: 2-terminal setup







Simulation characteristics:

- # of atoms/orbitals: 2240/16000

cell size: **32.03x80.07x34** Å³



Ge(001) surface addressed by a single tip: 3-terminal setup



Ge(001) surface addressed by two tips: 4-terminal setup

