Formalization of: Existence and Uniqueness of Homotopy Lifts & Universal Covering Spaces

Joint work:

(Tony, Daniel), (Amogh, Pranali, Brandon), (Gary, Shruthi)

Topology Study Group

June 16, 2023

Thm 1 (Existence and Uniqueness of Homotopy Lifts) Let X, Y, \tilde{X}

be topological spaces. Let maps be defined as follows

$$F: Y \times [0,1] \xrightarrow{cont.} X$$


$$p: \tilde{X} \xrightarrow{cont.} X$$

hp: p is a covering map

$$\tilde{F}_0: Y \times \{0\} \xrightarrow{cont.} X$$

$$h: F|_{Y\times\{0\}} = p\circ \tilde{F}_0$$

$$\vdash \exists ! \tilde{F} : Y \times [0,1] \xrightarrow{cont.} \tilde{X}, (p \circ \tilde{F} = F) \wedge (\tilde{F}|_{Y \times \{0\}}) = \tilde{F}_0)$$



Thm 2 (Uniqueness of Homotopy Lifts, General Case) Let X, Z, \tilde{X}

be topological spaces. Let maps be defined as follows

$$F: Z \xrightarrow{cont.} X$$
$$p: \tilde{X} \xrightarrow{cont.} X$$


hp: p is a covering map

 $H_1, H_2: Z \xrightarrow{cont.} \tilde{X}$

 $hH: p \circ H_1 = p \circ H_2 = F$

 $hconn: \forall z \in Z, \exists x \in ConnComp(z), H_1(x) = H_2(x)$

$$\vdash H_1 = H_2$$



Thm 3 (Existence of the Universal Covering)

Let X be a topological space satisfying the conditions

- Path-connected
- Locally path-connected
- Semilocally simply connected

Then there exists a covering space

$$\pi_{\infty}:\widetilde{X_{\infty}}\to X$$

which is simply connected (i.e. $\pi_1(\widetilde{X}_{\infty})=0$) and satisfies a universal property



Let X be a topological space satisfying the conditions

- Path-connected
- Locally path-connected
- Semilocally simply connected

Then there exists a basis of X whose elements are both path-connected and semilocally simply connected.

Lifting this basis constructs a basis for the topology of \widetilde{X}_{∞}