



Introduction to The LIGO Project

Brandon B. Miller

Allstate


9/1/2016

Classical Electrodynamics

ELECTRODYNAMICS

$$\square A^\alpha = \frac{4\pi}{c} J^\alpha$$

Radiation from an Accelerating Charge



Classical Electrodynamics vs General Relativity

ELECTRODYNAMICS


$$\square A^\alpha = \frac{4\pi}{c} J^\alpha$$

**Radiation from an
Accelerating Charge**


GENERAL RELATIVITY

$$\square h_{\mu\nu} = 0$$


**Radiation from an
Accelerating Mass?**



Chirps and Black Hole Binaries




A “*Chirp*” from a black hole binary in one dimension



How does “*Gravitational Radiation*” Manifest Itself?

Waves go straight through matter.

Space between objects warped and stretched



We need to measure the distances between things and look for changes and oscillations

VERY tiny signal, $\sim 10^{-23}$



Hulse and Taylor

The LIGO Project



[

The LSC/VSC Detector Network

]



September 14th, 2015

[


September 14th, 2015 – Success!

]

[The Problem – Reverse Engineer the Parameters from the Signal]

Not all Chrips are created equal!

15 independent numbers affect the shape




[What if there's noise? (there is)]

The real data (usually) looks something like this:

We need a way to make a quantitative statement about what parameters are the “best fit” while propagating through the uncertainty due to *noise*

A Bayesian Approach



Our main tool is thus the Bayesian Posterior:




$$P(X|Y) = \frac{P(Y|X)P(X)}{P(Y)}$$

Bayes Theorem: Our Application

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \longrightarrow \text{Think of a *Gaussian*!}$$




Bayes Theorem: Our Application




Problem: Posterior Narrow, Multimodal, and 15 Dimensional!

You can miss the peak, badly! Direct computation hard


Function $y = \sin(x^5)$ For Different Sample Densities




[One Solution: MCMC Sampling]




[One Solution: MCMC Sampling]




[One Solution: MCMC Sampling]



[One Solution: MCMC Sampling]





[One Solution: MCMC Sampling]



One Solution: MCMC Sampling


**Can take a few days to converge
Depending on wave template, hardware**



My Solution: GPU-based Direct Computation


Marginalizes out all the extrinsic parameters with a Large, brute force Monte-Carlo integration

Massively Parallel, scalable to multiple GPUs
Fully custom CUDA/PYCUDA



Results vs Fastest CPU Algorithm

Tesla K40M Sample Throughput



A Brief Word About Why We Care

Low Latency Electromagnetic Follow-Up

Get the angles and orientations quickly and you can tell EM observatories where to look

