

# MATH 114 Assignment 3

Brandon Tsang

January 30, 2020

1. Find a set of vectors, each with a norm of 5, such that the sum of the vectors is  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

Two vectors that satisfy these conditions are  $\vec{a} = \begin{bmatrix} \sqrt{\frac{99}{4}} \\ \frac{1}{2} \end{bmatrix}$  and  $\vec{b} = \begin{bmatrix} -\sqrt{\frac{99}{4}} \\ \frac{1}{2} \end{bmatrix}$ .

Both vectors have norm 5:

$$\begin{aligned}\|\vec{a}\| &= \sqrt{\left(\sqrt{\frac{99}{4}}\right)^2 + \left(\frac{1}{2}\right)^2} \\ &= \sqrt{\frac{99}{4} + \frac{1}{4}} \\ &= \sqrt{\frac{100}{4}} \\ &= \sqrt{25} \\ &= 5\end{aligned}$$

$$\begin{aligned}\|\vec{b}\| &= \sqrt{\left(-\sqrt{\frac{99}{4}}\right)^2 + \left(\frac{1}{2}\right)^2} \\ &= \sqrt{\frac{99}{4} + \frac{1}{4}} \\ &= \sqrt{\frac{100}{4}} \\ &= \sqrt{25} \\ &= 5\end{aligned}$$

And they sum to  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ :

$$\begin{aligned}\begin{bmatrix} \sqrt{\frac{99}{4}} \\ \frac{1}{2} \end{bmatrix} + \begin{bmatrix} -\sqrt{\frac{99}{4}} \\ \frac{1}{2} \end{bmatrix} &= \begin{bmatrix} \sqrt{\frac{99}{4}} - \sqrt{\frac{99}{4}} \\ \frac{1}{2} + \frac{1}{2} \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ 1 \end{bmatrix}\end{aligned}$$

2. (a) Consider the vectors  $\vec{u} = \begin{bmatrix} 2 \\ 3 \\ 0 \\ 5 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \end{bmatrix}$ .

i. Find a vector  $\vec{x}$  such that  $\vec{x}$  is orthogonal to both  $\vec{u}$  and  $\vec{v}$ .