

CMSC 207- Lecture 27

CHAPTER 9: Counting and Probability (9.7)

Dr. Ahmed Tarek

Think of Theorem 9.5.1 as a general template.

Theorem 9.5.1

The number of subsets of size r (or r -combinations) that can be chosen from a set of n elements, $\binom{n}{r}$, is given by the formula

$$\binom{n}{r} = \frac{P(n, r)}{r!} \quad \text{first version}$$

or, equivalently,

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} \quad \text{second version}$$

where n and r are nonnegative integers with $r \leq n$.

Regardless of what nonnegative numbers are placed in the boxes, if the number in the lower box is no greater than the number in the top box, then

$$\binom{\bullet}{\bullet} = \frac{\bullet!}{\bullet!(\bullet - \bullet)!}.$$

Example: Use Theorem 9.5.1 to show that for all integers

$n \geq 0$:

$$\binom{n}{n} = 1 \tag{9.7.1}$$

$$\binom{n}{n-1} = n, \quad \text{if } n \geq 1 \tag{9.7.2}$$

$$\binom{n}{n-2} = \frac{n(n-1)}{2}, \quad \text{if } n \geq 2. \tag{9.7.3}$$

Example 1 – *Solution*

$$\binom{n}{n} = \frac{n!}{n!(n-n)!}$$

$$= \frac{1}{0!}$$

$$= 1 \quad \text{since } 0! = 1 \text{ by definition}$$

$$\binom{n}{n-1} = \frac{n!}{(n-1)!(n-(n-1))!}$$

$$= \frac{n \cdot \cancel{(n-1)!}}{\cancel{(n-1)!}(n-n+1)!}$$

$$= \frac{n}{1}$$

$$= n$$

Example 1 – *Solution*

$$\binom{n}{n-2} = \frac{n!}{(n-2)!(n-(n-2))!}$$

$$= \frac{n \cdot (n-1) \cdot \cancel{(n-2)!}}{\cancel{(n-2)!} 2!}$$

$$= \frac{n(n-1)}{2}$$


Example 2

Deduce the formula $\binom{n}{r} = \binom{n}{n-r}$ for all nonnegative integers n and r with $r \leq n$, by interpreting it as saying that a set A with n elements has exactly as many subsets of size r as it has subsets of size $n - r$.

Example 2 – *Solution*

Observe that any subset of size r can be specified either by saying which r elements lie in the subset or by saying which $n - r$ elements lie outside the subset.

A , A Set with n Elements



Any subset B with r elements completely determines a subset, $A - B$, with $n - r$ elements.

Example 2 – *Solution*

Suppose A has k subsets of size r : B_1, B_2, \dots, B_k .
Then each B_i can be paired up with exactly one
set of size $n - r$, namely its complement $A - B_i$ as
shown below.

Subsets of Size r Subsets of Size $n - r$

$$B_1 \longleftrightarrow A - B_1$$

$$B_2 \longleftrightarrow A - B_2$$

\vdots

\vdots

$$B_k \longleftrightarrow A - B_k$$

Example 2 – *Solution*

All subsets of size r are listed in the left-hand column, and all subsets of size $n - r$ are listed in the right-hand column.

The number of subsets of size r equals the number of subsets of size $n - r$, and so

$$\binom{n}{r} = \binom{n}{n-r}.$$

Pascal's Formula

It relates the value of $\binom{n+1}{r}$ to the values of $\binom{n}{r-1}$ and $\binom{n}{r}$. Specifically, it says that

$$\binom{n+1}{r} = \binom{n}{r-1} + \binom{n}{r}$$

whenever n and r are positive integers with $r \leq n$.

This formula makes it easy to compute higher combinations in terms of lower ones: If all the values of $\binom{n}{r}$ are known, then the values of $\binom{n+1}{r}$ can be computed for all r such that $0 < r \leq n$.

Use Pascal's triangle to compute the values of

$$\binom{6}{2} \quad \text{and} \quad \binom{6}{3}.$$

Solution:

$$\binom{6}{2} = \binom{5}{1} + \binom{5}{2} = 5 + 10 = 15$$

$$\binom{6}{3} = \binom{5}{2} + \binom{5}{3} = 10 + 10 = 20.$$

Pascal's Formula

Pascal's formula can be derived by two entirely different arguments. One is algebraic; it uses the formula for the number of r -combinations obtained in Theorem 9.5.1. The other is combinatorial; it uses the definition of the number of r -combinations as the number of subsets of size r taken from a set with a certain number of elements.

Theorem 9.7.1 Pascal's Formula

Let n and r be positive integers and suppose $r \leq n$. Then

$$\binom{n+1}{r} = \binom{n}{r-1} + \binom{n}{r}.$$

Example 4 – *Deriving New Formulas from Pascal's Formula*

Use Pascal's formula to derive a formula for $\binom{n+2}{r}$ in terms of values of $\binom{n}{r}$, $\binom{n}{r-1}$ and $\binom{n}{r-2}$.

Assume n and r are nonnegative integers and $2 \leq r \leq n$.

Solution:

By Pascal's formula,

$$\binom{n+2}{r} = \binom{n+1}{r-1} + \binom{n+1}{r}.$$

Example 4 – *Solution*

Now apply Pascal's formula to $\binom{n+1}{r-1}$ and $\binom{n+1}{r}$ and substitute into the above to obtain

$$\binom{n+2}{r} = \left[\binom{n}{r-2} + \binom{n}{r-1} \right] + \left[\binom{n}{r-1} + \binom{n}{r} \right].$$

Combining the two middle terms gives

$$\binom{n+2}{r} = \binom{n}{r-2} + 2 \binom{n}{r-1} + \binom{n}{r}$$

for all nonnegative integers n and r such that

$$2 \leq r \leq n.$$

The Binomial Theorem

In algebra, a sum of two terms, such as $a + b$, is called a **binomial**.

The ***binomial theorem*** gives an expression for the powers of a **binomial** $(a + b)^n$, for each positive integer n , and all real numbers a and b .

Theorem 9.7.2 Binomial Theorem

Given any real numbers a and b and any nonnegative integer n ,

$$\begin{aligned}(a + b)^n &= \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \\ &= a^n + \binom{n}{1} a^{n-1} b^1 + \binom{n}{2} a^{n-2} b^2 + \cdots + \binom{n}{n-1} a^1 b^{n-1} + b^n.\end{aligned}$$

The Binomial Theorem

Note that the second expression equals the first
Because $\binom{n}{0} = 1$ and $\binom{n}{n} = 1$, $b^0 = 1$ and $a^{n-n} = 1$.
for all nonnegative integers n , provided
that

- **Definition**

For any real number a and any nonnegative integer n , the **nonnegative integer powers of a** are defined as follows:

$$a^n = \begin{cases} 1 & \text{if } n = 0 \\ a \cdot a^{n-1} & \text{if } n > 0 \end{cases}$$

The Binomial Theorem

If n and r are nonnegative integers and $r \leq n$, then $\binom{n}{r}$ is called a **binomial coefficient** because it is one of the coefficients in the expansion of the binomial expression $(a + b)^n$.

Example 5 – *Substituting into the Binomial Theorem*

Expand the following expressions using the binomial theorem:

a. $(a + b)^5$ **b.** $(x - 4y)^4$

Solution:

$$\begin{aligned}\mathbf{a.} \quad (a + b)^5 &= \sum_{k=0}^5 \binom{5}{k} a^{5-k} b^k \\ &= a^5 + \binom{5}{1} a^{5-1} b^1 + \binom{5}{2} a^{5-2} b^2 + \binom{5}{3} a^{5-3} b^3 + \binom{5}{4} a^{5-4} b^4 + b^5 \\ &= a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5\end{aligned}$$

Example 5 – *Solution*

b. Observe that $(x - 4y)^4 = (x + (-4y))^4$.

So let $a = x$ and $b = (-4y)$, and substitute into the binomial theorem.

$$\begin{aligned}(x - 4y)^4 &= \sum_{k=0}^4 \binom{4}{k} x^{4-k} (-4y)^k \\&= x^4 + \binom{4}{1} x^{4-1} (-4y)^1 + \binom{4}{2} x^{4-2} (-4y)^2 + \binom{4}{3} x^{4-3} (-4y)^3 + (-4y)^4 \\&= x^4 + 4x^3(-4y) + 6x^2(16y^2) + 4x^1(-64y^3) + (256y^4) \\&= x^4 - 16x^3y + 96x^2y^2 - 256xy^3 + 256y^4\end{aligned}$$

In-class Assignment #1

- Use the **Binomial Theorem** to expand:

$$(x + 1/x)^5$$

Example 7 – *Using a Combinatorial Argument to Derive the Identity*

According to Theorem 6.3.1, a set with n elements has 2^n subsets.

Theorem 6.3.1

For all integers $n \geq 0$, if a set X has n elements, then $\mathcal{P}(X)$ has 2^n elements.

Apply this fact to give a combinatorial argument to justify the identity

$$\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \binom{n}{3} + \cdots + \binom{n}{n} = 2^n.$$

Example 7 – *Solution*

Suppose S is a set with n elements. Then every subset of S has some number of elements k , where k is between 0 and n . It follows that the total number of subsets of S , $N(\mathcal{P}(S))$, can be expressed as the following sum:

$$\left[\begin{array}{c} \text{number of} \\ \text{subsets} \\ \text{of } S \end{array} \right] = \left[\begin{array}{c} \text{number of} \\ \text{subsets of} \\ \text{size 0} \end{array} \right] + \left[\begin{array}{c} \text{number of} \\ \text{subsets of} \\ \text{size 1} \end{array} \right] + \cdots + \left[\begin{array}{c} \text{number of} \\ \text{subsets of} \\ \text{size } n \end{array} \right].$$

Now the number of subsets of size k of a set with n elements is $\binom{n}{k}$.

Example 7 – *Solution*

Hence,

$$S = \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{n}$$

the number of subsets of S .

But by Theorem 6.3.1, S has 2^n subsets. Hence

$$\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \binom{n}{3} + \cdots + \binom{n}{n} = 2^n.$$

Example 8 – *Using the Binomial Theorem to Simplify a Sum*

Express the following sum in **closed form** (without using a summation symbol and without using an ellipsis \cdots):

$$\sum_{k=0}^n \binom{n}{k} 9^k$$

Solution:

When the number 1 is raised to any power, the result is still 1. Thus

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k} 9^k &= \sum_{k=0}^n \binom{n}{k} 1^{n-k} 9^k \\ &= (1 + 9)^n \quad \text{by the binomial theorem with } a = 1 \text{ and } b = 9 \\ &= 10^n. \end{aligned}$$

In-class Assignment #2

- For all integers $n \geq 0$, show that:

$$3^n = C(n, 0) + 2C(n, 1) + 2^2C(n, 2) + \dots + 2^nC(n, n)$$

- Find the coefficient of x^7 in the binomial expansion of: $(2x + 3)^{10}$