

Secondary Particle Showers from Hadron Absorber Interactions

Branton DeMoss
University of Colorado Boulder
High Energy Physics Group

August 12, 2015

1 Introduction

Proton pulses from Fermilab’s Main Injector interact with a solid graphite target to produce kaons and pions. Two focusing horns direct these products down the main decay pipe aimed at the near and far detectors. Any particles that have not decayed by the end of the pipe are stopped via interaction with the hadron absorber. This note discusses the secondary effects of interactions with the hadron absorber, such as potential secondary showers that could interfere with measurement statistics. In short, we’d like to assess whether the hadron absorber is an unaccounted-for particle gun.

2 New Geometry

Recent ¹ DUNE design documents have called for “scallop” to be placed in the absorber hall region. These are foot-thick aluminum blocks with spherical cuts taken out. Additionally, there is a proposal to add a foot thick solid block of aluminum to the front of the absorber region, the so-called “spoiler”.




Figure 2.1: The current configuration of the absorber

Our first task was to implement the new geometry 2.1 into G4LBNE, a GEANT4 implementation of the DUNE experiment managed by Fermilab. The

new geometry was implemented according mostly to ², with some supplementary information from Jan Boissevain’s 3D model.

Most of the absorber hall region was remade to ease the implementation of the new geometry. In addition to the modifications to geometry construction source files, we created some new inputs to aid in muon and pion tracking. If our changes are added to the master branch of G4LBNE, the commands

```
/LBNE/output/doPionVertexTracking bool
/LBNE/output/doMuonVertexTracking bool
```

will add an output Ntuple to your results file with plenty of useful information.

3 Contamination Studies


Behind the absorber hall is the “Muon alcove”, inside of which are various instruments to observe stopped and through-going muons, in order to correlate muon and neutrino flux data. Therefore, if muon production rates inside of absorber volumes are not insignificant, these absorber-born muons (which are generally poor proxies for neutrinos compared to muons created in the decay pipe) will make muon/neutrino correlation more difficult. Of interest, then, is the percent of all forward going muons created in absorber volumes, and their momenta distribution.

3.1 Muon Distributions


As expected, the muon profile at the tracking plane downstream of the absorber is fairly tight. If, instead of plotting the complete 3.1a distribution above, we cut muons by birth position and plot only those born in absorber volumes, we find the distribution of

¹LBNE-doc-10092


²LBNE-doc-10095




(a) Complete muon profile at tracking plane



(b) Muon profile at tracking plane as a function of r^2



(c) Profile of absorber born muons



(d) Ratio of 3.1b to 3.1c

Figure 3.1: Muon profile distributions

muons born in the hadron absorber has a seemingly similar spread.

Taking the ratio of the total muon profile 3.1b to the absorber muon profile 3.1c we get a ratio plot that increases in variance with radius. This increase of variance with radius suggests that the absorber born muon distribution will be a difficult background in determining overall muon distribution statistics

3.2 Muon counting methods (1)

Our original method of counting muons used the detectors downstream of the absorber in the standard G4LBNE implementation. When a muon hits the detectors we get its birth position using the built-in

```
const G4ThreeVector& GetVertexPosition() const;
//See G4Track.hh for reference
```

then, we simply divide the number of muons born in absorber volumes by the total number of muons that hit the tracking plane. Using this method, we find that the muon contamination from absorber volumes is, depending on specific run settings, around 3%.

3.3 Muon counting methods (2)

After preliminary results with the first method, there was confusion when Paul LeBrun shared his slides with us ³, showing results closer to 9% muon contamination, albeit with his own version of the new geometry (scallop and spoiler). Dr. LeBrun's method of counting (hereafter referred to as method 2) takes advantage of two tracking planes, respectively placed before and after the hadron absorber, to determine

³LBNE-doc-11203

which muons are born in the absorber. Simply, if a muon hits the plane downstream of the absorber but not the upstream plane, it must have been born in the absorber. We would expect to see results identical to our own with this method. As the results were significantly different, in-depth verification of the two methods had to be performed to determine the true value.

3.4 Counting Verification

To check the results of method 2, we coded an independent version of LeBrun’s double tracking plane analysis method (github). After a great deal of debugging and dealing with subtle volumetric errors, we confirm both methods give around 3% muon contamination, with up to a $\frac{1}{2}\%$ variance depending on specific geometry configurations.

3.5 Weighted counts

Also of interest to our studies is the degree of pion contamination from the absorber region. Here we would like to introduce a different view of what constitutes an “important” particle. Because muons are directly measured in the Muon Alcove, how their production rates change matters, neutrino production notwithstanding. Pions, however, are not directly measured, so we only consider those which are “important” with regards to how they contribute to neutrino flux at the detectors. If we weight a pion detection event with that pion’s eventual contribution to neutrino flux at the near detector ⁴, we get the following:




Figure 3.2: Pion births weighted with neutrino flux

⁴In reality we are actually counting neutrino events, weighting them with their likelihood to hit the near detector, and grabbing the parent particle birth position and type to make our “important” pion graph.

There is a clear spike in “important” pion births beginning at around 220 meters, which is the start of the hadron absorber. Integrating this spike and dividing by the total pion count integral, we find that the percent of “important” pion contamination from the hadron absorber is around 0.5%.

If we weight muon births by near detector neutrino flux, we find that there is not nearly as significant a spike at the absorber hall:




Figure 3.3: Muon births weighted with neutrino flux

Using the same pion contamination analysis as above, we find that the percent of “important” muon contamination from the absorber regions is around 0.3%.

Note the difference in total muon contamination (which the Muon Alcove will measure) and the “important” muon contamination.

3.6 Momentum Distributions

Both the pre-absorber 6.1 and absorber-born 6.2 muons follow a steep drop off in z-momentum at the tracking plane after passing through absorber volumes (found in Appendix).

Taking the ratio of the pre-absorber 6.1 to absorber born 6.2 muons yields:




Figure 3.4: Ratio of pre-absorber to absorber muon momentum distributions

The momentum distribution ratio 3.4 shows that for any given momentum, the percent of muons measured at that momentum level is less than 3%, often far less.

4 Neutrino Flux

Of the utmost importance for this analysis is determining how neutrino flux changes with the addition of the hadron absorber. Especially, we would like to know if absorber-born neutrinos affect the near and far detection rates differently, and if so, by how much.

Our first graph 3.5 is a plot of the flux of various neutrino flavors. Comparison to similar plots⁵ made for DUNE science documents shows that there are not severe differences with the new absorber geometry.

Looking into how specific neutrino detection rates change reveals that the ratio of near to far absorber-born counts does indeed change:

Regular Neutrino Mode	ν_μ	anti- ν_μ	ν_e	anti- ν_e
Horn current = 200kA				
Percent of Near Detector neutrino flux from absorber volumes	0.33%	3.32%	0.91%	2.61%
Percent of Far Detector neutrino flux from absorber volumes	0.13%	1.26%	0.41%	1.03%




Figure 3.5: Neutrino flux of various flavors at the near detector.

⁵[https://sharepoint.fnal.gov/project/lbne/LBNE Work/science doc pdfs/chapter_3_optim.pdf](https://sharepoint.fnal.gov/project/lbne/LBNE%20Work/science%20doc%20pdfs/chapter_3_optim.pdf) at figure 3.18

As you can see, the percent contamination of absorber-born neutrinos at the near detector is more than double that of the far detector.

Plotting the neutrino flux at the far detector, we see a similar flux distribution.

Anti Neutrino Mode	ν_μ	$\bar{\nu}_\mu$	ν_e	$\bar{\nu}_e$
Horn current = -200kA				
Percent of Near Detector neutrino flux from absorber volumes	3.47%	0.31%	4.13%	0.60%
Percent of Far Detector neutrino flux from absorber volumes	1.29%	0.12%	1.68%	0.27%




Figure 4.1: Neutrino flux of various flavors at the far detector.

The overall flux at the near 3.5 and far 4.1 detectors are useful, but of greater relevance to this study is the percent of flux of a given flavor from the absorber volumes, as a function of energy.

These plots 4.2 (found on next page) show how absorber-born neutrino count rates compare to the total neutrino count rate at the near and far detectors, as a function of energy.


4.1 Antineutrino mode

If we run our simulation with a negative current in the focusing horns, this is known as "antineutrino mode", as this will select the opposite sign particles for focusing, which will favor antineutrino production. The effects from the absorber are, as before:


5 Conclusions

Muon contamination from the hadron absorber is not a problem that can be swept under the rug. Muon detection groups should take these statistics into account when doing their analysis at the Alcove.


Furthermore, neutrino flux contributions from absorber volumes contributes differently but not insubstantially to the near and far detectors. The ratios 4.2 of absorber born neutrinos should be used by detector groups to aid in neutrino counting statistics, and will hopefully help to explain some anomalous-seeming spikes.




(a) Muon neutrinos



(b) Anti muon neutrinos



(c) Electron neutrinos



(d) Anti electron neutrinos

Figure 4.2: Ratio of absorber born to total neutrino flux at the detectors. Graphed by flavor. Far detector in red.

6 Appendix




Figure 6.1: Pre-absorber muon momentum distribution




Figure 6.2: Momentum distribution of absorber muons