

# Astronomy 400B Lecture 5: Stellar Orbits

Brant Robertson

February, 2015

## 1 Motion Under Gravity

Newton's Law of Gravity point mass  $M$  attracts another mass  $m$  separated by distance  $\mathbf{r}$ , causing a change in momentum  $m\mathbf{v}$  of:

$$\frac{d}{dt}(m\mathbf{v}) = -\frac{GmM}{r^3}\mathbf{r} \quad (1)$$

where  $G$  is Newton's gravitational constant. For an  $N$ -body system, we have

$$\frac{d}{dt}(m_i\mathbf{v}_i) = -\sum_{j \neq i} \frac{Gm_i m_j}{|\mathbf{x}_i - \mathbf{x}_j|^3}(\mathbf{x}_i - \mathbf{x}_j) \quad (2)$$

This equation can be re-written as

$$\frac{d}{dt}(m\mathbf{v}_i) = -m\nabla\Phi(\mathbf{x}_i) \quad (3)$$

where

$$\Phi(\mathbf{x}_i) = -\sum_{j \neq i} \frac{Gm_j}{|\mathbf{x}_i - \mathbf{x}_j|} \quad (4)$$

is the gravitational potential supplied by the point mass distribution at positions  $\mathbf{x}_i$ . Note we have chosen to define the potential such that  $\Phi(x) \rightarrow 0$  as  $x \rightarrow \infty$  but this is arbitrary. Note that

$$\nabla = \left[ \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right] \quad (5)$$

### 1.1 Continuous Matter Distributions

Now consider a continuous distribution of matter density  $\rho(\mathbf{x})$ . The potential generated by  $\rho(\mathbf{x})$  is given by

$$\Phi(\mathbf{x}) = -\int \frac{G\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} d^3\mathbf{x}' \quad (6)$$

Note that the integral is performed over  $\mathbf{x}'$ . The force  $\mathbf{F}$  per unit mass is

$$\mathbf{F}(\mathbf{x}) = -\nabla\Phi(\mathbf{x}) = -\int \frac{G\rho(\mathbf{x}')(\mathbf{x} - \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^3} d^3\mathbf{x}' \quad (7)$$

### 1.2 Poisson's Equation

Take Equation 6 and apply the Laplacian operator

$$\nabla^2 \equiv \nabla \cdot \nabla = \left[ \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right] \quad (8)$$

to both sides. Remembering that the operator acts on  $\mathbf{x}$  and not  $\mathbf{x}'$ , we have

$$\nabla^2 \Phi(\mathbf{x}) = - \int G\rho(\mathbf{x}') \nabla^2 \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) d^3 \mathbf{x}'. \quad (9)$$

We can evaluate this by noting that

$$\nabla \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right), \nabla^2 \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) = 0. \quad (10)$$

So we conclude that outside of a very small region around  $\mathbf{x}$ ,  $\nabla^2 \Phi(\mathbf{x}) = 0$ . Let's take a spherical region  $S(\epsilon)$  of radius  $\epsilon$  centered on  $\mathbf{x}$ . In proceeding, let's note that

$$\nabla^2 f(|\mathbf{x} - \mathbf{x}'|) = \nabla_{\mathbf{x}'}^2 f(|\mathbf{x} - \mathbf{x}'|) \quad (11)$$

for any function  $f(|\mathbf{x} - \mathbf{x}'|)$ . If we take  $\epsilon$  to be small enough such that  $\rho(\mathbf{x}) \approx$  a constant, then we can write

$$\begin{aligned} \nabla^2 \Phi(\mathbf{x}) &\approx -G\rho(\mathbf{x}) \int_{S(\epsilon)} \nabla^2 \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) d^3 \mathbf{x}' \\ &= -G\rho(\mathbf{x}) \int_{S(\epsilon)} \nabla_{\mathbf{x}}^2 \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) dV'. \end{aligned} \quad (12)$$

Now we get to use the *divergence* theorem

$$\int \nabla^2 f dV = \oint \nabla f \cdot dS, \quad (13)$$

which allows us to write Equation 12 as

$$-G\rho(\mathbf{x}) \int_{S(\epsilon)} \nabla_{\mathbf{x}}^2 \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) dV' = -G\rho(\mathbf{x}) \oint_S (\epsilon) \nabla \left( \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) \quad (14)$$