

# Snowmass2021 - Letter of Interest

## *Roman Space Telescope Strong Lensing Probes of Dark Matter Substructure*

**Thematic Areas:** (check all that apply ☐/☒)

- ☐ (CF1) Dark Matter: Particle Like
- ☐ (CF2) Dark Matter: Wavelike
- ☒ (CF3) Dark Matter: Cosmic Probes
- ☐ (CF4) Dark Energy and Cosmic Acceleration: The Modern Universe
- ☐ (CF5) Dark Energy and Cosmic Acceleration: Cosmic Dawn and Before
- ☐ (CF6) Dark Energy and Cosmic Acceleration: Complementarity of Probes and New Facilities
- ☐ (CF7) Cosmic Probes of Fundamental Physics
- ☐ (Other) *[Please specify frontier/topical group]*

**Contact Information:** (authors listed after the text)

Submitter Name/Institution: Brant Robertson (University of California, Santa Cruz)

Collaboration: Nancy Grace Roman Formulation Science Working Group

Contact Email: brant@ucsc.edu

**Abstract:** *Nancy Grace Roman Space Telescope*, formerly known as the *Wide Field Infrared Survey Telescope*, is scheduled to launch in the mid-2020's and will provide a multi-band, high-resolution view of the cosmos. Equipped with a mirror the size of *Hubble Space Telescope* and a  $0.3 \text{ deg}^2$  infrared-sensitive camera  $\sim 100\times$  larger than *Hubble/Wide Field Camera 3*, *Roman* will efficiently and sensitively map large areas of the sky with  $\sim 0.1''$  resolution. This resolution will enable the discovery of  $\sim 1000$  gravitationally-lensed quasars, which in turn provide detailed constraints on the properties of dark matter substructures. We review the promise of *Roman* for achieving dark matter constraints through strong lensing, both as stand-alone lensing experiment and in collaboration with other facilities.

*Nancy Grace Roman Space Telescope (Roman)* is a 2.4m space telescope scheduled to be launched in the mid-2020's by the United States National Aeronautics and Space Administration (NASA)<sup>1,2</sup>. Formerly known as the *Wide Field Infrared Survey Telescope*, *Roman* will undertake an ambitious set of mission surveys including a large area cosmology survey ( $\sim 1600 \text{ deg}^2$ ) with multiband imaging ( $H \sim 26.7$  AB flux limit) and slitless grism spectroscopy ( $\sim 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$  line sensitivity at  $\lambda \sim 1.5 \mu\text{m}$ ).

While the cosmology mission surveys are designed to constrain structure formation and the expansion history through weak lensing and baryon acoustic oscillations, *Roman* can test our picture for small-scale dark matter structure formation more directly through gravitational lensing. The cold dark matter plus cosmological constant ( $\Lambda$ CDM) paradigm forms one of the pillars of our models for the origin and evolution of cosmic structures. While remarkably successful at matching observations on large scales, there have been persistent observational challenges to the cold, collisionless dark matter expectations on dwarf-galaxy scales. These “small-scale controversies” may simply stem from a poor understanding of the baryonic processes involved in galaxy formation, or indicate more complex dark sector physics<sup>3</sup>.

Detailed testing of the standard paradigm on small scales remains one of the most pressing issues in cosmology. Numerical simulations in  $\Lambda$ CDM predict a rich spectrum of substructure in galaxy halos. Small fluctuations in the galaxy-scale lensing potential caused by these substructures should result in measurable “flux anomalies” in the magnifications of quadruply-lensed quasar images<sup>4</sup>. While discrepancies between the observed flux ratios and those predicted by a smooth lens model may have been found in quasar lenses<sup>5–7</sup>, the small size of current samples limits our understanding ( $\sim 56$  quad lenses<sup>8</sup>, even fewer with high-resolution imaging). According to predictions of the occurrence of multiply-imaged strong-lensed quasars in wide-area surveys<sup>9</sup>, the *Roman* cosmology survey will revolutionize this field by increasing the sample of quad lenses by more than  $20\times$  to  $\sim 1000$  objects.

The ability to use strong-lensed quasars as probes of the small-scale structure of dark matter halos can be powerful for constraining the nature of dark matter itself. The quasar strong-lensing method is potentially sensitive to the presence of any dark matter component that may suppress small-scale power. Subhalos in the host galaxy lens and in structures along the line of sight<sup>10</sup> affect the image positions and flux ratios. By modeling how the mass function of subhalos, which is directly controlled by the small-scale power spectrum, influences the lensing signal, the presence of a warm dark matter component can be constrained. For instance, Gilman et al. 2020<sup>11</sup> used space-based imaging<sup>12</sup> of only 8 multiply-lensed quasars to place 95% confidence limits of  $m_{DM} > 5.2 \text{ keV}$  assuming thermal relics. With increased samples of dozens of quad-image strong lenses with high-resolution imaging, constraints can be achieved<sup>13</sup> on sterile neutrino dark matter models with masses  $m_{DM} \sim 7 \text{ keV}$  required to explain the 3.5keV line seen in *XMM-Newton* and *Chandra* x-ray data<sup>14–16</sup>.

Over the last decade, methods have been developed to use extended lensed sources, in addition to the point-like AGN, to identify substructures through distortions of lensed images. Extended sources are imaged into arcs and rings that probe a larger volume in the lens halo, increasing the number of subhalos that can be sensed. They also can be drawn from larger populations of background sources, typical galaxies rather than bright AGN. Vegetti & Koopman (2009)<sup>17</sup> described a method for identifying substructures with high-resolution optical/IR imaging of extended galaxies, with Vegetti et al. (2012)<sup>18</sup> reporting a detection of a dark substructure in a gravitational lens at  $z = 0.9$ . In related work, Hezaveh et al. (2013, 2016a)<sup>19;20</sup> described the use of (sub-)millimeter spectral lines from dusty lensed galaxies to search for subhalos in the lenses. The wide survey area of the *Roman* cosmology survey, along with its combination of multi-band imaging and spectroscopy, will provide many avenues for identifying large samples of lensed galaxies for such substructure searches. By combining space-based and ground-based surveys<sup>21–23</sup>, *Roman* can contribute by extending strong-lensing probes of dark matter to halo inner density profile slope measurements<sup>24;25</sup>, group-to-cluster scale lenses<sup>26</sup>, and statistical measures of the matter power spectrum<sup>27</sup>.

The ability of *Roman* to provide improved constraints on the small-scale matter power spectrum through lensing is significantly enhanced by its high-resolution, space-based, multiband imaging and grism spectroscopy. The *Roman* cosmology survey is expected to have  $Y$ ,  $J$ , and  $H$  filter images, along with additional coverage in the  $F184$  filter that extends to  $\lambda \approx 2\mu\text{m}$ . The high-resolution imagery will help combat potential systematic issues with interpreting flux anomalies in the presences of baryonic structures like disks<sup>28</sup>, and help resolve uncertainties about the potential impact of central baryonic components on the survivability and presence of dark substructure<sup>29;30</sup>. This array of near-infrared filters will reduce the effect of dust attenuation from the lensing galaxies on the source images, improving the sensitivity to multiply imaged lenses. With the high spatial resolution of the grism spectroscopy, *Roman* gains a complementary probe using the flux ratio of strong quasar lines such as  $\text{O}_{\text{III}}$  to constrain substructure<sup>31</sup>.

Using *Roman*, astronomers will explore the substructure science achievable from the mission cosmology surveys and various guest investigator and observer programs.

## References

- [1] Spergel, D. *et al.* Wide-Field Infrared Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA 2015 Report. *arXiv e-prints* arXiv:1503.03757 (2015). [1503.03757](#).
- [2] Akeson, R. *et al.* The Wide Field Infrared Survey Telescope: 100 Hubbles for the 2020s. *arXiv e-prints* arXiv:1902.05569 (2019). [1902.05569](#).
- [3] Bullock, J. S. & Boylan-Kolchin, M. Small-Scale Challenges to the  $\Lambda$ CDM Paradigm. *ARAA* **55**, 343–387, DOI: [10.1146/annurev-astro-091916-055313](#) (2017). [1707.04256](#).
- [4] Metcalf, R. B. & Madau, P. Compound Gravitational Lensing as a Probe of Dark Matter Substructure within Galaxy Halos. *ApJ* **563**, 9–20, DOI: [10.1086/323695](#) (2001). [astro-ph/0108224](#).
- [5] Mao, S. & Schneider, P. Evidence for substructure in lens galaxies? *MNRAS* **295**, 587–594, DOI: [10.1046/j.1365-8711.1998.01319.x](#) (1998). [astro-ph/9707187](#).
- [6] Dalal, N. & Kochanek, C. S. Direct Detection of Cold Dark Matter Substructure. *ApJ* **572**, 25–33, DOI: [10.1086/340303](#) (2002). [astro-ph/0111456](#).
- [7] Metcalf, R. B. & Zhao, H. Flux Ratios as a Probe of Dark Substructures in Quadruple-Image Gravitational Lenses. *ApJL* **567**, L5–L8, DOI: [10.1086/339798](#) (2002). [astro-ph/0111427](#).
- [8] Lemon, C. *Gravitationally Lensed Quasar Database* (2020).
- [9] Oguri, M. & Marshall, P. J. Gravitationally lensed quasars and supernovae in future wide-field optical imaging surveys. *MNRAS* **405**, 2579–2593, DOI: [10.1111/j.1365-2966.2010.16639.x](#) (2010). [arXiv:1001.2037](#).
- [10] Hsueh, J. W. *et al.* SHARP - VII. New constraints on the dark matter free-streaming properties and substructure abundance from gravitationally lensed quasars. *MNRAS* **492**, 3047–3059, DOI: [10.1093/mnras/stz3177](#) (2020). [1905.04182](#).
- [11] Gilman, D. *et al.* Warm dark matter chills out: constraints on the halo mass function and the free-streaming length of dark matter with eight quadruple-image strong gravitational lenses. *MNRAS* **491**, 6077–6101, DOI: [10.1093/mnras/stz3480](#) (2020). [1908.06983](#).
- [12] Nierenberg, A. M. *et al.* Double dark matter vision: twice the number of compact-source lenses with narrow-line lensing and the WFC3 grism. *MNRAS* **492**, 5314–5335, DOI: [10.1093/mnras/stz3588](#) (2020). [1908.06344](#).
- [13] Despali, G., Lovell, M., Vegetti, S., Crain, R. A. & Oppenheimer, B. D. The lensing properties of subhaloes in massive elliptical galaxies in sterile neutrino cosmologies. *MNRAS* **491**, 1295–1310, DOI: [10.1093/mnras/stz3068](#) (2020). [1907.06649](#).
- [14] Bulbul, E. *et al.* Detection of an Unidentified Emission Line in the Stacked X-Ray Spectrum of Galaxy Clusters. *ApJ* **789**, 13, DOI: [10.1088/0004-637X/789/1/13](#) (2014). [1402.2301](#).
- [15] Boyarsky, A., Franse, J., Iakubovskiy, D. & Ruchayskiy, O. Checking the Dark Matter Origin of a 3.53 keV Line with the Milky Way Center. *Phys. Rev. Lett.* **115**, 161301, DOI: [10.1103/PhysRevLett.115.161301](#) (2015). [1408.2503](#).
- [16] Cappelluti, N. *et al.* Searching for the 3.5 keV Line in the Deep Fields with Chandra: The 10 Ms Observations. *ApJ* **854**, 179, DOI: [10.3847/1538-4357/aaaa68](#) (2018). [1701.07932](#).

- [17] Vegetti, S. & Koopmans, L. V. E. Bayesian strong gravitational-lens modelling on adaptive grids: objective detection of mass substructure in Galaxies. *MNRAS* **392**, 945–963, DOI: [10.1111/j.1365-2966.2008.14005.x](https://doi.org/10.1111/j.1365-2966.2008.14005.x) (2009). [arXiv:0805.0201](https://arxiv.org/abs/0805.0201).
- [18] Vegetti, S. *et al.* Gravitational detection of a low-mass dark satellite galaxy at cosmological distance. *Nature* **481**, 341–343, DOI: [10.1038/nature10669](https://doi.org/10.1038/nature10669) (2012). [arXiv:1201.3643](https://arxiv.org/abs/1201.3643).
- [19] Hezaveh, Y. D. *et al.* ALMA Observations of SPT-discovered, Strongly Lensed, Dusty, Star-forming Galaxies. *ApJ* **767**, 132, DOI: [10.1088/0004-637X/767/2/132](https://doi.org/10.1088/0004-637X/767/2/132) (2013). [arXiv:1303.2722](https://arxiv.org/abs/1303.2722).
- [20] Hezaveh, Y. D. *et al.* Detection of Lensing Substructure Using ALMA Observations of the Dusty Galaxy SDP.81. *ApJ* **823**, 37, DOI: [10.3847/0004-637X/823/1/37](https://doi.org/10.3847/0004-637X/823/1/37) (2016). [1601.01388](https://arxiv.org/abs/1601.01388).
- [21] Agnello, A. *et al.* DES meets Gaia: discovery of strongly lensed quasars from a multiplet search. *MNRAS* **479**, 4345–4354, DOI: [10.1093/mnras/sty1419](https://doi.org/10.1093/mnras/sty1419) (2018). [1711.03971](https://arxiv.org/abs/1711.03971).
- [22] Anguita, T. *et al.* The STRong lensing Insights into the Dark Energy Survey (STRIDES) 2016 follow-up campaign - II. New quasar lenses from double component fitting. *MNRAS* **480**, 5017–5028, DOI: [10.1093/mnras/sty2172](https://doi.org/10.1093/mnras/sty2172) (2018). [1805.12151](https://arxiv.org/abs/1805.12151).
- [23] Everett, W. B. *et al.* Millimeter-wave Point Sources from the 2500-square-degree SPT-SZ Survey: Catalog and Population Statistics. *arXiv e-prints* [arXiv:2003.03431](https://arxiv.org/abs/2003.03431) (2020). [2003.03431](https://arxiv.org/abs/2003.03431).
- [24] Gavazzi, R. *et al.* The Sloan Lens ACS Survey. IV. The Mass Density Profile of Early-Type Galaxies out to 100 Effective Radii. *ApJ* **667**, 176–190, DOI: [10.1086/519237](https://doi.org/10.1086/519237) (2007). [astro-ph/0701589](https://arxiv.org/abs/astro-ph/0701589).
- [25] Wong, K. C. *et al.* ALMA Observations of the Gravitational Lens SDP.9. *ApJL* **843**, L35, DOI: [10.3847/2041-8213/aa7d4a](https://doi.org/10.3847/2041-8213/aa7d4a) (2017). [1707.00702](https://arxiv.org/abs/1707.00702).
- [26] Jaelani, A. T. *et al.* Survey of Gravitationally lensed Objects in HSC Imaging (SuGOHI) - V. Group-to-cluster scale lens search from the HSC-SSP Survey. *MNRAS* **495**, 1291–1310, DOI: [10.1093/mnras/staa1062](https://doi.org/10.1093/mnras/staa1062) (2020). [2002.01611](https://arxiv.org/abs/2002.01611).
- [27] Diaz Rivero, A., Cyr-Racine, F.-Y. & Dvorkin, C. Power spectrum of dark matter substructure in strong gravitational lenses. *Phys. Rev. D* **97**, 023001, DOI: [10.1103/PhysRevD.97.023001](https://doi.org/10.1103/PhysRevD.97.023001) (2018). [1707.04590](https://arxiv.org/abs/1707.04590).
- [28] Hsueh, J. W. *et al.* SHARP - IV. An apparent flux-ratio anomaly resolved by the edge-on disc in B0712+472. *MNRAS* **469**, 3713–3721, DOI: [10.1093/mnras/stx1082](https://doi.org/10.1093/mnras/stx1082) (2017). [1701.06575](https://arxiv.org/abs/1701.06575).
- [29] Garrison-Kimmel, S. *et al.* Not so lumpy after all: modelling the depletion of dark matter subhaloes by Milky Way-like galaxies. *MNRAS* **471**, 1709–1727, DOI: [10.1093/mnras/stx1710](https://doi.org/10.1093/mnras/stx1710) (2017). [1701.03792](https://arxiv.org/abs/1701.03792).
- [30] Graus, A. S., Bullock, J. S., Boylan-Kolchin, M. & Nierenberg, A. M. Through a Smoother Lens: An expected absence of LCDM substructure detections from hydrodynamic and dark matter only simulations. *MNRAS* **480**, 1322–1332, DOI: [10.1093/mnras/sty1924](https://doi.org/10.1093/mnras/sty1924) (2018). [1710.11148](https://arxiv.org/abs/1710.11148).
- [31] Nierenberg, A. M. *et al.* Probing dark matter substructure in the gravitational lens HE 0435-1223 with the WFC3 grism. *MNRAS* **471**, 2224–2236, DOI: [10.1093/mnras/stx1400](https://doi.org/10.1093/mnras/stx1400) (2017). [1701.05188](https://arxiv.org/abs/1701.05188).

**Authors:**

Brant Robertson (UC Santa Cruz)