

Class test 2

Test duration is for 75 minutes and 10 minutes extra for submission. Answer all. Write the answer as instructed

in the question. After the examination submit rough copy to signalsystemspkd@gmail.com, give roll number as the file name.

...

Points: /40

1. A signal $x(t)$ is given by the following equation. choose the most correct answer for its Fourier transform $X(j\omega)$

(2/2 Points)

$x(t) = \sum_{n=-\infty}^{+\infty} \delta(t - nT)$, represent a periodic train of delta functions

☐ $X(j\omega) = 1$

☐ $X(j\omega) = \infty$

☐ $\sum_{n=-\infty}^{+\infty} \cos(\omega_0 n)$, $\omega_0 = \frac{2\pi}{T}$

☒ $X(j\omega) = \omega_0 \sum_{n=-\infty}^{+\infty} \delta(\omega - \omega_0 n)$, $\omega_0 = \frac{2\pi}{T}$



2. A periodic signal is given by $x(t) = 1 - (4/T)t$, $0 < t < T/2$ and $x(t) = (4/T)t - 3$, $T/2 < t < T$ and

have 0-th, 1st, 2nd and 3rd Fourier series coefficients a_0 , a_1 , a_2 and a_3 respectively.

Choose the most correct option

(4/4 Points)

☐ $a_0 = 0, a_1 = 0.202, a_2 = 0, a_3 = 0.81$

☐ $a_0 = 0.202, a_1 = 0.81, a_2 = 0, a_3 = 0$

☒ $a_0 = 0, a_1 = 0.81, a_2 = 0, a_3 = 0.09$ ✓



3. $h[n]$ and $x[n]$ is given by the following equation. $y[n]$ is the output of convolution of $h[n]$ and $x[n]$.

Find $y[14]$, value of $y[n]$ at $n=14$. Convolution property of DTFT may be used to compute $y[14]$.

$\delta[n]$ in the equation is the digital delta function. Give the answer rounded up to two decimal place

(0/4 Points)

$$h[n] = \delta[n] - \frac{\sin\left(\frac{\pi n}{8}\right)}{\pi n}, \quad x[n] = \cos[\pi n] + \sin\left[\frac{\pi n}{7} + \frac{1}{2}\right];$$

0.92

Correct answers: 0.479, 0.48, .48

4. Name *



5. A periodic signal $x(t)$ is given by the following equation. Find the 0-th Fourier series coefficient a_0 .

Write your answer rounded to two decimal places.

(4/4 Points)

$$x(t) = t^2, \quad -\pi < t < \pi \text{ and } x(t + 2\pi) = x(t) \text{ i.e. period of } x(t) \text{ is } 2\pi$$

3.29



6. An LTI system is given by the following difference equation. Its impulse response is given by $h[n]$.

Find the value of $h[n]$ at $n=1$. Write your answer rounded up to two decimal places.

(4/4 Points)

$$y[n] - \frac{1}{2}y[n-1] = x[n] - \frac{1}{4}x[n-1]$$

0.25 ✓

7. A signal $x(t)$ is given by the following equation has first four Fourier coefficients starting 0th coefficient as a_0 , a_1 , a_2 and a_3 . Choose the most correct option.

(4/4 Points)

$$x(t) = 1 + 2 \sin\left(\frac{7\pi}{3}t\right) - 3 \cos\left(\frac{7\pi}{3}t\right) + \sin(7\pi t) + \cos\left(\frac{28\pi}{3}t\right)$$

☐ $a_0 = \frac{1}{2}, a_1 = 1, a_2 = -\frac{j}{2}, a_3 = -\left(\frac{3}{2} + j1\right)$

☒ $a_0 = 1, a_1 = -\left(\frac{3}{2} + j1\right), a_2 = -\frac{j}{2}, a_3 = \frac{1}{2}$ ✓

☐ $a_0 = 1, a_1 = -\left(\frac{3}{2} + j1\right), a_2 = -\frac{j}{2}, a_3 = 0$



8. A system is given by a difference equation as given below. Determine whether the system is

stable or unstable, choose the most correct option


(0/2 Points)

$$y[n] - \frac{11}{6}y[n-1] - \frac{1}{2}y[n-2] + \frac{1}{3}y[n-3] = x[n] + 2x[n-2]$$

☒ stable

☐ unstable ✓

9. Roll Number *



10. A signal $x(t)$ is given by the following equation. Calculate the absolute value of the a_3 , fourth Fourier series coefficient. Write your answer rounded to two decimal places.
(0/4 Points)

$$x(t) = -1, -1 < t < 0; x(t) = 1, 0 < t < 1 \text{ and } x(t+2) = x(t). \text{ Find } |a_3|$$

0.11

Correct answers: 0.212,0.21

11. A signal $x[n]$ is given by the following equation, choose the most correct answer.
(2/2 Points)

$$x[n] = \cos(\sqrt{2} \pi n)$$

- ☐ Both discrete time Fourier series and discrete time Fourier transform exists for $x[n]$
☒ only discrete time Fourier transform exists for $x[n]$ ✓
☐ only discrete time Fourier series exists for $x[n]$

12. An impulse response $h[n]$ is given by the following equation. To check stability one has to evaluate the sum S given as follows. Find S . DTFT summation rule may be used. Write the answer rounded to two decimal places.
(4/4 Points)

$$h[n] = \left\{ n^2 \left(-\frac{1}{2}\right)^n + 2 \left(\frac{1}{4}\right)^n \right\} u[n]; S = \sum_{n=-\infty}^{+\infty} |h[n]|$$

8.67 ✓

13. An LTI system has an impulse response $h(t)$ and corresponding output $y(t)$ given by the following equations. Also the equation of $x(t)$ is given. Find the value of the constant K in $x(t)$.

(4/4 Points)

$$h(t) = e^{-2t}u(t), y(t) = \{e^{-2t} - e^{-3t}\}u(t), x(t) = e^{Kt}u(t)$$

-3 ✓

14. A signal $x(t)$ is given by the following equation. Find $x(jw)$ at $w=2$. Write the answer rounded up to two decimal places

(2/2 Points)

$$x(t) = e^{-2|t|}, \text{ find } X(jw) \text{ at } w = 2$$

0.5 ✓

This content is created by the owner of the form. The data you submit will be sent to the form owner. Microsoft is not responsible for the privacy or security practices of its customers, including those of this form owner. Never give out your password.

Powered by Microsoft Forms | [Privacy and cookies](#) | [Terms of use](#)