

Exercises

- 5.1.1 Defects occur along the length of a filament at a rate of $\lambda = 2$ per foot.
- Calculate the probability that there are no defects in the first foot of the filament.
 - Calculate the conditional probability that there are no defects in the second foot of the filament, given that the first foot contained a single defect.
- 5.1.2 Let $p_k = \Pr\{X = k\}$ be the probability mass function corresponding to a Poisson distribution with parameter λ . Verify that $p_0 = \exp\{-\lambda\}$, and that p_k may be computed recursively by $p_k = (\lambda/k)p_{k-1}$.
- 5.1.3 Let X and Y be independent Poisson distributed random variables with parameters α and β , respectively. Determine the conditional distribution of X , given that $N = X + Y = n$.
- 5.1.4 Customers arrive at a service facility according to a Poisson process of rate λ customer/hour. Let $X(t)$ be the number of customers that have arrived up to time t .
- What is $\Pr\{X(t) = k\}$ for $k = 0, 1, \dots$?
 - Consider fixed times $0 < s < t$. Determine the conditional probability $\Pr\{X(t) = n + k | X(s) = n\}$ and the expected value $E[X(t)X(s)]$.
- 5.1.5 Suppose that a random variable X is distributed according to a Poisson distribution with parameter λ . The parameter λ is itself a random variable, exponentially distributed with density $f(x) = \theta e^{-\theta x}$ for $x \geq 0$. Find the probability mass function for X .
- 5.1.6 Messages arrive at a telegraph office as a Poisson process with mean rate of 3 messages per hour.
- What is the probability that no messages arrive during the morning hours 8:00 A.M. to noon?
 - What is the distribution of the time at which the first afternoon message arrives?
- 5.1.7 Suppose that customers arrive at a facility according to a Poisson process having rate $\lambda = 2$. Let $X(t)$ be the number of customers that have arrived up to time t . Determine the following probabilities and conditional probabilities:
- $\Pr\{X(1) = 2\}$.
 - $\Pr\{X(1) = 2 \text{ and } X(3) = 6\}$.
 - $\Pr\{X(1) = 2 | X(3) = 6\}$.
 - $\Pr\{X(3) = 6 | X(1) = 2\}$.
- 5.1.8 Let $\{X(t); t \geq 0\}$ be a Poisson process having rate parameter $\lambda = 2$. Determine the numerical values to two decimal places for the following probabilities:
- $\Pr\{X(1) \leq 2\}$.
 - $\Pr\{X(1) = 1 \text{ and } X(2) = 3\}$.
 - $\Pr\{X(1) \geq 2 | X(1) \geq 1\}$.
- 5.1.9 Let $\{X(t); t \geq 0\}$ be a Poisson process having rate parameter $\lambda = 2$. Determine the following expectations:
- $E[X(2)]$.
 - $E[X(1)^2]$.
 - $E[X(1)X(2)]$.

Problems

- 5.1.1 Let ξ_1, ξ_2, \dots be independent random variables, each having an exponential distribution with parameter λ . Define a new random variable X as follows: If $\xi_1 > 1$, then $X = 0$; if $\xi_1 \leq 1$ but $\xi_1 + \xi_2 > 1$, then set $X = 1$; in general, set $X = k$ if

$$\xi_1 + \dots + \xi_k \leq 1 < \xi_1 + \dots + \xi_k + \xi_{k+1}.$$

Show that X has a Poisson distribution with parameter λ . (Thus, the method outlined can be used to simulate a Poisson distribution.)

Hint: $\xi_1 + \dots + \xi_k$ has a gamma density

$$f_k(x) = \frac{\lambda^k x^{k-1}}{(k-1)!} e^{-\lambda x} \quad \text{for } x > 0.$$

Condition on $\xi_1 + \dots + \xi_k$ and use the law of total probability to show

$$\Pr\{X = k\} = \int_0^1 [1 - F(1-x)] f_k(x) dx,$$

where $F(x)$ is the exponential distribution function.

- 5.1.2** Suppose that minor defects are distributed over the length of a cable as a Poisson process with rate α , and that, independently, major defects are distributed over the cable according to a Poisson process of rate β . Let $X(t)$ be the number of defects, either major or minor, in the cable up to length t . Argue that $X(t)$ must be a Poisson process of rate $\alpha + \beta$.
- 5.1.3** The *generating function* of a probability mass function $p_k = \Pr\{X = k\}$, for $k = 0, 1, \dots$ is defined by

$$g_X(s) = E[s^X] = \sum_{k=0}^{\infty} p_k s^k \quad \text{for } |s| < 1.$$

Show that the generating function for a Poisson random variable X with mean μ is given by

$$g_X(s) = e^{-\mu(1-s)}.$$

- 5.1.4** (Continuation) Let X and Y be independent random variables, Poisson distributed with parameters α and β , respectively. Show that the generating function of their sum $N = X + Y$ is given by

$$g_N(s) = e^{-(\alpha+\beta)(1-s)}.$$

Hint: Verify and use the fact that the generating function of a sum of independent random variables is the product of their respective generating functions. See Chapter 3, Section 3.9.2.

- 5.1.5** For each value of $h > 0$, let $X(h)$ have a Poisson distribution with parameter λh . Let $p_k(h) = \Pr\{X(h) = k\}$ for $k = 0, 1, \dots$. Verify that

$$\lim_{h \rightarrow 0} \frac{1 - p_0(h)}{h} = \lambda, \quad \text{or } p_0(h) = 1 - \lambda h + o(h);$$

$$\lim_{h \rightarrow 0} \frac{p_1(h)}{h} = \lambda, \quad \text{or } p_1(h) = \lambda h + o(h);$$

$$\lim_{h \rightarrow 0} \frac{p_2(h)}{h} = 0, \quad \text{or } p_2(h) = o(h).$$

Here $o(h)$ stands for any remainder term of order less than h as $h \rightarrow 0$.

- 5.1.6** Let $\{X(t); t \geq 0\}$ be a Poisson process of rate λ . For $s, t > 0$, determine the conditional distribution of $X(t)$, given that $X(t+s) = n$.

- 5.1.7** Shocks occur to a system according to a Poisson process of rate λ . Suppose that the system survives each shock with probability α , independently of other shocks, so that its probability of surviving k shocks is α^k . What is the probability that the system is surviving at time t ?

- 5.1.8** Find the probability $\Pr\{X(t) = 1, 3, 5, \dots\}$ that a Poisson process having rate λ is odd.

- 5.1.9** Arrivals of passengers at a bus stop form a Poisson process $X(t)$ with rate $\lambda = 2$ per unit time. Assume that a bus departed at time $t = 0$ leaving no customers behind. Let T denote the arrival time of the next bus. Then, the number of passengers present when it arrives is $X(T)$. Suppose that the bus arrival time T is independent of the Poisson process and that T has the uniform probability density function

$$f_T(t) = \begin{cases} 1 & \text{for } 0 \leq t \leq 1, \\ 0 & \text{elsewhere.} \end{cases}$$

- (a) Determine the conditional moments $E[X(T)|T=t]$ and $E[\{X(T)\}^2|T=t]$.
 (b) Determine the mean $E[X(T)]$ and variance $\text{Var}[X(T)]$.

Exercises

- 5.3.1 A radioactive source emits particles according to a Poisson process of rate $\lambda = 2$ particles per minute. What is the probability that the first particle appears after 3 min?
- 5.3.2 A radioactive source emits particles according to a Poisson process of rate $\lambda = 2$ particles per minute.
- (a) What is the probability that the first particle appears some time after 3 min but before 5 min?
 - (b) What is the probability that exactly one particle is emitted in the interval from 3 to 5 min?
- 5.3.3 Customers enter a store according to a Poisson process of rate $\lambda = 6$ per hour. Suppose it is known that only a single customer entered during the first hour. What is the conditional probability that this person entered during the first 15 min?
- 5.3.4 Let $X(t)$ be a Poisson process of rate $\xi = 3$ per hour. Find the conditional probability that there were two events in the first hour, given that there were five events in the first 3 h.
- 5.3.5 Let $X(t)$ be a Poisson process of rate θ per hour. Find the conditional probability that there were m events in the first t hours, given that there were n events in the first T hours. Assume $0 \leq m \leq n$ and $0 < t < T$.
- 5.3.6 For $i = 1, \dots, n$, let $\{X_i(t); t \geq 0\}$ be independent Poisson processes, each with the same parameter λ . Find the distribution of the first time that at least one event has occurred in every process.
- 5.3.7 Customers arrive at a service facility according to a Poisson process of rate λ customers/hour. Let $X(t)$ be the number of customers that have arrived up to time t . Let W_1, W_2, \dots be the successive arrival times of the customers. Determine the conditional mean $E[W_5 | X(t) = 3]$.
- 5.3.8 Customers arrive at a service facility according to a Poisson process of rate $\lambda = 5$ per hour. Given that 12 customers arrived during the first two hours of service, what is the conditional probability that 5 customers arrived during the first hour?
- 5.3.9 Let $X(t)$ be a Poisson process of rate λ . Determine the cumulative distribution function of the gamma density as a sum of Poisson probabilities by first verifying and then using the identity $W_r \leq t$ if and only if $X(t) \geq r$.