

**Topic:** Identity matrices**Question:** Which identity matrix is  $I_3$ ?**Answer choices:**

A  $I_3 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

B  $I_3 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

C  $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

D  $I_3 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$



**Solution: C**

We always call the identity matrix  $I$ , and it's always a square matrix, like  $2 \times 2$ ,  $3 \times 3$ ,  $4 \times 4$ , etc. For that reason, it's common to abbreviate  $I_{2 \times 2}$  as just  $I_2$ , or  $I_{3 \times 3}$  as just  $I_3$ , etc. So  $I_3$  is the  $3 \times 3$  identity matrix.

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



**Topic:** Identity matrices

**Question:** Which identity matrix can be multiplied by  $A$  (in other words,  $IA$ ), if  $A$  is a  $2 \times 4$  matrix?

**Answer choices:**

A  $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

B  $I_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

C  $I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

D  $I_4 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$



**Solution: A**

Start by setting up the equation  $IA = A$ . Next, substitute the dimensions for  $A$  into the equation.

$$I \cdot A = A$$

$$I \cdot 2 \times 4 = 2 \times 4$$

Break down the dimensions of the identity matrix as rows  $\times$  columns.

$$R \times C \cdot 2 \times 4 = 2 \times 4$$

In order to be able to multiply matrices, we need the same number of columns in the first matrix as we have rows in the second matrix. So the identity matrix must have 2 columns.

$$R \times 2 \cdot 2 \times 4 = 2 \times 4$$

And the dimensions of the resulting matrix come from the rows of the first matrix and the columns of the second matrix. So the identity matrix must have 2 rows.

$$2 \times 2 \cdot 2 \times 4 = 2 \times 4$$

Therefore, the identity matrix we need is  $I_2$ .

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



**Topic:** Identity matrices

**Question:** If we want to find  $IA$ , which identity matrix should we use, and what is the product?

$$A = \begin{bmatrix} -3 & 7 & 1 \\ 2 & 8 & 4 \end{bmatrix}$$

**Answer choices:**

A      Use  $I_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ , and the product is  $IA = \begin{bmatrix} 2 & 8 & 4 \\ -3 & 7 & 1 \end{bmatrix}$

B      Use  $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ , and the product is  $IA = \begin{bmatrix} -3 & 7 & 1 \\ 2 & 8 & 4 \end{bmatrix}$

C      Use  $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , and the product is  $IA = \begin{bmatrix} -3 & 7 & 1 \\ 2 & 8 & 4 \end{bmatrix}$

D      Use  $I_3 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ , and the product is  $IA = \begin{bmatrix} 2 & 8 & 4 \\ -3 & 7 & 1 \end{bmatrix}$



**Solution: B**

Matrix  $A$  is a  $2 \times 3$  matrix, and we need to find  $IA$ . We also know that  $IA$  will be  $2 \times 3$ . So we'll set up an equation of dimensions.

$$I \cdot A = A$$

$$I \cdot 2 \times 3 = 2 \times 3$$

$$R \times C \cdot 2 \times 3 = 2 \times 3$$

For matrix multiplication to be valid, we need the same number of columns in the first matrix as we have rows in the second matrix.

$$R \times 2 \cdot 2 \times 3 = 2 \times 3$$

The dimensions of the result are given by the rows from the first matrix, and columns from the second matrix.

$$2 \times 2 \cdot 2 \times 3 = 2 \times 3$$

So the identity matrix is  $2 \times 2$ , which means it's  $I_2$ .

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Then the product of  $I_2$  and matrix  $A$  is

$$IA = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -3 & 7 & 1 \\ 2 & 8 & 4 \end{bmatrix}$$

$$IA = \begin{bmatrix} 1(-3) + 0(2) & 1(7) + 0(8) & 1(1) + 0(4) \\ 0(-3) + 1(2) & 0(7) + 1(8) & 0(1) + 1(4) \end{bmatrix}$$



$$IA = \begin{bmatrix} -3 & 7 & 1 \\ 2 & 8 & 4 \end{bmatrix}$$

