ASTR/ATOC 5540 name: Baylee Bordwell

## **Problem Set 1**

1. In final plot.

2. The FD-1 approximation first displays larger, sporadic error which decreases until about  $\Delta x = 10^{-8.5}$ , when the error in the approximation begins to grow linearly. The smallest error in f'(x0) is about  $10^{-9}$ , and it occurs at approximately  $\Delta x = 10^{-8.5}$ . This is the effect of the roundoff error diminishing as the dominant source of error in the estimate, and the discretionary error taking over, as can be seen when the discretionary error is added into the plot.

3.

$$f(x0 + \Delta x) = f(x0) + \Delta x f'(x0) + \frac{\Delta x^2}{2} f''(x0) + \cdots$$

$$f'(x0) = \frac{f(x0 + \Delta x) - f(x0)}{\Delta x} + \frac{\Delta x}{2} f''(x0) + \cdots$$

Discretionary error D is approximated as the first neglected term,  $D \approx \frac{\Delta x}{2} f' \prime (x0)$ .

4.  $f(x0 + \Delta x)$  can be seen above,

$$f(x - \Delta x) = f(x0) - \Delta x f'(x0) + \frac{\Delta x^2}{2} f''(x0) + \cdots$$

$$f'_{FD-2}(x0) = \frac{f(x0 + \Delta x) - f(x0 - \Delta x)}{2\Delta x}$$


$$= \frac{1}{2\Delta x} (f(x0) + \Delta x f'(x0) + \frac{\Delta x^2}{2} f''(x0) + \dots - (f(x0) - \Delta x f'(x0) + \frac{\Delta x^2}{2} f''(x0) + \dots))$$

$$= f'(x0) + \frac{\Delta x^2}{6}f'''(x0) + \cdots$$

FD-2 is special in comparison to FD-1 because it removes all terms even in powers of  $\Delta x$  from the Taylor expansion, resulting in a smaller discretionary error,  $D \approx \frac{\Delta x^2}{6} f'''(x0)$ 

- 5. The FD-2 approximation behaves similarly to the FD-1, but with a consistently lower absolute error.
- 6. FD-1 is a first order approximation because we lose all terms after the first derivative in a Taylor expansion of f, while FD-2 is a second order approximation because we lose all terms after the second derivative in a Taylor expansion of f. If we want the lowest error at the lowest computational cost, FD-2 is better to use because we can use a larger  $\Delta x$  and achive a lower error than with FD-1.

ASTR/ATOC 5540 name: Baylee Bordwell



```
_1 #!/usr/bin/python3.4
3 import numpy as np
4 from matplotlib import pyplot as plt
  def first_order_finite_diff(f, x, delta_x):
      return (f(x+delta_x)-f(x))/delta_x
  def second_order_finite_diff(f, x, delta_x):
      return (f(x+delta_x)-f(x-delta_x))/(2*delta_x)
11
def plot_FD_error(f, dfdx, x0, ax, min_delta_x=-10, max_delta_x=10, ddfdxx=
     None, dddfdxxx=None):
      ""Function will find and plot the error in the FD approximation ""
14
15
      #Finding the error in the FD approximation
16
      delta_x = np.logspace(min_delta_x, max_delta_x, base = 10.0)
18
      FD_1_error = np.abs(first_order_finite_diff(f, x0, delta_x)-dfdx(x0))
19
      #print(FD_1_error) Default code
20
21
      FD_2_error = np.abs(second_order_finite_diff(f, x0, delta_x)-dfdx(x0))
23
24
      #Setting up the plot
25
      #ax.loglog(delta_x, FD_1_error, label='FD-1') Default code
26
      if ddfdxx != None:
          d_{err} = np. abs(delta_x/2.*ddfdxx(x0))
28
          ax.loglog(delta_x, d_err, 'r-o', label='FD-1 d_err')
29
      if dddfdxxx != None:
30
          d_{err2} = np.abs(delta_x **2/6.*dddfdxxx(x0))
31
          ax.loglog(delta_x, d_err2, 'g-o', label='FD-2 d_err')
      ax.loglog(delta_x, FD_1_error, 'b-o', label='FD-1')
33
      ax.loglog(delta_x, FD_2_error, 'k-o', label='FD-2')
34
      ax.set_xlabel(r'$\Delta x$')
35
      ax.set_ylabel('Absolute error in FD approximation')
36
      ax.legend(loc="lower right")
37
38
39
40
  if __name__=="__main__":
41
42
      figure_1 = plt.figure()
43
      ax = figure_1 \cdot add_subplot(1,1,1)
      #Defining the particulars of the function to be approximated
      f = np.sin
47
      dfdx = np.cos
```

ASTR/ATOC 5540 name: Baylee Bordwell

```
ddfdxx = lambda x: -1*np.sin(x)
dddfdxxx = lambda x: -1*np.cos(x)
x0 = .25

#plot_FD_error(f, dfdx, x0, ax) Default code
#plot_FD_error(f, dfdx, x0, ax, -20,1) Problem 2
#plot_FD_error(f, dfdx, x0, ax, -20,1, ddfdxx) Problem 3
#plot_FD_error(f, dfdx, x0, ax, -20,1, ddfdxx, dddfdxxx) Problem 4
plot_FD_error(f, dfdx, x0, ax, -15,0, ddfdxx, dddfdxxx)
figure_1.savefig("hw1_fig1.png")
plt.show()
```