

Section 12: Hypothesis testing and p values

STOR 155.02, Spring '21

updated 2021-04-13

What you will learn

- is in the title

Resources

- Textbook: 5.3

Eyes on the schedule

Four lectures remain

- April 13, 20, 27, May 4

Final project due May 11, 3p.m. sharp

- **no excuses for lateness**
- **except** by approved university exam excuses or exceptional circumstances
- assigned one week prior

Formula: pct (decimal) CI

a. Find the low quantile z^*

$$P(Z \leq -z^*) = \frac{1 - \text{pct}}{2}$$

Excel: `NORM.S.INV(.5*(1-pct))` and Python: `stats.norm.ppf(.5*(1-pct))`

with pct in decimals, e.g. 0.025 for 95% CI

b. or find the high quantile (same z^*)

$$P(Z \geq z^*) = 1 - \frac{1 - \text{pct}}{2}$$

c. Calculate the sample point estimates

$$\text{low} = \hat{m}_n - z^* \frac{\hat{s}_n}{\sqrt{n}}, \quad \text{high} = \hat{m}_n + z^* \frac{\hat{s}_n}{\sqrt{n}}$$

Shortcut: `NORM.INV(.5*(1-pct), mhat, shat/sqrtn)` for low

Shortcut: `stats.norm.interval(pct, mhat, shat/sqrtn)` both

need spear *and* net!

thank your classmate for making everything right



Hypothesis testing

Starter example

Adults in Turkey

- **random sample**
- mean caloric intake of **3500 calories / day**
- with s.d. of 500 calories

Sample of 100 Ankara University students

- **random sample**
- average intake of **3450 calories / day**

(Source: FAO, 2006-08 via Wikipedia)

Conclude Ankara U. student average calories is different from Turkish population average?

- could happen if **mean for Ankara U. students is in reality smaller**
- or if we got a different average **just because of randomness in sampling**

Assume averages are the same

What is the probability we would get the sample average 3450 or farther away from 3500 from a random sample size 100?

We have point estimates for the true mean (3500) and standard deviation (500).

By the CLT we know the distribution of the sample mean \bar{X}_{100} is approximately Normal $(3500, \frac{500}{\sqrt{100}})$,

$$\begin{aligned} P\left(\frac{|\bar{X}_{100} - 3500|}{50} > \frac{|3450 - 3500|}{50}\right) &\approx P(Z > 1 \text{ or } Z < -1) \\ &= 2P(Z > 1) \approx 0.32 \end{aligned}$$

That's a pretty high **chance a random sample of size 100 would have average like the one we saw, or smaller!**

Why the 2 above? Area under bell curve for $z < -1$ is the same as for $z > 1$.

Hypothesis test recipe

Model

Data $x_1 \dots x_n$ sampled randomly from an unknown population r.v. X , with true mean m and s.d. s .

Research question

Do our data give strong evidence that

$$m \neq m_0$$

for some number m_0 ?

Could also consider asking a one-sided question, $m < m_0$ or $m > m_0$.

Point estimates

\hat{s}_n, \hat{m}_n are point estimates for s, m based on the data, e.g. \hat{m}_n is the sample mean of $x_1 \dots x_n$.

H_0 'null hypothesis', H_1 'alternative'

- H_0 is the statement you want to reject (if possible)
- using the hypothesis test

$$H_0 : m = m_0, \quad H_1 : m \neq m_0$$

or can consider $H_1 : m < m_0$,
 $H_1 : m > m_0$.

in this class **will always be about true mean m being equal to some other number m_0**

Will try to **'reject H_0 in favor of H_1 '**

Make confidence interval for \bar{X}_n assuming H_0 is true

$$\text{low} = \hat{m}_n - z^* \frac{\hat{s}_n}{\sqrt{n}}, \quad \text{high} = \hat{m}_n + z^* \frac{\hat{s}_n}{\sqrt{n}}$$

If $m_0 > \text{high}$ or $m_0 < \text{low}$, **reject H_0 in favor of H_1**

Which pct to choose for CI?

Rejection with higher pct confidence interval gives stronger evidence

'Significance level' of hypothesis test

$\alpha = \text{test significance level} = 1 - \text{pct used in CI}$

intuitively

Rejecting a hypothesis test with α level means if we redid this experiment we would **incorrectly reject H_0 only α percent of the time**

'incorrectly reject' means our test tells us to reject H_0 when in fact it is true

Example

$\alpha = 0.05$ level corresponds to 0.95 CI



Revisit example

True mean and s.d. daily caloric intake for Ankara students are the unknown numbers m, s .

Hypotheses

$$m_0 = 3500$$

$$H_0 : m = 3500, \quad H_1 : m \neq 3500$$

Confidence interval pct = 0.95

Assume we get the same standard deviation for Ankara students, $\hat{s}_n = 500$.

$$\text{low} = 3450 - z^* \frac{500}{\sqrt{100}}, \quad \text{high} = 3450 + z^* \frac{500}{\sqrt{100}}$$

rounding to 3 decimals

$$(\text{low}, \text{high}) = (2470.018, 4429.982)$$

Q: Reject H_0 ?

Q: Significance level α ?

Q: If you consider a CI with $\text{pct} = 0.98$ would your answer change?

p-values

Definition

"The p-value is the probability of observing data at least as favorable to the alternative hypothesis as our current data set, **if the null hypothesis is true.**"

-- textbook ch 5.3.4

Translation

$$H_0 : m = m_0, \quad H_1 : m \neq m_0$$

p-value is the probability of observing data that is at least $|\hat{m}_n - m_0|$ away from m_0 as what we saw...

meaning chance that if we redrew our sample we would see a sample mean at least as different from m_0 as our current estimate \hat{m}_n

Calculation

'two-sided test'

$$H_1 : m \neq m_0$$

$$p = P \left(\frac{|\bar{X}_n - m_0|}{\hat{s}_n / \sqrt{n}} > \frac{|\hat{m}_n - m_0|}{\hat{s}_n / \sqrt{n}} \right) = 2P \left(Z < -\frac{|\hat{m}_n - m_0|}{\hat{s}_n / \sqrt{n}} \right)$$

'one-sided test'

$$H_1 : m < m_0$$

$$p = P \left(\frac{\bar{X}_n - m_0}{\hat{s}_n / \sqrt{n}} < \frac{\hat{m}_n - m_0}{\hat{s}_n / \sqrt{n}} \right) = P \left(Z < \frac{\hat{m}_n - m_0}{\hat{s}_n / \sqrt{n}} \right)$$

Often you'll see cutoff value referred to as the '**z-statistic**'. These are for the two-sided or one-sided tests:

$$\frac{|\hat{m}_n - m_0|}{\hat{s}_n / \sqrt{n}} \text{ or } \frac{\hat{m}_n - m_0}{\hat{s}_n / \sqrt{n}}$$

Accept-reject hypothesis test using p values

$$H_0 : m = m_0, \quad H_1 : m \neq m_0$$

$\alpha =$ test significance level

reject H_0 if

$$p \leq \alpha$$

This setup holds for one-sided tests too.

Flip it: Significance in terms of p-value

p-value is the largest significance level at which you could reject the null

How small a significance level / p-value is good enough?

Very much depends on the context

choose smaller α when you really don't want to reject incorrectly

Higgs boson discovery:

$$\alpha = 3 \times 10^{-7} \approx 1/3,500,000$$

Scientific American blog post

generic standards you will see

$$\alpha = 0.05, 0.1$$

But why? Should instead think carefully about what makes sense for the question.

Example for poll

It is suspected that patients treated with a certain drug have systolic blood pressures (SBP) that are lower on average than 115, which is the mean for people not treated.

A random sample of 20 patients is treated with the drug. Their sample average SBP is 112 and the sample s.d. is 10.

PollEv.com/brendanbrown849

poll closes at _____

Five more minutes

