Anisotropic Casimir stress on rectangular flat T^3 and a no-attractor result for an instantaneous Bianchi-I closure

Brendon Boyd[1, *]

[1] *Independent Researcher, Burswood, Western Australia, Australia*[†]

(Dated: Version 0.2.0-rc1 — October 1, 2026)

Compact spatial topology can source anisotropic vacuum stress. For a free massless scalar on a rectangular flat three-torus we test an already-renormalized periodic lattice formula for Casimir energy density and directional pressures against cutoff convergence, cubic isotropy, thermodynamic pressure derivatives, conformal trace, dimensional scaling, and an independent heat-kernel/Mellin evaluator at four cubic and noncubic points. We then transport the static shape-dependent stress through a declared instantaneous constitutive closure with a^{-4} dilution on a positive de Sitter biaxial Bianchi-I background. In the bounded scan reported here, this closure produces only *transient* passages near the historically proposed ratio $H_t/H_p = 13/12$, with no quasi-plateau or late-time attractor. This excludes a maintained ratio within the tested closure; it is not a general no-go theorem for a quantum field evolved and renormalized on a dynamical compact spacetime. We do not claim a resolution of the Hubble tension, a derivation of the galactic acceleration scale, or a completed cubic cosmology.

I. INTRODUCTION

Compact flat three-manifolds admit mode spectra that generically produce direction-dependent vacuum expectation values of the stress tensor [1–3]. That fact is distinct from any claim that a fixed numerical ratio of directional Hubble rates is a free parameter-free prediction of topology.

A historically attractive packaging of the latter idea is the cycle-counting assignment $H_t/H_p = 13/12$ together with the numerical identification $H_t = 72.97\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$ from a Planck anchor. Those statements are *not* conclusions of the present paper. Here we isolate two controlled questions that can be addressed with the declared lattice formula and biaxial closure integrator (gate package CBR-001):

1. Does the implemented periodic-scalar stress pass its internal numerical and thermodynamic controls and depend on the shape moduli?
2. When that static stress is transported by the declared instantaneous closure on biaxial expansion, does $H_t/H_p = 13/12$ appear as a free-field attractor, a quasi-plateau, or only as a transient crossing (if at all)?

The answers are: the implemented nonzero-mode static stress passes the internal identity checks, agrees with an independent evaluator at relative order 10^{-6}, and is shape dependent; and the bounded closure calculation yields only transient crossings of 13/12, with every valid candidate returning to isotropy by late expansion.

* brendon.boyd@itsm-cosmology.org
ORCID: 0009-0007-4177-2612
† GitHub: `github.com/brendohxd/ITSM-...`
Zenodo: doi:10.5281/zenodo.18808348
Gate: `Analysis/Casimir/CBR-001/`

Companion work on present-epoch scale matching and weak-field matching invariants [4] records independent no-go results for a different geometric packaging; the Casimir calculation below does not rely on those withdrawn shortcuts and does not restore them.

II. RECTANGULAR LATTICE CASIMIR STRESS

A. Mode sum and pressures

For a massless scalar with periodic lengths (L_1, L_2, L_3), CBR-001 takes as input the already-renormalized rectangular-periodic Epstein-lattice expression developed in Ref. [3]:

$$\rho_{\mathrm{Cas}} = -\frac{\hbar c}{2\pi^2} \sum_{\mathbf{n}\in\mathbb{Z}^3\setminus\{0\}} \frac{1}{\left[(n_1L_1)^2 + (n_2L_2)^2 + (n_3L_3)^2\right]^2}. \tag{1}$$

Directional pressures follow thermodynamically,

$$p_i = -\frac{1}{L_jL_k}\frac{\partial E_{\mathrm{Cas}}}{\partial L_i}, \qquad E_{\mathrm{Cas}} = L_1L_2L_3\rho_{\mathrm{Cas}}. \tag{2}$$

In code and tables we report dimensionless lattice units with the overall $\hbar c$ factor restored only when matching cosmological amplitudes. Figure 1 recalls the rectangular fundamental domain (not an embedded doughnut T^2).

B. Stage-1 validation (cubic point)

At the cube $L_1 = L_2 = L_3 = 1$, the direct cutoff sequence at quadratic extrapolation order gives $\rho = -0.837535893$, while an independent heat-kernel/Mellin evaluation gives $\rho = -0.837536911$. Their relative discrepancy is 1.22×10^{-6}. We therefore quote

$$\rho = -0.837537, \qquad p_i = \rho/3 = -0.279179 \tag{3}$$




FIG. 1. Rectangular flat T^3 fundamental domain with independent edge lengths L_p and L_t (biaxial case $L_2 = L_3 = L_t$). This is the geometry of the lattice sum, not a Planck-safe cubic cosmology and not an embedded solid torus.

TABLE I. Cubic Stage-1 diagnostics (lattice units, $L_i = 1$). Extrapolated ρ uses a polynomial retained through $1/N^2$, including a fitted $1/N$ term; validation residuals are at mode cutoff $N = 120$.

Quantity	Value
ρ (direct, order 2)	-0.837535893
ρ (heat kernel/Mellin)	-0.837536911
p_i (quoted cube value)	-0.279179
max FD pressure rel. error	4.8×10^{-10}
conformal-trace residual	$\sim 10^{-15}$
cube isotropy error	$\sim 10^{-16}$

in lattice units rather than retaining unsupported extra digits. The direct run has conformal-trace residual $|\rho - \sum_i p_i| \sim 10^{-15}$ at $N = 120$ modes per edge, finite-difference pressure errors $\sim 5 \times 10^{-10}$, and cubic isotropy at machine precision (Table I). The Stage-1 package reports `STATUS: PASS`.

C. Independent noncubic benchmark and sensitivity

The heat-kernel/Mellin implementation does not call the direct lattice-sum routine. It evaluates the Epstein sum through a Jacobi-theta product and obtains pressures by differentiating the total energy. Table II compares it with the registered direct extrapolation. The maximum normalized pressure discrepancy over these points is 7.42×10^{-7}, while relative trace residuals are below 5.7×10^{-12}. This verifies the nonzero-mode topological stress, not the state-dependent isotropic zero mode retained in a full compact-space treatment [5].

TABLE II. Independent heat-kernel/Mellin benchmark. The final column is the relative energy-density discrepancy.

(L_1, L_2, L_3)	$\rho_{\rm HK}$	$\rho_{\rm dir}$	rel.
$(1, 1, 1)$	-0.8375369	-0.8375359	1.22×10^{-6}
$(1, 1.25, 1.25)$	-0.4788673	-0.4788668	1.20×10^{-6}
$(1, 1, 2)$	-0.4362267	-0.4362262	1.08×10^{-6}
$(0.8, 1.1, 1.4)$	-0.7519731	-0.7519723	1.13×10^{-6}

A preregistered release-sensitivity suite varied the cube extrapolation order and fitting window and re-ran Stage 3B at twice the temporal sampling with rtol $= 10^{-12}$ and atol $= 10^{-16}$. All four static precision checks passed. All three registered dynamic shapes retained the `TRANSIENT_CROSSING` classification and no attractor appeared. One threshold amplitude shifted by 8.74×10^{-5} relative, however, exceeding the deliberately stringent preregistered 10^{-8} bound; the suite therefore records `FAIL` (18/19), not a rewritten pass. The qualitative no-attractor conclusion is stable, but threshold amplitudes are not quoted as precision predictions.

D. Stage-2 biaxial shape scan

Let $r = L_t/L_p$. CBR-001 Stage 2 holds $L_p = 1$ while varying $L_t = rL_p$; it does not hold the mean scale fixed. Stage 3 converts this raw table to fixed mean scale using the homogeneous $r^{8/3}$ stress rescaling before constructing its source. The stress $\rho(r)$, $p_p(r)$, $p_t(r)$ and the anisotropy $\Delta p \equiv p_t - p_p$ are smooth functions of shape; Δp changes sign across the cube $r = 1$ (Fig. 2). Representative points (lattice units) include

$$r = 1: \qquad \rho = -0.8375, \qquad \Delta p \simeq 0, \tag{4}$$

$$r \simeq 1.26: \qquad \rho = -0.4702, \qquad \Delta p \simeq +0.247, \tag{5}$$

where the second line is in the raw fixed-L_p Stage-2 convention. Compact *shape-dependent* anisotropic stress is therefore a concrete control of the implemented static formula. It is *not* a universal cycle-counting coefficient independent of the mode sum.

III. BIAXIAL INSTANTANEOUS-STRESS CLOSURE

A. Geometry and equations

For the biaxial Bianchi-I line element

$$\mathrm{d}s^2 = -\mathrm{d}t^2 + a_p^2\,\mathrm{d}x^2 + a_t^2(\mathrm{d}y^2 + \mathrm{d}z^2), \tag{6}$$

define

$$H = \frac{H_p + 2H_t}{3}, \qquad \delta = H_t - H_p, \qquad r = \frac{a_t}{a_p}. \tag{7}$$




FIG. 2. Stage-2 biaxial anisotropy $\Delta p = p_t - p_p$ versus shape ratio $r = L_t/L_p$. The cubic point $r = 1$ is isotropic; anisotropy is shape moduli dependent.

The shear system is the biaxial specialization of Bianchi-I [6], with $\kappa = 8\pi G$,

$$\dot{\delta} + 3H\delta = \kappa(p_t - p_p), \qquad \dot{r} = r\delta, \tag{8}$$

and Hamiltonian constraint

$$3H^2 - \frac{\delta^2}{3} = \kappa\rho_{\text{total}}. \tag{9}$$

Stage 3A treats the verified Casimir tensor as a small radiation-like perturbation ($\propto a^{-4}$) on a positive de Sitter background and integrates in $N = \ln a$. It does not evolve quantum-field mode functions on the time-dependent Bianchi-I geometry or renormalize $\langle T_{\mu\nu}\rangle$ there. The dimensionless source-scale ratio ϵ absorbs $\hbar c/L_\star^4$ relative to the de Sitter density scale (CBR-001 Stage 3A convention). No present ITSM derivation fixes this ratio. It validates the engine (perturbative limit, constraint and continuity residuals) but does *not* by itself establish any preferred Hubble ratio.

B. The Stage-3B ratio test

Let

$$x = \frac{\delta}{H}, \qquad q = \frac{H_t}{H_p} = \frac{1 + x/3}{1 - 2x/3}. \tag{10}$$

The historical target $q = 13/12$ corresponds to $x_\star = 3/38$. The small-source fixed-background profile $x(N) = S(e^{-3N} - e^{-4N})$ peaks at $N = \ln(4/3)$ with maximum $(27/256)S$; merely reaching $x_\star$ already requires $S \geq 128/171 \simeq 0.7485$, outside the strict small-perturbation regime.

TABLE III. Stage-3B classification at threshold amplitudes for the 13/12 search. "Regime" labels the peak Casimir energy fraction (perturbative / marginal / nonperturbative).

r_0	class	$\epsilon_\star$	regime
1.01	no crossing	—	—
1.05	no crossing	—	—
1.10	transient	2.55	nonpert.
1.25	transient	1.34	nonpert.
1.50	transient	0.676	nonpert.
2.00	transient	0.278	nonpert.
3.00	transient	0.089	marginal
4.00	invalid	—	—




FIG. 3. Representative Stage-3B trajectories of $q = H_t/H_p$ versus N. The horizontal line marks 13/12. Passages, when present, are brief; late evolution returns to isotropy.

A nonlinear root search over initial shapes $r_0 \in \{1.01, 1.05, 1.1, 1.25, 1.5, 2, 3, 4\}$ and source-scale ratios spanning seven decades finds target-reaching thresholds for five of eight shapes. Classification vocabulary:

- `NO_CROSSING`: peak q stays below 13/12;
- `TRANSIENT_CROSSING`: touch or brief crossing;
- `QUASI_PLATEAU`: dwell ≥ 1 e-fold within 1% without attractor criteria;
- `ATTRACTOR`: dwell ≥ 5 e-folds within 1%, nearby trajectories converge, and no subsequent decay toward $q = 1$.

Results: zero attractors, zero quasi-plateaus, five transient crossings, two no-crossings, and one trajectory initialized at the upper interpolation boundary whose outward drive leaves the supported domain (Table III; Fig. 3). Dwell times within 1% of the target are only $\sim$ 0.27–0.30 e-folds. Every valid candidate returns to $q \to 1$ by $N = 20$, consistent with the declared a^{-4} source becoming negligible. The tested nearby trajectories also decay toward isotropy; none satisfy the registered attractor criteria.




FIG. 4. Stage-3B threshold amplitudes $\epsilon_\star(r_0)$ for first touch of $q = 13/12$. Near-cubic shapes require large source-scale ratios; no threshold defines an attractor.

C. Kinematic warning

Because $\mathrm{d}\ln r/\mathrm{d}N = x$, a constant $q \neq 1$ requires continuous evolution of the shape ratio; it is not a fixed-torus property. A persistent anisotropic state would have to be a *driven* solution with an additional constitutive stress, not a free Casimir attractor. For a prescribed $x(N)$, the required anisotropic pressure is

$$\Pi_{\mathrm{req}} = \frac{H^2}{\kappa}\Big[\frac{\mathrm{d}x}{\mathrm{d}N} + x\big(3 + \tfrac{\mathrm{d}\ln H}{\mathrm{d}N}\big)\Big]. \tag{11}$$

In de Sitter at fixed $x = 3/38$ this is $\Pi_{\mathrm{req}} = 9H^2/(38\kappa)$. The free Casimir anisotropy scales as

$$\Pi_{\mathrm{Cas}} \propto a^{-4}\, r^{8/3}\big[\widehat{p}_t(r) - \widehat{p}_p(r)\big] \tag{12}$$

and, under the stated positive-H, bounded-shape instantaneous closure, cannot maintain Π_{req} once the source redshifts away. Any future positive claim of a persistent 13/12-class ratio must derive that extra stress (open programme CBR-002) without inserting the target into a coefficient by hand.

IV. HISTORICAL PACKAGING (EXPLICITLY NOT RESTORED)

Cycle counting that assigns a fixed fraction 1/12 per compact direction and promotes $H_t/H_p = 13/12$ to a free-field prediction, together with the numerical map $H_p = 67.36\,\mathrm{km\,s^{-1}\,Mpc^{-1}} \Rightarrow H_t = 72.97\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$, is a packaging of the *question* this paper tests—not a result we endorse. The lattice stress is shape dependent; free-field backreaction does not lock the ratio; and CMB topology bounds on cubic E_1 cells [7, 8] are a separate observational sector that this rectangular free-field simulation does not claim to satisfy.

We likewise do not claim that rectangular Casimir stress derives a galactic acceleration scale, fixes weak-field matching coefficients, or resolves the Hubble tension [9, 10].

V. LIMITATIONS AND OPEN ROUTES

- Free massless scalar only; interacting or multi-field sectors untested.
- The renormalized static lattice expression is supplied as input. A local independent heat-kernel/Mellin evaluation now agrees at four points, but an external replication and a full-domain error analysis remain outstanding.
- The time-dependent calculation is an instantaneous a^{-4} closure, not a mode-function evolution or curved-spacetime renormalization of $\langle T_{\mu\nu}\rangle$. Particle production, curvature counterterms, trace-anomaly and memory terms are not calculated.
- Rectangular moduli; twisted flat spaces (E_2, E_3) not scanned.
- Positive de Sitter testbed with Casimir as a small perturbation—not a Casimir-dominated universe.
- The preregistered sensitivity suite is complete but not all-pass: one threshold-amplitude stability check missed its 10^{-8} bound, although all classifications and the no-attractor conclusion were unchanged.
- Persistent anisotropy requires a derived driven or memory stress (CBR-002), not free-field packaging.
- CMB compatibility of any particular (L_i) is not established here.

These limitations are research boundaries, not research bans: driven constitutive stress, alternate topologies, and richer field content remain open under an honest claim ladder.

VI. CONCLUSIONS

A free massless scalar on rectangular flat T^3 yields an internally checked and independently benchmarked nonzero-mode, shape-dependent anisotropic Casimir stress. When that static stress is transported as an instantaneous a^{-4} source on biaxial de Sitter expansion, the ratio $H_t/H_p = 13/12$ appears at most as a brief transient at large source-scale ratios, never as a quasi-plateau or attractor in the bounded search reported here. The tested closure therefore cannot sustain the historically proposed anisotropic expansion or justify a map to $H_t = 72.97\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$. Public code and stage summaries under `Analysis/Casimir/CBR-001/` define the

Stage-1–3B pipeline. The current release candidate remains on publication hold pending deferred external review and resolution or explicit editorial acceptance of the declared dynamical-state limitations. Portable SHA-256 anchors for the repaired candidate are recorded in `CBR001_CHECKSUMS.md`; these anchors identify artifacts and do not constitute a physics pass.

CODE AND DATA AVAILABILITY

The complete Stage-1–3B implementation, frozen CSV/JSON outputs, independent literature-control receipt, and preregistered sensitivity contract and receipt are available in the public companion repository at `https://github.com/brendohxd/ITSM-Integrated-Toroidal-Syntropic-Model`. Hashes identify exact artifacts but do not constitute a physics pass.

DECLARATION ON THE USE OF ARTIFICIAL-INTELLIGENCE TOOLS

Artificial-intelligence-assisted tools were used for editorial drafting, source-code assistance, and checks of symbolic and numerical consistency. The author independently reviewed the manuscript, equations, cited sources, and generated artifacts and accepts full responsibility for the content.

ACKNOWLEDGMENTS

Computations use the repository environment `itsm_env` and the CBR-001 gate package. Claim hygiene for related geometric packaging is recorded in companion note [4].

Appendix A: Reproducibility anchors

Table IV lists SHA-256 prefixes for portable CBR-001 CSV/JSON artifacts underlying the Stage-1–3B numbers in the main text (branch `recovery/v12-core-architecture`). CSV writers use a fixed LF record terminator and JSON output paths are relative to the summary directory. Re-running the Stage scripts after intentional code changes will alter digests; any published revision must update this table.

Full digests and reproduction commands appear in the companion `CBR001_CHECKSUMS.md` ledger.

[1] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, 1982).
[2] M. Lachièze-Rey and J.-P. Luminet, Cosmic topology, Physics Reports **254**, 135 (1995).
[3] A. Edery, Multidimensional cut-off technique, odd-dimensional Epstein zeta functions and Casimir energy of massless scalar fields, Journal of Physics A: Mathematical and General **39**, 685 (2006).
[4] B. Boyd, Present-epoch scale matching, the weak-field invariant $c_{\rm obs}$, and no-go results for a cubic t^3 geometric story (2026), companion reconstruction note; working draft under papers/P1-Scale-Matching-Reconstruction/.
[5] A. Negro *et al.*, Quantum signatures of cosmic topology: How casimir backreaction transmits isotropy violation, arXiv:2603.12319 (2026), arXiv:2603.12319.
[6] G. F. R. Ellis and M. A. H. MacCallum, A class of homogeneous cosmological models, Communications in Mathematical Physics **12**, 108 (1969).
[7] Planck Collaboration, Planck 2015 results. XVIII. background geometry and topology of the Universe, Astronomy & Astrophysics **594**, A18 (2016).
[8] Y. Akrami *et al.*, Promise of future searches for cosmic topology, Physical Review Letters **132**, 171501 (2024).
[9] Planck Collaboration, Planck 2018 results. VI. cosmological parameters, Astronomy & Astrophysics **641**, A6 (2020).
[10] A. G. Riess *et al.*, A comprehensive measurement of the local value of the Hubble constant with 1 km s^{-1} Mpc^{-1} uncertainty from the Hubble space telescope and the SH0ES team, Astrophysical Journal Letters **934**, L7 (2022).

TABLE IV. SHA-256 of CBR-001 artifacts used in this draft.

Artifact	SHA-256 (prefix)
`cbr001_stage1.csv`	`57847F382BAF...`
`cbr001_stage2_scan.csv`	`FE648629545F...`
`cbr001_stage3b_thresholds.csv`	`94B0AE4C6381...`
`cbr001_stage3b_summary.json`	`9A5E80D2F713...`