

Problem Set 8

Complex Analysis

Bennett Rennier
bennett@brennier.com

December 7, 2018

Ex 1

- a) Let U be a simply-connected domain and $f : U \rightarrow \mathbb{C} \setminus \{0\}$ be analytic. Let $g : U \rightarrow \mathbb{C}$ be an analytic function with $e^g = f$. Let $z, w \in U$ with $f(z) = f(w)$. Let $\gamma : [a, b] \rightarrow U$ be a piecewise C^1 curve with $\gamma(a) = z$, $\gamma(b) = w$. Show that $g(z) - g(w)$ is the $2\pi i$ times the winding number of $f \circ \gamma$ about 0.
- b) Let V be a simply-connected domain, $a_1, \dots, a_k \in V$ distinct points, and set $U = V \setminus \{a_1, \dots, a_k\}$. Let $\delta > 0$ be small enough so that $\overline{B_\delta(a_j)}$ are disjoint. For $j = 1, \dots, k$, let γ_j be $\partial B_\delta(a_j)$ traversed once counterclockwise. Given an analytic $f : U \rightarrow \mathbb{C} \setminus \{0\}$, show that f has a logarithm if and only if $n(f \circ \gamma_j, 0) = 0$ for $j = 1, \dots, k$. [Hint: It suffices to show that for any cycle Γ in U , we have that $\int_\Gamma \frac{f'}{f} dz = 0$. Reduce to the case of a general cycle Γ to a cycle which is the sum of the γ_j .]

Proof.

- a) We see that

$$n(f \circ \gamma; 0) = \frac{1}{2\pi i} \int_{f \circ \gamma} \frac{1}{s} ds = \frac{1}{2\pi i} \int_0^1 \frac{f'(\gamma(s))\gamma'(s)}{f(\gamma(s))} ds = \frac{1}{2\pi i} \int_\gamma \frac{f'(s)}{f(s)} ds = \frac{1}{2\pi i} (g(z) - g(w))$$

which proves that $g(z) - g(w)$ is $2\pi i$ times $n(f \circ \gamma; 0)$ as desired.

- b) Suppose f has a logarithm, that is, a holomorphic function g such that $e^g = f$. Since each γ_j is a closed curve, we can apply part (a) to get that

$$n(f \circ \gamma_j; 0) = \frac{1}{2\pi i} (g(\gamma_j(0)) - g(\gamma_j(1))) = 0$$

for each γ_j . Now conversely, suppose that $n(f \circ \gamma_j; 0) = 0$ for $j = 1, \dots, k$ and let γ be an arbitrary cycle in U . If we let $\Gamma = \gamma - \sum_i n(\gamma; a_i) \gamma_i$, we see that

$$n(\Gamma; a_j) = n(\gamma; a_j) - \sum_i n(\gamma; a_j) n(\gamma_i; a_j) = n(\gamma; a_j) - \sum_i n(\gamma; a_j) \delta_{ij} = n(\gamma; a_j) - n(\gamma; a_j) = 0.$$

Since Γ is contained in U and its winding number around each a_j is zero, this means that Γ

is homologous to zero on U . As f is never zero, we get by the Cauchy Integral Theorem that

$$\begin{aligned} 0 &= \int_{\Gamma} \frac{f'}{f} dz = \int_{\gamma} \frac{f'}{f} dz - \sum_i n(\gamma; a_i) \int_{\gamma_i} \frac{f'}{f} dz \\ &= \int_{\gamma} \frac{f'}{f} dz - \sum_i n(\gamma; a_i) \int_{f \circ \gamma_i} \frac{1}{s} ds \\ &= \int_{\gamma} \frac{f'}{f} dz - \sum_i n(\gamma; a_j) n(f \circ \gamma; 0) = \int_{\gamma} \frac{f'}{f} dz. \end{aligned}$$

Since $\int_{\gamma} \frac{f'}{f} dz = 0$ for any cycle γ in U , path integrals in U are path-independent. After fixing some $a \in U$, this allows use to define the logarithm

$$g(z) = \int_{\gamma_{a,z}} \frac{f'(s)}{f(s)} ds$$

where $\gamma_{a,z}$ is any path from a to z . □

Ex 2

- a) Suppose that $U, V \subseteq \mathbb{C}$ are open and that $f : U \rightarrow V$ is a conformal map. Let z_n be a sequence in U . Show that $z_n \rightarrow \partial U$ if and only if $f(z_n) \rightarrow \partial V$.
- b) Let $A = \{z \in \mathbb{C} : 0 < |z| < 1\}$ and $B = \{z \in \mathbb{C} : 1 < |z| < 2\}$. Show that there is no conformal map $f : A \rightarrow B$.

Proof.

- a) Let $z_n \rightarrow \partial U$; that is z_n eventually escapes any compact subset of U . Let $K \subseteq V$ be a compact set. Since f^{-1} is continuous, we have that $f^{-1}(K)$ is a compact subset of U . Thus, for some N , we have that for all $n \geq N$, $z_n \notin f^{-1}(K)$. But this means that for all $n \geq N$, $f(z_n) \notin K$. Since K was an arbitrary compact set, we have that $f(z_n) \rightarrow \partial V$.

The converse is proved similarly. Suppose that $f(z_n) \rightarrow \partial V$. Let $K \subseteq U$ be a compact set. As f is continuous, we know that $f(K)$ is a compact subset of V . Thus, for some N , for all $n \geq N$, $f(z_n) \notin f(K)$. This means that for all $n \geq N$, we have that $z_n \notin K$. Since K was an arbitrary compact set, we have that $z_n \rightarrow \partial U$.

- b) Suppose that $f : A \rightarrow B$ was a conformal map. Since $\text{Im}(f) = B$, we know that f is bounded. In particular, f is bounded near 0, which means that 0 is a removable singularity of f . This proves that we can extend f holomorphically to a function $\tilde{f} : \mathbb{D} \rightarrow B$. By part (a), if $z_n \rightarrow 0 \in \partial A$, then $f(z_n) \rightarrow \partial B$. This proves that $\tilde{f}(0) \in \partial B$. However, this means that $\text{Im}(\tilde{f}) = B \cup \{\tilde{f}(0)\}$, where $\tilde{f}(0) \in \partial B$. However, this image cannot be open, contradicting the open mapping theorem. Thus, there can be no conformal map $f : A \rightarrow B$. □

Ex 3 Suppose that f is entire and satisfies $\lim_{z \rightarrow \infty} f(z) = \infty$. Show that f is a polynomial.

Proof. Let $g : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ be given by $g(z) = f(1/z)$. We see that if g had an essential singularity at 0, then by Casorati–Weierstrass, for any $z \in \mathbb{C}$ there is a sequence $z_n \rightarrow 0$ such that $g(z_n) \rightarrow z$. However, we see that $\lim_{z \rightarrow 0} g(z) = \lim_{z \rightarrow \infty} f(z) = \infty$, contradicting Casorati–Weierstrass. Thus, g does not have an essential singularity at 0, meaning $g(z) = z^{-n}h(z)$ for some holomorphic function

h . Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ and $h(z) = \sum_{k=0}^{\infty} b_k z^k$ be the analytic representations of f and h . This gives us that

$$\sum_{k=-\infty}^0 a_{-k} x^k = \sum_{k=0}^{\infty} a_k x^{-k} = f(1/z) = g(z) = \frac{h(z)}{z^n} = \sum_{k=0}^{\infty} b_k x^{k-n} = \sum_{k=-n}^{\infty} b_{k+n} x^k.$$

By Exercise 5, Laurant series are unique, so it must be that $a_{-k} = b_{k+n}$ for all k . In particular, this means that $a_k = 0$ for all $k \geq n$. Thus, $f(z) = \sum_{k=0}^n a_k z^k$, proving that f is a polynomial. \square

Ex 4 Suppose that $f : \mathbb{C} \rightarrow \mathbb{C}$ is injective and entire. Define $g : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ by $g(z) = f(1/z)$.

- a) Show that the singularity of g at 0 is not an essential singularity.
- b) Show that $f(z) = az + b$ for some $a, b \in \mathbb{C}$.
- c) Show that if $h : \mathbb{C}_{\infty} \rightarrow \mathbb{C}_{\infty}$ is injective and analytic, then h is a linear fractional transformation.

Proof.

- a) Let D be the punctured disk and let E be the set $\{z : |z| > 1\}$. Suppose that 0 was an essential singularity of g . By Cassorati-Weierstrass, we have that $g(D)$ is dense in \mathbb{C} . This means that $f(1/D) = f(E)$ is dense in \mathbb{C} . But $f(D)$ is open by the Open Mapping Theorem, so it must be that $f(D) \cap f(E) \neq \emptyset$ contradicting the injectivity of f .
- b) By part (a), we know that 0 is either a removable singularity or a pole. If 0 were a removable singularity, then we could extend g to be defined at 0, but this is equivalent to extending f at infinity to be some complex number. But this implies that f is bounded around infinity, meaning f must be constant by Liouville's Theorem, contradicting injectivity of f . Thus, 0 must be a pole of f . This means that $\lim_{z \rightarrow \infty} f(z) = \lim_{z \rightarrow 0} g(z) = \infty$. By Ex 3, this implies that f is a polynomial. But polynomials of degree greater than 2 are not injective, so it must be that $f(z) = az + b$.
- c) If $\infty \notin \text{Im}(f)$, then $f|_{\mathbb{C}}$ is an entire function such that $\lim_{z \rightarrow \infty} f(z) = c \in \mathbb{C}$. This implies that f is bounded around infinity, meaning f must be constant by Liouville's Theorem, contradicting the injectivity of f .

If $h(\infty) = \infty$, then by injectivity, $h|_{\mathbb{C}} : \mathbb{C} \rightarrow \mathbb{C}$ is an injective, entire function such that $\lim_{z \rightarrow \infty} h|_{\mathbb{C}} = h(\infty) = \infty$. By part (b), it must be that $h|_{\mathbb{C}}(z) = az + b$. By continuity, we can conclude that $h(z) = az + b$.

Now suppose that $h(c) = \infty$ where $c \neq \infty$. Let $\phi(z) = \frac{cz}{z-c}$ which is a mobius transformation such that $c \mapsto \infty$ and $\infty \mapsto c$. If we then define $g = h \circ \phi$, then g is injective (as its the composition of injective functions) and $g(\infty) = h(\phi(\infty)) = h(c) = \infty$. By the previous paragraph, this means that $g(z) = az + b$. Since the inverse of ϕ is $\phi^{-1}(z) = \frac{-cz}{z-c}$, we get that

$$h(z) = (h \circ \phi \circ \phi^{-1})(z) = (g \circ \phi^{-1})(z) = g\left(\frac{-cz}{z-c}\right) = a \frac{-cz}{z-c} + b = \frac{(b-ac)z - bc}{z-c}$$

which shows that h is a linear fractional as desired. \square

Ex 5 Let $r, R \in [0, +\infty]$ and let $A_{r,R} = \{z \in \mathbb{C} : r < |z| < R\}$. Suppose that (a_n) and (b_n) are sequences of complex numbers such that

$$\sum_{n=-\infty}^{\infty} a_n z^n = \sum_{n=-\infty}^{\infty} b_n z^n \quad \text{for all } z \in A_{r,R}.$$

Prove that $a_n = b_n$ for all n .

Proof. Combining the terms to one side we get that

$$\sum_{n=-\infty}^{\infty} (a_n - b_n) z^n = 0 \quad \text{for all } z \in A_{r,R}.$$

Since this Laurent series converges to the holomorphic function 0 on $A_{r,R}$, we can use the Cauchy Integral Formula to find the coefficients of this power series as

$$a_n - b_n = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z^{n+1}} dz = \frac{1}{2\pi i} \int_{\gamma} \frac{0}{z^{n+1}} dz = 0$$

where γ is a path in $A_{r,R}$ that traverses around 0 once counterclockwise. This proves that $a_n = b_n$ for all n . \square

Ex 6

- a) For each of the following functions f has an isolated singularity at $z = 0$. Determine the type of singularity.

i) $f(z) = \frac{\cos(z)}{z}$

ii) $f(z) = \frac{\log(z+1)}{z^2}$

iii) $f(z) = z \sin(1/z)$

iv) $f(z) = \frac{1}{1-e^z}$

- b) Let $f(z) = \frac{1}{z(z-1)(z-2)}$. Give the Laurent expansion of f in each of the following annuli: $A_{0,1}$, $A_{1,2}$, $A_{2,\infty}$.

- c) Show that $f(z) = \tan(z)$ is analytic in \mathbb{C} except for simple poles at $z = \frac{\pi}{2} + n\pi$ for each integer n . Determine the singular part of f at each of these poles.

Proof.

- a) i) Using the Taylor series for cosine, we get that

$$f(z) = \cos(z)/z = \sum_{k=0}^{\infty} \frac{(-1)^k z^{2k}}{(2k)!} \cdot \frac{1}{z} = \sum_{k=0}^{\infty} \frac{(-1)^k z^{2k-1}}{(2k)!}.$$

We see then that f has a pole of order one at 0 and that the singular part of f is z^{-1} .

- ii) Using the Taylor series for log, we get that

$$f(z) = \frac{\log z + 1}{z^2} = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} z^k}{k z^2} = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} z^{k-2}}{k} = \sum_{k=0}^{\infty} \frac{(-1)^k z^{k-1}}{k+1}.$$

We see then that f has a pole of order one at 0 and that the singular part of f is z^{-1} .

iii) Using the Taylor series for \sin , we get that

$$f(z) = \sin(1/z) \cdot z = \sum_{k=0}^{\infty} \frac{(-1)^k z^{-(2k+1)}}{(2k+1)!} \cdot z = \sum_{k=0}^{\infty} \frac{(-1)^k z^{-2k}}{(2k+1)!}.$$

Since this Laurent series has infinitely many non-zero terms in its singular part, we get that the singularity of f at 0 is an essential singularity.

iv) Since $1 - e^z = \sum_{n=1}^{\infty} -\frac{z^n}{n!}$ has a zero of order 1 at $z = 0$, we know that f must have a pole of order 1 at $z = 0$. Since this pole is simple, we can calculate the residue as

$$\text{res}_0(f) = \lim_{z \rightarrow 0} z f(z) = \lim_{z \rightarrow 0} \frac{z}{1 - e^z} = \lim_{z \rightarrow 0} -\frac{1}{e^z} = -1.$$

This means the singular part of f is $-x^{-1}$.

b) First, we use partial fractions to obtain that

$$\frac{1}{z(z-1)(z-2)} = \frac{1/2}{z} + \frac{-1}{z-1} + \frac{1/2}{z-2} = \frac{1/2}{z} + \frac{1}{1-z} - \frac{1/2}{2-z}$$

Now we see that for $0 < |z| < 1$

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n$$

and for $0 < |z| < 2$ we have that

$$-\frac{1/2}{2-z} = -\frac{1}{4} \left(\frac{1}{1-z/2} \right) = -\frac{1}{4} \sum_{n=0}^{\infty} \frac{z^n}{2^n} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+2}}.$$

This means the Laurent series for $A_{0,1}$ is

$$f(z) = \frac{1/2}{z} + \sum_{n=0}^{\infty} z^n - \sum_{n=0}^{\infty} \frac{z^n}{2^{n+2}} = \frac{1/2}{z} + \sum_{n=0}^{\infty} \left(1 - \frac{1}{2^{n+2}} \right) z^n = \sum_{n=-1}^{\infty} \left(1 - \frac{1}{2^{n+2}} \right) z^n.$$

Now for $|z| > 1$, we have that

$$\frac{1}{1-z} = -\frac{1}{z-1} = -\frac{1}{z} \frac{1}{(1-1/z)} = -\frac{1}{z} \sum_{n=0}^{\infty} z^{-n} = -\frac{1}{z} \sum_{n=-\infty}^0 z^n = -\sum_{n=-\infty}^0 z^{n-1} = -\sum_{n=-\infty}^{-1} z^n$$

This means the Laurent series for $A_{1,2}$ is

$$f(z) = \frac{1/2}{z} - \sum_{n=-\infty}^{-1} z^n - \sum_{n=0}^{\infty} \frac{z^n}{2^{n+2}}.$$

And finally, for $|z| > 2$, we have that

$$-\frac{1/2}{2-z} = \frac{1}{2z} \cdot \frac{1}{1-2/z} = \frac{1}{2z} \sum_{n=0}^{\infty} 2^n z^{-n} = \frac{1}{2z} \sum_{n=-\infty}^0 2^{-n} z^n = \sum_{n=-\infty}^0 2^{-n-1} z^{n-1} = \sum_{n=-\infty}^{-1} 2^{-n-2} z^n.$$

This means the Laurent series for $A_{2,\infty}$ is

$$f(z) = \frac{1/2}{z} - \sum_{n=-\infty}^{-1} z^n - \sum_{n=-\infty}^{-1} 2^{-n-2} z^n = \frac{1/2}{z} - \sum_{n=-\infty}^{-1} \left(1 - \frac{1}{2^{n+2}} \right) z^n.$$

- c) The zeros of $\cos(z)$ are the points $z = \frac{\pi}{2} + n\pi$ for each integer n . Since $\frac{d}{dz} \cos(z) = -\sin(z)$ and the zeros of sine (which are $n\pi$ for each integer n) are distinct from our original zeros, it must be that the zeros of $\cos(z)$ are of order 1. Again, as the zeros of sine are distinct from cosine, they don't cancel out in $f(z) = \frac{\sin(z)}{\cos(z)}$. This implies that the poles of $\tan(z)$ are at the points $\frac{\pi}{2} + n\pi$ and that these poles are simple. Since the poles are simple, we can calculate the residue as

$$\text{res}_{\frac{\pi}{2}+n\pi}(f) = \lim_{z \rightarrow \frac{\pi}{2}+n\pi} (z - \frac{\pi}{2} - n\pi) \frac{\sin(z)}{\cos(z)} = \lim_{z \rightarrow \frac{\pi}{2}+n\pi} \frac{\sin(z) + (z - \frac{\pi}{2} - n\pi) \cos(z)}{-\sin(z)} = -1.$$

This means that the singular part of each of these poles is simple $-x^{-1}$. □