A gap is not a graviton: strong-coupling exclusions for lattice gravity in first-order form

Tristan Romanov[1] and Brent Hartshorn

[1] *Independent Researcher, San Francisco, CA, USA*[*]

(Dated: Draft, 31 October 2026)

In Yang–Mills theory a mass gap is the goal; in gravity it is a diagnosis. We prove, in Lean 4 with Mathlib, that a family of reflection-positive lattice theories whose transfer-matrix gap is bounded below in physical units along a continuum tuning has no continuum limit with a massless mode, and that a massless mode makes the two-point function decay more slowly than any exponential. Combined with the strong-coupling theorems of lattice Yang–Mills for every compact simple group, this excludes the strong-coupling phase of Euclidean lattice gravity written as a gauge theory of the Lorentz group $SO(4)$ (for $\beta_{\mathrm{W}} < 4/29$) or of the de Sitter group $SO(5)$ (for $\beta_{\mathrm{W}} < 15/58$, without compensator) from being gravitational: these phases have a unique state, a positive transfer matrix (so no ghost states), and a gap. We also prove in Lean that an antiscreening gravitational correction to the Yang–Mills β-function preserves asymptotic freedom while a screening one produces an interacting ultraviolet fixed point; that on the 600-cell a gauge-fixing forest can lower the single-link Dobrushin partner count only from 10 to 9, so $\beta_{\mathrm{W}} < 4/9$ is optimal there; and we give two exactly solvable checks: two-dimensional Yang–Mills on a Brownian disk, where the area law becomes $d_F(1 + gL\sqrt{C_F})e^{-gL\sqrt{C_F}}$ in the quantum boundary length L, and the two-dimensional causal-triangulation transfer matrix, whose spectrum is geometric with a lattice gap of order the spacing. Nothing here identifies a gravitational phase or constructs a continuum theory.

I. INTRODUCTION

Four-dimensional lattice Yang–Mills theory has a unique state and a transfer-matrix gap at strong coupling for every compact simple gauge group [1], and the gap problem in physical units is reduced there to a single cube estimate. Gravity is the next force to add [2]. The present note starts from one observation: the graviton is massless, so in any approach where gravity is a lattice theory with a transfer matrix, a gap in physical units is a proof that the phase is *not* gravitational. The strong-coupling machinery of [1] produces such gaps at explicit couplings, and some lattice formulations of gravity are gauge theories of compact groups to which it applies as stated.

The formulations are first-order (Palatini) gravity, whose connection is an $SO(4)$ gauge field in Euclidean signature, and MacDowell–Mansouri gravity [3], an $SO(5)$ gauge field broken to $SO(4)$ by a compensating vector [4]. Monte Carlo simulations of such models find a vanishing tetrad at strong coupling [5], and causal and Euclidean dynamical triangulations have non-geometric phases [6]; none of this is proved. The pure Lorentz sector of a Wilson plaquette action generates $R_{\mu\nu}R^{\mu\nu}$ [7, 8], the curvature-squared sector of quadratic gravity, whose spin-2 ghost is conjectured to be confined at strong coupling [9, 10].

All Lean statements are in the repository `romanov360/quantum-gravity`, directory `QuantumGravity/`, built against Mathlib (Lean v4.33.0), without `sorry`, with the standard axioms only; numerical checks are in `python/qg/`.

[*] romanov360@gmail.com

II. A PHYSICAL GAP EXCLUDES A MASSLESS MODE

Let ρ_a be the finite spectral measure, in lattice units, of the transfer matrix at spacing a for a spatially smeared gauge-invariant field, so that the temporal two-point function is $S_a(t) = \int e^{-Et/a}\, d\rho_a(E)$. A gap m in physical units along the tuning means $\rho_a((-\infty, ma)) = 0$.

Theorem 1 (`physicalGap_ratio_le`, `limit_ratio_of_physicalGap`)**.** *For $0 \le t_1 \le t_2$, $S_a(t_2) \le e^{-m(t_2-t_1)} S_a(t_1)$ for every a, and if $Z_a S_a(t_i) \to L_i$ along any filter for positive renormalisations Z_a, then $L_2 \le e^{-m(t_2-t_1)} L_1$.*

A massless particle in a reflection-positive theory gives the smeared field spectral weight at arbitrarily small energies: a packet contains momenta $|p| < \varepsilon$ and $E = |p|$. We call a finite measure σ on $[0, \infty)$ *massless* if $\sigma([0, \varepsilon)) > 0$ for every $\varepsilon > 0$.

Theorem 2 (`massless_not_expDecay`)**.** *If σ is massless, then for no $\Delta > 0$ and C does $\int e^{-Et} d\sigma(E) \le Ce^{-\Delta t}$ hold for all $t \ge 1$.*

Idea. $\int e^{-Et} d\sigma \ge \sigma([0, \Delta/2))\, e^{-\Delta t/2}$, and $Ce^{-\Delta t/2} \to 0$. □

Theorem 3 (`no_massless_limit_of_physicalGap`)**.** *If a family of lattice theories has a gap $m > 0$ in physical units along the tuning and its renormalised two-point functions converge at every $t > 0$ to $\int e^{-Et} d\sigma(E)$ for a finite measure σ on $[0, \infty)$, then σ is not massless.*

The other alternative, a gap bounded below in *lattice* units, is the no-go theorem of [1]: every renormalised two-point function then vanishes in the limit. A graviton therefore needs $\Delta_{\rm lat}(a)/a \to 0$.

III. FIRST-ORDER LATTICE GRAVITY AT STRONG COUPLING

The theorem of [1] on every compact simple group gives, in the forest gauge, a unique free-boundary state, exponential clustering and a transfer-matrix gap for $\beta_{\rm W} = N^2\beta < 2N\,\mathrm{Ric}/29$, where N is the dimension of the representation in the Wilson action and $\mathrm{Ric} = c/4$ when the Killing form is c times the trace form. The constants are quadratic-form identities on a Cartan subalgebra, proved for all Cartan elements in `GravityGroups.lean`:

group (rep.)	role	c	Ric	range of $\beta_{\rm W}$
$SU(2)$ (2)	self-dual Spin(4)	4	1	4/29
$SO(4)$ (4)	Lorentz, tetrad index	2	1/2	4/29
$SO(5)$ (5)	de Sitter (MM)	3	3/4	15/58
$Sp(2)$ (4)	de Sitter, spinor	6	3/2	12/29

$SO(4)$ is semisimple; its two factors have the same Ricci constant, which is all the Bakry–Émery step uses.

Corollary 4. *The Euclidean lattice Lorentz gauge theory with weight* $\exp(4\beta\sum_p \mathrm{Tr}_4 U_p)$, $U \in SO(4)$, *has for* $\beta_{\rm W} = 16\beta < 4/29$ *a unique infinite-volume state, exponential clustering and a transfer-matrix gap. Its transfer matrix is a positive operator (reflection positivity of the Wilson action), so its Hilbert space has no negative-norm states. If the gap persists in physical units along a tuning, the limit has no massless mode in any channel (Theorem 3).*

The centre element $-1 \in SO(4)$ acts as -1 on the vector representation, so vector Wilson loops have an area law at strong coupling [11]. The fundamental character coefficient (`lorentz_strong.py`) is $a_F(\beta_{\rm W}) = \beta_{\rm W}/16 + O(\beta_{\rm W}^3)$ (for $SU(2)$: $I_2(\beta_{\rm W})/I_1(\beta_{\rm W}) = \beta_{\rm W}/4 + \dots$), so the leading-order string tension at $\beta_{\rm W} = 4/29$ is $-\log a_F = 4.75$ in lattice units.

Remark. In the frozen-tetrad model the curvature-squared sector is therefore ghost-free and gapped at strong coupling; this is the fixed-spacing analogue of ghost confinement in quadratic gravity [9, 10]. Whether a continuum limit of this lattice model is quadratic gravity is not addressed.

For MacDowell–Mansouri gravity the plaquette-pair action and the compensator change the influence count; the programme is in `docs/G2-pregeometric-phase.md`.

IV. GRAVITY AND ASYMPTOTIC FREEDOM

Above the Planck scale the Yang–Mills coupling runs as $dg/dt = -bg^3 - fg$, with f a gravitational coefficient whose sign is scheme dependent but non-negative in symmetry-preserving computations [12]. In $u = 1/g^2$, $u(t) = (u_0 + b/f)e^{2ft} - b/f$.

Theorem 5 (`RunningWithGravity.lean`). *(i) u solves $du/dt = 2b + 2fu$. (ii) If $f > 0$, $b \ge 0$, $u_0 \ge 0$ then $u(t) \ge u_0 + 2bt$: the coupling is never larger than without gravity, and $g^2 \to 0$. (iii) If $f < 0 < b$, then $g^2 \to -f/b > 0$, an interacting ultraviolet fixed point. (iv) If $f > 0$, every $u_* \ge u_0$ is attained at $t_* = \log\frac{u_*+b/f}{u_0+b/f}/(2f) \ge 0$.*

Item (iv) is the statement of [1] that the intermediate coupling window is entered at a spacing-independent scale, with the gravitational correction included.

V. YANG–MILLS ON CURVED LATTICES

The single-link Dobrushin count of [1] is local: on any 2-complex with a gauge-fixing forest F, a free link e has $P(e) = \sum_{p\ni e}(\#\text{free links of } p-1)$ partners and $SU(2)$ is unique for $\beta_{\rm W} < 4/\max_e P(e)$. The search in `forest_partners.py` reproduces $P = 12$ on $\mathbb{Z}^4$ and $P = 7$ for the period-3 forest on $\mathbb{Z}^3$, and finds by exhaustion that period-2 forests on $\mathbb{Z}^3$ cannot go below 8.

Proposition 6 (`cell600_nine_optimal`). *On the 600-cell (120 vertices, 720 edges, every edge in 5 triangles) every forest leaves some free link with at least 9 partners, and a forest with* $\max P = 9$ *exists. The optimal single-link $SU(2)$ range is $\beta_{\rm W} < 4/9$.*

Proof. $P(e) = 10 - n(e)$, where $n(e)$ is the number of fixed links sharing a triangle with e. A forest has at most 119 links, each a neighbour of 10 links, so $\sum_{e\notin F} n(e) \le 1190 < 2\cdot 601$; some free link has $n(e) \le 1$. A greedy forest attains 9. □

For closed triangulations a forest has $V-1$ links among $E \approx 6V$, so gauge fixing helps little and the range is governed by the maximal edge degree; a uniform statement over dynamical triangulations needs a degree cap.

VI. TWO EXACTLY SOLVABLE CHECKS

a. YM_2 on a Brownian disk. Two-dimensional Yang–Mills depends on the surface only through its area [13, 14]; the free Brownian disk of perimeter L has area density $L^3(2\pi A^5)^{-1/2}e^{-L^2/2A}$ [15]. Hence $\mathbb{E}[e^{-sA}] = (1+L\sqrt{2s})e^{-L\sqrt{2s}}$ (an inverse-Gamma Laplace transform via $K_{3/2}$), and the Haar-averaged boundary Wilson loop $d_F e^{-g^2 A C_F/2}$ becomes

$$\langle\!\langle W_F \rangle\!\rangle_L = d_F\,(1 + gL\sqrt{C_F})\,e^{-gL\sqrt{C_F}},$$

a perimeter law in quantum length with tension $\sqrt{2\sigma_F}$. Since $\int Z(U,A)\,dU = 1$ for every A, quenched and annealed averages agree. On the sphere each representation shifts the cosmological constant by $g^2 C_\lambda/2$, and only the

trivial one is non-analytic at $\mu = 0$: the string susceptibility exponent $-1/2$ is unchanged. Monte Carlo checks are in `ym2_brownian.py`.

b. 2D causal triangulations. The one-step transfer matrix $M(l_1, l_2) = g^{l_1+l_2-1}\binom{l_1+l_2-1}{l_1-1}$ [16] has, numerically to ten digits, an exactly geometric spectrum $\lambda_{n+1}/\lambda_n = (1-s)/(1+s)$, $s = \sqrt{1-4g^2}$; with $g = \frac{1}{2}e^{-a^2\Lambda}$ the lattice gap is $2\,\mathrm{artanh}\, s \sim 2\sqrt{2\Lambda}\, a$. This is a solvable instance of the scaling that [1] requires of a theory with a continuum gap.

VII. COSMOLOGICAL BILLIARDS

Near a spacelike singularity, vacuum gravity becomes a billiard for the hyperbolic Kac–Moody algebra AE_3, and eleven-dimensional supergravity one for E_{10} [17]. `Billiards.lean` checks by `decide` the leading principal minors $2, 3, -2$ of AE_3 and $2, 3, \ldots, 10, -1$ of E_{10} (one sign change: Lorentzian signature), and `bkl_billiard.py` that the BKL era lengths are the continued-fraction digits of the initial u, with Gauss–Kuzmin statistics.

VIII. WHAT IS NOT CLAIMED

No result here identifies a gravitational phase, constructs a continuum limit, or concerns Lorentzian signature. The strong-coupling statements are at fixed lattice spacing; Theorem 3 needs the gap along a tuning, which is not proved for any of the models above. The open problems are listed in `OPEN.md`.

[1] D. Bhattacharjee, U. Bhattacharya, J. M. Lockwood, T. Romanov, *Explicit lattice Yang–Mills mass gap for every compact simple group reducing the continuum gap to one cube*, Zenodo 10.5281/zenodo.23444338 (2026), code at github.com/creelie/4Y-M.

[2] B. Hartshorn, *Quantum Gravity Lab*, github.com/brentharts/Quantum_Gravity_Lab (2026).

[3] S. W. MacDowell, F. Mansouri, Phys. Rev. Lett. **38**, 739 (1977).

[4] K. S. Stelle, P. C. West, Phys. Rev. D **21**, 1466 (1980).

[5] S. Catterall, D. Ferrante, arXiv:0912.5525.

[6] J. Ambjørn, R. Loll, arXiv:2401.09399.

[7] M. A. Zubkov, arXiv:1308.3406.

[8] *Gravitational Wilson loop in discrete quantum gravity*, arXiv:0907.2652.

[9] B. Holdom, J. Ren, Phys. Rev. D **93**, 124030 (2016).

[10] R. Liu, J. Quintin, N. Afshordi, Phys. Rev. Lett. **136**, 111501 (2026), arXiv:2510.18733.

[11] K. Osterwalder, E. Seiler, Ann. Phys. **110**, 440 (1978).

[12] S. Folkerts, D. F. Litim, J. M. Pawlowski, Phys. Lett. B **709**, 234 (2012), arXiv:1101.5552.

[13] A. Sengupta, Mem. Amer. Math. Soc. **126** (1997).

[14] T. Lévy, Mem. Amer. Math. Soc. **166** (2003).

[15] J. Bettinelli, G. Miermont, Ann. Inst. H. Poincaré Probab. Statist. **53**, 1244 (2017).

[16] J. Ambjørn, R. Loll, Nucl. Phys. B **536**, 407 (1998).

[17] T. Damour, M. Henneaux, H. Nicolai, Class. Quant. Grav. **20**, R145 (2003).