THE SUGGESTED CLINICAL PATHWAY FOR REMICADE®

PREPARATION 1 10 minutes Greet and escort patient Give Medication Guide to patient* Weigh patient and record. Take initial vital signs and record Place patient in reclining/IV chair **Assess patient** for therapy administration Review charts/notes and verify appropriate patient screening, including TB test Obtain/review physician's order Calculate drug dose or verify dose ordered Gather and set-up IV equipment 10 Wash hands; observe universal precautions **PREPARATION 2** 10 minutes **Assess** patient's arm for infusion site Prep venipuncture site Perform venipuncture with appropriate infusion access device **Check blood return** for site patency Connect tubing to IV site, secure connection junction Stabilize access device with dressing or tape Start infusing saline solution Assess access site for any redness or swelling Reassess patient for problems 10 Reconstitute/mix drug per package insert or notify pharmacy **ADMINISTRATION** 125 minutes This is an example of an infusion rate schedule Start infusion with REMICADE® as follows 10 mL/h x 15 minutes 20 mL/h x 15 minutes 40 mL/h x 15 minutes 80 mL/h x 15 minutes 150 mL/h x 30 minutes 250 mL/h x 30 minutes or until completely infused **Check and record** vital signs every 15 to 30 minutes during the infusion Monitor patient with every rate adjustment during the infusion **COMPLETION** 30 minutes Flush IV line with saline to ensure REMICADE® is completely infused Continue to monitor patient after infusion with REMICADE® is completely infused Remove venous access and observe site Provide patient with education materials Complete patient discharge **POST-INFUSION** 5 minutes

Dispose of waste materials

Clean and straighten room and equipment
Complete notes and documentation



^{*}Patients or their caregivers should be given the Medication Guide for REMICADE® and provided the opportunity to read it and discuss questions prior to each treatment session.