

MONTE CARLO METHOD FOR CALCULATING UNCERTAINTY IN OXYGEN ABUNDANCE FROM STRONG-LINE FLUX MEASUREMENTS

AUTHOR ORDER TO BE DETERMINED: MARYAM MODJAZ¹, FEDERICA B. BIANCO¹, SEUNG MAN OH^{1,2}, DAVID FIERROZ¹, YUQIAN LIU¹, LISA KEWLEY^{3,4}

Draft version March 12, 2015

ABSTRACT

MODIFY & FINALIZE AT THE END: We present an open-source Python code for the determination of the strong-emission-line estimators of oxygen abundance in the standard scales, based on the original IDL-code in Kewley & Dopita (2002). The standard strong line Metallicity scales and diagnostics IMPROVE have been used to estimate metal abundance through emission line ratios. Here we introduce a Monte Carlo resampling of these methods in order to better characterize an oxygen abundance confidence region. We output median values, 16th and 84th percentile confidence regions, for various standard metallicity diagnostics, and, when possible, for reddening $E(B-V)$. We produce Monte Carlo parameter distributions for the oxygen abundance and when possible for reddening $E(B-V)$. We test our code on emission lines measurements from a sample of galaxies ($z < 0.15$) and compare our metallicity results with those from previous methods. We show that our metallicity estimates are consistent with previous methods but yields smaller uncertainties. The code is open source and can be found at www.github.com/nyusgroup/ (add repo and DOI).

1. INTRODUCTION

Small amounts of carbon, oxygen, nitrogen, sulfur and iron and other elements provide a splash of color to the otherwise dominating greyscale of hydrogen and helium in the stars and gas of galaxies. Nevertheless, even this minute presence of heavy elements (all elements heavier than H and He, also called metals or collectively metallicity) is important for many areas of astrophysics. For example, Johnson & Li (2012) suggest that if it was not for the relatively high metallicity level in our Solar System, planet formation may not have been possible. With Z representing the mass fraction of metals, for our own Sun the value is measured to be $Z=0.0153$ (Caffau et al. 2011), though there are others who suggest a lower solar metallicity of $Z = 0.0134$ in particular because of oxygen (Asplund et al. 2009; Grevesse et al. 2010)⁵. Furthermore, when properly observed and estimated, metallicity measurements of galaxies can tightly constrain models of galaxy formation and evolution (e.g., Kewley & Ellison 2008 and references therein), as well as shed light on the metallicity dependence and production conditions for different types of SNe and long-duration GRBs (e.g., Modjaz et al. 2008; Levesque et al. 2010; Anderson et al. 2010; Modjaz et al. 2011; Kelly & Kirshner 2012; Sanders et al. 2012; Lunnan et al. 2014; Leloudas et al. 2014), to just name a few examples.

Metals are produced in the cores of stars during their fusion life cycle but also during the extreme conditions

of stellar explosions. For example, the majority of iron comes from thermonuclear explosions (SN Ia) while nearly all of oxygen is the result of core collapse (SN Ib, Ic, & II). Since new stars are born from the clouds these explosions produce, metallicity will increase with each passing generation.

However, for almost all astronomical objects, metallicity cannot be measured directly. The oxygen abundance in the gas-phase is the canonical choice of metallicity indicator for interstellar medium (ISM) studies, since oxygen is the most abundant metal and only weakly depleted onto dust grains (in contrast to refractory elements such as Mg, Si, Fe, with Fe, being depleted by more than a factor of 10 in Orion; see Simón-Díaz & Stasińska 2011). The oxygen abundance⁶ is expressed as $12 + \log_{10}(\frac{O}{H})$, where O and H represent the number of Oxygen and Hydrogen atoms, respectively. Importantly, oxygen exhibits very strong nebular lines in the optical wavelength range of HII regions (e.g., Pagel et al. 1979; Osterbrock 1989; Tremonti et al. 2004), which can be measured. Thus, many different diagnostic techniques, relying on different lines, have been developed (e.g., Kewley & Dopita 2002; Pettini & Pagel 2004; Kobulnicky & Kewley 2004; Kewley & Ellison 2008), which are discussed in the next section.

1.1. The different oxygen abundance diagnostics

Here we present a brief overview of the various observational methods for measuring the gas-phase oxygen abundance - however, for a full discussion with all the caveats we encourage the reader to read the reviews by e.g. Stasińska (2002); Kewley & Ellison (2008); Moustakas et al. (2010); Stasińska (2010); Dopita et al. (2013); Blanc et al. (2015). The so-called “classical” way to estimate the oxygen abundance is the electron temperature (T_e) method, which estimates the electron tempera-

¹ Center for Cosmology and Particle Physics, New York University, 4 Washington Place, New York, NY 10003, USA

² NYU Abu Dhabi PO Box 129188 Abu Dhabi, UAE

³ Australian National University, Research School for Astronomy & Astrophysics, Mount Stromlo Observatory, Cotter Road, Weston, ACT 2611, Australia

⁴ Institute of Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

⁵ Note that these abundances refer to the current abundances in the Sun, which are lower than the value with which the Sun was formed 4.56 Gyr ago, since diffusion at the bottom of the convection zone has decreased metallicity over time (Grevesse et al. 2010).

⁶ We note that in many cases in the literature, including here, the terms metallicity and oxygen abundance are used interchangeably.

ture and density of the nebula using a number of oxygen lines with different ionization states, including the auroral [OIII] $\lambda 4363$ line, to then directly estimate the OII and OIII abundances to obtain the total oxygen abundance, after correcting for the unseen stages of ionization. However, the auroral [OIII] $\lambda 4363$ line is very weak, even in low-metallicity environments, and saturates at higher metallicity (since at higher metallicities the cooling is dominated by the Oxygen NIR fine structure lines) – thus, other methods had to be developed that use other, stronger lines in the spectra of HII regions. These are called *strong-line methods* and are the subject of this manuscript. Strong-line methods can be categorized into two types: theoretical methods, that rely on calibrating various observed line ratios using photoionization methods (basically theoretically simulating HII regions, using stellar model atmospheres, stellar population synthesis and photoionization models) and empirical ones, that calibrate various observed strong line ratios using observed T_e -based metallicities. While historically there have been large systematic offsets between the T_e method and the strong line methods, Dopita et al. (2013) demonstrated that the T_e method gives the same results as the strong line methods, if the energy distribution of the electrons in the HII regions is assumed to not be a simple Maxwell-Boltzmann distribution (as assumed in prior works), but a more realistic κ distribution, as observed in solar system astrophysical plasma. They also find that the effect of the κ distribution on the strong-line methods is minor.

For theoretical strong-line method, a ratio of oxygen line fluxes to $H\beta$, referred to as $R23$, is commonly used to determine the metallicity of galaxies:

$$R23 = \frac{[\text{OII}]\lambda 3727 + [\text{OIII}]\lambda 4959, \lambda 5007}{H\beta}$$

(Pagel et al. 1979). The drawback of this method is that it is double-valued with metallicity, and thus other line ratios need to be used to break the degeneracy between the high values (“upper branch”) and the low values (“lower branch”) of the $R23$ metallicities. Furthermore, Kewley & Dopita (2002) showed the importance of ionization parameter, which can be physically understood to correspond to the maximum velocity of an ionized front that can be driven by the local radiation field of hot massive stars that are ionizing the ISM gas. This ionization parameter needs to be taken into account in the various strong-line methods, as HII regions at the same metallicity but with different ionization parameters produce different line strengths. Calibrations of $R23$ by McGaugh (1991) (hereafter M91) and by Kewley & Dopita (2002) (hereafter KD02) use different theoretical photoionization models and take the ionization parameter into account, while other calibrations such as of Zaritsky et al. (1994) (hereafter Z94) do not. Thus, Z94 is mostly valid for only metal-rich galaxies. M91 and KD02 use an iterative process to break the $R23$ degeneracy (KD02 uses different ratios $[\text{NII}]/[\text{OII}]$ and $[\text{NII}]/H\alpha$) and to also constrain the ionization parameter q in order to arrive at the metallicity estimate.

As to empirical strong-line methods, the most commonly used one is that by Pettini & Pagel (2004) (hereafter PP04). PP04 used HII regions with T_e -based

metallicities to derive empirical fits to strong-line ratios, and introduce the line ratios of $([\text{NII}]/H\beta(N2))$ and $([\text{OIII}]/H\beta)/([\text{NII}]/H\alpha(O3N2))$ as metallicity diagnostics. Since PP04-N2 employs two closely spaced lines ($H\alpha$ and NII), which are not affected by stellar absorption, nor uncertain reddening, and are easily observed in one simple spectroscopic setup, it has become an often-used scale, at least for low- z SN host galaxy studies (see metal-analysis by e.g., Sanders et al. 2012; Modjaz 2012; Leloudas et al. 2014). However, it is important to mention that this scale has a number of short-comings: it does not take into account the impact of the ionization parameter, it was initially derived based on only 137 extragalactic HII regions, and the nitrogen emission line employed saturates at high metallicity (**CHECK!**), and thus this method may not be well-suited for high-metallicity galaxies. An updated calibration by Marino et al. (2013) based on many more T_e -based metallicities (almost three time larger than that of PP04) derives a significantly shallower slope between $O3N2$ index and oxygen abundance than the PP04 calibration.

As it can be seen, each scale has different advantages and disadvantages and should be used in different metallicity regimes (see detailed discussion in e.g., Kewley & Dopita 2002; Stasińska 2002; Kewley & Ellison 2008; Moustakas et al. 2010; Dopita et al. 2013; Blanc et al. 2015). Thus, this open-source code outputs the oxygen abundance in the main 7⁷ metallicity scales (of the KD02 diagnostic has four outputs and the PP04 diagnostic has two outputs). While there is a long-standing debate about which diagnostic to use, as there are systematic metallicity offsets between different methods (recombination lines vs. strong-line method vs. “direct” T_e method, see the above sources), **the relative metallicity trends can be considered robust, if the analysis is performed self-consistently in the same scale, and trends are seen across different scales (Kewley & Ellison 2008; Moustakas et al. 2010)**. Note however, that while there are conversion values between different scales (Kewley & Ellison 2008), they apply for large data sets, since those conversion values were derived based on ten thousands of SDSS galaxies, and thus should be used with caution (or not at all) for smaller samples. In addition, one should note that there is a debate about the value of the solar oxygen abundance (Asplund et al. 2009; Caffau et al. 2011), such that the absolute oxygen calibration is still uncertain.

Here we introduce the open-source python code `””`. pro .. In § 2 we describe our method, the input and output values of the code. In § 4, we compare our method of obtaining abundance uncertainties to previous methods in the literature.

2. DESCRIPTION OF METALLICITY CODE

For computing oxygen abundances, we use the iterative code by Kewley & Dopita (2002), which has been updated in Kewley & Ellison (2008) and reflects .. **LISA: YOUR INPUT HERE: what is the update??** which was initially written in IDL. We translated the code into python, and added new features, most importantly the capability of obtaining uncertainties on the metallicity outputs via Monte Carlo resampling, and made it open

⁷ as of version v1.0, Spring 2015




FIG. 1.— WILL UPDATE WITH NEW FIGURES BASED ON A PUBLISHED SN WITH SIGNFICANT E(B-V) AND WRITE MORE TEXT. Metallicity and reddening (E(B-V)) parameter distributions based on example data (namely, emission line data of the HII regions at the position of SN 2008D, published in Modjaz et al. (2011), based on $N=20,000$ samples. The median value is shown with the dashed lines, while the 16th till 84th confidence region is shaded in tan color. The shown metallicity scales are by Kewley & Dopita (2002), updated by Kewley & Ellison (2008) (KD02combupdated), Pettini & Pagel (2004), using OIII and NII (PP04-O3N2) and McGaugh (1991) (M91). These are the same plots that are outpitted by the code, which produces such plots for all scales (namely those of **sLIST THEM ALL**). The legend also includes the ID number of

source on GitHub, as we explain below.

2.1. Input and Output of code

The input of the code is a set of spectral line fluxes. We assume that the observed emission lines to be used to indicate metallicity originate in HII regions and are not due to non-thermal excitation by e.g., AGN or interstellar shocks from SNe or stellar winds. Tests to exclude data contaminated by such non-thermal sources should be executed using the recommended line ratios by e.g., Baldwin et al. 1981; Kauffmann et al. 2003; Kewley et al. 2006 *prior to running this code*. Furthermore these lines should have all the correct calibration (at least correct relative calibration) and *should have a signal-to-noise ratio (S/N) of at least 3*. The latter is important for the success of the Monte Carlo resampling technique as described below.

Emission line flux values are fed into our Python implementation as in the original IDL code by Kewley & Dopita (2002), hereafter referred to as IDLKD02. The inputs are emission line flux values and their uncertainties for the following lines: $H\alpha$, $H\beta$, [OI] 6300, [OII] 3727, [OIII] 4959, [OIII] 5007, [NII] 6584, [SII] 6717, [SII] 6731. SIII 9532, and SIII 9096 can be used to calculate $S23$, but

are currently not used to calculate metallicity with any of the diagnostics enabled by this code. The line fluxes are to be stored in an ascii file, and the measurement errors in a separate ascii files. (consult the README.md⁸ in the GitHub repository for details about the input format, and sample files). **CHECK AT THE END WITH CODE!** If the fluxes for the specified lines are not available, the entry should be set to 'NaN' and the out-putted oxygen abundance will be calculated only for metallicity scales that use the provided line fluxes. In absence of measurement errors the flux errors entry should be set to 0.0 (and the code will not generate confidence intervals, as explained later in this Section).

As part of the code, the inputted line fluxes are corrected for reddening by using the observed Balmer decrement, for which $H\alpha$ and $H\beta$ flux values need to be provided. We assume case B recombination, and thus the standard value of 2.86 as the intrinsic $H\alpha/H\beta$ ratio (Osterbrock 1989), and apply the standard Galactic reddening law with $R_V = 3.1$ (Cardelli et al. 1989). However, the user can choose other extinction laws and R_V values,

⁸ https://github.com/nyusngroup/MC_Metalicity/blob/master/README.md

if desired, given the code’s open-source nature. If the input measurements are already de-reddened, the user can easily disable the reddening correction. If either $H\alpha$ or $H\beta$ are not provided the reddening correction cannot be implemented, the user is notified, and has the option to proceed with the calculation with uncorrected line fluxes.

As output, we obtain metallicity values and their uncertainties in various metallicity scales. The user can choose which of the following calibrations to calculate:

- **M91** (McGaugh 1991),
- **Z94** (Zaritsky et al. 1994),
- **C01**: a diagnostic based on $R23$, $C01_R23$, and one based on $[NII]/[SII]$, $C01_N2S2$ (Charlot & Longhetti 2001),
- **P01** (Pilyugin 2001),
- **D02** (Denicoló et al. 2002),
- **KD02**: 4 computations using $R23$ ($KD02_R23$), the $[NII]/[OII]$ ratio ($KD02_N2O2$), the $[NII]/H\alpha$ ratio ($KD02_N2O2$), and a combined method that chooses the optimal method given the input line fluxes ($KD02comb$) (Kewley & Dopita 2002),
- **PP04**: 2 computations based on the $[NII]/H\alpha$ ratio and $[OIII]/H\beta$ respectively ($PP04_N2$, $PP04_O3N2$) **CHECK WITH FINAL CODE - ALSO CHECK whether P01 is P01 or P05 by looking at the code** (Pettini & Pagel 2004).
- **pyqz**: In addition, if the user has installed the publicly available **pyqz**⁹ Python Module, based on Dopita et al. (2013), $[NII]$, $[SII]$, $[OIII]$, $H\alpha$, and $H\beta$ lines are fed to the **pyqz** module, which produces up to 8 emission line ratio diagnostics for $12+\log(O/H)$, each using two of the line ratios $[NII]/[SII]$, $[NII]/H\alpha$, $[OIII]/[SII]$, and $[OIII]/H\beta$. The **pyqz** module assumes a κ -distribution of energy for the electrons in HII regions, as suggested by Nicholls et al. (2012). Our code sets the κ parameter to 20, which is the value that Dopita et al. (2013) found best resolves the inconsistencies between oxygen abundance values derived from the strong-line method vs. the “direct” T_e method¹⁰.

All of the above calibrations, except for the **pyqz** results, are calculated in the original IDLKD02 code, and are discussed in detail in Kewley & Dopita (2002); Kewley & Ellison (2008).

If the line fluxes necessary for specific scales are not provided, the output metallicities will default to ‘NaN’, if the errors in the measurements are not provided, the code will specify that it cannot create a measurement distribution and determine a confidence interval, but calculate the nominal metallicity.

The value of $E(B-V)$ and $\log(R23)$ are output, together with the metallicity diagnostics. While parameters, such

as the ionization parameter q and the electron density (using the $SiIII$ lines) are computed, as long as the necessary lines are provided, they are not out-putted in the current version of our code – however, the reader can easily modify the code to suite their needs, given it’s an open-source code.

2.2. Computing Uncertainties

The novel aspect of our work is that for every set of input line measurements we introduce a Monte Carlo (MC) resampling method to obtain iterations via random sampling within the measurement errors, and thus we obtain a robust result for error estimation (e.g., Efron 1979; Hastie et al. 2009; Andrae 2010).

Given a data set with error bars from which certain parameters are estimated, Monte Carlo resampling generates synthetic data samples drawing from a given distribution. Here we draw synthetic data from a Gaussian distribution centered on each measured line flux value, with standard deviation corresponding to the measurement error. The implicit assumption is made, of course, that the line flux error is Gaussian distributed in nature¹¹.

For each metallicity scale, for each of N values chosen randomly within the relevant emission line distributions we run the calculations that derives the metallicity. This effectively simulates conducting multiple experiments when repeating observations is impractical or impossible, as in the case of the emission line flux data, and thus generates alternative data sets. The sample size N is set by the user, and one should expect an appropriate value of N to be a few 1000s, depending on the metallicity scale chosen and measurement errors (for example $N = 2,000$ is determined to be sufficient for our example data, as shown below **WE NEED TO SHOW IT YET**, and we provide tools to assure the sample size is sufficiently large). A distribution of parameter estimates for the oxygen abundance is generated for each scale, the metallicity and its confidence region are calculated, and the results are binned and visualized in a histogram (see Section 2.2.1)¹². This is done for each scale the user chooses to calculate. The fiftieth (50%) percentile, i.e. the median, is reported as the “true” metallicity value, and the 16th and 84th percentiles of the distribution as its confidence region.

This MC resampling approach takes into account the impact of the uncertain reddening (due to the uncertainties in the measurement of the $H\alpha$ and $H\beta$ fluxes), when the option for de-reddened metallicities is chosen. For each synthetic set of measurements a new reddening value is calculated based on the resampled $H\alpha$ and $H\beta$ fluxes, and used to compute the de-reddened metallicity value, thus the derived distribution of metallicity values takes into account the uncertain reddening. As part of the output, a probability distribution plot for $E(B-V)$ is provided as well (first panel in figure 1, along with confidence intervals derived using the same method as

¹¹ Users may wish to provides their own probability distribution for the emission line uncertainties, and modify the code to suite their needs.

¹² However, note this method is a conservative approach, since it overestimates the intrinsic metallicity uncertainty, as we are centering this error distribution on the measured values instead of the (unknown) true values (Andrae 2010).

⁹ <https://datacommons.anu.edu.au:8443/DataCommons/item/anudc:5037>

¹⁰ The user may wish to modify the value of κ by editing the code.

for the metallicity measurements. If either the $H\alpha$ or the $H\beta$ flux is not provided, no reddening correction can be applied. The computed metallicity will not be reddening-corrected and the $E(B-V)$ output will be set to zero.

Figure 1 shows the metallicity estimate distribution for three representative scales (M91, PP04_O3N2, and KD02comb), and for the reddening parameter $E(B-V)$ - similar plots that are out-putted by our code for all scales as listed above (not all shown here). Although the input distributions are Gaussian, the metallicity distributions are not, for two reasons: first, since the metallicities are computed based on log values of line flux ratios, symmetric error bars in linear space will translate into asymmetric error bars in log space; and second, some metallicity computations are non-linear, and (especially those that include R_{23}) sometimes bimodal since they choose upper vs lower branch to break degeneracy.

Since the metallicity distributions are rarely Gaussian, the percentiles we report cannot be expressed in terms of σ values. In determining the confidence region for asymmetric and multi-modal distributions, there are broadly three approaches (e.g., Andrae 2010): choosing a symmetric interval, the shortest interval, or a *central* interval. With the central method we determine the confidence interval by choosing the left and right boundaries such that the region outside the confidence interval on each side contains 16% of the total distribution - in analogy to the one-sigma-interval of a Gaussian distribution. This ensures that the algorithm finds the proper boundaries even for asymmetric, non-Gaussian distributions, and in the case of multiple peaks. In summary, the output for the measured value corresponds to the fiftieth (50%) percentile, while the lower error bar corresponds to the 50th-16th percentile and the upper error bar corresponds to 84th-50th of the metallicity estimate distribution.

The distributions for the D02 scale include the uncertainty in the fit parameters: the oxygen abundance in this scale is generated as $12 + \log_{10}(\frac{O}{H}) = 9.12 \pm 0.05 + (0.73 \pm 0.10) \text{ NII}$ (Denicoló et al. 2002). The parameters of the fit are generated as the sum of the nominal parameters (9.12 and 0.73) and a Gaussianly distributed random value centered on zero, and within a standard deviation of 0.05 and 0.10, respectively.

We note that our code does not output the *systematic* uncertainty of each scale, which, for example, are ~ 0.07 dex (or 0.14 dex **CHECK**) for PP04-O2N2. Thus systematic errors can be as large, if not larger, than the statistical errors. However, if all metallicity measurements are in the *same* scale and only *relative* comparisons are made, then the systematic error does not have any impact (by definition!).

2.2.1. Visual diagnostics

In order for the user to check the validity of a measurement, and to better understand the distribution, we provide two visualizations: for each metallicity scale calculated, for each input set, we generate a histogram of the output distribution (Figure 1 and 2), and for each set of input lines we generate a *box-and-whiskers* plot (hereafter *boxplot*, for short) summarizing the result of all scales calculated (Figure 3).

Choosing the binning size for a histogram is not a trivial task. Hogg (2008) describes various data analysis recipes for selecting a histogram bin size. Too many bins will result in many empty bins and an “over-fit” histogram, while too few bins may miss features of the distribution. By default, we use *Knuth’s Method* to choose the number of bins N_{bins} for each histogram. Knuth’s method optimizes a Bayesian fitness function across fixed-width bins (Knuth 2006). Additionally, however, we enable a number of binning options from which the user can choose, including: the square root of the number of bins, *Rice rule* ($N_{\text{bins}} = 2\sqrt[3]{N}$, e.g., Hastie et al. 2009), *Doane’s formula* ($N_{\text{bins}} = 1 + \log_2 N + \log_2 \left(1 + \text{Kurt} \sqrt{(N/6)}\right)$, where Kurt is the third moment of the distribution, Doane 1976¹³), and the full Bayesian solution, known as Bayesian Blocks, which optimizes a fitness function across an arbitrary configuration of bins, such that the bins are of variable size (Scargle et al. 2013). The implementation of the latter method requires the *astroML* python package to be installed on the user’s system (Vanderplas et al. 2012¹⁴). If the *astroML* package is not found, the code will default to Knuth’s Rule. As mentioned, Knuth’s method implies an optimization. In cases in which the convergence of this minimization takes too long (or if the number of bins after the minimization is $N_{\text{bins}}/\sqrt{N} > 5$ or $N_{\text{bins}}/\sqrt{N} < 1/3$) the code will revert to Rice rule. Some methods may be computationally prohibitive with a very large sample size, or very little computational power, such as the Bayesian methods that rely on optimization (Knuth’s method, and the Bayesian Block method) in which case the user may choose to use Doane’s formula, Rice rule, or even the square root of the number of samples, to choose the bin size for the histograms.




FIG. 2.— **FINISH**

Lastly, the user can generate and visualize a *Kernel Density* if the *sklearn* package is installed. The Kernel Density of the distribution is then calculated via *KD*

¹³ Doane (1976) attempted to address the issue of finding the proper number of bins for the histogram of a skewed distribution. Several version of the so-called Doane’s formula can be found in the literature. Our formula can be found, for example, in ?

¹⁴ <https://github.com/astroML/astroML>

Tree with a top-hat function, as explained in the `sklearn` package documentation¹⁵. The results will then show both a histogram, with N_{bins} chosen via Knuth’s method, as well as the distribution Kernel Density, as shown in Figure 2.




FIG. 3.— **FINISH**

The boxplot summarizes the result from each scale the user chooses to calculate. For each scale the median of the $12 + \log_{10}(\frac{O}{H})$ distribution is plotted as a black horizontal line. The height of the corresponding box represents the 25th percentile of the $12 + \log_{10}(\frac{O}{H})$ distribution. The bars represent the maximum and minimum value of the distribution, excluding outliers. The outliers, that are plotted as circles, are defined as all data points farther than $1.5 \times \text{IQR}$, where IQR is the *interquartile range* (and the length of the box) from the 25th percentile (i.e. from either end of the box). The Solar oxygen abundance is indicated in this plot for comparison: a gray box

shows a range of estimated values for Solar oxygen abundances, from $12 + \log_{10}(\frac{O}{H}) = 8.69$ (Asplund et al. 2009) to $12 + \log_{10}(\frac{O}{H}) = 8.76$ (Caffau et al. 2011). Notice that only the diagnostics requested by the user have a slot in the plot (in the example in Figure 2 the computed scales are M91, the PP04 scales, and the KD02 scales. However these slot exists on the plot whether the diagnostic can be produced or not, i.e. if the set of input lines does not allow a requested scale to be calculated an empty column will be generated in this plot in correspondence of said metallicity scale.

2.3. Visual diagnostic to assure completeness

The user can choose the number of samples to be generated. Of course the reliability of the metallicity estimates depends crucially on the sample size being sufficiently large to properly characterize the distribution of metallicity values. It is however not trivial to decide when N is sufficiently large. As soon as N is large enough, and the distribution is well characterized, adding synthetic data will not change its shape. Consider a cumulative distribution for a metallicity scale, D02, for example, which has noise from the measurement errors, as well as from the error in the fit parameters, or KD02, which is a non linear combination of the input line flux values. We plot the cumulative distribution for a randomly selected subsample of 10% of the datapoints in the distribution, a subsample of 25%, 50%, 75%, and for the entire distribution. If 1/10th of the points were a sufficiently large Monte Carlo sample, the cumulative distributions would appear smooth, and most importantly overlap. In Figure 4 we show these cumulative plots for a distribution generated with $N = 200$, $N = 2,000$, and $N = 20,000$, for D02 and KD02comb. The cumulative distributions for the subsets of the $N = 200$ simulation are different, and noisy, indicating that 200 data points is not a sufficiently large dataset. Conversely, at $N = 20,000$ the distributions are indistinguishable, indicating that $N = 2,000$ (the smaller subset plotted) is large enough. The noise in the $N = 2,000$ cumulative distribution is small, indicating that a value of N somewhere between 200 and 2,000 will characterize the distribution appropriately. In the light of this we chose, conservatively, $N = 2000$ to run our simulations for this dataset. However, we remind the user that the appropriate number will depend on the diagnostic, and on the input data. The way in which the lines are combined, which is in many cases non-linear, will affect the shape of the distribution, and thus the number of datapoints that are needed to fully characterize it.

2.4. performance and benchmarks

2.5. Availability

The source code is published under MIT-licensed on GitHub¹⁶. At this time the code is released under DOI NUMBER HERE as version 1.0: NAME HERE. Project details, step-wise tutorials, and further information can be found in the project README¹⁷. Development is done in Linux and OS X. The package requires standard

¹⁵ <http://scikit-learn.org/stable/modules/density.html>

¹⁶ [repoURLhere](#)
¹⁷ [README.mdURL](#)




FIG. 4.— TOBD

python packages, such as `numpy`, `scipy`, `pylab`, and additional features are enabled if the packages `astroml`, `skitlearn`, and `pyqz` are installed, but the packages are *not* required. Contact the authors to be included in a mailing list and be notified about critical changes.

3. COMPARISON TO PRIOR UNCERTAINTY COMPUTATION AND OTHER WORKS

A previous method for determining the uncertainty in the oxygen abundance (as used in Modjaz et al. 2008; Kewley et al. 2010; Rupke et al. 2010; Modjaz et al. 2011) was an *analytic* approach of propagating the emission-line flux uncertainties: it found the maximum and minimum abundances via maximizing and minimizing, respectively, the various line ratios by adding/subtracting to the measured line values their uncertainties. For comparison we computed the metallicities and their errors in both ways (both analytic and using our current MC resampling method) for 3 representative scales. We plot our results and the residuals in Fig.5, which shows a number of important points: i) The metallicity reported as the 50th percentile of the metallicity parameter distribution from the MC resampling method is completely consistent with the analytically derived metallicity - well within the respective error bars - and thus, the prior published results still stand (unsurprisingly, since our code, aside for the calculation of the confidence interval, uses the same algorithms developed for IDLKD02). ii) The MC resampling method has smaller error bars than the analytic method, especially for the scales of M91 and KD02. This is easily understandable, since the

analytic method assumes the worst-case-scenario, as it basically yields 2 metallicity parameter draws (the "minimum" and "maximum") which are in the tail of the full metallicity probability distribution. However, the MC resampling method is the more appropriate method as it empirically characterizes the full parameter estimation distribution.

3.1. Comparison with other works

The field of SN host metallicity studies has been rapidly developing as these kinds of studies may be crucial avenues for constraining the progenitor systems of different kinds of explosions - however, a few of the works do not even compute errors and others not show how they compute their statistical errors or (e.g., Anderson et al. (2010); Leloudas et al. (2011); Sanders et al. (2012); Leloudas et al. (2014)). While Sanders et al. (2012) compute the line flux errors from a Markov-Chain Monte Carlo fitting of a gaussian to the emission lines, they only mention in passing that they propagate the line flux uncertainties into the metallicity measurements, but do not describe how **this is a little harsh, right? i think mentioning it without explaining is generally the standard for iid errors -j Maryam: " How would you rephrase it then? "**

In contrast, the general metallicity field has considered in detail how to estimate the uncertainties in measured metallicities- however, none of those codes are open-source and many of them are for specific scales which were chosen by the authors: Moustakas et al. (2010) also use MC resampling to estimate the metallicity uncertain-




FIG. 5.— **FINISH**. Comparison of metallicity estimation between the analytic method and our Monte Carlo resampling method (top) and their residuals (bottom) for 3 different metallicity scales. Flux measurements come from 19 galaxies previously measured in Modjaz et al. (2011). To add asymmetric errors in quadrature we use $residual_{min} = \sqrt{x_{max}^2 + y_{min}^2}$ and $residual_{max} = \sqrt{x_{min}^2 + y_{max}^2}$

ties (in their case using 500 trials which seems to lead to a Gaussian distribution) but only do this for two scale, KK04 and Pilyugin & Thuan (2005). For computing the metallicities of the SDSS star forming galaxies, Tremonti et al. (2004) fit a combination of stellar population synthesis and CLOUDY (citation) photoionization models to the observed strong emission lines [OII], $H\beta$, [OIII], $H\alpha$, NII and SII and report the median of the metallicity likelihood distribution as the metallicity estimate, with

the width of the distribution giving the 1σ (Gaussian) error. However, this constitutes their own scale (the T04 scale).

In the last stages of preparing this manuscript Blanc et al. (2015) was published. Blanc et al. (2015) employ Bayesian inference for doing something similar to Tremonti et al. (2004) - they use Bayesian inference to derive the joint and marginalized posterior probability density functions for metallicity Z and ionization parameter q given a set of observed line fluxes and an input photoionization model. They provide a publicly available IDL implementation of their method named *IZI* (inferring metallicities (Z) and ionization parameters) on the author's web site.

4. CONCLUSIONS

FINISH. We hope that this open-access code will be used in many different fields where gas-phase metallicities are important, including in the emerging field of SN and GRB host galaxies, where either it is not described how they got or no error bars are computed (e.g., Lunnan et al. 2014). Given its public-access nature, the users are free to include any new metallicity diagnostics, e.g., that suggested by Dopita et al. (2013).

M. Modjaz is supported in parts by the NSF CAREER award AST-1352405 and by NSF award AST-1413260. FBB is supported by a *James Arthur Fellowship* at the NYU-Center for Cosmology and Particle Physics. Some plots are produced with public code DOI:10.5281/zenodo.15419 available at https://github.com/fedhere/residuals_pylib. This research made use of NASA Astrophysics Data System; the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Anderson, J. P., Covarrubias, R. A., James, P. A., Hamuy, M., & Haberman, S. M. 2010, MNRAS, 407, 2660
- Andrae, R. 2010, ArXiv e-prints, arXiv:1009.2755
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, PASP, 93, 5
- Blanc, G. A., Kewley, L., Vogt, F. P. A., & Dopita, M. A. 2015, ApJ, 798, 99
- Caffau, E., Ludwig, H.-G., Steffen, M., Freytag, B., & Bonifacio, P. 2011, Sol. Phys., 268, 255
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245
- Charlot, S., & Longhetti, M. 2001, MNRAS, 323, 887
- Denicoló, G., Terlevich, R., & Terlevich, E. 2002, MNRAS, 330, 69
- Doane, D. P. 1976, The American Statistician, 30, 181
- Dopita, M. A., Sutherland, R. S., Nicholls, D. C., Kewley, L. J., & Vogt, F. P. A. 2013, ApJS, 208, 10
- Efron, R. 1979, Ann. Stat., 7, 1
- Grevesse, N., Asplund, M., Sauval, A. J., & Scott, P. 2010, Ap&SS, 328, 179
- Hastie, T., Tibshirani, R., & Friedman, J. 2009, The Elements of Statistical Learning: Data Mining, Inference, and Prediction (Springer Science+Business Media, New York)
- Hogg, D. W. 2008, ArXiv e-prints, arXiv:0807.4820
- Johnson, J. L., & Li, H. 2012, ApJ, 751, 81
- Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, MNRAS, 346, 1055
- Kelly, P. L., & Kirshner, R. P. 2012, ApJ, 759, 107
- Kewley, L. J., & Dopita, M. A. 2002, ApJS, 142, 35
- Kewley, L. J., & Ellison, S. L. 2008, ApJ, 681, 1183
- Kewley, L. J., Groves, B., Kauffmann, G., & Heckman, T. 2006, MNRAS, 372, 961
- Kewley, L. J., Rupke, D., Zahid, H. J., Geller, M. J., & Barton, E. J. 2010, ApJ, 721, L48
- Knuth, K. H. 2006, ArXiv Physics e-prints, physics/0605197
- Kobulnicky, H. A., & Kewley, L. J. 2004, ApJ, 617, 240
- Leloudas, G., Gallazzi, A., Sollerman, J., et al. 2011, A&A, 530, A95
- Leloudas, G., Schulze, S., Kruehler, T., et al. 2014, ArXiv e-prints, arXiv:1409.8331
- Levesque, E. M., Berger, E., Kewley, L. J., & Bagley, M. M. 2010, AJ, 139, 694
- Lunnan, R., Chornock, R., Berger, E., et al. 2014, ApJ, 787, 138
- Marino, R. A., Rosales-Ortega, F. F., Sánchez, S. F., et al. 2013, A&A, 559, A114
- McGaugh, S. S. 1991, ApJ, 380, 140
- Modjaz, M. 2012, in IAU Symposium, Vol. 279, IAU Symposium, 207–211
- Modjaz, M., Kewley, L., Bloom, J. S., et al. 2011, ApJ, 731, L4
- Modjaz, M., Kewley, L., Kirshner, R. P., et al. 2008, AJ, 135, 1136
- Moustakas, J., Kennicutt, Jr., R. C., Tremonti, C. A., et al. 2010, ApJS, 190, 233
- Nicholls, D. C., Dopita, M. A., & Sutherland, R. S. 2012, ApJ, 752, 148

- Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galaxies* (Mill Vallery: University Science Books)
- Pagel, B. E. J., Edmunds, M. G., Blackwell, D. E., Chun, M. S., & Smith, G. 1979, *MNRAS*, 189, 95
- Pettini, M., & Pagel, B. E. J. 2004, *MNRAS*, 348, L59
- Pilyugin, L. S. 2001, *A&A*, 369, 594
- Pilyugin, L. S., & Thuan, T. X. 2005, *ApJ*, 631, 231
- Rupke, D. S. N., Kewley, L. J., & Chien, L.-H. 2010, *ApJ*, 723, 1255
- Sanders, N. E., Soderberg, A. M., Levesque, E. M., et al. 2012, *ApJ*, 758, 132
- Scargle, J. D., Norris, J. P., Jackson, B., & Chiang, J. 2013, *ApJ*, 764, 167
- Simón-Díaz, S., & Stasińska, G. 2011, *A&A*, 526, A48+
- Stasińska, G. 2002, *ArXiv Astrophysics e-prints*, astro-ph/0207500
- Stasińska, G. 2010, in *IAU Symposium*, Vol. 262, IAU Symposium, ed. G. R. Bruzual & S. Charlot, 93–96
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898
- Vanderplas, J., Connolly, A., Ivezić, Ž., & Gray, A. 2012, in *Conference on Intelligent Data Understanding (CIDU)*, 47–54
- Zaritsky, D., Kennicutt, Jr., R. C., & Huchra, J. P. 1994, *ApJ*, 420, 87

APPENDIX

MINIMUM CODE VALIDATION

WORK WITH FED - and any other code specific things