

Addressing Confounding and Continuous Exposure Measurement Error Using Corrected Score Functions

Brian Richardson

Department of Biostatistics,
University of North Carolina at Chapel Hill

August 5, 2025



Scan for slides

Acknowledgements

Bryan Blette, PhD



Vanderbilt University

Peter Gilbert, PhD



University of Washington
Medical Center


Michael Hudgens, PhD



UNC Chapel Hill


This research was supported by the U.S. Public Health Service Grant AI068635, the National Institute of Allergy And Infectious Diseases of the National Institutes of Health (NIH) under Award Number R37 AI054165, and the National Institute of Environmental Health Sciences of the NIH under Award Number T32

Motivation: HVTN 505 trial



- **HVTN 505 trial:** trial of a preventive HIV vaccine

Motivation: HVTN 505 trial



- **HVTN 505 trial:** trial of a preventive HIV vaccine
- Stopped early after reaching predetermined cutoffs for efficacy futility [2]

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE

 IDSA
Infectious Diseases Society of America

 hivma
hiv medicine association

 OXFORD

Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. James,¹ Kristen W. Cohen,¹ Nicole Frahm,¹ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Beutley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1c} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients [3, 1, 4]

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE

 IDSA
Infectious Diseases Society of America

 hivma
hiv medicine association

 OXFORD

Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. James,¹ Kristen W. Cohen,¹ Nicole Frahm,¹ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients [3, 1, 4]
- Is there a **causal** relationship between these biomarkers and HIV?

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE

 IDSA
Infectious Diseases Society of America

 hivma
hiv medicine association

 OXFORD

Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. James,¹ Kristen W. Cohen,¹ Nicole Frahm,¹ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients [3, 1, 4]
- Is there a **causal** relationship between these biomarkers and HIV?
- Biomarker-HIV relationship is **confounded**

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE

IDSA
Infectious Diseases Society of America

hivma
hiv medicine association

oxford

Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. James,¹ Kristen W. Cohen,¹ Nicole Frahm,¹ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Beutley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients [3, 1, 4]
- Is there a **causal** relationship between these biomarkers and HIV?
- Biomarker-HIV relationship is **confounded**
- Biomarkers are **measured with error**

Goal

To estimate the **causal effect** of a continuous exposure on an outcome when

Goal

To estimate the **causal effect** of a continuous exposure on an outcome when
(i) the exposure-outcome relationship is potentially **confounded**

Goal

To estimate the **causal effect** of a continuous exposure on an outcome when

- (i) the exposure-outcome relationship is potentially **confounded**
- (ii) the exposure is **measured with error**

What Do We Mean by Causal Effect?

“What would be the risk of HIV if, possibly counter to fact, somebody were to have biomarker level \mathbf{a} ?”


What Do We Mean by Causal Effect?

“What would be the risk of HIV if, possibly counter to fact, somebody were to have biomarker level a ?”

a
biomarker

What Do We Mean by Causal Effect?

“What would be the risk of HIV if, possibly counter to fact, somebody were to have biomarker level \mathbf{a} ?”



What Do We Mean by Causal Effect?

“What would be the risk of HIV if, possibly counter to fact, somebody were to have biomarker level \mathbf{a} ?”



What Do We Mean by Causal Effect?

“What would be the risk of HIV if, possibly counter to fact, somebody were to have biomarker level \mathbf{a} ?”




Estimand (**dose-response surface**): $\eta(\mathbf{a}) \equiv E\{Y(\mathbf{a})\}$ for $\mathbf{a} \in \mathcal{A}$


Confounding




Confounding



Confounding



Measurement Error



Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$
- potential outcome: $Y(\mathbf{a})$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$
- potential outcome: $Y(\mathbf{a})$
- observed outcome: Y

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$
- potential outcome: $Y(\mathbf{a})$
- observed outcome: Y
- confounders: $\mathbf{L} = (L_1, L_2, \dots, L_p)$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$
- potential outcome: $Y(\mathbf{a})$
- observed outcome: Y
- confounders: $\mathbf{L} = (L_1, L_2, \dots, L_p)$

Observe: iid copies of $(Y_i, \mathbf{L}_i, \mathbf{A}_i^*)$

Notation

- true exposure: $\mathbf{A} = (A_1, \dots, A_m)$
- measured exposure: $\mathbf{A}^* = (A_1^*, \dots, A_m^*) = \mathbf{A} + \boldsymbol{\epsilon}$
- measurement error: $\boldsymbol{\epsilon}$
- potential outcome: $Y(\mathbf{a})$
- observed outcome: Y
- confounders: $\mathbf{L} = (L_1, L_2, \dots, L_p)$

Observe: iid copies of $(Y_i, \mathbf{L}_i, \mathbf{A}_i^*)$

Estimand: dose-response surface $\eta(\mathbf{a}) \equiv E\{Y(\mathbf{a})\}$ for $\mathbf{a} \in \mathcal{A}$

Assumptions

(i) **causal consistency**: $Y = Y(\mathbf{a})$ when $\mathbf{A} = \mathbf{a}$

Assumptions

- (i) **causal consistency:** $Y = Y(\mathbf{a})$ when $\mathbf{A} = \mathbf{a}$
- (ii) **conditional exchangeability:** $Y(\mathbf{a}) \perp\!\!\!\perp \mathbf{A} | \mathbf{L}$ for all $\mathbf{a} \in \mathcal{A}$

Assumptions

- (i) **causal consistency**: $Y = Y(\mathbf{a})$ when $\mathbf{A} = \mathbf{a}$
- (ii) **conditional exchangeability**: $Y(\mathbf{a}) \perp\!\!\!\perp \mathbf{A} | \mathbf{L}$ for all $\mathbf{a} \in \mathcal{A}$
- (iii) **positivity**: $f_{\mathbf{A}|\mathbf{L}}(\mathbf{a}|\mathbf{l}) > 0$ for all \mathbf{l} such that $f_{\mathbf{L}}(\mathbf{l}) > 0$ and for all $\mathbf{a} \in \mathcal{A}$

Assumptions

- (i) **causal consistency**: $Y = Y(\mathbf{a})$ when $\mathbf{A} = \mathbf{a}$
- (ii) **conditional exchangeability**: $Y(\mathbf{a}) \perp\!\!\!\perp \mathbf{A} | L$ for all $\mathbf{a} \in \mathcal{A}$
- (iii) **positivity**: $f_{\mathbf{A}|L}(\mathbf{a}|l) > 0$ for all l such that $f_L(l) > 0$ and for all $\mathbf{a} \in \mathcal{A}$
- (iv) **classical additive measurement error**: $\epsilon \sim \mathcal{N}_m(0, \Sigma_{me})$ and $\epsilon \perp\!\!\!\perp (Y, L, \mathbf{A})$

Assumptions

- (i) **causal consistency**: $Y = Y(\mathbf{a})$ when $\mathbf{A} = \mathbf{a}$
- (ii) **conditional exchangeability**: $Y(\mathbf{a}) \perp\!\!\!\perp \mathbf{A} | \mathbf{L}$ for all $\mathbf{a} \in \mathcal{A}$
- (iii) **positivity**: $f_{\mathbf{A}|\mathbf{L}}(\mathbf{a}|\mathbf{l}) > 0$ for all \mathbf{l} such that $f_{\mathbf{L}}(\mathbf{l}) > 0$ and for all $\mathbf{a} \in \mathcal{A}$
- (iv) **classical additive measurement error**: $\epsilon \sim \mathcal{N}_m(0, \Sigma_{me})$ and $\epsilon \perp\!\!\!\perp (Y, \mathbf{L}, \mathbf{A})$
- (v) **known or estimable measurement error variance** Σ_{me}

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

- 3 classical methods:

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

- 3 classical methods:

- 1 G-Formula

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

- 3 classical methods:

- ① G-Formula
- ② Inverse Probability Weighting (IPW)

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

- 3 classical methods:

- ① G-Formula
- ② Inverse Probability Weighting (IPW)
- ③ Doubly Robust Method (DR)

Addressing Confounding Alone

How can we address confounding in the absence of measurement error?

- 3 classical methods:
 - 1 G-Formula
 - 2 Inverse Probability Weighting (IPW)
 - 3 Doubly Robust Method (DR)
- These can all be framed as **M-estimators**

Crash Course on M-Estimation

Score Function: a function of the observed data and the parameter of interest

$$\underbrace{\psi_0}_{\text{score fun.}} (\quad ; \quad)$$

Crash Course on M-Estimation

Score Function: a function of the observed data and the parameter of interest

$$\underbrace{\psi_0}_{\text{score fun.}} \left(\underbrace{Y, L, A}_{\text{data}} ; \quad \right)$$

Crash Course on M-Estimation

Score Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi_0}_{\text{score fun.}} \left(\underbrace{Y, L, A}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

Crash Course on M-Estimation

Score Function: a function of the observed data and the parameter of interest

$$\underbrace{\psi_0}_{\text{score fun.}} \left(\underbrace{Y, L, A}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

Crash Course on M-Estimation

Score Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi_0}_{\text{score fun.}} \left(\underbrace{Y, L, A}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

$$E\left\{ \Psi_0(Y, L, A; \underbrace{\theta_0}_{\text{true}}) \right\} = \mathbf{0}.$$

Crash Course on M-Estimation (cont'd)

- Given a score function $\boldsymbol{\psi}_0$ and observed data $\{(Y_i, \mathbf{L}_i, \mathbf{A}_i) : i = 1, \dots, n\}$, we can find an estimator $\hat{\boldsymbol{\theta}}$ as the solution to

Crash Course on M-Estimation (cont'd)

- Given a score function $\boldsymbol{\psi}_0$ and observed data $\{(Y_i, \mathbf{L}_i, \mathbf{A}_i) : i = 1, \dots, n\}$, we can find an estimator $\hat{\boldsymbol{\theta}}$ as the solution to

$$\sum_{i=1}^n \boldsymbol{\psi}_0(\underbrace{Y_i, \mathbf{L}_i, \mathbf{A}_i}_{\text{observed}}; \boldsymbol{\theta}) = \mathbf{0}.$$

Crash Course on M-Estimation (cont'd)

- Given a score function $\boldsymbol{\psi}_0$ and observed data $\{(Y_i, \mathbf{L}_i, \mathbf{A}_i) : i = 1, \dots, n\}$, we can find an estimator $\hat{\boldsymbol{\theta}}$ as the solution to

$$\sum_{i=1}^n \boldsymbol{\psi}_0(\underbrace{Y_i, \mathbf{L}_i, \mathbf{A}_i}_{\text{observed}}; \boldsymbol{\theta}) = \mathbf{0}.$$

- $\hat{\boldsymbol{\theta}}$ is consistent and asymptotically normal and has a simple variance estimator [6].

G-Formula

- 1 fit the **outcome model** $\mu(\mathbf{L}, \mathbf{A}; \beta) \equiv E(Y|\mathbf{L}, \mathbf{A})$

G-Formula

- 1 fit the **outcome model** $\mu(\mathbf{L}, \mathbf{A}; \beta) \equiv E(Y|\mathbf{L}, \mathbf{A})$
- 2 estimate the dose-response curve by marginalizing over the distribution of confounders: $\hat{\eta}(\mathbf{a}) = n^{-1} \sum_{i=1}^n \mu(\mathbf{L}_i, \mathbf{a}; \hat{\beta})$

G-Formula

- 1 fit the **outcome model** $\mu(\mathbf{L}, \mathbf{A}; \beta) \equiv E(Y|\mathbf{L}, \mathbf{A})$
- 2 estimate the dose-response curve by marginalizing over the distribution of confounders: $\hat{\eta}(\mathbf{a}) = n^{-1} \sum_{i=1}^n \mu(\mathbf{L}_i, \mathbf{a}; \hat{\beta})$

This can be expressed as an M-estimator with score function

$$\Psi_{0-GF}(Y, \mathbf{L}, \mathbf{A}; \theta_{GF}) = \begin{bmatrix} \{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\} \partial_{\beta} \mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{bmatrix}$$

- 1 obtain/estimate **standardized propensity score weights**

$$SW(L, A) = \frac{f_A(A)}{f_{A|L}(A|L)}$$

- 1 obtain/estimate **standardized propensity score weights**

$$SW(\mathbf{L}, \mathbf{A}) = \frac{f_{\mathbf{A}}(\mathbf{A})}{f_{\mathbf{A}|\mathbf{L}}(\mathbf{A}|\mathbf{L})}$$

- 2 use weighted observations to estimate the dose-response curve $\eta(\mathbf{a}; \gamma)$

IPW

- 1 obtain/estimate **standardized propensity score weights**

$$SW(\mathbf{L}, \mathbf{A}) = \frac{f_{\mathbf{A}}(\mathbf{A})}{f_{\mathbf{A}|\mathbf{L}}(\mathbf{A}|\mathbf{L})}$$

- 2 use weighted observations to estimate the dose-response curve $\eta(\mathbf{a}; \gamma)$

This can be expressed as an M-estimator with score function

$$\psi_{0-IPW}(Y, \mathbf{L}, \mathbf{A}; \theta_{IPW}) = \begin{bmatrix} \psi_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A}) \{Y - \eta(\mathbf{A}; \gamma)\} \partial_{\gamma} \eta(\mathbf{A}; \gamma) \end{bmatrix}$$

- 1 obtain/estimate **standardized propensity score weights** $SW(L, A)$

DR

- 1 obtain/estimate **standardized propensity score weights** $SW(L, A)$
- 2 use weighted observations to estimate the **outcome model** $\mu(L, A; \beta) \equiv E(Y|L, A)$

DR

- ① obtain/estimate **standardized propensity score weights** $SW(L, A)$
- ② use weighted observations to estimate the **outcome model** $\mu(L, A; \beta) \equiv E(Y|L, A)$
- ③ estimate the dose-response curve by marginalizing over the distribution of confounders

DR

- 1 obtain/estimate **standardized propensity score weights** $SW(\mathbf{L}, \mathbf{A})$
- 2 use weighted observations to estimate the **outcome model** $\mu(\mathbf{L}, \mathbf{A}; \beta) \equiv E(Y|\mathbf{L}, \mathbf{A})$
- 3 estimate the dose-response curve by marginalizing over the distribution of confounders

This can be expressed as an M-estimator with score function

$$\boldsymbol{\psi}_{0-DR}(Y, \mathbf{L}, \mathbf{A}; \boldsymbol{\theta}_{DR}) = \begin{bmatrix} \boldsymbol{\psi}_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A})\{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\}\partial_{\beta}\mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{bmatrix}$$

DR

- 1 obtain/estimate **standardized propensity score weights** $SW(\mathbf{L}, \mathbf{A})$
- 2 use weighted observations to estimate the **outcome model** $\mu(\mathbf{L}, \mathbf{A}; \beta) \equiv E(Y|\mathbf{L}, \mathbf{A})$
- 3 estimate the dose-response curve by marginalizing over the distribution of confounders

This can be expressed as an M-estimator with score function

$$\boldsymbol{\psi}_{0-DR}(Y, \mathbf{L}, \mathbf{A}; \boldsymbol{\theta}_{DR}) = \begin{bmatrix} \boldsymbol{\psi}_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A})\{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\}\partial_{\beta}\mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{bmatrix}$$

Doubly robust* to models for $\mu(\mathbf{L}, \mathbf{A}; \beta)$ and $f_{\mathbf{A}|\mathbf{L}}(\mathbf{A}|\mathbf{L})$.

Addressing Confounding and Measurement Error

Can we just substitute \mathbf{A}^* for \mathbf{A} and find the solution to

$$\sum_{i=1}^n \psi_0(Y_i, L_i, \underbrace{A_i^*}_{\text{mismeasured}}; \theta) = \mathbf{0}?$$

Addressing Confounding and Measurement Error

Can we just substitute \mathbf{A}^* for \mathbf{A} and find the solution to

$$\sum_{i=1}^n \boldsymbol{\psi}_0(Y_i, L_i, \underbrace{\mathbf{A}_i^*}_{\text{mismeasured}}; \boldsymbol{\theta}) = \mathbf{0}?$$

No! This leads to bias in $\hat{\boldsymbol{\theta}}$ because

$$E\{\boldsymbol{\psi}_0(Y, L, \mathbf{A}^*; \boldsymbol{\theta}_0)\} \neq \mathbf{0}.$$

Addressing Confounding and Measurement Error

Can we just substitute \mathbf{A}^* for \mathbf{A} and find the solution to

$$\sum_{i=1}^n \psi_0(Y_i, L_i, \underbrace{\mathbf{A}_i^*}_{\text{mismeasured}}; \theta) = \mathbf{0}?$$

No! This leads to bias in $\hat{\theta}$ because

$$E\{\psi_0(Y, L, \mathbf{A}^*; \theta_0)\} \neq \mathbf{0}.$$

We need a new score function ψ_{CS} such that

$$E\left\{ \underbrace{\psi_{CS}}_{\text{new score fun.}} (Y, L, \underbrace{\mathbf{A}^*}_{\text{mismeasured}}; \theta_0) \right\} = \mathbf{0}.$$

Corrected Score Functions

Given the “oracle” score function Ψ_0 , the “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) [5]:

Corrected Score Functions

Given the “oracle” score function Ψ_0 , the “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) [5]:

$$\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) = E[\operatorname{Re}\{\Psi_0(Y, L, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, L, \mathbf{A}^*]$$

Corrected Score Functions

Given the “oracle” score function Ψ_0 , the “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) [5]:

$$\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) = E[\text{Re}\{\Psi_0(Y, L, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, L, \mathbf{A}^*]$$

- 1 Add additional *imaginary* measurement error $i\tilde{\epsilon}$ to the mismeasured exposure

Corrected Score Functions

Given the “oracle” score function Ψ_0 , the “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) [5]:

$$\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) = E[\operatorname{Re}\{\Psi_0(Y, L, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, L, \mathbf{A}^*]$$

- 1 Add additional *imaginary* measurement error $i\tilde{\epsilon}$ to the mismeasured exposure
- 2 Keep only the real part of the resulting complex-valued function

Corrected Score Functions

Given the “oracle” score function Ψ_0 , the “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) [5]:

$$\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) = E[\operatorname{Re}\{\Psi_0(Y, L, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, L, \mathbf{A}^*]$$

- 1 Add additional *imaginary* measurement error $i\tilde{\epsilon}$ to the mismeasured exposure
- 2 Keep only the real part of the resulting complex-valued function
- 3 Take the expectation over the additional measurement error $\tilde{\epsilon}$.

Corrected Score Functions (contd)

Under certain conditions, the corrected score function Ψ_{CS} is then unbiased, meaning

$$E\{\Psi_{CS}(Y, L, \underbrace{A^*}_{\text{mismeasured}}; \theta_0)\} = \mathbf{0}.$$

Corrected Score Functions (contd)

Under certain conditions, the corrected score function Ψ_{CS} is then unbiased, meaning

$$E\{\Psi_{CS}(Y, L, \underbrace{A^*}_{\text{mismeasured}}; \theta_0)\} = \mathbf{0}.$$

The G-Formula, IPW, and DR score functions all satisfy these conditions, and so can be “corrected:”

$$\Psi_{0-GF} \longrightarrow \Psi_{CS-GF}$$

$$\Psi_{0-IPW} \longrightarrow \Psi_{CS-IPW}$$

$$\Psi_{0-DR} \longrightarrow \Psi_{CS-DR}$$

Monte Carlo Corrected Score Functions

- Sometimes we can find a closed-form algebraic expression for

$$\boldsymbol{\Psi}_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \boldsymbol{\theta}) = \underbrace{\mathbb{E}[\operatorname{Re}\{\boldsymbol{\Psi}_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\boldsymbol{\epsilon}}; \boldsymbol{\theta})\} \mid Y, \mathbf{L}, \mathbf{A}^*]}_{\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\}}.$$

Monte Carlo Corrected Score Functions

- Sometimes we can find a closed-form algebraic expression for

$$\boldsymbol{\Psi}_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \boldsymbol{\theta}) = \underbrace{\mathbb{E}[\text{Re}\{\boldsymbol{\Psi}_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\boldsymbol{\epsilon}}; \boldsymbol{\theta})\} \mid Y, \mathbf{L}, \mathbf{A}^*]}_{\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\}}.$$

- Alternatively, we can approximate this expectation with Monte Carlo replicates

Monte Carlo Corrected Score Functions

- Sometimes we can find a closed-form algebraic expression for

$$\boldsymbol{\Psi}_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \boldsymbol{\theta}) = \underbrace{\mathbb{E}[\operatorname{Re}\{\boldsymbol{\Psi}_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\boldsymbol{\epsilon}}; \boldsymbol{\theta})\} \mid Y, \mathbf{L}, \mathbf{A}^*]}_{\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\}}.$$

- Alternatively, we can approximate this expectation with Monte Carlo replicates

$$\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\} \approx B^{-1} \sum_{b=1}^B f(\tilde{\boldsymbol{\epsilon}}_b)$$

Monte Carlo Corrected Score Functions

- Sometimes we can find a closed-form algebraic expression for

$$\boldsymbol{\Psi}_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \boldsymbol{\theta}) = \underbrace{\mathbb{E}[\operatorname{Re}\{\boldsymbol{\Psi}_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\boldsymbol{\epsilon}}; \boldsymbol{\theta})\} \mid Y, \mathbf{L}, \mathbf{A}^*]}_{\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\}}.$$

- Alternatively, we can approximate this expectation with Monte Carlo replicates

$$\begin{aligned}\mathbb{E}\{f(\tilde{\boldsymbol{\epsilon}})\} &\approx B^{-1} \sum_{b=1}^B f(\tilde{\boldsymbol{\epsilon}}_b) \\ \implies \boldsymbol{\Psi}_{MCCS}^B(Y, \mathbf{L}, \mathbf{A}^*; \boldsymbol{\theta}) &= B^{-1} \sum_{b=1}^B \operatorname{Re}\{\boldsymbol{\Psi}_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\boldsymbol{\epsilon}}_b; \boldsymbol{\theta})\}\end{aligned}$$

Simulation Setting

$$L \sim \mathcal{N}(0, 0.36)$$




Simulation Setting

$$L \sim \mathcal{N}(0, 0.36)$$




$$\mathbf{A}|L \sim \mathcal{N}_2(\mathbf{0}, I)$$

Simulation Setting




Simulation Setting


$$\epsilon \sim \mathcal{N}_2(\mathbf{0}, \sigma_{me}^2 \mathbf{I})$$



$$L \sim \mathcal{N}(0, 0.36)$$




$$\mathbf{A}|L \sim \mathcal{N}_2(\mathbf{0}, \mathbf{I})$$




$$Y|L, \mathbf{A} \sim \mathcal{N}(A_1 + A_2 + L, 1)$$

Simulation Setting



Simulation Setting



Simulation Setting

- implies a dose-response surface of $\eta(\mathbf{a}; \gamma) = \gamma_0 + \underbrace{\gamma_1}_{\text{estimand}} a_1 + \gamma_2 a_2$

Simulation Setting

- implies a dose-response surface of $\eta(\mathbf{a}; \gamma) = \gamma_0 + \underbrace{\gamma_1}_{\text{estimand}} a_1 + \gamma_2 a_2$
- sample size $n = 800$

Simulation Setting

- implies a dose-response surface of $\eta(\mathbf{a}; \gamma) = \gamma_0 + \underbrace{\gamma_1}_{\text{estimand}} a_1 + \gamma_2 a_2$
- sample size $n = 800$
- 2 estimators compared:


Simulation Setting

- implies a dose-response surface of $\eta(\mathbf{a}; \boldsymbol{\gamma}) = \gamma_0 + \underbrace{\gamma_1}_{\text{estimand}} a_1 + \gamma_2 a_2$
- sample size $n = 800$
- 2 estimators compared:
 - ① Naive IPW (ignores measurement error)


Simulation Setting

- implies a dose-response surface of $\eta(\mathbf{a}; \boldsymbol{\gamma}) = \gamma_0 + \underbrace{\gamma_1}_{\text{estimand}} a_1 + \gamma_2 a_2$
- sample size $n = 800$
- 2 estimators compared:
 - 1 Naive IPW (ignores measurement error)
 - 2 Corrected Score IPW


Simulation Results: Estimator




Simulation Results: Estimator



Simulation Results: Confidence Interval



Simulation Results: Confidence Interval



Application: HVTN 505 Trial

- **two exposures** (both log-transformed):

Application: HVTN 505 Trial

- **two exposures** (both log-transformed):
 - (i) antibody-dependent cellular phagocytosis (**ADCP**)

Application: HVTN 505 Trial

- **two exposures** (both log-transformed):
 - (i) antibody-dependent cellular phagocytosis (**ADCP**)
 - (ii) recruitment of $\text{Fc}\gamma\text{RIIa}$ of the H131-Con S gp140 protein (**RII**)

Application: HVTN 505 Trial

- **two exposures** (both log-transformed):
 - (i) antibody-dependent cellular phagocytosis (**ADCP**)
 - (ii) recruitment of $\text{Fc}\gamma\text{RIIa}$ of the H131-Con S gp140 protein (**RII**)
- **case-cohort sampling**: biomarkers only measured in stratified random sample of controls


Application: HVTN 505 Trial

- **two exposures** (both log-transformed):
 - (i) antibody-dependent cellular phagocytosis (**ADCP**)
 - (ii) recruitment of $\text{Fc}\gamma\text{RIIa}$ of the H131-Con S gp140 protein (**RII**)
- **case-cohort sampling**: biomarkers only measured in stratified random sample of controls
- **covariates**: age, race, BMI, behavior risk, CD4-P, and CD8-P

Application: HVTN 505 Trial


- **two exposures** (both log-transformed):
 - (i) antibody-dependent cellular phagocytosis (**ADCP**)
 - (ii) recruitment of $\text{Fc}\gamma\text{RIIa}$ of the H131-Con S gp140 protein (**RII**)
- **case-cohort sampling**: biomarkers only measured in stratified random sample of controls
- **covariates**: age, race, BMI, behavior risk, CD4-P, and CD8-P
- **analysis**: DR estimator with a log-linear outcome model

Application: HVTN 505 Trial



Exposure values


Application: HVTN 505 Trial



Mismex: Causal Inference for Mismeasured Exposures



Paper in *Biometrics*



GitHub R package

References

- [1] Youyi Fong, Xiaoying Shen, Vicki C Ashley, Aaron Deal, Kelly E Seaton, Chenchu Yu, Shannon P Grant, Guido Ferrari, Allan C deCamp, Robert T Bailer, et al. Modification of the association between T-cell immune responses and human immunodeficiency virus type 1 infection risk by vaccine-induced antibody responses in the HVTN 505 trial. *The Journal of Infectious Diseases*, 217(8):1280–1288, 2018.
- [2] Scott M Hammer, Magdalena E Sobieszczyk, Holly Janes, Shelly T Karuna, Mark J Mulligan, Doug Grove, Beryl A Koblin, Susan P Buchbinder, Michael C Keefer, Georgia D Tomaras, et al. Efficacy trial of a DNA/rAd5 HIV-1 preventive vaccine. *New England Journal of Medicine*, 369(22):2083–2092, 2013.
- [3] Holly E Janes, Kristen W Cohen, Nicole Frahm, Stephen C De Rosa, Brittany Sanchez, John Hural, Craig A Magaret, Shelly Karuna, Carter Bentley, Raphael Gottardo, et al. Higher T-cell responses induced by DNA/rAd5 HIV-1 preventive vaccine are associated with lower HIV-1 infection risk in an efficacy trial. *The Journal of Infectious Diseases*, 215(9):1376–1385, 2017.
- [4] Scott D Neidich, Youyi Fong, Shuying S Li, Daniel E Geraghty, Brian D Williamson, William Chad Young, Derrick Goodman, Kelly E Seaton, Xiaoying Shen, Sheetal Sawant, et al. Antibody Fc effector functions and IgG3 associate with decreased HIV-1 risk. *Journal of Clinical Investigation*, 129(11):4838–4849, 2019.
- [5] Steven J Novick and Leonard A Stefanski. Corrected score estimation via complex variable simulation extrapolation. *Journal of the American Statistical Association*, 97(458):472–481, June 2002. ISSN 0162-1459, 1537-274X. doi: 10.1198/016214502760047005. URL <http://www.tandfonline.com/doi/abs/10.1198/016214502760047005>.
- [6] Leonard A Stefanski and Dennis D Boos. The calculus of M-estimation. *The American Statistician*, 56(1):29–38, 2002.

Appendix

Appendix: Corrected Score Functions

- Suppose the oracle score function is **conditionally unbiased**, meaning

$$E\{\Psi_0(Y, L, \mathbf{A}; \theta) | \mathbf{A}\} = 0.$$

- Define the corrected score function as

$$\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) = E \left[\text{Re} \left\{ \Psi_0(Y, L, \tilde{\mathbf{A}}; \theta) \right\} \mid Y, L, \mathbf{A}^* \right],$$

where $\tilde{\mathbf{A}} = \mathbf{A}^* + i\tilde{\epsilon}$, $i = \sqrt{-1}$, $\text{Re}(\cdot)$ denotes the real component of a complex number, and $\tilde{\epsilon} \sim \mathcal{N}(\mathbf{0}, \Sigma_{me})$.

- Then

$$\begin{aligned} E\{\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) \mid Y, L, \mathbf{A}\} &= \Psi_0(Y, L, \mathbf{A}; \theta) \\ \implies E[E\{\Psi_{CS}(Y, L, \mathbf{A}^*; \theta) \mid Y, L, \mathbf{A}\}] &= E\{\Psi_0(Y, L, \mathbf{A}; \theta)\} \\ \implies E\{\Psi_{CS}(Y, L, \mathbf{A}^*; \theta)\} &= 0 \end{aligned}$$