

Causal Inference Methods for Infectious Disease Prevention Trials

Brian Richardson

Department of Biostatistics, University of North Carolina at Chapel Hill

March 25, 2026



Scan for slides

Outline

Outline

- **Project 2:** Physical Distancing to Reduce Transmission of Influenza-Like-Illness on College Campuses: the eX-FLU Trial

Outline

- **Project 2:** Physical Distancing to Reduce Transmission of Influenza-Like-Illness on College Campuses: the eX-FLU Trial
- **Project 3:** Causal Inference from Cluster Randomized Trials with Differential Nonresponse

Outline

- **Project 2:** Physical Distancing to Reduce Transmission of Influenza-Like-Illness on College Campuses: the eX-FLU Trial
- **Project 3:** Causal Inference from Cluster Randomized Trials with Differential Nonresponse
- **Project 1:** Addressing Confounding and Continuous Exposure Measurement Error Using Corrected Score Functions

Project 2: Physical Distancing to Reduce Transmission of Influenza-Like-Illness on College Campuses: the eX-FLU Trial

Brian Richardson, Allison Aiello, Michael Hudgens

- **eX-FLU**: trial to evaluate a physical distancing intervention on a college campus during flu season ([Aiello et al., 2016](#); [Zivich et al., 2020](#))

Design and methods of a social network isolation study for reducing respiratory infection transmission: The eX-FLU cluster randomized trial

Allison E. Aiello^{a,*}, Amanda M. Simanek^{b,1}, Marisa C. Eisenberg^c, Alison R. Walsh^c, Brian Davis^c, Erik Volz^{d,1}, Caroline Cheng^c, Jeanette J. Rainey^e, Amra Uzicanin^e, Hongjiang Gao^e, Nathaniel Osgood^f, Dylan Knowles^f, Kevin Stanley^f, Kara Tarter^c, Arnold S. Monto^c

eX-FLU Trial

- **eX-FLU**: trial to evaluate a physical distancing intervention on a college campus during flu season ([Aiello et al., 2016](#); [Zivich et al., 2020](#))
- **Intervention**: encouragement to isolate in dorm for three days upon developing symptoms of **influenza-like illness** (ILI)

Design and methods of a social network isolation study for reducing respiratory infection transmission: The eX-FLU cluster randomized trial

Allison E. Aiello^{a,*}, Amanda M. Simanek^{b,1}, Marisa C. Eisenberg^c, Alison R. Walsh^c, Brian Davis^c, Erik Volz^{d,1}, Caroline Cheng^c, Jeanette J. Rainey^e, Amra Uzicanin^e, Hongjiang Gao^e, Nathaniel Osgood^f, Dylan Knowles^f, Kevin Stanley^f, Kara Tarter^c, Arnold S. Monto^c


eX-FLU Trial

- **eX-FLU**: trial to evaluate a physical distancing intervention on a college campus during flu season ([Aiello et al., 2016](#); [Zivich et al., 2020](#))
- **Intervention**: encouragement to isolate in dorm for three days upon developing symptoms of **influenza-like illness** (ILI)
- **Central question**: does the encouragement-to-isolate intervention reduce transmission of ILI?


Design and methods of a social network isolation study for reducing respiratory infection transmission: The eX-FLU cluster randomized trial

Allison E. Aiello^{a,*}, Amanda M. Simanek^{b,1}, Marisa C. Eisenberg^c, Alison R. Walsh^c, Brian Davis^c, Erik Volz^{d,1}, Caroline Cheng^c, Jeanette J. Rainey^e, Amra Uzicanin^e, Hongjiang Gao^e, Nathaniel Osgood^f, Dylan Knowles^f, Kevin Stanley^f, Kara Tarter^c, Arnold S. Monto^c


Transmission of Influenza-Like-Illness




Transmission of Influenza-Like-Illness




Transmission of Influenza-Like-Illness




Transmission of Influenza-Like-Illness




Example I: Intervention Affects Network and Transmission




Example I: Intervention Affects Network and Transmission




Example I: Intervention Affects Network and Transmission




Example II: Intervention Affects Network, Not Transmission




Example II: Intervention Affects Network, Not Transmission




Example II: Intervention Affects Network, Not Transmission




Example III: Intervention Affects Transmission, Not Network






Example III: Intervention Affects Transmission, Not Network



Example III: Intervention Affects Transmission, Not Network



Central Question: Does the Intervention Affect Transmission of ILI?




In Examples I and III, the answer is yes

eX-FLU Observed Data

- Baseline randomization assignments


$\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for

$Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$




eX-FLU Observed Data

- Baseline randomization assignments
 $\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for
 $Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$
 - ▶ randomization distribution $r(\mathbf{z})$




eX-FLU Observed Data

- Baseline randomization assignments
 $\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for
 $Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$
 - ▶ randomization distribution $r(\mathbf{z})$
- At weeks $k = 1, \dots, \tau$, we observe:



eX-FLU Observed Data

- Baseline randomization assignments
 $\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for
 $Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$
 - ▶ randomization distribution $r(\mathbf{z})$
- At weeks $k = 1, \dots, \tau$, we observe:
 - ▶ networks $\mathbf{A}^k = [A_{ij}^k]$, for
 $A_{ij}^k = \mathbb{1}(\text{students } i, j \text{ in contact at week } k)$



	1	2	3	4	5	6
1	0	1	0	1	1	0
2	1	0	0	0	1	0
3	0	0	0	0	1	0
4	1	0	0	0	1	0
5	1	1	1	1	0	0
6	0	0	0	0	0	0

eX-FLU Observed Data

- Baseline randomization assignments

$\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for

$Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$

- ▶ randomization distribution $r(\mathbf{z})$


- At weeks $k = 1, \dots, \tau$, we observe:

- ▶ networks $\mathbf{A}^k = [A_{ij}^k]$, for

$A_{ij}^k = \mathbb{1}(\text{students } i, j \text{ in contact at week } k)$


- ▶ infections $\mathbf{Y}^k = (Y_1^k, \dots, Y_n^k)$, for

$Y_i^k = \mathbb{1}(\text{student } i \text{ infected at week } k)$




eX-FLU Observed Data

- Baseline randomization assignments
 $\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for
 $Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$
 - ▶ randomization distribution $r(\mathbf{z})$
- At weeks $k = 1, \dots, \tau$, we observe:
 - ▶ networks $\mathbf{A}^k = [A_{ij}^k]$, for
 $A_{ij}^k = \mathbb{1}(\text{students } i, j \text{ in contact at week } k)$
 - ▶ infections $\mathbf{Y}^k = (Y_1^k, \dots, Y_n^k)$, for
 $Y_i^k = \mathbb{1}(\text{student } i \text{ infected at week } k)$
- network history: $\bar{\mathbf{A}} = \{\mathbf{A}^k\}_{k=1}^{\tau}$




...




eX-FLU Observed Data


- Baseline randomization assignments
 $\mathbf{Z} = (Z_1, \dots, Z_n) \in \mathcal{Z}$, for
 $Z_i = \mathbb{1}(\text{student } i \text{ gets intervention})$
 - ▶ randomization distribution $r(\mathbf{z})$
- At weeks $k = 1, \dots, \tau$, we observe:
 - ▶ networks $\mathbf{A}^k = [A_{ij}^k]$, for
 $A_{ij}^k = \mathbb{1}(\text{students } i, j \text{ in contact at week } k)$
 - ▶ infections $\mathbf{Y}^k = (Y_1^k, \dots, Y_n^k)$, for
 $Y_i^k = \mathbb{1}(\text{student } i \text{ infected at week } k)$
- network history: $\overline{\mathbf{A}} = \{\mathbf{A}^k\}_{k=1}^{\tau}$
- infection history: $\overline{\mathbf{Y}} = \{\mathbf{Y}^k\}_{k=1}^{\tau}$



eX-FLU Observed Data



eX-FLU Observed Data




Nodes

- uninfected student
- ✕ infected student

(93 out of 579 students with at least one infection)

eX-FLU Observed Data



Nodes


● uninfected student

× infected student

intervention group

control group

eX-FLU Observed Data




Nodes

- uninfected student
- × infected student
- intervention group
- control group

Edges

- between uninfected
- including infected control
- including infected intervention

eX-FLU Observed Data



eX-FLU Potential Outcomes

eX-FLU Potential Outcomes

Potential outcomes are defined for the networks and ILI infections:

$$\overline{\mathbf{A}}(\mathbf{z}), \quad \overline{\mathbf{Y}}(\mathbf{z}) \quad \text{for } \mathbf{z} \in \mathcal{Z}$$

eX-FLU Potential Outcomes

Potential outcomes are defined for the networks and ILI infections:

$$\overline{\mathbf{A}}(\mathbf{z}), \quad \overline{\mathbf{Y}}(\mathbf{z}) \quad \text{for } \mathbf{z} \in \mathcal{Z}$$

- $\overline{\mathbf{A}}(\mathbf{z})$: sequence of networks that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$

eX-FLU Potential Outcomes

Potential outcomes are defined for the networks and ILI infections:

$$\overline{\mathbf{A}}(\mathbf{z}), \quad \overline{\mathbf{Y}}(\mathbf{z}) \quad \text{for } \mathbf{z} \in \mathcal{Z}$$

- $\overline{\mathbf{A}}(\mathbf{z})$: sequence of networks that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$
- $\overline{\mathbf{Y}}(\mathbf{z})$: sequence of infection statuses that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$

eX-FLU Potential Outcomes

Potential outcomes are defined for the networks and ILI infections:

$$\overline{\mathbf{A}}(\mathbf{z}), \quad \overline{\mathbf{Y}}(\mathbf{z}) \quad \text{for } \mathbf{z} \in \mathcal{Z}$$

- $\overline{\mathbf{A}}(\mathbf{z})$: sequence of networks that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$
- $\overline{\mathbf{Y}}(\mathbf{z})$: sequence of infection statuses that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$
- Both of these depend on the entire vector $\mathbf{z}$
 - ▶ potential outcomes for one student depend on intervention assignment of other students (**interference**)
([Hudgens and Halloran, 2008](#))

eX-FLU Potential Outcomes

Potential outcomes are defined for the networks and ILI infections:

$$\overline{\mathbf{A}}(\mathbf{z}), \quad \overline{\mathbf{Y}}(\mathbf{z}) \quad \text{for } \mathbf{z} \in \mathcal{Z}$$


- $\overline{\mathbf{A}}(\mathbf{z})$: sequence of networks that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$
- $\overline{\mathbf{Y}}(\mathbf{z})$: sequence of infection statuses that **would** occur if students **were** encouraged to isolate according to $\mathbf{z}$
- Both of these depend on the entire vector $\mathbf{z}$
 - ▶ potential outcomes for one student depend on intervention assignment of other students (**interference**)
([Hudgens and Halloran, 2008](#))
- Assume **causal consistency**:

$$\overline{\mathbf{A}} = \overline{\mathbf{A}}(\mathbf{Z}) \text{ and } \overline{\mathbf{Y}} = \overline{\mathbf{Y}}(\mathbf{Z})$$

eX-FLU Null Hypotheses

$$H_0^Y : \overline{Y}(\mathbf{z}) = \overline{Y}(\tilde{\mathbf{z}}) \text{ for all } \mathbf{z}, \tilde{\mathbf{z}} \in \mathcal{Z}$$

(“the intervention has no effect on the infections”)




eX-FLU Null Hypotheses

$$H_0^Y : \bar{Y}(z) = \bar{Y}(\tilde{z}) \text{ for all } z, \tilde{z} \in \mathcal{Z}$$

(“the intervention has no effect on the infections”)

$$H_0^A : \bar{A}(z) = \bar{A}(\tilde{z}) \text{ for all } z, \tilde{z} \in \mathcal{Z}$$

(“the intervention has no effect the networks”)



eX-FLU Null Hypotheses

$$H_0^Y : \bar{Y}(z) = \bar{Y}(\tilde{z}) \text{ for all } z, \tilde{z} \in \mathcal{Z}$$


(“the intervention has no effect on the infections”)

$$H_0^A : \bar{A}(z) = \bar{A}(\tilde{z}) \text{ for all } z, \tilde{z} \in \mathcal{Z}$$

(“the intervention has no effect the networks”)

$$H_0^\# = H_0^Y \cap H_0^A : \bar{Y}(z) = \bar{Y}(\tilde{z}) \text{ and } \bar{A}(z) = \bar{A}(\tilde{z}) \text{ for all } z, \tilde{z} \in \mathcal{Z}$$

(“the intervention has no effect on the infections or on the networks”)



eX-FLU Null Hypotheses

$$H_0^Y : \overline{\mathbf{Y}}(\mathbf{z}) = \overline{\mathbf{Y}}(\tilde{\mathbf{z}}) \text{ for all } \mathbf{z}, \tilde{\mathbf{z}} \in \mathcal{Z}$$

(“the intervention has no effect on the infections”)

$$H_0^A : \overline{\mathbf{A}}(\mathbf{z}) = \overline{\mathbf{A}}(\tilde{\mathbf{z}}) \text{ for all } \mathbf{z}, \tilde{\mathbf{z}} \in \mathcal{Z}$$

(“the intervention has no effect the networks”)

$$H_0^\# = H_0^Y \cap H_0^A : \overline{\mathbf{Y}}(\mathbf{z}) = \overline{\mathbf{Y}}(\tilde{\mathbf{z}}) \text{ and } \overline{\mathbf{A}}(\mathbf{z}) = \overline{\mathbf{A}}(\tilde{\mathbf{z}}) \text{ for all } \mathbf{z}, \tilde{\mathbf{z}} \in \mathcal{Z}$$

(“the intervention has no effect on the infections or on the networks”)

Previous analyses of the eX-FLU trial ([Alexandria et al., 2023](#)) tested $H_0^\#$, but no analyses have tested H_0^Y

Testing $H_0^\#$

“How unlikely are the observed data under $H_0^\#$?”

Testing $H_0^\#$

“How unlikely are the observed data under $H_0^\#$?” We can answer this using a **randomization test** ([Fisher, 1936](#); [Alexandria et al., 2023](#); [Zhang and Zhao, 2023](#); [Ritzwoller et al., 2024](#))

Testing $H_0^\#$

“How unlikely are the observed data under $H_0^\#$?” We can answer this using a **randomization test** ([Fisher, 1936](#); [Alexandria et al., 2023](#); [Zhang and Zhao, 2023](#); [Ritzwoller et al., 2024](#))

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** ([Fisher, 1936](#); [Alexandria et al., 2023](#); [Zhang and Zhao, 2023](#); [Ritzwoller et al., 2024](#))

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** ([Fisher, 1936](#); [Alexandria et al., 2023](#); [Zhang and Zhao, 2023](#); [Ritzwoller et al., 2024](#))

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$			
$\mathbf{z}_2$			
$\dots$			
$\mathbf{z}_{ \mathcal{Z} }$			

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = ?$	$T(\mathbf{z}_1, ?, ?)$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = ?$	$T(\mathbf{z}_1, ?, ?)$
$\dots$	$\dots$	$\dots$	$\dots$
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$T(\mathbf{z}_{ \mathcal{Z} }, ?, ?)$

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$
 - ▶ This is possible since $\bar{\mathbf{A}}(\mathbf{z}), \bar{\mathbf{Y}}(\mathbf{z})$ are **imputable** under $H_0^\sharp$

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$

Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$
 - ▶ This is possible since $\bar{\mathbf{A}}(\mathbf{z}), \bar{\mathbf{Y}}(\mathbf{z})$ are **imputable** under $H_0^\sharp \implies H_0^\sharp$ is **sharp**

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{A}}$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$


Testing $H_0^\sharp$

“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$
 - ▶ This is possible since $\bar{\mathbf{A}}(\mathbf{z}), \bar{\mathbf{Y}}(\mathbf{z})$ are **imputable** under $H_0^\sharp \implies H_0^\sharp$ is **sharp**
- Compute a p-value: $\rho^\sharp = F_T(T|H_0^\sharp)$, where

$$F_T(t|H_0^\sharp) = \Pr(T \leq t|H_0^\sharp) = \sum_{\mathbf{z} \in \mathcal{Z}} r(\mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}}) \leq t\}$$

is the **cumulative distribution function** (CDF) of T



Testing $H_0^\sharp$


“How unlikely are the observed data under $H_0^\sharp$?” We can answer this using a **randomization test** (Fisher, 1936; Alexandria et al., 2023; Zhang and Zhao, 2023; Ritzwoller et al., 2024)

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under $H_0^\sharp$
 - ▶ This is possible since $\bar{\mathbf{A}}(\mathbf{z}), \bar{\mathbf{Y}}(\mathbf{z})$ are **imputable** under $H_0^\sharp \implies H_0^\sharp$ is **sharp**
- Compute a p-value: $\rho^\sharp = F_T(T|H_0^\sharp)$, where

$$F_T(t|H_0^\sharp) = \Pr(T \leq t|H_0^\sharp) = \sum_{\mathbf{z} \in \mathcal{Z}} r(\mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}}) \leq t\}$$

is the **cumulative distribution function** (CDF) of T

- Reject $H_0^\sharp$ if $\rho^\sharp \leq 0.05$



Testing $H_0^\sharp$

- This test of $H_0^\sharp$ controls the **type I error rate** exactly:

$$\Pr(\rho^\sharp \leq \alpha | H_0^\sharp) \leq \alpha$$

for any $\alpha \in [0, 1]$, for any sample size, and for any choice of test statistic

Testing $H_0^\sharp$

- This test of $H_0^\sharp$ controls the **type I error rate** exactly:

$$\Pr(\rho^\sharp \leq \alpha | H_0^\sharp) \leq \alpha$$

for any $\alpha \in [0, 1]$, for any sample size, and for any choice of test statistic


- The **power** of this test:

$$\Pr(\rho^\sharp \leq \alpha | H_1^\sharp)$$

depends on the choice of test statistic


Choice of Test Statistic

- 93 (out of 579) students with at least one infection



Choice of Test Statistic


- 93 (out of 579) students with at least one infection
- Bold edges represent possible transmission events



Choice of Test Statistic

- 93 (out of 579) students with at least one infection
- Bold edges represent possible transmission events
 - ▶ Proportion of possible transmission events attributable to students in the intervention group:

$$T = \frac{\text{number of yellow edges}}{\text{total number of edges}} = \frac{\sum_{k=2}^{\tau} \sum_{i=1}^n \sum_{j \neq i} Z_i Y_i^{k-1} A_{ij}^{k-1}}{\sum_{k=2}^{\tau} \sum_{i=1}^n \sum_{j \neq i} Y_i^{k-1} A_{ij}^{k-1}} = 0.359$$



Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$
- Find the distribution of T under H_0^Y

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$
- Find the distribution of T under H_0^Y

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$			
$\mathbf{z}_2$			
$\dots$			
$\mathbf{z}_{ \mathcal{Z} }$			

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \overline{\mathbf{A}}, \overline{\mathbf{Y}})$
- Find the distribution of T under H_0^Y

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\overline{\mathbf{A}}(\mathbf{z}_1) = ?$	$\overline{\mathbf{Y}}(\mathbf{z}_1) = ?$	$T(\mathbf{z}_1, ?, ?)$
$\mathbf{z}_2$	$\overline{\mathbf{A}}(\mathbf{z}_2) = ?$	$\overline{\mathbf{Y}}(\mathbf{z}_1) = ?$	$T(\mathbf{z}_1, ?, ?)$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\overline{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$\overline{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$T(\mathbf{z}_{ \mathcal{Z} }, ?, ?)$

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under H_0^Y

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, ?, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, ?, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, ?, \bar{\mathbf{Y}})$

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under H_0^Y
 - ▶ This is not possible since $\bar{\mathbf{A}}(\mathbf{z})$ are **not imputable** under H_0^Y

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, ?, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, ?, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, ?, \bar{\mathbf{Y}})$

Testing H_0^Y

“How unlikely are the observed data under H_0^Y ?”

- Choose a test statistic $T = T(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}})$
- Find the distribution of T under H_0^Y
 - ▶ This is not possible since $\bar{\mathbf{A}}(\mathbf{z})$ are **not imputable** under $H_0^Y \implies H_0^Y$ is **nonsharp**

Randomization	Networks	Infections	Test Statistic
$\mathbf{z}_1$	$\bar{\mathbf{A}}(\mathbf{z}_1) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_1) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_1, ?, \bar{\mathbf{Y}})$
$\mathbf{z}_2$	$\bar{\mathbf{A}}(\mathbf{z}_2) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_2) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_2, ?, \bar{\mathbf{Y}})$
...	...	...	...
$\mathbf{z}_{ \mathcal{Z} }$	$\bar{\mathbf{A}}(\mathbf{z}_{ \mathcal{Z} }) = ?$	$\bar{\mathbf{Y}}(\mathbf{z}_{ \mathcal{Z} }) = \bar{\mathbf{Y}}$	$T(\mathbf{z}_{ \mathcal{Z} }, ?, \bar{\mathbf{Y}})$

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**
 - ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**
 - ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$
 - ▶ Then

$$F_T(t|H_0^Y) = \Pr(T \leq t|H_0^Y) = \sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}} r(\mathbf{z}) q(\bar{\mathbf{a}}, \mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{a}}, \bar{\mathbf{Y}}) \leq t\}$$

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**
 - ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$
 - ▶ Then

$$F_T(t|H_0^Y) = \Pr(T \leq t|H_0^Y) = \sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}} r(\mathbf{z}) q(\bar{\mathbf{a}}, \mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{a}}, \bar{\mathbf{Y}}) \leq t\}$$

- ▶ p-value: $\rho^Y = F_T(T|H_0^Y)$

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**
 - ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$
 - ▶ Then

$$F_T(t|H_0^Y) = \Pr(T \leq t|H_0^Y) = \sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}} r(\mathbf{z}) q(\bar{\mathbf{a}}, \mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{a}}, \bar{\mathbf{Y}}) \leq t\}$$

- ▶ p-value: $\rho^Y = F_T(T|H_0^Y)$
- ▶ If/when q is unknown, estimate it

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y
- **Solution:** model the potential networks as **stochastic**
 - ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$
 - ▶ Then

$$F_T(t|H_0^Y) = \Pr(T \leq t|H_0^Y) = \sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}} r(\mathbf{z}) q(\bar{\mathbf{a}}, \mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{a}}, \bar{\mathbf{Y}}) \leq t\}$$

- ▶ p-value: $\rho^Y = F_T(T|H_0^Y)$
- ▶ If/when q is unknown, estimate it
 - ★ E.g., using a **Separable Temporal Exponential Random Graph Model**

Testing H_0^Y

- **Problem:** H_0^Y is nonsharp
 $\implies$ we don't know $F_T(t|H_0^Y)$ and thus cannot compute a p-value testing H_0^Y

- **Solution:** model the potential networks as **stochastic**

- ▶ Define $q(\bar{\mathbf{a}}, \mathbf{z}) = \Pr\{\bar{\mathbf{A}}(\mathbf{z}) = \bar{\mathbf{a}}\}$, the **probability mass function** (PMF) of $\bar{\mathbf{A}}(\mathbf{z})$
- ▶ Then

$$F_T(t|H_0^Y) = \Pr(T \leq t|H_0^Y) = \sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}} r(\mathbf{z}) q(\bar{\mathbf{a}}, \mathbf{z}) \mathbb{1}\{T(\mathbf{z}, \bar{\mathbf{a}}, \bar{\mathbf{Y}}) \leq t\}$$

- ▶ p-value: $\rho^Y = F_T(T|H_0^Y)$
- ▶ If/when q is unknown, estimate it
 - ★ E.g., using a **Separable Temporal Exponential Random Graph Model**
- ▶ If/when the double sum $\sum_{\mathbf{z} \in \mathcal{Z}} \sum_{\bar{\mathbf{a}} \in \mathcal{A}}$ cannot be computed exactly, approximate it using a Monte-Carlo method

Testing H_0^Y

Testing Procedure:

- 1 Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$

Testing H_0^Y

Testing Procedure:

- 1 Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- 2 Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B do:

Testing H_0^Y

Testing Procedure:

- ① Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- ② Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B **do**:
 - ① generate randomization vector $\mathbf{z}_b$ from $r(\mathbf{z})$

Testing H_0^Y

Testing Procedure:

- ① Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- ② Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B **do**:
 - ① generate randomization vector $\mathbf{z}_b$ from $r(\mathbf{z})$
 - ② generate networks $\bar{\mathbf{a}}_b$ from $q(\bar{\mathbf{a}}, \mathbf{z}_b; \hat{\theta})$

Testing H_0^Y

Testing Procedure:

- ① Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- ② Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B **do**:
 - ① generate randomization vector $\mathbf{z}_b$ from $r(\mathbf{z})$
 - ② generate networks $\bar{\mathbf{a}}_b$ from $q(\bar{\mathbf{a}}, \mathbf{z}_b; \hat{\theta})$
 - ③ compute the test statistic $T_b = T(\mathbf{z}_b, \bar{\mathbf{a}}_b, \mathbf{Y})$

Testing H_0^Y

Testing Procedure:

- ① Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- ② Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B do:
 - ① generate randomization vector $\mathbf{z}_b$ from $r(\mathbf{z})$
 - ② generate networks $\bar{\mathbf{a}}_b$ from $q(\bar{\mathbf{a}}, \mathbf{z}_b; \hat{\theta})$
 - ③ compute the test statistic $T_b = T(\mathbf{z}_b, \bar{\mathbf{a}}_b, \mathbf{Y})$
- ③ compute the plug-in p-value testing H_0^Y :

$$\hat{\rho}_B^Y = B^{-1} \sum_{b=1}^B \mathbb{1}(T_b \leq T)$$

Testing H_0^Y

Testing Procedure:

- ① Estimate the unknown parameter θ in the PMF $q(\bar{\mathbf{a}}, \mathbf{z}; \theta)$ of $\bar{\mathbf{A}}(\mathbf{z})$
- ② Approximate $F_T(t|H_0^Y)$ using B Monte-Carlo replicates:
for $b = 1$ to B do:
 - ① generate randomization vector $\mathbf{z}_b$ from $r(\mathbf{z})$
 - ② generate networks $\bar{\mathbf{a}}_b$ from $q(\bar{\mathbf{a}}, \mathbf{z}_b; \hat{\theta})$
 - ③ compute the test statistic $T_b = T(\mathbf{z}_b, \bar{\mathbf{a}}_b, \mathbf{Y})$
- ③ compute the plug-in p-value testing H_0^Y :

$$\hat{\rho}_B^Y = B^{-1} \sum_{b=1}^B \mathbb{1}(T_b \leq T)$$


Proposition: This procedure will **asymptotically** control the type I error rate

eX-FLU: Hypothesis Test Results

- Test statistic (proportion of possible transmission events attributable to students in the intervention group):
 $T = 0.359$.

eX-FLU: Hypothesis Test Results


- Test statistic (proportion of possible transmission events attributable to students in the intervention group):
 $T = 0.359$.



- Encouragement to isolate affects the social network and/or influenza-like illness ($\rho_B^\# = 0.0466$)

eX-FLU: Hypothesis Test Results

- Test statistic (proportion of possible transmission events attributable to students in the intervention group):
 $T = 0.359$.



- Encouragement to isolate affects the social network and/or influenza-like illness ($\rho_B^\# = 0.0466$)
- Encouragement to isolate specifically affects influenza-like illness ($\hat{\rho}_B^Y = 0.0367$)

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
 - ★ Using mismeasured networks does not hurt the type I error, but may hurt power
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
 - ★ Using mismeasured networks does not hurt the type I error, but may hurt power
 - ★ Use Bluetooth data from eX-FLU sub-study
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
 - ★ Using mismeasured networks does not hurt the type I error, but may hurt power
 - ★ Use Bluetooth data from eX-FLU sub-study
 - ▶ Focus on **estimation** as opposed to **hypothesis testing**
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
 - ★ Using mismeasured networks does not hurt the type I error, but may hurt power
 - ★ Use Bluetooth data from eX-FLU sub-study
 - ▶ Focus on **estimation** as opposed to **hypothesis testing**
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Summary of Project 2

- Developed a new randomization-based inference procedure
 - ▶ Established theoretical properties (asymptotic control of Type I error)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **eX-FLU trial** to find a protective effect of an encouragement-to-isolate intervention on a college campus
- **Future directions:**
 - ▶ Account for **measurement error** in the self-reported social networks
 - ★ Using mismeasured networks does not hurt the type I error, but may hurt power
 - ★ Use Bluetooth data from eX-FLU sub-study
 - ▶ Focus on **estimation** as opposed to **hypothesis testing**
 - ▶ **Design future trials** with independent clusters
 - ★ can allow **identification** of more causal estimands
- **Status:** under revision at *Proceedings of the National Academy of Sciences*

Project 3 : Causal Inference from Cluster Randomized Trials with Differential Nonresponse

Brian Richardson, Bonnie Shook-Sa, Michael Hudgens

PopART Trial

- **PopART (HPTN 071) Trial:** large **cluster randomized trial** designed to evaluate a combination **HIV prevention intervention** in Zambia and South Africa ([Hayes et al., 2014](#))

Hayes et al. *Trials* 2014, **15**:57
<http://www.trialsjournal.com/content/15/1/57>



STUDY PROTOCOL

Open Access

HPTN 071 (PopART): Rationale and design of a cluster-randomised trial of the population impact of an HIV combination prevention intervention including universal testing and treatment – a study protocol for a cluster randomised trial

Richard Hayes¹, Helen Ayles^{2,3}, Nulda Beyers⁴, Kalpana Sabapathy^{1*}, Sian Floyd¹, Kwame Shanaube³, Peter Bock⁴, Sam Griffith⁵, Ayana Moore⁵, Deborah Watson-Jones², Christophe Fraser⁶, Sten H Vermund⁷, Sarah Fidler⁸ and The HPTN 071 (PopART) Study Team

PopART Trial

- **PopART (HPTN 071) Trial:** large **cluster randomized trial** designed to evaluate a combination **HIV prevention intervention** in Zambia and South Africa ([Hayes et al., 2014](#))
- **Intervention:**
 - ▶ Arm A: annual home-based HIV testing, promotion of medical male circumcision for HIV-negative men, and offer of immediate ART for those testing HIV-positive
 - ▶ Arm B: same as Arm A, but ART initiation following national guidelines
 - ▶ Arm C: standard of care

Hayes et al. *Trials* 2014, **15**:57
<http://www.trialsjournal.com/content/15/1/57>



STUDY PROTOCOL

Open Access

HPTN 071 (PopART): Rationale and design of a cluster-randomised trial of the population impact of an HIV combination prevention intervention including universal testing and treatment – a study protocol for a cluster randomised trial

Richard Hayes¹, Helen Ayles^{2,3}, Nulda Beyers⁴, Kalpana Sabapathy^{1*}, Sian Floyd¹, Kwame Shanaube³, Peter Bock⁴, Sam Griffith⁵, Ayana Moore⁵, Deborah Watson-Jones², Christophe Fraser⁶, Sten H Vermund⁷, Sarah Fidler⁸ and The HPTN 071 (PopART) Study Team

PopART Trial

- **PopART (HPTN 071) Trial:** large **cluster randomized trial** designed to evaluate a combination **HIV prevention intervention** in Zambia and South Africa ([Hayes et al., 2014](#))
- **Intervention:**
 - ▶ Arm A: annual home-based HIV testing, promotion of medical male circumcision for HIV-negative men, and offer of immediate ART for those testing HIV-positive
 - ▶ Arm B: same as Arm A, but ART initiation following national guidelines
 - ▶ Arm C: standard of care
- **Outcome:** HIV Incidence

Hayes et al. *Trials* 2014, **15**:57
<http://www.trialsjournal.com/content/15/1/57>



STUDY PROTOCOL

Open Access

HPTN 071 (PopART): Rationale and design of a cluster-randomised trial of the population impact of an HIV combination prevention intervention including universal testing and treatment – a study protocol for a cluster randomised trial

Richard Hayes¹, Helen Ayles^{2,3}, Nulda Beyers⁴, Kalpana Sabapathy^{1*}, Sian Floyd¹, Kwame Shanaube³, Peter Bock⁴, Sam Griffith⁵, Ayana Moore⁵, Deborah Watson-Jones², Christophe Fraser⁶, Sten H Vermund⁷, Sarah Fidler⁸ and The HPTN 071 (PopART) Study Team

PopART Trial

- **Initial Findings:** no protective effect of the Arm A intervention ([Hayes et al., 2019](#))
 - ▶ “unanticipated and not consistent with the data on viral suppression”



Effect of Universal Testing and Treatment on HIV Incidence — HPTN 071 (PopART)

R.J. Hayes, D. Donnell, S. Floyd, N. Mandla, J. Bwalya, K. Sabapathy, B. Yang, M. Phiri, A. Schaap, S.H. Eshleman, E. Piwowar-Manning, B. Kosloff, A. James, T. Skalland, E. Wilson, L. Emel, D. Macleod, R. Dunbar, M. Simwinga, N. Makola, V. Bond, G. Hoddinott, A. Moore, S. Griffith, N. Deshmane Sista, S.H. Vermund, W. El-Sadr, D.N. Burns, J.R. Hargreaves, K. Hauck, C. Fraser, K. Shanaube, P. Bock, N. Beyers, H. Ayles, and S. Fidler, for the HPTN 071 (PopART) Study Team

PopART Trial

- **Initial Findings:** no protective effect of the Arm A intervention ([Hayes et al., 2019](#))
 - ▶ “unanticipated and not consistent with the data on viral suppression”
- **Later Analyses:** identified possibly differential nonresponse that could bias trial results ([Shook-Sa et al., 2025](#))
 - ▶ Males were less likely to respond (i.e., have data collected)
 - ▶ The intervention appeared more effective among males



Effect of Universal Testing and Treatment on HIV Incidence — HPTN 071 (PopART)

R.J. Hayes, D. Donnell, S. Floyd, N. Mandla, J. Bwalya, K. Sabapathy, B. Yang, M. Phiri, A. Schaap, S.H. Eshleman, E. Piwowar-Manning, B. Kosloff, A. James, T. Skalland, E. Wilson, L. Emel, D. Macleod, R. Dunbar, M. Simwinga, N. Makola, V. Bond, G. Hoddinott, A. Moore, S. Griffith, N. Deshmane Sista, S.H. Vermund, W. El-Sadr, D.N. Burns, J.R. Hargreaves, K. Hauck, C. Fraser, K. Shanaube, P. Bock, N. Beyers, H. Ayles, and S. Fidler, for the HPTN 071 (PopART) Study Team

Goal

To estimate the **causal effect** of a cluster-level exposure on an outcome under **differential nonresponse**

Goal

To estimate the **causal effect** of a cluster-level exposure on an outcome under **differential nonresponse**

- i.e., nonresponse that depends on missing covariates and on treatment

Goal

To estimate the **causal effect** of a cluster-level exposure on an outcome under **differential nonresponse**


- i.e., nonresponse that depends on missing covariates and on treatment

(also under outcome censoring)


(Non)Participation




(Non)Participation




(Non)Participation




(Non)Participation




(Non)Participation



(Non)Participation



(Non)Participation




(Non)Response




(Non)Response




(Non)Response




(Non)Response




(Non)Response



(Non)Response




(Non)Response




PopART Observed Data


Control Treatment



PopART Observed Data




PopART Observed Data




- $\mathbf{X}_i$: cluster covariates

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator
- Y_{ij} : outcome

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator
- Y_{ij} : outcome

PopART Observed Data




- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator
- Y_{ij} : outcome

PopART Observed Data



- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator
- Y_{ij} : outcome

PopART Observed Data



- $\mathbf{X}_i$: cluster covariates
- A_i : cluster treatment
- S_{ij} : trial membership
- R_{ij} : response indicator
- $\mathbf{W}_{ij}$: individual covariates
- C_{ij} : censoring indicator
- Y_{ij} : outcome

Causal Estimand

- **Potential Outcome:** $Y_{ij}(a)$: potential outcome for individual j in cluster i , under cluster-level exposure a

Causal Estimand

- **Potential Outcome:** $Y_{ij}(a)$: potential outcome for individual j in cluster i , under cluster-level exposure a
- **Expected Potential Outcome:** $\eta(a) = E\{Y_{ij}(a)\}$

Causal Estimand

- **Potential Outcome:** $Y_{ij}(a)$: potential outcome for individual j in cluster i , under cluster-level exposure a
- **Expected Potential Outcome:** $\eta(a) = E\{Y_{ij}(a)\}$
- **Causal Contrasts:**

$$RD = \eta(1) - \eta(0),$$

Causal Estimand

- **Potential Outcome:** $Y_{ij}(a)$: potential outcome for individual j in cluster i , under cluster-level exposure a
- **Expected Potential Outcome:** $\eta(a) = E\{Y_{ij}(a)\}$
- **Causal Contrasts:**

$$RD = \eta(1) - \eta(0), \quad RR = \frac{\eta(1)}{\eta(0)},$$


Causal Estimand

- **Potential Outcome:** $Y_{ij}(a)$: potential outcome for individual j in cluster i , under cluster-level exposure a
- **Expected Potential Outcome:** $\eta(a) = E\{Y_{ij}(a)\}$
- **Causal Contrasts:**

$$RD = \eta(1) - \eta(0), \quad RR = \frac{\eta(1)}{\eta(0)}, \quad OR = \frac{\eta(1)/\{1 - \eta(1)\}}{\eta(0)/\{1 - \eta(0)\}}$$

Identification


- Under assumptions*, the estimand $\eta(a)$ can be identified using either:



Identification

- Under assumptions*, the estimand $\eta(a)$ can be identified using either:
- G-formula:

$$\eta(a) = E \{E(Y_{ij} | A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0, \mathbf{X}_i, \mathbf{W}_{ij}) | S_{ij} = 0\}$$




Identification

- Under assumptions*, the estimand $\eta(a)$ can be identified using either:
- G-formula:

$$\eta(a) = E \{ E(Y_{ij} | A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0, \mathbf{X}_i, \mathbf{W}_{ij}) | S_{ij} = 0 \}$$

- Inverse Probability Weighting:

$$\eta(a) = E \left\{ \frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}{\Pr(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i, \mathbf{W}_{ij})} \right\}$$



Estimation: G-Formula

- 1 Fit the **outcome regression model** among **uncensored responders**

$$\mu_a(\mathbf{x}, \mathbf{w}) = E(Y_{ij} | R_{ij} = 1, C_{ij} = 0, A_i = a, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})$$

Estimation: G-Formula

- 1 Fit the **outcome regression model** among **uncensored responders**

$$\mu_a(\mathbf{x}, \mathbf{w}) = \underbrace{E(Y_{ij})}_{\text{expected outcome}} \mid R_{ij} = 1, C_{ij} = 0, A_i = a, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})$$

Estimation: G-Formula

- 1 Fit the **outcome regression model** among **uncensored responders**

$$\mu_a(\mathbf{x}, \mathbf{w}) = \underbrace{E(Y_{ij})}_{\text{expected outcome}} \mid \underbrace{R_{ij} = 1, C_{ij} = 0}_{\text{among uncensored responders}}, A_i = a, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})$$

Estimation: G-Formula

- 1 Fit the **outcome regression model** among **uncensored responders**

$$\mu_a(\mathbf{x}, \mathbf{w}) = \underbrace{E(Y_{ij})}_{\text{expected outcome}} \mid \underbrace{R_{ij} = 1, C_{ij} = 0,}_{\text{among uncensored responders}} \underbrace{A_i = a, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given treatment and covariates}}$$

Estimation: G-Formula

- 1 Fit the **outcome regression model** among **uncensored responders**

$$\mu_a(\mathbf{x}, \mathbf{w}) = \underbrace{E(Y_{ij})}_{\text{expected outcome}} \mid \underbrace{R_{ij} = 1, C_{ij} = 0}_{\text{among uncensored responders}} \underbrace{A_i = a, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given treatment and covariates}}$$

- 2 Marginalize the estimator $\hat{\mu}_a(\mathbf{X}, \mathbf{W})$ over the distribution of covariates in the **auxiliary data**:

$$\hat{\eta}^{GF}(a) = \underbrace{\frac{1}{n_{\text{aux}}} \sum_{i=1}^m \sum_{j=1}^{n_i} \mathbb{1}(S_{ij} = 0) \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\text{sample mean in auxiliary data}}$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w} \right)$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

- 2 Take the **weighted mean outcome**

$$\hat{\eta}^{IPW}(a) = \frac{1}{n^{\text{trial}}} \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}{\hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

- 2 Take the **weighted mean outcome**

$$\hat{\eta}^{IPW}(a) = \frac{1}{n^{\text{trial}}} \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\overbrace{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}^{\text{outcomes among uncensored responders with treatment } a}}{\hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

- 2 Take the **weighted mean outcome**

$$\hat{\eta}^{IPW}(a) = \frac{1}{n^{\text{trial}}} \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\overbrace{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}^{\text{outcomes among uncensored responders with treatment } a}}{\underbrace{\hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\text{inverse probability weights}}}$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

- 2 Take the **weighted mean outcome**

$$\hat{\eta}^{IPW}(a) = \underbrace{\frac{1}{n^{\text{trial}}}}_{\text{trial size}} \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\overbrace{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}^{\text{outcomes among uncensored responders with treatment } a}}{\underbrace{\hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\text{inverse probability weights}}}$$

Estimation: Inverse Probability Weighting

- 1 Fit the **joint propensity score model**

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr \left(\underbrace{(A_i = a, R_{ij} = 1, C_{ij} = 0)}_{\text{treatment } a \text{ and uncensored response}} \mid \underbrace{\mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}}_{\text{given covariates}} \right)$$

- 2 Take the **weighted mean outcome**

$$\hat{\eta}^{IPW}(a) = \underbrace{\frac{1}{\hat{n}^{\text{trial}}}}_{\text{Hájek estimator}} \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\overbrace{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0) Y_{ij}}^{\text{outcomes among uncensored responders with treatment } a}}{\underbrace{\hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\text{inverse probability weights}}}$$

Joint Propensity Scores

$$\pi_a(\mathbf{x}, \mathbf{w}) = \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})$$

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \times \Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})\end{aligned}$$

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})\end{aligned}$$

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})\end{aligned}$$

Response and Treatment Mechanism: $\pi_a^{AR}(\mathbf{x}, \mathbf{w})$

$$= \Pr(A_i = 1, R_{ij} = 1) \frac{f_{\mathbf{X}, \mathbf{W} | A, R}(\mathbf{x}, \mathbf{w} | a, 1)}{f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w})}$$

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})\end{aligned}$$

Response and Treatment Mechanism: $\pi_a^{AR}(\mathbf{x}, \mathbf{w})$

$$= \Pr(A_i = 1, R_{ij} = 1) \frac{f_{\mathbf{X}, \mathbf{W} | A, R}(\mathbf{x}, \mathbf{w} | a, 1)}{f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w})}$$

- Estimate this ratio of densities following an approach in [Sugiyama et al. \(2010\)](#)

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \underbrace{\Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^C(\mathbf{x}, \mathbf{w})}\end{aligned}$$

Response and Treatment Mechanism: $\pi_a^{AR}(\mathbf{x}, \mathbf{w})$

$$= \Pr(A_i = 1, R_{ij} = 1) \frac{f_{\mathbf{X}, \mathbf{W} | A, R}(\mathbf{x}, \mathbf{w} | a, 1)}{f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w})}$$

- Estimate this ratio of densities following an approach in [Sugiyama et al. \(2010\)](#)

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \underbrace{\Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^C(\mathbf{x}, \mathbf{w})}\end{aligned}$$

Response and Treatment Mechanism: $\pi_a^{AR}(\mathbf{x}, \mathbf{w})$

$$= \Pr(A_i = 1, R_{ij} = 1) \frac{f_{\mathbf{X}, \mathbf{W} | A, R}(\mathbf{x}, \mathbf{w} | a, 1)}{f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w})}$$

Censoring Mechanism: $\pi_a^C(\mathbf{x}, \mathbf{w})$

- Estimate this ratio of densities following an approach in [Sugiyama et al. \(2010\)](#)

Joint Propensity Scores

$$\begin{aligned}\pi_a(\mathbf{x}, \mathbf{w}) &= \Pr(A_i = a, R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w}) \\ &= \underbrace{\Pr(A_i = a, R_{ij} = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^{AR}(\mathbf{x}, \mathbf{w})} \times \underbrace{\Pr(C_{ij} = 0 | A_i = a, R_{ij} = 1, \mathbf{X}_i = \mathbf{x}, \mathbf{W}_{ij} = \mathbf{w})}_{\equiv \pi_a^C(\mathbf{x}, \mathbf{w})}\end{aligned}$$

Response and Treatment Mechanism: $\pi_a^{AR}(\mathbf{x}, \mathbf{w})$

$$= \Pr(A_i = 1, R_{ij} = 1) \frac{f_{\mathbf{X}, \mathbf{W} | A, R}(\mathbf{x}, \mathbf{w} | a, 1)}{f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w})}$$

- Estimate this ratio of densities following an approach in Sugiyama et al. (2010)

Censoring Mechanism: $\pi_a^C(\mathbf{x}, \mathbf{w})$

- Estimate using responders in the trial (for whom $A_i, \mathbf{X}_i, \mathbf{W}_{ij}, C_{ij}$ are available)

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a
- 3 AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \left[\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{n}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})} \{Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})\} + \frac{\mathbb{1}(S_{ij} = 0)}{n^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij}) \right]$$

Estimation: Augmented Inverse Probability Weighting

- ① Fit the **outcome regression model** μ_a
- ② Fit the **joint propensity score model** π_a
- ③ AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \left[\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{n}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})} \{Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})\} + \frac{\mathbb{1}(S_{ij} = 0)}{n^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij}) \right]$$

$\Rightarrow$ **Double robustness** to parametric model specifications for μ_a, π_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a
- 3 AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \left[\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{n}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})} \{Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})\} + \underbrace{\frac{\mathbb{1}(S_{ij} = 0)}{n^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\hat{\eta}^{GF}(a)} \right]$$

$\Rightarrow$ **Double robustness** to parametric model specifications for μ_a, π_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a
- 3 AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \left[\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{n}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})} \underbrace{\{Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})\}}_{\rightarrow 0} + \underbrace{\frac{\mathbb{1}(S_{ij} = 0)}{\hat{n}^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\hat{\eta}^{GF}(a)} \right]$$

$\Rightarrow$ **Double robustness** to parametric model specifications for μ_a, π_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a
- 3 AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \underbrace{\left[\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{n}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})} \{Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})\} + \frac{\mathbb{1}(S_{ij} = 0)}{n^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij}) \right]}_{\hat{\eta}^{IPW}(a)}$$

$\Rightarrow$ **Double robustness** to parametric model specifications for μ_a, π_a

Estimation: Augmented Inverse Probability Weighting

- 1 Fit the **outcome regression model** μ_a
- 2 Fit the **joint propensity score model** π_a
- 3 AIPW estimator: $\hat{\eta}^{AIPW}(a) =$

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \left[\underbrace{\frac{\mathbb{1}(A_i = a, S_{ij} = 1, R_{ij} = 1, C_{ij} = 0)}{\hat{\eta}^{\text{trial}} \hat{\pi}_a(\mathbf{X}_i, \mathbf{W}_{ij})}}_{\hat{\eta}^{IPW}(a)} \{ Y_{ij} - \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij}) \} + \underbrace{\frac{\mathbb{1}(S_{ij} = 0)}{n^{\text{aux}}} \hat{\mu}_a(\mathbf{X}_i, \mathbf{W}_{ij})}_{\rightarrow 0} \right]$$

$\Rightarrow$ **Double robustness** to parametric model specifications for μ_a, π_a

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\quad ; \quad \right)$$

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \quad \right)$$

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

$$E\{\underbrace{\Psi(\mathbf{D}_{ij}; \theta_0)}_{\text{true}}\} = \mathbf{0}$$

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

$$E\{\underbrace{\Psi(\mathbf{D}_{ij}; \theta_0)}_{\text{true}}\} = \mathbf{0}$$

The M-estimator $\hat{\theta}$ is the solution to

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

$$E\{\underbrace{\Psi(\mathbf{D}_{ij}; \theta_0)}_{\text{true}}\} = \mathbf{0}$$

The M-estimator $\hat{\theta}$ is the solution to

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \Psi(\mathbf{D}_{ij}; \theta) = \mathbf{0}$$

Crash Course on M-Estimation

Estimating (Score) Function: a function of the observed data and the parameter of interest

$$\underbrace{\Psi}_{\text{score fun.}} \left(\underbrace{\mathbf{D}_{ij}}_{\text{data}} ; \underbrace{\theta}_{\text{param.}} \right)$$

that is **unbiased**, meaning

$$E\{\underbrace{\Psi(\mathbf{D}_{ij}; \theta_0)}_{\text{true}}\} = \mathbf{0}$$

The M-estimator $\hat{\theta}$ is the solution to

$$\sum_{i=1}^m \sum_{j=1}^{n_i} \Psi(\mathbf{D}_{ij}; \theta) = \mathbf{0}$$

$\hat{\theta}$ is consistent and asymptotically normal and has a simple variance estimator ([Stefanski and Boos, 2002](#))

Inference: Estimating Equations

Each of the proposed estimators can be expressed as an **M-estimator**:

Inference: Estimating Equations

Each of the proposed estimators can be expressed as an **M-estimator**:

$\hat{\eta}^{GF}$ has estimating function

$$\Psi_{GF}(\mathbf{D}_{ij}; \theta_{GF}) = \begin{bmatrix} \Psi_{\mu}(\mathbf{D}_{ij}, \beta) \\ \frac{1}{p^{\text{aux}}} \mathbb{1}(S_{ij} = 0) \mu_0(\mathbf{X}_i, \mathbf{W}_{ij}; \beta) - \eta(0) \\ \frac{1}{p^{\text{aux}}} \mathbb{1}(S_{ij} = 1) \mu_1(\mathbf{X}_i, \mathbf{W}_{ij}; \beta) - \eta(1) \end{bmatrix},$$

where

- Ψ_{μ} is an estimating function for fitting the outcome model $\mu_a(\mathbf{x}, \mathbf{w}; \beta)$
- $\theta_{GF} = [\beta, \eta(0), \eta(1)]$
- $p^{\text{aux}} = n^{\text{aux}}/n$

Inference: Estimating Equations

Each of the proposed estimators can be expressed as an **M-estimator**:

$\hat{\eta}^{IPW}$ has estimating function

$$\Psi_{IPW}(\mathbf{D}_{ij}; \theta_{IPW}) = \begin{bmatrix} \Psi_{\pi}(\mathbf{D}_{ij}, \gamma) \\ \frac{1}{p^{\text{trial}}} \frac{\mathbb{1}(A_i=0, S_{ij}=1, R_{ij}=1, C_{ij}=0) Y_{ij}}{\hat{\pi}_0(\mathbf{x}_i, \mathbf{w}_{ij}; \gamma)} - \eta(0) \\ \frac{1}{p^{\text{trial}}} \frac{\mathbb{1}(A_i=1, S_{ij}=1, R_{ij}=1, C_{ij}=0) Y_{ij}}{\hat{\pi}_1(\mathbf{x}_i, \mathbf{w}_{ij}; \gamma)} - \eta(1) \end{bmatrix},$$

where

- Ψ_{π} is an estimating function for fitting the joint propensity score model $\pi_a(\mathbf{x}, \mathbf{w}; \gamma)$
- $\theta_{IPW} = [\gamma, \eta(0), \eta(1)]$
- $p^{\text{trial}} = n^{\text{trial}}/n$

Inference: Estimating Equations

Each of the proposed estimators can be expressed as an **M-estimator**:

$\hat{\eta}^{AIPW}$ has estimating function

$$\Psi_{AIPW}(\mathbf{D}_{ij}; \theta_{AIPW}) = \begin{bmatrix} \Psi_{\mu}(\mathbf{D}_{ij}, \beta) \\ \Psi_{\pi}(\mathbf{D}_{ij}, \gamma) \\ \frac{\mathbb{1}(A_i=0, S_{ij}=1, R_{ij}=1, C_{ij}=0)}{p^{\text{trial}} \pi_0(\mathbf{X}_i, \mathbf{W}_{ij}; \gamma)} \{Y_{ij} - \mu_0(\mathbf{X}_i, \mathbf{W}_{ij}; \beta)\} + \frac{\mathbb{1}(S_{ij}=0)}{p^{\text{aux}}} \mu_0(\mathbf{X}_i, \mathbf{W}_{ij}; \beta) - \eta(0) \\ \frac{\mathbb{1}(A_i=1, S_{ij}=1, R_{ij}=1, C_{ij}=0)}{p^{\text{trial}} \pi_1(\mathbf{X}_i, \mathbf{W}_{ij}; \gamma)} \{Y_{ij} - \mu_1(\mathbf{X}_i, \mathbf{W}_{ij}; \beta)\} + \frac{\mathbb{1}(S_{ij}=0)}{p^{\text{aux}}} \mu_1(\mathbf{X}_i, \mathbf{W}_{ij}; \beta) - \eta(1) \end{bmatrix},$$

where


$$\bullet \theta_{AIPW} = [\beta, \gamma, \eta(0), \eta(1)]$$

Simulation Study



Simulation Study

$$X \sim N(0,1)$$




Simulation Study

$$X \sim N(0,1)$$

$$W_1 \sim \text{Bernoulli}(0.5)$$

$$W_2 \sim N(0,1)$$




Simulation Study

$$X \sim N(0,1)$$

$$W_1 \sim \text{Bernoulli}(0.5)$$

$$W_2 \sim N(0,1)$$




Treatment marginally protective,
but harmful when $W_1 = 1$

Simulation Study

$$X \sim N(0,1)$$

$$W_1 \sim \text{Bernoulli}(0.5)$$

$$W_2 \sim N(0,1)$$




Treatment marginally protective,
but harmful when $W_1 = 1$


Invited Trial Sample,
size $n^{trial} \in \{500, 5000\}$




Simulation Study




Simulation Study




Simulation Study




Simulation Study









$\hat{RD}$




RD




RD



RD





RB





Auxiliary and Trial Sample Sizes

Inconsistent if μ_a is incorrect














consistent if π_a
is correct














Estimand ● $\eta(0)$ ▲ $\eta(1)$ ■ RD + RR

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014](#), [2019](#))

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014](#), [2019](#))
- Auxiliary data:

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014, 2019](#))
- Auxiliary data:
 - ▶ 2016 Zambia Population-based HIV Impact Assessment survey (ZAMPHIA) ([Ministry of Health Zambia, 2019](#))

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014, 2019](#))
- Auxiliary data:
 - ▶ 2016 Zambia Population-based HIV Impact Assessment survey (ZAMPHIA) ([Ministry of Health Zambia, 2019](#))
 - ▶ 2017 South African National HIV Prevalence, Incidence, Behaviour and Communication Survey V (SABSSMV) ([Simbayi et al., 2021](#))

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014, 2019](#))
- Auxiliary data:
 - ▶ 2016 Zambia Population-based HIV Impact Assessment survey (ZAMPHIA) ([Ministry of Health Zambia, 2019](#))
 - ▶ 2017 South African National HIV Prevalence, Incidence, Behaviour and Communication Survey V (SABSSMV) ([Simbayi et al., 2021](#))
 - ▶ Both restricted to match age range, provinces, urbanicity, and HIV status of the PopART trial

PopART Application

- Trial data: PopART trial ([Hayes et al., 2014, 2019](#))
- Auxiliary data:
 - ▶ 2016 Zambia Population-based HIV Impact Assessment survey (ZAMPHIA) ([Ministry of Health Zambia, 2019](#))
 - ▶ 2017 South African National HIV Prevalence, Incidence, Behaviour and Communication Survey V (SABSSMV) ([Simbayi et al., 2021](#))
 - ▶ Both restricted to match age range, provinces, urbanicity, and HIV status of the PopART trial


	PopART Trial Responders				Auxiliary Data
	Overall <i>N</i> = 29,130	A <i>N</i> = 9,594	B <i>N</i> = 10,235	C (Control) <i>N</i> = 9,301	<i>N</i> = 5,076
Sex and Male Circumcision					
Female	19,494 (68%)	6,547 (69%)	6,828 (68%)	6,119 (68%)	54%
Male (Circumcised)	4,190 (15%)	1,403 (15%)	1,516 (15%)	1,271 (14%)	22%
Male (Uncircumcised)	4,847 (17%)	1,474 (16%)	1,725 (17%)	1,648 (18%)	24%

PopART Application


- Trial data: PopART trial ([Hayes et al., 2014, 2019](#))
- Auxiliary data:
 - ▶ 2016 Zambia Population-based HIV Impact Assessment survey (ZAMPHIA) ([Ministry of Health Zambia, 2019](#))
 - ▶ 2017 South African National HIV Prevalence, Incidence, Behaviour and Communication Survey V (SABSSMV) ([Simbayi et al., 2021](#))
 - ▶ Both restricted to match age range, provinces, urbanicity, and HIV status of the PopART trial

	PopART Trial Responders				Auxiliary Data
	Overall <i>N</i> = 29,130	A <i>N</i> = 9,594	B <i>N</i> = 10,235	C (Control) <i>N</i> = 9,301	<i>N</i> = 5,076
Sex and Male Circumcision					
Female	19,494 (68%)	6,547 (69%)	6,828 (68%)	6,119 (68%)	54%
Male (Circumcised)	4,190 (15%)	1,403 (15%)	1,516 (15%)	1,271 (14%)	22%
Male (Uncircumcised)	4,847 (17%)	1,474 (16%)	1,725 (17%)	1,648 (18%)	24%


PopART Results




PopART Results



PopART Results



PopART Results




Summary of Project 3

- Developed estimators to adjust for differential nonresponse and outcome censoring in cluster RCT
 - ▶ Established theoretical properties (consistency, asymptotic normality, double robustness of AIPW)
 - ▶ Demonstrated empirical performance through simulations
- Applied the method to the **PopART trial**, along with auxiliary data from national surveys
- **Future Directions:**
 - ▶ show semiparametric efficiency of AIPW estimator
 - ▶ derive asymptotic results for AIPW estimator given nonparametric estimators of μ_a, π_a
- **Status:** in preparation to submit to *Annals of Applied Statistics*

Project 1 : Addressing Confounding and Continuous Exposure Measurement Error Using Corrected Score Functions


Brian Richardson, Brian Blette, Peter Gilbert, Michael Hudgens (2025)

Motivation: HVTN 505 trial



- **HVTN 505 trial:** trial of a preventive HIV vaccine

Motivation: HVTN 505 trial



- **HVTN 505 trial:** trial of a preventive HIV vaccine
- Stopped early after reaching predetermined cutoffs for efficacy futility ([Hammer et al., 2013](#))

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE

 IDSA
Infectious Diseases Society of America

 hivma
HIV Medicine Association

 OXFORD

Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. Janes,¹ Kristen W. Cohen,² Nicole Frahm,³ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients ([Janes et al., 2017](#); [Fong et al., 2018](#); [Neidich et al., 2019](#))

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE



Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. Janes,¹ Kristen W. Cohen,¹ Nicole Frahm,² Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,³ Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients ([Janes et al., 2017](#); [Fong et al., 2018](#); [Neidich et al., 2019](#))
- Is there a **causal** relationship between these biomarkers and HIV?

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE



Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. Janes,¹ Kristen W. Cohen,² Nicole Frahm,³ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients ([Janes et al., 2017](#); [Fong et al., 2018](#); [Neidich et al., 2019](#))
- Is there a **causal** relationship between these biomarkers and HIV?
- Biomarker-HIV relationship is **confounded**

Motivation: HVTN 505 trial

The Journal of Infectious Diseases

MAJOR ARTICLE



Higher T-Cell Responses Induced by DNA/rAd5 HIV-1 Preventive Vaccine Are Associated With Lower HIV-1 Infection Risk in an Efficacy Trial

Holly E. Janes,¹ Kristen W. Cohen,² Nicole Frahm,³ Stephen C. De Rosa,¹ Brittany Sanchez,¹ John Hural,¹ Craig A. Magaret,¹ Shelly Karuna,¹ Carter Bentley,¹ Raphael Gottardo,¹ Greg Finak,¹ Douglas Grove,² Mingchao Shen,¹ Barney S. Graham,³ Richard A. Koup,³ Mark J. Mulligan,⁴ Beryl Koblin,⁵ Susan P. Buchbinder,⁶ Michael C. Keefer,⁷ Elizabeth Adams,⁸ Chuka Anude,^{8a} Lawrence Corey,¹ Magdalena Sobieszczyk,^{1b} Scott M. Hammer,^{1b} Peter B. Gilbert,¹ and M. Juliana McElrath¹

- Several biomarkers correlated with HIV among vaccine recipients ([Janes et al., 2017](#); [Fong et al., 2018](#); [Neidich et al., 2019](#))
- Is there a **causal** relationship between these biomarkers and HIV?
- Biomarker-HIV relationship is **confounded**
- Biomarkers are **measured with error**

Goal

To estimate the **causal effect** of a continuous exposure on an outcome when

Goal

To estimate the **causal effect** of a continuous exposure on an outcome when


- (i) The exposure-outcome relationship is potentially **confounded**

Goal


To estimate the **causal effect** of a continuous exposure on an outcome when

- (i) The exposure-outcome relationship is potentially **confounded**
- (ii) The exposure is **measured with error**


Confounding




Confounding



Confounding



Measurement Error



Addressing Confounding Alone

- G-Formula
- IPW
- Doubly Robust Estimator

Addressing Confounding Alone

- G-Formula

$$\Psi_{0-GF}(Y, \mathbf{L}, \mathbf{A}; \theta_{GF}) = \begin{bmatrix} \{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\} \partial_{\beta} \mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{bmatrix}$$

- IPW
- Doubly Robust Estimator

Addressing Confounding Alone

- G-Formula

$$\Psi_{0-GF}(Y, \mathbf{L}, \mathbf{A}; \theta_{GF}) = \left[\begin{array}{c} \{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\} \partial_{\beta} \mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{array} \right]$$

- IPW

$$\Psi_{0-IPW}(Y, \mathbf{L}, \mathbf{A}; \theta_{IPW}) = \left[\begin{array}{c} \Psi_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A}) \{Y - \eta(\mathbf{A}; \gamma)\} \partial_{\gamma} \eta(\mathbf{A}; \gamma) \end{array} \right]$$

- Doubly Robust Estimator

Addressing Confounding Alone

- G-Formula

$$\psi_{0-GF}(Y, \mathbf{L}, \mathbf{A}; \theta_{GF}) = \left[\begin{array}{c} \{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\} \partial_{\beta} \mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{array} \right]$$

- IPW

$$\psi_{0-IPW}(Y, \mathbf{L}, \mathbf{A}; \theta_{IPW}) = \left[\begin{array}{c} \Psi_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A}) \{Y - \eta(\mathbf{A}; \gamma)\} \partial_{\gamma} \eta(\mathbf{A}; \gamma) \end{array} \right]$$

- Doubly Robust Estimator

$$\psi_{0-DR}(Y, \mathbf{L}, \mathbf{A}; \theta_{DR}) = \left[\begin{array}{c} \Psi_{PS}(\mathbf{L}, \mathbf{A}) \\ SW(\mathbf{L}, \mathbf{A}) \{Y - \mu(\mathbf{L}, \mathbf{A}; \beta)\} \partial_{\beta} \mu(\mathbf{L}, \mathbf{A}; \beta) \\ \eta(\mathbf{a}) - \mu(\mathbf{L}, \mathbf{a}; \beta) \end{array} \right]$$

Addressing Confounding and Measurement Error

Can we just substitute $\mathbf{A}^*$ for $\mathbf{A}$ and find the solution to

$$\sum_{i=1}^n \psi_0(Y_i, \mathbf{L}_i, \underbrace{\mathbf{A}_i^*}_{\text{mismeasured}}; \theta) = \mathbf{0}?$$

Addressing Confounding and Measurement Error

Can we just substitute $\mathbf{A}^*$ for $\mathbf{A}$ and find the solution to

$$\sum_{i=1}^n \psi_0(Y_i, \mathbf{L}_i, \underbrace{\mathbf{A}_i^*}_{\text{mismeasured}}; \theta) = \mathbf{0}?$$

No! This leads to bias in $\hat{\theta}$ because

$$\mathbb{E}\{\psi_0(Y, \mathbf{L}, \mathbf{A}^*; \theta_0)\} \neq \mathbf{0}.$$

Addressing Confounding and Measurement Error

Can we just substitute $\mathbf{A}^*$ for $\mathbf{A}$ and find the solution to

$$\sum_{i=1}^n \underbrace{\psi_0(Y_i, \mathbf{L}_i, \mathbf{A}_i^*)}_{\text{mismeasured}}; \theta) = \mathbf{0}?$$

No! This leads to bias in $\hat{\theta}$ because

$$E\{\psi_0(Y, \mathbf{L}, \mathbf{A}^*; \theta_0)\} \neq \mathbf{0}.$$

We need a new score function ψ_{CS} such that

$$E\{\underbrace{\psi_{CS}}_{\text{new score fun.}}(Y, \mathbf{L}, \underbrace{\mathbf{A}^*}_{\text{mismeasured}}; \theta_0)\} = \mathbf{0}.$$

Corrected Score Functions

Given the “oracle” score function Ψ_0 , a “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) ([Novick and Stefanski, 2002](#)):

Corrected Score Functions

Given the “oracle” score function Ψ_0 , a “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) ([Novick and Stefanski, 2002](#)):

$$\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta) \equiv E[\text{Re}\{\Psi_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, \mathbf{L}, \mathbf{A}^*]$$

Corrected Score Functions

Given the “oracle” score function Ψ_0 , a “corrected score” function Ψ_{CS} can be created following Novick and Stefanski (2002) ([Novick and Stefanski, 2002](#)):

$$\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta) \equiv E[\text{Re}\{\Psi_0(Y, \mathbf{L}, \mathbf{A}^* + i\tilde{\epsilon}; \theta)\} | Y, \mathbf{L}, \mathbf{A}^*]$$

The G-Formula, IPW, and DR score functions can all be corrected in this way

$$\Psi_{0-GF} \longrightarrow \Psi_{CS-GF}$$


$$\Psi_{0-IPW} \longrightarrow \Psi_{CS-IPW}$$

$$\Psi_{0-DR} \longrightarrow \Psi_{CS-DR}$$

Summary of Project 1



Paper in *Biometrics* (2025)



GitHub R package

Mismex: Causal Inference for Mismeasured Exposures

Acknowledgments

- add photos

Thank you! Questions?

References I

- Allison E. Aiello, Amanda M. Simanek, Marisa C. Eisenberg, Alison R. Walsh, Brian Davis, Erik Volz, Caroline Cheng, Jeanette J. Rainey, Amra Uzicanin, Hongjiang Gao, Nathaniel Osgood, Dylan Knowles, Kevin Stanley, Kara Tarter, and Arnold S. Monto. Design and methods of a social network isolation study for reducing respiratory infection transmission: The eX-FLU cluster randomized trial. *Epidemics*, 15:38–55, June 2016. ISSN 17554365. doi: 10.1016/j.epidem.2016.01.001. URL <https://linkinghub.elsevier.com/retrieve/pii/S1755436516000025>.
- Shaina J. Alexandria, Michael G. Hudgens, and Allison E. Aiello. Assessing Intervention Effects in a Randomized Trial Within a Social Network. *Biometrics*, 79(2):1409–1419, June 2023. ISSN 0006-341X, 1541-0420. doi: 10.1111/biom.13606. URL <https://academic.oup.com/biometrics/article/79/2/1409-1419/7513977>.
- R. A. Fisher. The design of experiments. *Nature*, 137(3459):252–254, February 1936. ISSN 0028-0836, 1476-4687. doi: 10.1038/137252a0.
- Youyi Fong, Xiaoying Shen, Vicki C Ashley, Aaron Deal, Kelly E Seaton, Chenchen Yu, Shannon P Grant, Guido Ferrari, Allan C deCamp, Robert T Bailer, et al. Modification of the association between T-cell immune responses and human immunodeficiency virus type 1 infection risk by vaccine-induced antibody responses in the HVTN 505 trial. *The Journal of Infectious Diseases*, 217(8):1280–1288, 2018.
- Scott M Hammer, Magdalena E Sobieszczyk, Holly Janes, Shelly T Karuna, Mark J Mulligan, Doug Grove, Beryl A Koblin, Susan P Buchbinder, Michael C Keefer, Georgia D Tomaras, et al. Efficacy trial of a DNA/rAd5 HIV-1 preventive vaccine. *New England Journal of Medicine*, 369(22):2083–2092, 2013.
- Richard Hayes, Helen Ayles, Nulda Beyers, Kalpana Sabapathy, Sian Floyd, Kwame Shanaube, Peter Bock, Sam Griffith, Ayana Moore, Deborah Watson-Jones, Christophe Fraser, Sten H Vermund, Sarah Fidler, and The HPTN 071 (PopART) Study Team. Hptn 071 (popart): Rationale and design of a cluster-randomised trial of the population impact of an hiv combination prevention intervention including universal testing and treatment – a study protocol for a cluster randomised trial. *Trials*, 15(1):57, December 2014. ISSN 1745-6215. doi: 10.1186/1745-6215-15-57.

References II

- Richard J. Hayes, Deborah Donnell, Sian Floyd, Nomtha Mandla, Justin Bwalya, Kalpana Sabapathy, Blia Yang, Mwelwa Phiri, Ab Schaap, Susan H. Eshleman, Estelle Piwowar-Manning, Barry Kosloff, Anelet James, Timothy Skalland, Ethan Wilson, Lynda Emel, David Macleod, Rory Dunbar, Musonda Simwinga, Nozizwe Makola, Virginia Bond, Graeme Hoddinott, Ayana Moore, Sam Griffith, Nirupama Deshmane Sista, Sten H. Vermund, Wafaa El-Sadr, David N. Burns, James R. Hargreaves, Katharina Hauck, Christophe Fraser, Kwame Shanaube, Peter Bock, Nulda Beyers, Helen Ayles, and Sarah Fidler. Effect of universal testing and treatment on hiv incidence — hptn 071 (popart). *New England Journal of Medicine*, 381(3):207–218, 2019. ISSN 0028-4793, 1533-4406. doi: 10.1056/NEJMoA1814556.
- Michael G Hudgens and M. Elizabeth Halloran. Toward causal inference with interference. *Journal of the American Statistical Association*, 103(482):832–842, 2008. ISSN 0162-1459, 1537-274X. doi: 10.1198/016214508000000292.
- Holly E Janes, Kristen W Cohen, Nicole Frahm, Stephen C De Rosa, Brittany Sanchez, John Hural, Craig A Magaret, Shelly Karuna, Carter Bentley, Raphael Gottardo, et al. Higher T-cell responses induced by DNA/rAd5 HIV-1 preventive vaccine are associated with lower HIV-1 infection risk in an efficacy trial. *The Journal of Infectious Diseases*, 215(9):1376–1385, 2017.
- Ministry of Health Zambia. Zambia population-based hiv impact assessment (zamphia) 2016. *Lusaka: Ministry of Health*, 2019.
- Scott D Neidich, Youyi Fong, Shuying S Li, Daniel E Geraghty, Brian D Williamson, William Chad Young, Derrick Goodman, Kelly E Seaton, Xiaoying Shen, Sheetal Sawant, et al. Antibody Fc effector functions and IgG3 associate with decreased HIV-1 risk. *Journal of Clinical Investigation*, 129(11):4838–4849, 2019.
- Steven J Novick and Leonard A Stefanski. Corrected score estimation via complex variable simulation extrapolation. *Journal of the American Statistical Association*, 97(458):472–481, June 2002. ISSN 0162-1459, 1537-274X. doi: 10.1198/016214502760047005. URL <http://www.tandfonline.com/doi/abs/10.1198/016214502760047005>.
- David M. Ritzwoller, Joseph P. Romano, and Azeem M. Shaikh. Randomization inference: Theory and applications, 2024.

References III

- Bonnie E. Shook-Sa, Paul N. Zivich, Stephen R. Cole, Nora E. Rosenberg, Michael G. Hudgens, Deborah J. Donnell, Sizulu Moyo, Khangelani Zuma, Helen Ayles, Peter Bock, Joseph J. Eron, Richard J. Hayes, and Jessie K. Edwards. Examining the effect of universal testing and treatment strategies for hiv prevention in zambia and south africa: generalizing the results of the hptn 071 (popart) trial. *Journal of the International AIDS Society*, 28(12):e70062, December 2025. ISSN 1758-2652, 1758-2652. doi: 10.1002/jia2.70062.
- L. Simbayi, Khangelani Zuma, S. Moyo, E. Marinda, M. Mabaso, S. Ramlagan, A. North, N. Mohlabane, C. Dietrich, and I. Naidoo. South african national hiv prevalence, incidence, behaviour and communication survey, 2017. *Cape Town: HSRC Press*, 2021.
- Leonard A Stefanski and Dennis D Boos. The calculus of m-estimation. *The American Statistician*, 56(1):29–38, February 2002. ISSN 0003-1305, 1537-2731. doi: 10.1198/000313002753631330.
- M. Sugiyama, T. Suzuki, and Takafumi Kanamori. Density ratio estimation: A comprehensive review. *RIMS Kokyuroku*, pages 10–31, 01 2010.
- Yao Zhang and Qingyuan Zhao. What is a randomization test? *Journal of the American Statistical Association*, 118(544):2928–2942, October 2023. ISSN 0162-1459, 1537-274X. doi: 10.1080/01621459.2023.2199814.
- P.N. Zivich, M.C. Eisenberg, A.S. Monto, A. Uzicanin, R. S. Baric, T. P. Sheahan, J. J. Rainey, H. Gao, and A. E. Aiello. Transmission of viral pathogens in a social network of university students: the eX-FLU study. *Epidemiology and Infection*, 148:e267, 2020. ISSN 0950-2688, 1469-4409. doi: 10.1017/S0950268820001806. URL https://www.cambridge.org/core/product/identifier/S0950268820001806/type/journal_article.

Supplement to Project 2: Physical Distancing to Reduce Transmission of Influenza-Like-Illness on College Campuses: the eX-FLU Trial

Brian Richardson, Allison Aiello, Michael Hudgens

Choice of Test Statistic

Test statistics: “proportion of possible transmission events attributable to students in the intervention group:”

$$T^{**}(\mathbf{Z}, \bar{\mathbf{A}}, \bar{\mathbf{Y}}) = \frac{\sum_{k=2}^{\tau} \sum_{i=1}^n \sum_{j>i} Z_i E_{ijk}^{**}}{\sum_{k=2}^{\tau} \sum_{i=1}^n \sum_{j>i} E_{ijk}^{**}}.$$

	From Infected to Infected	From Infected
Contact at $k - 1$	$E_{ijk}^{11} = Y_i^{k-1} A_{ij}^{k-1} Y_j^k$	$E_{ijk}^{12} = Y_i^{k-1} A_{ij}^{k-1}$
Contact at k	$E_{ijk}^{21} = Y_i^{k-1} A_{ij}^k Y_j^k$	$E_{ijk}^{22} = Y_i^{k-1} A_{ij}^k$
Contact at $k - 1$ or k	$E_{ijk}^{31} = Y_i^{k-1} (A_{ij}^{k-1} \vee A_{ij}^k) Y_j^k$	$E_{ijk}^{32} = Y_i^{k-1} (A_{ij}^{k-1} \vee A_{ij}^k)$
Contact at $k - 1$ and k	$E_{ijk}^{41} = Y_i^{k-1} (A_{ij}^{k-1} * A_{ij}^k) Y_j^k$	$E_{ijk}^{42} = Y_i^{k-1} (A_{ij}^{k-1} * A_{ij}^k)$


Table: Definitions of a possible transmission event E_{ijk} from student i to student j at time k . Y_i^k is an indicator for student i being infected at week k , A_{ij}^k is an indicator for students i and j being in contact at week k , and $a \vee b$ denotes the maximum of a and b .

Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)

Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)




Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)

An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$



Exponential Family Random Graph Models


Goal: model a network (**A**) given covariates (**X**)

An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients
- $\mathbf{g}(\mathbf{a}, \mathbf{x})$: sufficient statistics



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients
- $\mathbf{g}(\mathbf{a}, \mathbf{x})$: sufficient statistics
 - ▶ number of edges



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients
- $\mathbf{g}(\mathbf{a}, \mathbf{x})$: sufficient statistics
 - ▶ number of edges
 - ▶ number of edges touching treated students



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients
- $\mathbf{g}(\mathbf{a}, \mathbf{x})$: sufficient statistics
 - ▶ number of edges
 - ▶ number of edges touching treated students
 - ▶ number of edges touching infected students



Exponential Family Random Graph Models

Goal: model a network (**A**) given covariates (**X**)


An **ERGM** assumes:

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})}$$

- normalizing constant:

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

- $\mathcal{A}$: space of possible networks
- θ : model coefficients
- $\mathbf{g}(\mathbf{a}, \mathbf{x})$: sufficient statistics
 - ▶ number of edges
 - ▶ number of edges touching treated students
 - ▶ number of edges touching infected students
 - ▶ number of edges touching treated $\times$ infected students



ERGM Model Formulation

$$\Pr_{\theta}(\mathbf{A} = \mathbf{a} | \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}}{\kappa(\theta, \mathcal{A}, \mathbf{x})},$$

$$\kappa(\theta, \mathcal{A}, \mathbf{x}) = \sum_{\mathbf{a} \in \mathcal{A}} \exp\{\theta \cdot \mathbf{g}(\mathbf{a}, \mathbf{x})\}$$

For example, in the simulation study and eX-FLU application,

$$\mathbf{g}(\mathbf{a}, \mathbf{x}) = \begin{bmatrix} \# \text{ edges} \\ \# \text{ edges touching a treated node} \\ \# \text{ edges touching an infected node} \\ \# \text{ edges touching a treated and infected node} \\ \# \text{ edges between roommate pairs} \end{bmatrix} = \begin{bmatrix} \sum_{i,j} a_{ij} \\ \sum_{i,j} a_{ij}(z_i + z_j) \\ \sum_{i,j} a_{ij}(y_i + y_j) \\ \sum_{i,j} a_{ij}(z_i y_i + z_j y_j) \\ \sum_{i,j} a_{ij} \mathbb{1}(i, j \text{ roommates}) \end{bmatrix}$$

ERGM Change Statistic Model Formulation

- **Change statistic:** $\delta_{\mathbf{g}}(\mathbf{a}, \mathbf{x})_{ij} = \mathbf{g}(\mathbf{a}_{ij}^+, \mathbf{x}) - \mathbf{g}(\mathbf{a}_{ij}^-, \mathbf{x})$ is the change in network statistic that would occur if a_{ij} were changed from 0 to 1
 - ▶ where $\mathbf{a}_{ij}^+$ and $\mathbf{a}_{ij}^-$ represent the network $\mathbf{a}$ with dyad a_{ij} set to 1 or 0, respectively
- Then the equivalent ERGM specification is

$$\text{logit}\{\Pr_{\theta}(A_{ij} = 1 | \mathbf{A}_{ij}^C = \mathbf{a}_{ij}^C, \mathbf{X} = \mathbf{x})\} = \theta^k \delta_{\mathbf{g}}(\mathbf{a}, \mathbf{x})_{ij}$$

- ▶ where $\mathbf{A}_{ij}^C$ represents all dyads in $\mathbf{A}$ except A_{ij}
- **Interpretation of θ :** the change in conditional log-odds of the network associated with a one-unit increase in the corresponding component of $\mathbf{g}(\mathbf{a}, \mathbf{x})$ resulting from switching a particular dyad A_{ij} from 0 to 1 and leaving the rest of the network fixed at $\mathbf{A}_{ij}^C$

Dyadic Independence EGRMs


- **Dyadic independence term:** a component g of $\mathbf{g}$ in an ERGM for which the corresponding change statistic $\delta_g(\mathbf{a}, \mathbf{x})_{ij}$ can be calculated for any i, j without knowing $\mathbf{a}$
 - ▶ For example, if $g(\mathbf{a}, \mathbf{x}) = \sum_{i,j} a_{ij}(Z_i + Z_j)$ counts the number of edges touching treated nodes, then $\delta_g(\mathbf{a}, \mathbf{x})_{ij} = z_i + z_j$ doesn't depend on $\mathbf{a}$
- **Dyadic independence ERGM:** an ERGM with only dyadic independence terms
 - ▶ replace $\delta_g(\mathbf{a}, \mathbf{x})_{ij}$ with $\delta_g(\mathbf{x})_{ij}$ and write the model as

$$\text{logit}\{\Pr(A_{ij} = 1 | \mathbf{X} = \mathbf{x})\} = \theta \cdot \delta_g(\mathbf{x})_{ij}$$

- **Interpretation of θ :** the change in log-odds of the network associated with a one-unit increase in the corresponding component of $\mathbf{g}(\mathbf{a}, \mathbf{x})$ resulting from switching a particular dyad A_{ij} from 0 to 1

Separable Temporal Exponential Family Random Graph Models

A **STERGM** assumes that, at each time step $k = 1, \dots, \tau - 1$:




Separable Temporal Exponential Family Random Graph Models

A **STERGM** assumes that, at each time step $k = 1, \dots, \tau - 1$:

- New edges form according to a **formation** ERGM

$$\Pr_{\theta^+}(\mathbf{A}^{k+1} = \mathbf{a}^{k+1} | \mathbf{A}^k = \mathbf{a}^k, \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta^+ \cdot \mathbf{g}^+(\mathbf{a}^{k+1}, \mathbf{x})\}}{\kappa\{\theta^+, \mathcal{A}^+(\mathbf{a}^k), \mathbf{x}\}}$$




Separable Temporal Exponential Family Random Graph Models

A **STERGM** assumes that, at each time step $k = 1, \dots, \tau - 1$:

- New edges form according to a **formation** ERGM
- Old edges persist according to a **persistence** ERGM

$$\Pr(\mathbf{A}^{k-} = \mathbf{a}^{k-} | \mathbf{A}^k = \mathbf{a}^k, \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta^- \cdot \mathbf{g}^-(\mathbf{a}^{k-}, \mathbf{x})\}}{\kappa\{\theta^-, \mathcal{A}^-(\mathbf{a}^k), \mathbf{x}\}}.$$




Separable Temporal Exponential Family Random Graph Models

A **STERGM** assumes that, at each time step $k = 1, \dots, \tau - 1$:

- New edges form according to a **formation** ERGM
- Old edges persist according to a **persistence** ERGM
- The network at time step $k + 1$ is the result of formation and persistence

$$\mathbf{A}^{k+1} = \underbrace{\mathbf{A}^k}_{\text{previous network}} \cup \underbrace{(\mathbf{A}^{k+} - \mathbf{A}^k)}_{\text{new edges formed}} - \underbrace{(\mathbf{A}^k - \mathbf{A}^{k-})}_{\text{old edges not persisting}}$$



STERGM Model Formulation

- **Formation model** is an ERGM conditional on only adding edges:

$$\Pr_{\theta^+}(\mathbf{A}^{k+1} = \mathbf{a}^{k+1} | \mathbf{A}^k = \mathbf{a}^k, \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta^+ \cdot \mathbf{g}^+(\mathbf{a}^{k+1}, \mathbf{x})\}}{\kappa\{\theta^+, \mathcal{A}^+(\mathbf{a}^k), \mathbf{x}\}}$$

► $\mathcal{A}^+(\mathbf{a})$: space of possible networks that can be formed by adding edges to $\mathbf{a}$

- **Persistence model** is an ERGM conditional on only removing edges:

$$\Pr_{\theta^-}(\mathbf{A}^{k+1} = \mathbf{a}^{k+1} | \mathbf{A}^k = \mathbf{a}^k, \mathbf{X} = \mathbf{x}) = \frac{\exp\{\theta^- \cdot \mathbf{g}^-(\mathbf{a}^{k+1}, \mathbf{x})\}}{\kappa\{\theta^-, \mathcal{A}^-(\mathbf{a}^k), \mathbf{x}\}}$$

► $\mathcal{A}^-(\mathbf{a})$: space of possible networks that can be formed by removing edges to $\mathbf{a}$

- A **STERGM** assumes the network at time $k + 1$ is then the result of applying the changes in $\mathbf{A}^{k+}$ and $\mathbf{A}^{k-}$ to $\mathbf{A}^k$:


$$\mathbf{A}^{k+1} = \underbrace{\mathbf{A}^k}_{\text{previous network}} \cup \underbrace{(\mathbf{A}^{k+} - \mathbf{A}^k)}_{\text{new edges formed}} - \underbrace{(\mathbf{A}^k - \mathbf{A}^{k-})}_{\text{old edges not persisting}}$$

Separability of STERGMs

- 1 $\mathbf{A}^{k+} \perp\!\!\!\perp \mathbf{A}^{k-} | \mathbf{A}^k$, i.e., the formation and persistence processes are conditionally independent given the network at time k


Separability of STERGMs

- ① $\mathbf{A}^{k+} \perp\!\!\!\perp \mathbf{A}^{k-} | \mathbf{A}^k$, i.e., the formation and persistence processes are conditionally independent give the network at time k



Separability of STERGMs

- 1 $\mathbf{A}^{k+} \perp\!\!\!\perp \mathbf{A}^{k-} | \mathbf{A}^k$, i.e., the formation and persistence processes are conditionally independent given the network at time k
- 2 the parameter space for $\theta = (\theta^+, \theta^-)$ is the product of the parameter spaces for θ^+ and θ^-



Simulation Setup

- $n \in \{24, 48\}$ students were equally divided into two residence halls, each with six clusters of equal size
- Within each residence hall group, the clusters were randomized using a 50:50 allocation to either the intervention group or the control group
- Five pairs of students were randomly selected to be roommates, meaning they had contact with each other each week
- Baseline ($k = 1$) face-to-face contacts were simulated between each pair of students with probability 0.5
- Baseline infection statuses were simulated for each student with probability 0.5
- Social networks were simulated over the remaining $\tau \in \{5, 10\}$ weeks according to a STERGM with both formation and persistence models including edge count, intervention assignment Z , infection status Y , $Z \times Y$ interaction, and an offset to force constant edges between roommates
- Infection statuses were simulated for each student i at each week $k \in \{2, \dots, \tau\}$ with probability $\Pr(Y_i^k = 1 | \bar{\mathbf{A}}^{k-1}, \bar{\mathbf{Y}}^{k-1}) = h_i(\mathbf{S}^{k-1})$ depending on the vector $\mathbf{S}^{k-1} = \mathbf{A}^{k-1} \mathbf{Y}^{k-1}$ of counts of infected contacts for each student at week $k - 1$.

Simulation Setup

Three scenarios:

Null	Formation Model	Persistence Model	Infection Probability
Hypothesis	Parameters θ^+	Parameters θ^-	Function h_i
$H_0^\sharp$	$(-0.5, 0, 0, 0)$	$(-0.5, 0, 0, 0)$	$h_i(\mathbf{s}) = 0.5$
$\overline{H_0^A} \cap H_0^Y$	$(-0.2, 0, 0, -1)$	$(-0.2, 0, 0, -1)$	$h_i(\mathbf{s}) = 0.5$
$\overline{H_0^A} \cap \overline{H_0^Y}$	$(-0.2, 0, 0, -1)$	$(-0.2, 0, 0, -1)$	$h_i(\mathbf{s}) = \frac{n}{2} \text{expit}(s_i - \bar{s}) / \sum_{j=1}^n \text{expit}(s_j - \bar{s})$

Table: The null hypothesis refers to the null hypothesis that is true under the data generating process, formation model parameters are $\theta^+ = (\theta_{\text{edges}}^+, \theta_Z^+, \theta_Y^+, \theta_{ZY}^+)$, persistence model parameters are $\theta^- = (\theta_{\text{edges}}^-, \theta_Z^-, \theta_Y^-, \theta_{ZY}^-)$, the infection probability function $h_i(\mathbf{s})$ gives the probability of student i being infected at week k given the vector $\mathbf{S}^{k-1} = \mathbf{s} = (s_1, \dots, s_n)$ of infected contact counts for each student at week $k-1$, and $\bar{s} = n^{-1} \sum_{i=1}^n s_i$.

Three p-values:

- $\rho_B^\sharp$: testing the sharp null
- ρ_B^Y : testing H_0^Y using known q
- $\hat{\rho}_B^Y$: testing H_0^Y using estimated q

Simulation Study

Goal: use simulated data to investigate the performance of our testing procedure

Simulation Study

Goal: use simulated data to investigate the performance of our testing procedure

- $n \in \{24, 48\}$ students
- $\tau \in \{5, 10\}$ weeks
- 50:50 cluster randomization with 12 equal-sized clusters

Simulation Study

Goal: use simulated data to investigate the performance of our testing procedure


- $n \in \{24, 48\}$ students
- $\tau \in \{5, 10\}$ weeks
- 50:50 cluster randomization with 12 equal-sized clusters
- Three scenarios:
 - 1 $H_0^\sharp$: no effect of intervention on networks or infection
 - 2 $\overline{H}_0^A \cap H_0^Y$: no effect of intervention on infection
 - 3 $\overline{H}_0^A \cap \overline{H}_0^Y$: effect of intervention on both networks and infection

Simulation Study


Goal: use simulated data to investigate the performance of our testing procedure

- $n \in \{24, 48\}$ students
- $\tau \in \{5, 10\}$ weeks
- 50:50 cluster randomization with 12 equal-sized clusters
- Three scenarios:
 - 1 $H_0^\#$: no effect of intervention on networks or infection
 - 2 $\overline{H}_0^A \cap H_0^Y$: no effect of intervention on infection
 - 3 $\overline{H}_0^A \cap \overline{H}_0^Y$: effect of intervention on both networks and infection
- Three testing procedures:
 - 1 $\rho_B^\#$: testing $H_0^\#$
 - 2 ρ_B^Y : testing H_0^Y using known q
 - 3 $\hat{\rho}_B^Y$: testing H_0^Y using estimated q

Simulation Results: Empirical CDF of P-Values



eX-FLU: STERGM Results



Type I Error Control

- **Proposition 2.1:** Let T_N be a test statistic with CDF F_N under H_0 .
 - ① Under H_0 , $F_N(T_N)$ stochastically dominates a $\text{Uniform}(0, 1)$ distribution for any N .
 - ② If the test statistic T_N has a continuous limiting distribution, then $F_N(T_N) \rightarrow^d \text{Uniform}(0, 1)$.
- **Corollary 2.2:**
 - ① Under $H_0^\sharp$, the sharp null p-value $\rho_N^\sharp$ stochastically dominates a $\text{Uniform}(0, 1)$ distribution for any N .
 - ② Under H_0^Y , the oracle p-value ρ_N^Y stochastically dominates a $\text{Uniform}(0, 1)$ distribution for any N .
 - ③ If the test statistic T_N has a continuous limiting distribution, then $\rho_N^\sharp \rightarrow^d \text{Uniform}(0, 1)$ under $H_0^\sharp$ and $\rho_N^Y \rightarrow^d \text{Uniform}(0, 1)$ under H_0^Y .

Type I Error Control

- **Proposition 2.3:** Let $q(\mathbf{a}, \mathbf{z}, \theta) \equiv \Pr\{\mathbf{A}(\mathbf{z}) = \mathbf{a}; \theta\}$ denote the PMF of the distribution of stochastic potential networks $\mathbf{A}(\mathbf{z})$ at parameter value θ , let $\hat{\theta}_N$ denote the estimator of θ , and let $F_N(\cdot; \theta)$ denote the CDF of the test statistic T_N at θ , with limiting CDF $F(\cdot; \theta)$. Let θ_0 denote the true value of θ . Assume the following:

(A1) $\hat{\theta}_N \rightarrow^P \theta_0$

(A2) $F(t; \theta_0)$ is continuous in t on $\mathbb{R}$

(A3) there exists a $\delta_0 > 0$ such that

$$\sup_{\theta \in B_{\delta_0}(\theta_0)} \sup_{t \in \mathbb{R}} |F_N(t; \theta) - F(t; \theta)| \rightarrow 0$$

(A4) $F(t; \theta)$ is continuous in θ at θ_0 uniformly in t , i.e.,


$$\lim_{\theta \rightarrow \theta_0} \sup_{t \in \mathbb{R}} |F(t; \theta) - F(t; \theta_0)| = 0$$

Then the plug-in p-value $\hat{\rho}_N^Y$ converges in distribution to $\text{Uniform}(0, 1)$.

Type I Error Control


- **Proposition 2.4:** Let $\rho_N = F_N(T_N)$ for test statistic T_N and CDF F_N (not necessarily the true CDF of T_N). Let $T_N^* = h_N(T_N)$ for a sequence of deterministic, strictly increasing functions h_N . Define $F_N^*(t) = F_N\{h_N^{-1}(t)\}$, the (not necessarily true) CDF of the transformed test statistic, and let $\rho_N^* = F_N^*(T_N^*)$. Then $\rho_N^* = \rho_N$.
- **Corollary 2.5:** Let h_N be a sequence of deterministic, strictly increasing functions.
 - 1 If the hypotheses of Proposition 2.1 are met for a test statistic T_N , then the results also hold for $T_N^* = h_N(T_N)$.
 - 2 If the hypotheses of Proposition 2.3 are met for $T_N, F_N(\cdot; \theta)$, then the results also hold for $T_N^* = h_N(T_N), F_N^*(\cdot; \theta) = F_N\{h_N^{-1}(\cdot); \theta\}$.

Assumptions



Assumptions


- Design assumptions



Assumptions

- Design assumptions


- Cluster-randomization: $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$



Assumptions


- Design assumptions

- Cluster-randomization: $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$
- Independent random sampling of trial and auxiliary data: $S_{ij} \perp\!\!\!\perp (A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij}, C_{ij}, Y_{ij})$




Assumptions

- Design assumptions
 - ▶ **Cluster-randomization:** $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$
 - ▶ **Independent random sampling of trial and auxiliary data:** $S_{ij} \perp\!\!\!\perp (A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij}, C_{ij}, Y_{ij})$
- “Standard” causal inference assumptions



Assumptions

- Design assumptions
 - ▶ **Cluster-randomization:** $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$
 - ▶ **Independent random sampling of trial and auxiliary data:** $S_{ij} \perp\!\!\!\perp (A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij}, C_{ij}, Y_{ij})$
- “Standard” causal inference assumptions
 - ▶ **Causal consistency:** $Y_{ij} = Y_{ij}(a)$, $R_{ij} = R_{ij}(a)$, and $C_{ij} = C_{ij}(a)$ when $A_i = a$




Assumptions

- Design assumptions

- ▶ **Cluster-randomization:** $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$
- ▶ **Independent random sampling of trial and auxiliary data:** $S_{ij} \perp\!\!\!\perp (A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij}, C_{ij}, Y_{ij})$

- “Standard” causal inference assumptions

- ▶ **Causal consistency:** $Y_{ij} = Y_{ij}(a)$, $R_{ij} = R_{ij}(a)$, and $C_{ij} = C_{ij}(a)$ when $A_i = a$
- ▶ **Conditional exchangeability:** $Y_{ij}(a) \perp\!\!\!\perp A_i | \mathbf{X}_i, \mathbf{W}_{ij}$




Assumptions

- Design assumptions

- Cluster-randomization: $A_i \perp\!\!\!\perp (\mathbf{W}_{ij}, \dots, \mathbf{W}_{in_i}) | \mathbf{X}_i$
- Independent random sampling of trial and auxiliary data: $S_{ij} \perp\!\!\!\perp (A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij}, C_{ij}, Y_{ij})$


- “Standard” causal inference assumptions

- Causal consistency: $Y_{ij} = Y_{ij}(a)$, $R_{ij} = R_{ij}(a)$, and $C_{ij} = C_{ij}(a)$ when $A_i = a$
- Conditional exchangeability: $Y_{ij}(a) \perp\!\!\!\perp A_i | \mathbf{X}_i, \mathbf{W}_{ij}$
- Positivity of treatment assignment: $\Pr(A_i = a | \mathbf{X}_i = \mathbf{x}) > 0 \forall \mathbf{x}$ with positive density $f_{\mathbf{X}}(\mathbf{x}) > 0$.



Assumptions

- Additional assumptions




Assumptions

- Additional assumptions

- ▶ **Positivity of uncensored response:**

$$\Pr(R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, A_i = a, \mathbf{W}_{ij} = \mathbf{w}) > 0 \quad \forall \mathbf{x}, \mathbf{w} \text{ with } f_{\mathbf{X}, \mathbf{W}}(\mathbf{x}, \mathbf{w}) > 0$$




Assumptions

- Additional assumptions

- ▶ **Positivity of uncensored response:**

$$\Pr(R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, A_i = a, \mathbf{W}_{ij} = \mathbf{w}) > 0 \quad \forall \mathbf{x}, \mathbf{w} \text{ with } f_{\mathbf{x}, \mathbf{w}}(\mathbf{x}, \mathbf{w}) > 0$$

- ▶ **Conditional independence of potential outcome and response:** $Y_{ij}(a) \perp\!\!\!\perp R_{ij}(a) | A_i, \mathbf{X}_i, \mathbf{W}_{ij}$



Assumptions

- Additional assumptions


- **Positivity of uncensored response:**

$$\Pr(R_{ij} = 1, C_{ij} = 0 | \mathbf{X}_i = \mathbf{x}, A_i = a, \mathbf{W}_{ij} = \mathbf{w}) > 0 \quad \forall \mathbf{x}, \mathbf{w} \text{ with } f_{\mathbf{x}, \mathbf{w}}(\mathbf{x}, \mathbf{w}) > 0$$

- **Conditional independence of potential outcome and response:** $Y_{ij}(a) \perp\!\!\!\perp R_{ij}(a) | A_i, \mathbf{X}_i, \mathbf{W}_{ij}$

- **Conditional independence of potential outcome and censoring among responders:**

$$Y_{ij}(a) \perp\!\!\!\perp C_{ij}(a) | A_i, \mathbf{X}_i, \mathbf{W}_{ij}, R_{ij} = 1$$



Supplement to Project 3 : Causal Inference from Cluster Randomized Trials with Differential Nonresponse

Brian Richardson, Bonnie Shook-Sa, Michael Hudgens

Simulation Details

- $m = 20$ clusters
 - ▶ cluster treatment $\Pr(A = 1) = 0.5$
 - ▶ cluster covariate $X \sim N(0, 1)$
- invited trial sample of size $n^{\text{trial}} \in \{500, 5000\}$
- individual covariates $W_1 \sim \text{Bernoulli}(0.5)$, $W_2 \sim N(0, 1)$
- potential outcomes

$$Y(0) \sim \text{Bernoulli}\{\text{expit}(-0.5 + 0.25X_1 + 0.5W_1 + 0.5W_2)\},$$

$$Y(1) \sim \text{Bernoulli}\{\text{expit}(0 - 1W_1 - 0.5W_2)\}$$


$\implies$ treatment is marginally protective, but harmful when $W_1 = 0$

- observed outcomes $Y = (1 - A)Y(0) + AY(1)$
- response $R \sim \text{Bernoulli}\{\text{expit}(\alpha_0 - 2AW_1)\}$
 - $\implies$ marginal $\Pr(R = 1) = 0.5$, but higher response rate if $W_1 = 0$
- auxiliary sample of size $n^{\text{aux}} \in \{500, 5000\}$
 - ▶ same covariates W_1, W_2


Simulation Details

- naive and proposed versions of the g-formula, IPW, and AIPW estimators of $\eta(a)$
 - ▶ naive versions used trial data only, and ignored nonresponse
- Estimand: $RD = \eta(1) - \eta(0)$
- Estimator: $\widehat{RD} = \widehat{\eta}(1) - \widehat{\eta}(0)$
 - ▶ variance estimated using the delta method
- Outcome model specification
 - ▶ μ_a correctly modeled with logistic regression with covariates X, W_1, W_2 and their interaction with treatment A
 - ▶ μ_a mis-specified by excluding the covariate W_1 and its interaction with A
- Joint propensity score model specification
 - ▶ censoring mechanism π_a^C always correctly specified as logistic regression with covariates X, W_1, W_2 and their interaction with treatment A
 - ▶ response mechanism π_a^{AR} possibly mis-specified by incorrectly assuming $R_{ij} \perp\!\!\!\perp A_i | \mathbf{X}_i, \mathbf{W}_{ij}$


Simulation: Variance Estimation



Supplemental Analysis: Arms A/B vs Standard of Care



Supplemental Analysis: Arms A/B vs Standard of Care



Supplement to Project 1 : Addressing Confounding and Continuous Exposure Measurement Error Using Corrected Score Functions

Brian Richardson, Brian Blette, Peter Gilbert, Michael Hudgens

Corrected Score Functions: What?

- Suppose the oracle score function is **conditionally unbiased**, meaning

$$E\{\Psi_0(Y, \mathbf{L}, \mathbf{A}; \theta) | \mathbf{A}\} = \mathbf{0}.$$

- Define the corrected score function as

$$\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta) = E \left[\text{Re} \left\{ \Psi_0(Y, \mathbf{L}, \tilde{\mathbf{A}}; \theta) \right\} \mid Y, \mathbf{L}, \mathbf{A}^* \right],$$

where $\tilde{\mathbf{A}} = \mathbf{A}^* + i\tilde{\epsilon}$, $i = \sqrt{-1}$, $\text{Re}(\cdot)$ denotes the real component of a complex number, and $\tilde{\epsilon} \sim \mathcal{N}(\mathbf{0}, \Sigma_{me})$.

- Then

$$\begin{aligned} E\{\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta) \mid Y, \mathbf{L}, \mathbf{A}\} &= \Psi_0(Y, \mathbf{L}, \mathbf{A}; \theta) \\ \implies E[E\{\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta) \mid Y, \mathbf{L}, \mathbf{A}\}] &= E\{\Psi_0(Y, \mathbf{L}, \mathbf{A}; \theta)\} \\ \implies E\{\Psi_{CS}(Y, \mathbf{L}, \mathbf{A}^*; \theta)\} &= \mathbf{0} \end{aligned}$$

Corrected Score Functions: Why?

The key result ([Novick and Stefanski, 2002](#)) for corrected score functions is that, for a smooth enough function $\mathbf{f}$, the function $\tilde{\mathbf{f}}$ defined by

$$\tilde{\mathbf{f}}(\mathbf{A}^*) \equiv \mathbb{E}[\text{Re}\{\mathbf{f}(\mathbf{A}^* + i\tilde{\epsilon})\} | \mathbf{A}, \mathbf{A}^*]$$

does not depend on $\mathbf{A}$ and satisfies

$$\mathbb{E}\{\tilde{\mathbf{f}}(\mathbf{A}^*) | \mathbf{A}\} = \mathbf{f}(\mathbf{A}).$$

The proof of this for the special case $f(\mathbf{A}) = \exp(\mathbf{c}\mathbf{A}^T)$ is illustrative.

Corrected Score Functions: Why?

CLAIM: $E\{\tilde{f}(\mathbf{A}^*)|\mathbf{A}\} = f(\mathbf{A})$ for $f(\mathbf{A}) = \exp(\mathbf{c}\mathbf{A}^T)$.

Corrected Score Functions: Why?

CLAIM: $E\{\tilde{f}(\mathbf{A}^*)|\mathbf{A}\} = f(\mathbf{A})$ for $f(\mathbf{A}) = \exp(\mathbf{c}\mathbf{A}^T)$.

$$\begin{aligned}\tilde{f}(\mathbf{A}^*) &\equiv E(\text{Re}[\exp\{\mathbf{c}(\mathbf{A}^* + i\tilde{\epsilon})^T\}]|\mathbf{A}, \mathbf{A}^*) \\ &= \exp(\mathbf{c}\mathbf{A}^{*T}) \underbrace{\text{Re}[E\{\exp(i\mathbf{c}\tilde{\epsilon}^T)\}]}_{\text{normal c.f.}} = \exp(\mathbf{c}\mathbf{A}^{*T}) \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right)\end{aligned}$$

Corrected Score Functions: Why?

CLAIM: $E\{\tilde{f}(\mathbf{A}^*)|\mathbf{A}\} = f(\mathbf{A})$ for $f(\mathbf{A}) = \exp(\mathbf{c}\mathbf{A}^T)$.

$$\begin{aligned}\tilde{f}(\mathbf{A}^*) &\equiv E(\text{Re}[\exp\{\mathbf{c}(\mathbf{A}^* + i\tilde{\epsilon})^T\}]|\mathbf{A}, \mathbf{A}^*) \\ &= \exp(\mathbf{c}\mathbf{A}^{*T}) \underbrace{\text{Re}[E\{\exp(i\mathbf{c}\tilde{\epsilon}^T)\}]}_{\text{normal c.f.}} = \exp(\mathbf{c}\mathbf{A}^{*T}) \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \\ \implies E\{\tilde{f}(\mathbf{A}^*)|\mathbf{A}\} &= E\{\exp(\mathbf{c}\mathbf{A}^{*T})|\mathbf{A}\} \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \\ &= E\{\exp\{\mathbf{c}(\mathbf{A}^T + \epsilon^T)\}|\mathbf{A}\} \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \\ &= \exp(\mathbf{c}\mathbf{A}^T) \underbrace{E\{\exp(\mathbf{c}\epsilon^T)\}}_{\text{normal m.g.f.}} \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \\ &= \exp(\mathbf{c}\mathbf{A}^T) \exp\left(\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \exp\left(-\frac{1}{2}\mathbf{c}\boldsymbol{\Sigma}_{me}\mathbf{c}^T\right) \\ &= \exp(\mathbf{c}\mathbf{A}^T) = f(\mathbf{A})\end{aligned}$$