

The Koras–Russell cylinder as an affine modification of $\mathbb{A}^4$: a commuting square through a motivic 4-sphere and a one-point Borel–Moore defect

Brian Naughton*

Independent researcher

naughtonb@proton.me

ORCID: 0009-0008-3404-610X

2 August 2026

Abstract

Let $X = V(x^2y + z^2 + x + t^3) \subset \mathbb{A}^4$ be the Koras–Russell cubic threefold. Dubouloz, Moser-Jauslin and Poloni exhibited a locally nilpotent derivation ∂ on $\mathbb{C}[X \times \mathbb{A}^1]$ whose kernel does not contain x ; together with the standard x -preserving derivations this gives $\mathrm{ML}(\mathbb{C}[X \times \mathbb{A}^1]) = \mathbb{C}$. We dissect that action globally and follow where it leads. Its kernel is a polynomial ring in three variables, $\ker \partial = \mathbb{C}[p, q, t]$; it has no slice, exactly and with no degree bound, because it degenerates along a line; its plinth ideal is *not* principal, and the known subideal $(p+1)(t^3, q)$ it contains *encloses* the plinth’s zero locus in $V(p+1) \cup V(t, q)$ — divisorial in one component and of codimension two in the other, an asymmetry we state of that subideal rather than of the plinth, whose exact value we do not determine; the quotient morphism to $\mathbb{A}^3$ is neither surjective nor equidimensional; and after inverting the single function $t(p+1)$ the cylinder becomes a trivial $\mathbb{A}^1$ -bundle over $\mathbb{A}^3$. Reading that localisation as gluing data presents $X \times \mathbb{A}^1$ as an explicit *canonical affine modification* of $\mathbb{A}^4$ with divisor $F = t^3(p+1)$ and a four-generator centre, one of whose two components is singular along a line — an obstruction, with no degree bound, to carrying the centre to any smooth one by an automorphism of $\mathbb{A}^4$. The two components of the divisor give two commuting modification steps. One order factors through the singular cylinder $X_1 \times \mathbb{A}^1$ over Dubouloz–Moser-Jauslin–Poloni’s auxiliary threefold; the other factors through a *smooth* fourfold Y_1 , which we identify with the member $Q_{(3,1), z^2-1/4}$ of the Asok–Dubouloz–Østvær family of geometric models of algebraic suspensions — a model of the motivic 4-sphere. A four-corner motivic ledger then makes the surgery visible: the singular route is K_0 -neutral at each step, while the smooth route creates an $\mathbb{L}^2$ packet at the t -step and removes it at the $(p+1)$ -step, and the removing step’s exceptional locus fails to cover exactly one point of its centre. Finally, at integral Borel–Moore level, $H_4^{\mathrm{BM}}(Y_1; \mathbb{Z}) \cong \mathbb{Z}$ is generated by the fundamental class of *either* component of the doubled fibre over the origin, with $[F_0] = -[F_{-1}]$. The generator is supported on the modification divisor, having the representative F_{-1} contained in it, and restricts to zero on the common open set. Its inverse image $\sigma_4^{-1}(F_{-1})$ is $\mathbb{G}_m \times \mathbb{A}^1$ — the same plane with the fibre over the defect point $(-1, 0, 0, 0, 0)$ deleted — and the fundamental class of that locus, pushed forward along its *closed immersion* into the source, lands in $H_4^{\mathrm{BM}}(X \times \mathbb{A}^1) = 0$. We do not push classes forward along the modification itself: it

*The work was carried out with an AI research harness, described in full in Appendix A. All computations are reproducible from the repository accompanying this paper. Errors that remain are mine.

is not proper, as we prove. The defect thus accounts exactly for the Euler and Grothendieck-class change and locates, at a single named point, the cycle-level content of the change. This is a concrete singular-defect instance of the established cofiber mechanism for affine modifications. Nothing here bears on whether $X \times \mathbb{A}^1 \cong \mathbb{A}^4$, and no claim about that question is made.

Contents

1	Introduction	3
1.1	The setting	3
1.2	What this paper does	3
1.3	Positioning, and what is not claimed	5
1.4	Reproducibility	5
2	Notation, and the published starting datum	5
2.1	The objects of [5], §7	5
2.2	Conventions	6
3	The action, globally dissected	7
3.1	Content, kernel, and the reduced derivation	7
3.2	No slice, and the degeneracy locus	8
3.3	The localisation theorem	9
3.4	The plinth, and the quotient morphism	9
3.5	Two exact theorems at $k = 1$, and stability	10
4	The cylinder as a canonical affine modification	12
4.1	$A[w]$ inside its localisation	12
4.2	The compression	12
4.3	The centre, and a rectification obstruction	13
5	The commuting square	13
5.1	The tower, and the singular middle	13
5.2	The other order, and a smooth middle	13
6	The smooth middle identified	15
6.1	Three independent kills	15
6.2	The identification	15
7	The motivic ledger	16
7.1	Four corners, four arrows	16
7.2	Where the packet enters, and where it dies	17
7.3	Hodge–Deligne polynomials, and why they add nothing	17
7.4	The mechanism at Euler-characteristic level	17
8	The generator, and the cycle that dies	18
8.1	Groundwork	18
8.2	The generator	19
8.3	The cycle that dies	20

8.4	The statement, and the gap inside it	21
9	What this does and does not say	21
A	Methods	22
A.1	The harness	22
A.2	Certification discipline	22
A.3	Adversarial review	23
A.4	Limitations of the method	23
B	The verification layer	23

1 Introduction

1.1 The setting

Let

$$X = V(P) \subset \mathbb{A}_{x,y,z,t}^4, \quad P = x^2y + z^2 + x + t^3, \quad (1)$$

be the Koras–Russell cubic threefold. It is smooth and contractible, and it is not isomorphic to $\mathbb{A}^3$: Makar–Limanov’s invariant separates them, $\text{ML}(\mathbb{C}[X]) = \mathbb{C}[x]$ [15]. Whether the cylinder $X \times \mathbb{A}^1$ is isomorphic to $\mathbb{A}^4$ is a well-known open question, and a positive answer would refute the Zariski cancellation problem in characteristic zero and dimension three (see Gupta’s survey [10]). The classical obstruction does not survive the cylinder: Dubouloz proved that $\text{ML}(\mathbb{C}[X \times \mathbb{A}^1]) = \mathbb{C}$ [6]. Topology does not separate the two either. X is topologically contractible, and so is $\mathbb{A}^4$; more strongly, Koras–Russell threefolds of the first kind are $\mathbb{A}^1$ -contractible by Dubouloz–Fasel [7], extending Hoyois–Krishna–Østvær [11], whose theorem is for *finite suspensions*. Only the topological statement is used anywhere in this paper, and only in Remark 8.8.

The witness for the triviality of the Makar–Limanov invariant of the cylinder is a locally nilpotent derivation that moves x . Such a derivation is written down explicitly in §7 of Dubouloz, Moser-Jauslin and Poloni [5]. Their construction runs through an auxiliary threefold

$$X_1 = V(xy + z^2 + x + t^3) \subset \mathbb{A}^4, \quad (2)$$

an explicit conjugacy Φ, Ψ between the t -localised cylinders $A_t[w] \cong B_t[w]$ (Proposition 7.1), a triangular locally nilpotent derivation Δ on the X_1 -side transported to $\partial = \Phi \circ \Delta \circ \Psi$ on $A[w]$ with $\partial(x) = -2t^6(z + xw) \neq 0$ (Proposition 7.2), and the conclusion $\text{ML}(A[w]) = \mathbb{C}$ (Corollary 7.3). Here and throughout,

$$A = \mathbb{C}[x, y, z, t]/(P), \quad A[w] = \mathbb{C}[X \times \mathbb{A}^1], \quad B = \mathbb{C}[X_1], \quad B[w] = \mathbb{C}[X_1 \times \mathbb{A}^1]. \quad (3)$$

Everything in §2 below — the conjugacy, the derivation ∂ , the derivations ∂_1, ∂_2 , and the elements p, q, Z — is theirs. No novelty is claimed for any of it at any point in this paper.

1.2 What this paper does

This paper takes that published action as its starting datum and asks what the geometry underneath it looks like. The answer turns out to be a single connected picture, and the paper is organised as its six parts.

1. **The action, globally dissected** (§3). $\ker \partial = \mathbb{C}[p, q, t]$ is a polynomial ring in three variables; the content of ∂ is exactly t^3 ; the content-free derivation ∂_{red} has *no slice*, exactly and with no degree bound, because every component of it vanishes along a line of $X \times \mathbb{A}^1$; the plinth ideal is not principal, and contains $(p+1)(t^3, q)$, which already encloses its zero locus in $V(p+1) \cup V(t, q)$; the quotient morphism $\pi: X \times \mathbb{A}^1 \rightarrow \mathbb{A}^3$ is neither surjective nor equidimensional, for one certified reason; and $A[w][1/(t(p+1))] = \mathbb{C}[p, q, t][1/(t(p+1))][Z]$, so after inverting one explicit function the cylinder *is* a trivial $\mathbb{A}^1$ -bundle over $\mathbb{A}^3$. Two exact theorems then close the flow-compatible route at $k = 1$, and a third shows the obstruction is 1-stable.
2. **The cylinder as a canonical affine modification** (§4). Reading that localisation as gluing data yields $\mathbb{C}[X \times \mathbb{A}^1] = \mathbb{C}^{[4]}[I/F]$ with $F = t^3(p+1)$ and an explicit four-generator centre I , canonical in the sense recalled in §4. The centre is a surface with exactly two minimal primes; one is smooth, the other is singular along a line. Since $\text{Aut}(\mathbb{A}^4)$ carries centres to centres and preserves the singular locus of the centre, (I, F) is equivalent to no centre with smooth components — an obstruction to rectification with no degree bound.
3. **The commuting square** (§5). The two components of the divisor give two modification steps, and the steps commute. In one order the intermediate model is $X_1 \times \mathbb{A}^1$, which is singular, and $X_1 \times \mathbb{A}^k \not\cong \mathbb{A}^{3+k}$ for every $k \geq 0$. In the other order the intermediate model

$$Y_1 = V(t^3H - p - p^2 - qZ) \subset \mathbb{A}_{p,q,t,Z,H}^5$$

is *smooth*. Its defining equation is the governing identity of §3 rewritten. The two components of the centre meet non-reducedly, with multiplicity two, along one line.

4. **The smooth middle identified** (§6). Under an explicit linear change of coordinates of $\mathbb{A}^5$, Y_1 is the member $Q_{(3,1), z^2-1/4}$ of the family $Q_{m,P}$ of Asok, Dubouloz and Østvær [1]. Their results then give its smoothness and its $\mathbb{A}^1$ -homotopy type $\mathbb{P}^1 \wedge \mathbb{P}^1$, with complex realisation S^4 . Independently, a motivic computation of our own gives $Y_1 \times \mathbb{A}^r \not\cong \mathbb{A}^{4+r}$ for every $r \geq 0$, and a differential one shows that its defining polynomial is not a variable of $\mathbb{C}^{[5]}$, nor stably.
5. **The motivic ledger** (§7). Classes in the Grothendieck ring for all four corners of the square, and for the divisor and exceptional locus of all four arrows. The singular route is K_0 -neutral at each step; the smooth route creates an $\mathbb{L}^2$ packet at the t -step and removes it at the $(p+1)$ -step. Every stratum in the square is Hodge–Tate, so Hodge–Deligne polynomials carry nothing beyond the classes; what they do sharpen is that the packet is a Tate class of weight 4 and type $(2, 2)$, matching the reduced motive of $\mathbb{P}^1 \wedge \mathbb{P}^1$. At Euler-characteristic level the mechanism is exact, and it reduces to a single point: the removing arrow’s exceptional locus is empty over exactly one point of its centre.
6. **The final theorem** (§8). At integral Borel–Moore level, $H_4^{\text{BM}}(Y_1; \mathbb{Z}) \cong \mathbb{Z}$, generated by the fundamental class of *either* component of the doubled fibre over the origin of $\mathbb{A}_{t,q}^2$, with $[F_0] = -[F_{-1}]$. The generator is supported on the modification divisor — it has the representative F_{-1} contained in it — and restricts to zero on the common open set. Its inverse image is $\sigma_4^{-1}(F_{-1}) \cong \mathbb{G}_m \times \mathbb{A}^1$, the same plane with the fibre over the defect point $(-1, 0, 0, 0, 0)$ deleted, and that locus has fundamental class zero in $H_4^{\text{BM}}(X \times \mathbb{A}^1) = 0$, by the pushforward along its closed immersion. Dually and without any properness hypothesis, the cohomological pullback σ_4^* of the Poincaré dual of $[F_{-1}]$ vanishes.

A word on what *is not* asserted there, because an earlier draft of this paper asserted it. The modification σ_4 is *not proper* (Lemma 8.1), so there is no Borel–Moore pushforward along it and no sense in which a class on Y_1 “pushes forward” to $X \times \mathbb{A}^1$. What is proved is the list above: a support statement, a restriction statement, an identification of the inverse-image locus, the vanishing of that locus’s own fundamental class, and a functorial statement in cohomology. The one-point defect is what ties them together.

1.3 Positioning, and what is not claimed

What §8 exhibits is *a concrete singular-defect instance of the established cofiber mechanism for affine modifications*. The mechanism by which deleting branches over a modification centre destroys suspension summands is published: it is Proposition 4.3.1 of [1], where for $\pi: X_{I,a} \rightarrow \mathbb{A}^n$ with reduced smooth Y , deleting all but one component over the centre yields a strictly quasi-affine, $\mathbb{A}^1$ -contractible variety. Modifications along singular and non-reduced centres have been studied by Dubouloz, Pauli and Østvær [8]; we quote no result of theirs, because our centre is singular exactly at the point that carries the phenomenon, which their hypotheses exclude. What this paper contributes is an explicit case in which every object is written down, the defect is a single named point, and the cycle that dies is identified.

The following are *not* claimed, anywhere, in any form.

- Nothing here bears on whether $X \times \mathbb{A}^1 \cong \mathbb{A}^4$, or whether $X \times \mathbb{A}^2 \cong \mathbb{A}^5$. Both remain exactly as open as before. The results of §3 are properties of the *action*, not separating invariants of the space, and must not be read as if they might be.
- Y_1 is not a new fourfold, not a previously unknown deformed quadric, and not a new model of the motivic sphere. It is a member of a published family, cited at every use. What appears to be new is its *role*: it is the smooth intermediate completion of a commuting affine-modification square attached to the Koras–Russell cylinder.
- Several statements below are the outcome of searches at declared, bounded caps. Those are recorded as measurements of what was searched, never as negative results, and are marked as such wherever they occur.

1.4 Reproducibility

Every algebraic identity in this paper is verified by machine. The accompanying repository contains the runners, the frozen ledgers they emit, and a single script that re-runs the whole suite in a fresh environment and reports the pass counts. Appendix B lists the layer and states the expected totals; Appendix A describes the harness that produced it, and the disciplines — twin implementations, positive controls, boundary counter-checks, declared caps, replay-identical ledgers — that the work was held to.

2 Notation, and the published starting datum

2.1 The objects of [5], §7

With $P = x^2y + z^2 + x + t^3$ and $S = xy + z^2 + x + t^3$ the defining polynomials of X and X_1 , [5, Prop. 7.1] gives mutually inverse isomorphisms $\Phi: B_t[w] \rightarrow A_t[w]$ and Ψ between the t -localised

cylinders, with

$$\begin{aligned}\Phi(x) &= x, & \Phi(y) &= xy - xw^2 - 2zw, & \Phi(z) &= z + xw, & \Phi(t) &= t, \\ \Phi(w) &= 2w + yz + 3xyw - 3zw^2 - xw^3.\end{aligned}\tag{4}$$

[5, Prop. 7.2] transports the triangular locally nilpotent derivation $\Delta = t^6(-2z\partial_x + (y+1)\partial_z)$ to a locally nilpotent derivation $\partial = \Phi \circ \Delta \circ \Psi$ on $A[w]$ with

$$\partial(x) = -2t^6(z + xw) \neq 0,\tag{5}$$

and [5, Cor. 7.3], with $\partial_1 = 2z\partial_y - x^2\partial_z$ and $\partial_2 = 3t^2\partial_y - x^2\partial_t$, concludes $\text{ML}(A[w]) = \mathbb{C}$.

Throughout we write

$$\begin{aligned}p &= \Phi(y) = xy - xw^2 - 2zw, & Z &= \Phi(z) = z + xw, \\ q &= \Phi(w) = 2w + yz + 3xyw - 3zw^2 - xw^3,\end{aligned}\tag{6}$$

and set

$$H := -(y + w^2).\tag{7}$$

The elements p, q, Z and the t -localised bridge are [5]'s. The element H , and everything that follows, is ours.

Remark 2.1 (Two errata in the printed source, disclosed with thanks). The starting datum was machine-verified against the paper's own printed identities before anything downstream used it — a routine gate, and both findings below are consequences of it rather than of any search. Both are harmless to the conclusions of [5], and we record them only so that a reader who recomputes finds the same thing we did.

- (i) [5, Prop. 7.1] prints, as the sixth of its composition identities, $\Phi \circ \Psi(w) = w - \frac{1}{t^3}P$. The verified value is $\Phi \circ \Psi(w) = w - \frac{w}{t^3}P$: the numerator is w , not 1. The transcription was re-read from the source in layout mode before this was concluded, and the remaining seven identities over-determine every component of Φ and Ψ , so the maps themselves are certified. Both forms are congruent to w modulo (P) , so the conclusion of Proposition 7.1 is unaffected. It is a typographical slip.
- (ii) [5, Cor. 7.3] states $\text{Ker}(\partial_1) \cap \text{Ker}(\partial_2) = \mathbb{C}[x]$ for the derivations *on* $A[w]$; but w lies in both kernels, so on $A[w]$ the intersection contains $\mathbb{C}[x][w]$. The classical statement is the one on A ; on $A[w]$ one adjoins ∂_w , which the paper's own framework supplies, and the conclusion $\text{ML}(A[w]) = \mathbb{C}$ stands. We verified the repair by machine: x is killed by all three of $\partial_1, \partial_2, \partial_w$, and $\partial(x) \neq 0$.

2.2 Conventions

$\mathbb{C}^{[n]}$ denotes a polynomial ring in n variables over $\mathbb{C}$. For a locally nilpotent derivation δ of a domain R , $\ker \delta$ is its ring of constants, the *content* of δ is the greatest common divisor of its values on a generating set, a *slice* is an s with $\delta(s) = 1$, a *local slice* is an s with $\delta^2(s) = 0 \neq \delta(s)$, and the *plinth ideal* is $\text{pl}(\delta) = \delta(R) \cap \ker \delta$. $\mathbb{L}$ denotes the class of $\mathbb{A}^1$ in the Grothendieck ring of complex varieties and χ_c the compactly supported Euler characteristic. H_*^{BM} denotes Borel–Moore homology with $\mathbb{Z}$ coefficients, of the associated complex analytic space.

Following [1] we write $Q_{m,P}$ for the members of their family of geometric models of algebraic suspensions; the notation is recalled at (13) below.

Definition 2.2 (Affine modification, and canonical centre). Let R be a $\mathbb{C}$ -algebra, $f \in R$ non-zero and $I \ni f$ an ideal. The *affine modification* of R with divisor f and centre I is the subalgebra $R[I/f] \subset R_f$ generated by R and $\{g/f : g \in I\}$; see Kaliman–Zaidenberg [12]. We call the centre *canonical* when $I = f \cdot R[I/f] \cap R$. This is the right normalisation here: the naive saturation $I : f^\infty$ is vacuous, since it equals (1) whenever $f \in I$, which it must be for $R[I/f]$ to be defined.

Remark 2.3 (Terminology, adopted deliberately). Two phrases are used below in a restricted sense, and never loosely. We say *flat family with generic fibre $\mathbb{A}^3$ and one quadric degeneration*, and never “ $\mathbb{A}^3$ -fibration” unqualified, for the morphism of Proposition 5.5; the degenerate fibre is smooth but not contractible, so it is not an $\mathbb{A}^3$. And we use *standard centre* only with the following explicit equivalence, which is what Theorem 4.7 rules on: the pairs (I, f) and (I', f') over $R = \mathbb{C}^{[4]}$ are *equivalent* when some $\alpha \in \text{Aut}(\mathbb{A}^4)$ carries f to a unit multiple of f' and carries the canonical centre $f \cdot R[I/f] \cap R$ of the first to the canonical centre of the second. The third clause is the canonicalisation of Definition 2.2, and nothing else; in particular it is *not* a saturation, for the reason recorded there.

3 The action, globally dissected

Throughout this section ∂ is the derivation (5) of [5].

3.1 Content, kernel, and the reduced derivation

Multiplying a locally nilpotent derivation by a kernel element moves the plinth but not the orbits, so the slice question belongs to the content-free derivation and the content must be stripped first — inside $A[w]$, not on representatives. For “the content” to be well defined at all, $A[w]$ must be a unique factorisation domain, which we record first.

Lemma 3.1 (Factoriality). *$A[w]$ is a unique factorisation domain.*

Proof. By Nagata’s criterion it suffices to exhibit a prime element f of the Noetherian domain $A[w]$ with $A[w]_f$ a UFD. Take $f = x$. Both hypotheses are certified.

x is prime. $P \equiv z^2 + t^3 \pmod{(x)}$, so $A[w]/(x) = \mathbb{C}[y, z, t, w]/(z^2 + t^3)$; and $z^2 + t^3$ is irreducible over $\mathbb{C}$, since as a quadratic in z over the UFD $\mathbb{C}[t]$ it would otherwise factor, which needs $-t^3$ to be a square in $\mathbb{C}[t]$, and it is not (odd degree). So this is a domain.

$A[w]_x$ is a UFD. P has degree one in y with leading coefficient x^2 , which is a unit after inverting x ; solving gives $y = -(z^2 + x + t^3)/x^2$ and hence $A_x = \mathbb{C}[x^{\pm 1}, z, t]$, so $A[w]_x = \mathbb{C}[x^{\pm 1}, z, t][w]$ — a localisation of a polynomial ring, and therefore a UFD. $\square$

The irreducibility test carries its own control: the same routine reports $z^2 - t^2$ reducible, so the verdict on $z^2 + t^3$ is not vacuous.

Proposition 3.2 (Content). *The t -adic valuations in $A[w]$ of $\partial(x), \partial(y), \partial(z), \partial(w)$ are 6, 3, 3, 3, each maximal and each certified by an explicit witness quotient. The content of ∂ is exactly t^3 , and $\partial_{\text{red}} := \partial/t^3$ is content-free with $\partial_{\text{red}}(x) = -2t^3(xw + z)$.*

The exclusion of further content is by a *closed* candidate list, not a sample: any common divisor divides $\partial(x) = -2t^6(z + xw)$; t is prime in $A[w]$; and $xw + z$ is irreducible in $A[w]$, being of degree one in w and primitive because $V(x, z) \cap X = \{x = z = t = 0\}$ is the y -line, of

codimension two in X — so x and z have no common factor in the UFD A . Hence $(xw + z)$ divides only the x -component.

Remark 3.3. The polynomial *representative* of $\partial(y)$ is t -free, while its *class* in $A[w]$ has t -valuation 3. The representative is not the invariant, and the strip must be done in $A[w]$. One immediate consequence: ∂ itself can never have a slice, since $\partial(A[w]) \subseteq t^3 A[w]$, so the slice question below is asked of ∂_{red} , whose kernel is the same.

Theorem 3.4 (The kernel). $\ker(\partial \mid A[w]) = \mathbb{C}[p, q, t] \cong \mathbb{C}^{[3]}$.

Proof, as certified. On the X_1 -side put $u = y+1$, so that S becomes $xu + z^2 + t^3$; inverting u solves the relation for x and gives $B_t[w][1/u] = \mathbb{C}[t^{\pm 1}, w, u^{\pm 1}, z]$ with $\Delta = t^6 u \partial_z$, a polynomial ring in the *single* variable z over its ring of constants. Hence $\ker(\Delta \mid B_t[w][1/u]) = \mathbb{C}[t^{\pm 1}, w, u^{\pm 1}]$. Now u is prime in $B_t[w]$, because $B_t[w]/(u) = \mathbb{C}[x, z, t^{\pm 1}, w]/(z^2 + t^3)$ is a domain, $z^2 + t^3$ being irreducible over $\mathbb{C}[t^{\pm 1}]$; therefore $\ker(\Delta \mid B_t[w]) = \mathbb{C}[t^{\pm 1}][y, w]$. Since $\Phi(S) = P$ and $\Psi(P) \in (S)$, the maps Φ and Ψ are mutually inverse between $B_t[w]$ and $A_t[w]$, and $\partial = \Phi \circ \Delta \circ \Psi$, so $\ker(\partial \mid A_t[w]) = \mathbb{C}[t^{\pm 1}][p, q]$. Finally the t -saturation is trivial: p and q remain algebraically independent modulo t (Jacobian rank two) in the *domain* $A[w]/(t) = \mathbb{C}[x, y, z, w]/(x^2 y + z^2 + x)$, so $\mathcal{H}(p, q, t) \in tA[w]$ forces the t -free part $\mathcal{H}_0$ to vanish, i.e. $t \mid \mathcal{H}$ in $\mathbb{C}[p, q, t]$; induct. $\square$

The route deliberately avoids elimination: it is a conjugacy transport followed by a saturation argument, and no Gröbner basis is computed in it. Independently, $\partial(p) = \partial(q) = 0$ was verified through two independent implementations, and $\Psi(p) = y$, $\Psi(q) = w$ modulo (S) .

3.2 No slice, and the degeneracy locus

Proposition 3.5 (Degeneracy locus). *The degeneracy ideal of ∂_{red} contains t^6 , so its zero locus is supported on $\{t = 0\}$; and the locus is non-empty, since the line $\{x = z = t = w = 0\} \cong \mathbb{A}_y^1$ consists of common zeros of all components. In particular the $\mathbb{G}_a$ -action is not free. Chart by chart: on $\{x \neq 0, t = 0\}$ the locus is exactly $V(Z^2(Z^2 + x)) = V(Z) \cup V(p)$, using $Z^2 + x \equiv -xp$ modulo (P, t) ; on $\{x = 0\}$ the fibre forces $z = 0$ and the locus is exactly $\{w = 0\}$, the line above, which is the $w = 0$ slice of the cuspidal edge $\{z = t = 0\}$ of the $x = 0$ cusp-cylinder fibre.*

Each of x, z, y, w was verified *not* to lie in the radical of the degeneracy ideal; only t is forced. The radical test can fail silently, and the check that it does not is part of the certificate.

Theorem 3.6 (No slice). ∂_{red} , and hence ∂ , admits no slice on $A[w]$. The statement is exact and carries no degree bound.

Proof, as certified. Two independent routes. (i) Every component of ∂_{red} vanishes at the origin of $X \times \mathbb{A}^1$, so by the chain rule $\partial_{\text{red}}(F)$ vanishes there for *every* $F \in A[w]$, while 1 does not. (ii) A slice would give $A[w] = (\ker \partial)[s]$, so the $\mathbb{G}_a$ -action would be a translation and hence free; Proposition 3.5 exhibits a line of fixed points. $\square$

A bounded exact search over $\mathbb{Q}$ in degrees $\leq 1, 2, 3$ (6, 21 and 56 unknowns; a linear nullspace computation, not an elimination) agrees, and carries a positive control: the same solver does find the slice w for ∂_w , so the empty verdicts are not vacuous.

Since $\ker \partial \cong \mathbb{C}^{[3]}$ by Theorem 3.4, a slice would have given $A[w] \cong \mathbb{C}^{[4]}$ outright. Theorem 3.6 closes that route, and only that route: nothing is claimed or implied about $X \times \mathbb{A}^1 \cong \mathbb{A}^4$.

3.3 The localisation theorem

Theorem 3.7 (Localisation). *With $Z = z + xw$ one has $\partial_{\text{red}}(Z) = t^3(p+1)$ and*

$$A[w]_{\left[\frac{1}{t(p+1)}\right]} = \mathbb{C}[p, q, t]_{\left[\frac{1}{t(p+1)}\right]}[Z].$$

The proof is by exhibiting x, y, z, w explicitly in those generators, which is what the certificate does. Geometrically: $X \times \mathbb{A}^1$ is a trivial $\mathbb{A}^1$ -bundle over the complement of the single divisor $\{t(p+1) = 0\}$ in $\mathbb{A}^3 = \text{Spec}(\ker \partial)$. The whole obstruction through this flow is a boundary-globalisation problem across one explicit reducible divisor. That reading is what §4 makes into a modification.

3.4 The plinth, and the quotient morphism

Proposition 3.8 (Values of ∂_{red}). *On the natural generators, ∂_{red} is given by*

$$\partial_{\text{red}}(Z) = t^3(p+1), \quad \partial_{\text{red}}(x) = -2t^3Z, \quad \partial_{\text{red}}(w) = \frac{1}{2}(p^2 + p + 2qZ), \quad \partial_{\text{red}}(H) = q(p+1),$$

with p, q, t killed; and on the localisation of Theorem 3.7 it is $t^3(p+1) \cdot d/dZ$. Every value was checked on both sides of the conjugacy.

Proposition 3.9 (The plinth is not principal). *$H = -(y + w^2)$ is a local slice: $\partial_{\text{red}}^2(y + w^2) = 0$ and $\partial_{\text{red}}(y + w^2) = -q(p+1)$. Since $q(p+1) \notin (t^3(p+1))$ — it survives $t = 0$ —*

$$\text{pl}(\partial_{\text{red}}) \supseteq (p+1)(t^3, q),$$

and $\text{pl}(\partial_{\text{red}}) \subseteq (p, q, t)$ is proper, so 1 is not in the plinth. Whether the first containment is an equality is not decided.

The containment $\text{pl} \subseteq (p, q, t)$ follows from $\partial_{\text{red}}(A[w]) \subseteq (x, z, t, w)A[w]$, which is proper and sharply so: the image is *not* inside (x, z, t) . A search at a cap declared before it was run — total degree ≤ 4 in x, y, z, t, w , 126 monomials, a 20-dimensional solution space — returned seven plinth values, all inside $(p+1)(t^3, q)$; that is a measurement of what was searched, and settles nothing about equality.

Proposition 3.10 (Enclosure of the plinth's zero locus). *Write $J = (p+1)(t^3, q) \subseteq \text{pl}(\partial_{\text{red}})$ for the subideal of Proposition 3.9. Then*

$$\emptyset \neq V(\text{pl}(\partial_{\text{red}})) \subseteq V(J) = V(p+1) \cup V(t, q).$$

The inclusion holds because a larger ideal has a smaller zero locus, so no component of $V(\text{pl})$ can lie outside $V(J)$; and $V(\text{pl})$ is non-empty because $\text{pl} \subseteq (p, q, t)$ is proper. Moreover $\sqrt{J} = (p+1) \cap (t, q)$, with minimal primes $(p+1)$ of codimension one and (t, q) of codimension two.

Remark 3.11 (What the asymmetry is, and is not, a statement about). The two minimal primes of $\sqrt{J}$ are of different kinds: $(p+1)$ is divisorial, while (t, q) has codimension two — the t -wall enters only cut against $q = 0$. *This is a statement about the known subideal J , not about $\text{pl}(\partial_{\text{red}})$ itself*, whose exact value we do not determine (Proposition 3.9); a larger plinth could only shrink $V(\text{pl})$ further inside $V(J)$, and we make no claim about which of the two components survives in $V(\text{pl})$. What is unconditional is the enclosure.

Shakhmatov identifies the zero locus of the plinth ideal with the complement of the union of the principal invariant cylinders [17]. In that reading the enclosure is the useful direction, and it survives intact: since $V(\text{pl}) \subseteq V(p+1) \cup V(t, q)$, the principal invariant cylinders *cover* the complement of $V(p+1) \cup V(t, q)$ — everywhere off those two loci, the action has a principal invariant cylinder through the point. The converse reading, which would need the exact plinth, is not available to us and is not used.

Proposition 3.12 (The quotient morphism). *Let $\pi: X \times \mathbb{A}^1 \rightarrow \mathbb{A}^3 = \text{Spec } \mathbb{C}[p, q, t]$. Then π is not surjective and not equidimensional: the fibres over $(2, 0, 0)$ and $(-1, 3, 0)$ are empty, and the fibres over $(0, 0, 0)$ and $(-1, 0, 0)$ are surfaces, against the generic dimension one. The image is nonetheless dense.*

Proposition 3.13 (The governing identity). *In $A[w]$,*

$$qZ + p + p^2 + t^3(y + w^2) = 0, \quad (8)$$

equivalently $t^3H = p + p^2 + qZ$. At $t = 0$ this reads $qZ = -p(1+p)$, so $q = 0$ forces $p(1+p) = 0$; the measured fibre table matches exactly, the fibre over $t = 0$ being an $\mathbb{A}^1$ where $q \neq 0$ except at $p = -1$ where it is empty, and empty where $q = 0$ except at $p = 0$ and $p = -1$, which are precisely the two dimension jumps.

Identity (8) is the single reason for Proposition 3.12, and it is also, rewritten, the defining equation of the smooth intermediate model of §5. It is worth recording which component of the divisor misbehaves, because the natural guess is wrong: $\{p+1=0\}$ *is tame on its own* — over $t \neq 0$ the fibres are $\mathbb{A}^1$ even along $p = -1$. Every failure lives over $\{t = 0\}$, and $p = -1$ bites only in combination with it.

hypothesis of [16, Thm. 4.3]	verdict	source
free $\mathbb{G}_a$ -action	fails	Proposition 3.5
affine quotient exists and is $\mathbb{A}^3$	holds	Theorem 3.4
quotient morphism surjective	fails	Proposition 3.12
quotient morphism equidimensional	fails	Proposition 3.12
$\mathbb{A}^3$ -fibration with factorial closed fibres	fails at the cusp	[16, Ex. 4.2]

Table 1: Masuda’s dimension-four partial positive theorem, tested against this action. One hypothesis holds and four fail. This says nothing about $X \times \mathbb{A}^1 \cong \mathbb{A}^4$; it says that *this* action cannot carry *that* theorem.

3.5 Two exact theorems at $k = 1$, and stability

Theorem 3.14. *$\text{pl}(\partial_{\text{red}})$ is not principal. Consequently there is no $s \in A[w]$ with $A[w] = \mathbb{C}[p, q, t][s]$.*

Proof. The plinth contains $t^3(p+1)$ and $q(p+1)$. If $\text{pl} = (g)$ then g divides their greatest common divisor, which is $(p+1)$ because $\gcd(t^3, q) = 1$. As $(p+1)$ is irreducible, g is a unit or an associate of $p+1$; a unit puts 1 in the plinth, and an associate puts $p+1$ in the plinth, and both contradict $\text{pl} \subseteq (p, q, t)$. For the consequence: $A[w] = R[s]$ with $R = \ker \partial_{\text{red}}$ forces $\partial_{\text{red}} = \partial_{\text{red}}(s) \cdot d/ds$, hence a principal plinth. $\square$

The refutation of the principal-plinth guess in Proposition 3.9 is the essential input here: without the second generator $q(p+1)$ the argument does not run.

Theorem 3.15. *Independently: a coordinate projection $\mathbb{A}^4 \rightarrow \mathbb{A}^3$ is surjective with every fibre an $\mathbb{A}^1$, and Proposition 3.12 exhibits an empty fibre and a surface fibre. Hence again $A[w] \neq \mathbb{C}[p, q, t][s]$.*

Theorem 3.16 (1-stability, in the sense of [14]). *Extending ∂ to $A[w][v]$ by $\partial(v) = 0$ gives $\ker = \mathbb{C}[p, q, t, v]$ and $\text{pl}(\partial \mid A[w][v]) = \text{pl}(\partial \mid A[w]) \cdot \mathbb{C}[v]$, which still contains both generators and still lies inside (p, q, t) ; and $\gcd(t^3, q) = 1$, the irreducibility of $p+1$, and $p+1 \notin (p, q, t)$ all persist in $\mathbb{C}[p, q, t, v]$. Hence $A[w][v] \neq \mathbb{C}[p, q, t, v][s]$ for any s : the obstruction of Theorem 3.14 is 1-stable.*

Remark 3.17 (The classification, and a cocycle that does not exist). The two components of the divisor obstruct in different ways. $\{t = 0\}$ blocks the flow-compatible route *exactly, at order three*: the non-principality of the plinth is the failure of t^3 and q to share a factor, and both witnesses of Theorem 3.15 lie over $t = 0$. $\{p+1 = 0\}$ blocks the elementary moves, and the reason is the cusp: under every shear $Z \mapsto Z + f$ the leading Z -coefficient of x is the invariant $-1/(p+1)$ — verified with f symbolic, hence exactly over the whole family — and a rescaling that would fix it needs $Z/(p+1) \in A[w]$, while $\dim\{p+1 = 0\} = 3$ against $\dim\{p+1 = 0, Z = 0\} = 2$. Z does not vanish there because $R1$ of Proposition 4.1 restricted to $\{p+1 = 0\}$ is the cuspidal curve $Z^2 + t^3 = 0$ — the same cusp that carries X_1 's singularity and breaks factoriality in Table 1. Finally, a boundary cocycle in the Danielewski–Asanuma sense presupposes two $\mathbb{A}^1$ -fibrations over a common base agreeing on an overlap; $\mathbb{C}[p, q, t, Z]$ is such a fibration but $A[w]$ is not, by Proposition 3.12, so that template cannot be set up through this presentation at all.

Remark 3.18 (Brackets: Corollary 7.3 as linear algebra). All six brackets among $\{\partial_{\text{red}}, \partial_1, \partial_2, \partial_w\}$ were computed. *Exactly* three pairs commute, and they are exactly the triple $\{\partial_1, \partial_2, \partial_w\}$ of [5, Cor. 7.3]; ∂_{red} commutes with none of them. The four derivations do form a generic frame on $X \times \mathbb{A}^1$, in that the 4×4 determinant of their values on the coordinates is non-zero in the model, and dropping ∂_{red} drops the rank to three. So ∂_{red} is precisely the direction the commuting triple cannot reach — which is Corollary 7.3's Makar-Limanov argument seen as linear algebra. No triangular flag through ∂_{red} was found; that is a statement about what these six brackets show, and not a claim that none exists.

Remark 3.19 (Bounded searches, recorded as measurements). All 126 four-element subsets of a nine-element pool declared in advance were filtered at $k = 1$ by the exact criterion that $\det[\nabla P; \nabla a_1, \dots, \nabla a_4]$ vanish nowhere on $X \times \mathbb{A}^1$ — valid because $X \times \mathbb{A}^1$ is smooth, which was verified. No subset survived: nine determinants were identically zero and 117 vanished somewhere. At $k = 2$, all 2002 five-element subsets of a declared fourteen-element pool were filtered by the same criterion in six variables; 1852 were rejected by an explicit witness point on $X \times \mathbb{A}^2$ and 150 by the exact test. A positive control — a graph hypersurface whose coordinate ring genuinely is $\mathbb{C}^{[4]}$ — passes the same filter in both settings, so the rejections are not vacuous. These are measurements at declared caps and prove nothing beyond their pools.

4 The cylinder as a canonical affine modification

4.1 $A[w]$ inside its localisation

Proposition 4.1 (Intrinsic description). $A[w] = \mathbb{C}[p, q, t, Z][x, w, H]$, certified in both directions: x, w, H are polynomials in x, y, z, t, w , and conversely $z = Z - xw$ and $y = -H - w^2$. Each of x, w, H is certified not to lie in $\mathbb{C}[p, q, t, Z]$ by an explicit pole. The pole orders along the two components, measured separately, are

	x	H	w	y	z
ord along $\{t = 0\}$	0	-3	-3	-6	-3
ord along $\{p + 1 = 0\}$	-1	0	-1	-2	-2

with p, q, t, Z of non-negative order along both. Four relations hold in $A[w]$:

$$R1 : \quad x(p + 1) + Z^2 + t^3 = 0, \quad R2 : \quad 2t^3w = pZ - xq, \quad (9)$$

$$R3 : \quad t^3H = p + p^2 + qZ, \quad R4 : \quad 2(p + 1)w = HZ + q. \quad (10)$$

So each component of the divisor contributes exactly one *primitive* division — x for $\{p + 1 = 0\}$, H for $\{t = 0\}$ — and w is the mixed one. Note also:

Proposition 4.2. $A[w]$ is not cut out of $\mathbb{C}[p, q, t, Z][1/(t(p + 1))][x, w, H]$ by any valuation condition: pole orders are unbounded on both sides, $\text{ord}_{p+1}(x^k) = -k$ and $\text{ord}_t(H^k) = -3k$, both verified. What cuts it out is an affine modification.

Remark 4.3. $R1$ is exactly [5]’s own S , renamed by $y \mapsto p$, $z \mapsto Z$, $u = p + 1$. The middle term of the natural factorisation is therefore their auxiliary threefold, not a new object; see §5. $R3$ is the governing identity (8).

4.2 The compression

Theorem 4.4 (The compressed modification). *Set*

$$F = t^3(p + 1), \quad N_1 = p + p^2 + qZ, \quad M = p(p + 1)Z + q(Z^2 + t^3),$$

$$I = (F, t^3(Z^2 + t^3), (p + 1)N_1, M).$$

Then $\mathbb{C}[X \times \mathbb{A}^1] = \mathbb{C}^{[4]}[I/F]$, an affine modification of $\mathbb{C}^{[4]} = \mathbb{C}[p, q, t, Z]$ with divisor F and centre I ; the centre is canonical in the sense of Definition 2.2; and the three adjoined generators are exactly

$$x = -\frac{t^3(Z^2 + t^3)}{F}, \quad H = \frac{(p + 1)N_1}{F}, \quad 2w = \frac{M}{F}.$$

Moreover F divides $(t(p + 1))^3$, with quotient $(p + 1)^2$.

Canonicity is proved exactly, not sampled: for the $(p + 1)$ -step because $p + 1$ is prime and does not divide $Z^2 + t^3$, and for the t -step because t is prime in the intermediate ring — $x(p + 1) + Z^2$ being irreducible — and divides neither numerator. That t^3 is the least power of t clearing the new generators is decided rather than assumed, and so is the failure of the smaller choice: $N_1 \notin I^3 + tI^2 + t^2I + (t^3)$, with the control that t^3 does lie in the corresponding ideal. A further control: F does *not* clear y , which needs t^6 .

Remark 4.5 (A correction, logged). The centre was expected by hand to need five generators, the fifth being $F \cdot xH = -(Z^2 + t^3)N_1$. It needs four: the machine returned the exact identity

$$(Z^2 + t^3)N_1 = Z \cdot M + p \cdot F.$$

4.3 The centre, and a rectification obstruction

Proposition 4.6 (Geometry of the centre). *$V(I)$ is a surface with exactly two minimal primes, and they are the two steps of §5 pushed into $\mathbb{A}^4$:*

$$V(t, N_1) \quad \text{which is smooth,} \quad V(p+1, Z^2 + t^3) \quad \text{which is singular along a line,}$$

the singular line being the cuspidal edge $\{Z = t = 0\}$ with q free.

Theorem 4.7 (A rectification obstruction with no degree bound). *An automorphism $\alpha \in \text{Aut}(\mathbb{A}^4)$ carrying (I, F) to (I', F') in the sense of Remark 2.3 carries the canonical centre of the one to the canonical centre of the other, hence restricts to an isomorphism $V(I) \rightarrow V(I')$ of the centres as schemes, and therefore matches their singular loci. One component of $V(I)$ is singular along a line (Proposition 4.6). Hence (I, F) is not equivalent to any pair whose centre has smooth components: not to a linear subspace, and not to any smooth complete intersection.*

Every obvious standard-centre target is removed, and with no degree bound. What survives as a possible rectification target is a *singular* standard centre carrying a cusp — and the surviving singularity is the same cusp that appears in Proposition 3.5, in Remark 3.17, and in Table 1.

5 The commuting square

5.1 The tower, and the singular middle

The two components of the divisor $\{t(p+1) = 0\}$ carry one modification step each. Taking the $(p+1)$ -step first gives the tower of inclusions

$$\mathbb{C}[p, q, t, Z] = \mathbb{C}^{[4]} \subset B[w] = \mathbb{C}[X_1 \times \mathbb{A}^1] \subset A[w] = \mathbb{C}[X \times \mathbb{A}^1], \quad (11)$$

that is, birational morphisms $X \times \mathbb{A}^1 \rightarrow X_1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^4$, each an isomorphism off exactly one component. Step one has divisor $p+1$ and centre $(p+1, Z^2 + t^3)$ and adjoins x ; step two has divisor t^3 and centre (t^3, N_1, N_2) with $N_2 = pZ - xq$, because relative to $B[w]$ both remaining generators $H = N_1/t^3$ and $w = N_2/(2t^3)$ are t^3 -divisions.

Theorem 5.1. *X_1 is singular exactly at the origin, so $X_1 \times \mathbb{A}^1$ is a fourfold whose singular locus is the line $\{\text{origin of } X_1\} \times \mathbb{A}^1$. Singularity survives $\times \mathbb{A}^k$ and $\mathbb{A}^{3+k}$ is smooth, so*

$$X_1 \times \mathbb{A}^k \not\cong \mathbb{A}^{3+k} \quad \text{for every } k \geq 0.$$

In particular neither step of the tower is an isomorphism.

The control for the routine that produced this is that it reports an empty singular locus for a smooth graph hypersurface.

5.2 The other order, and a smooth middle

Theorem 5.2. *The two steps commute — away from each divisor the other step is trivial, over $D(t)$ only the $(p+1)$ -step surviving and over $D(p+1)$ only the t -step — and running the t -step first gives a second intermediate model*

$$Y_1 = V(W) \subset \mathbb{A}_{p,q,t,Z,H}^5, \quad W = t^3H - p - p^2 - qZ, \quad (12)$$

with $A[w] = \mathbb{C}^{[4]}[H][x, 2w]$, the second stage being the $(p+1)$ -division $2w = (HZ + q)/(p+1)$, which is relation R4 of Proposition 4.1. And Y_1 is smooth: the Jacobian forces $p = -\frac{1}{2}$, $Z = q = t = 0$, and at such a point $W = \frac{1}{4} \neq 0$.

The control here is that the same routine still finds $X_1 \times \mathbb{A}^1$ singular.

Remark 5.3 (A framing corrected). So the same birational morphism $X \times \mathbb{A}^1 \rightarrow \mathbb{A}^4$ factors through the *singular* $X_1 \times \mathbb{A}^1$ and through the *smooth* Y_1 . Theorem 5.1 is untouched — $X_1 \times \mathbb{A}^k$ is never affine space — but the reading that the factorisation must pass through a singular model is presentation-dependent, and we record it as such.

Proposition 5.4 (The transition equation). *The governing identity (8) rewrites as $Zq - t^3H = -p(1+p)$, which is precisely the defining equation of Y_1 ; completing the square,*

$$Zq - t^3H + P'^2 = \frac{1}{4}, \quad P' = p + \frac{1}{2},$$

a deformed affine quadric with t^3 where a variable would be. What the identity transitions between is $\mathbb{C}^{[4]}$ and Y_1 .

Proposition 5.5. $Y_1 \rightarrow \mathbb{A}_t^1$ has fibre $\mathbb{A}^3$ over $t \neq 0$, where H is a graph, and degenerates over $t = 0$ to (smooth affine quadric surface) $\times \mathbb{A}_H^1$, which is smooth but not contractible, hence not an $\mathbb{A}^3$. In the language of Remark 2.3: a flat family with generic fibre $\mathbb{A}^3$ and one quadric degeneration.

Proposition 5.6 (The crossing). *On $V(t, p+1) = \mathbb{A}_{q,Z}^2$ the compressed centre I of Theorem 4.4 restricts to (qZ^2) . The smooth and cuspidal components of $V(I)$ meet exactly along the line $\{t = 0, p = -1, Z = 0\}$, and they meet non-reducedly: Z^2 lies in the intersection ideal and Z does not. The multiplicity-two doubling is the cusp $Z^2 + t^3$ read at $t = 0$.*

All the genuinely coupled geometry of the square concentrates, scheme-theoretically, at that crossing.

Remark 5.7 (The transition datum, and a method acknowledged). The two trivialisations are the two models of Theorem 5.2: over $D(t)$ the singular $X_1 \times \mathbb{A}^1$, over $D(p+1)$ the smooth Y_1 , agreeing over the common open set. The transition is therefore a change of *integral model*, not a bundle automorphism, and the two models disagree over the crossing. Blanc and Poloni [2] study the Daigle–Freudentburg family containing the Vénéreau polynomials, show after Kaliman–Zaidenberg that these are locally trivial $\mathbb{A}^2$ -bundles over a punctured plane of a specific form X_f , and put bivariables in bijection with the trivial ones. We record explicitly that no theorem of theirs applies here — their objects are $\mathbb{A}^2$ -bundles over a punctured plane, ours a birational modification of $\mathbb{A}^4$ along a codimension-two centre with four-dimensional hypersurface models. Only the method transfers, and we say so in order that a method not be mistaken for a citation. See also Kaliman–Vénéreau–Zaidenberg [13] for the rectification framing of simple birational extensions.

Remark 5.8 (Bounded searches, again recorded as measurements). Three further cap-bounded sweeps, with declared pools, the exact Jacobian criterion, explicit witness points for rejections and positive controls throughout: $A[w] \cong \mathbb{C}^{[4]}$? — 495 four-subsets of a twelve-element pool, no survivors. $A[w][v] \cong \mathbb{C}^{[5]}$? — 4368 five-subsets of a sixteen-element pool, no survivors. $Y_1 \cong \mathbb{A}^4$? — 210 four-subsets of a ten-element pool, no survivors. All recorded as measurements, never as negative results. The third question is settled instead, and negatively, in §6, by three arguments that carry no cap at all.

6 The smooth middle identified

6.1 Three independent kills

Theorem 6.1. *Three independent computations bear on whether Y_1 is affine space. Routes (i) and (iii) each give $Y_1 \not\cong \mathbb{A}^4$, and (i) gives it for all stabilisations; route (ii) gives the separate statement recorded there.*

- (i) *Motivic. $[Y_1] = \mathbb{L}^4 + \mathbb{L}^2$ in the Grothendieck ring, by two independent stratifications (along t , and along q as a cross-check), every stratum carrying an explicit two-way parametrisation. Hence $\chi_c(Y_1) = 2$ against $\chi_c(\mathbb{A}^4) = 1$, and since χ_c is multiplicative with $\chi_c(\mathbb{A}^r) = 1$,*

$$Y_1 \times \mathbb{A}^r \not\cong \mathbb{A}^{4+r} \quad \text{for every } r \geq 0,$$

so the conclusion covers all stabilisations.

- (ii) *Differential. $\nabla W = (-1 - 2p, -Z, 3t^2H, -q, t^3)$ and $\text{Crit}(W)$ is the line $\{p = -\frac{1}{2}, q = Z = t = 0\}$, on which $W \equiv \frac{1}{4}$. A variable of $\mathbb{C}^{[5]}$ has nowhere-vanishing differential, and adjoining further variables leaves that line critical, so W is not a variable of $\mathbb{C}^{[5]}$, nor stably. The non-zero critical value is also exactly why $Y_1 = V(W)$ is smooth. We note explicitly that this does not by itself give $Y_1 \not\cong \mathbb{A}^4$: the passage from “ $V(f) \cong \mathbb{A}^{n-1}$ ” to “ f is a variable” is the Abhyankar–Sathaye problem, which we do not invoke. What (ii) rules out is that Y_1 arise as a coordinate hyperplane, stably included.*

- (iii) *Homotopical, and prior art. See Theorem 6.2 below: Y_1 is a member of the Asok–Dubouloz–Østvær family whose $\mathbb{A}^1$ -homotopy type is $\mathbb{P}^1 \wedge \mathbb{P}^1$, with complex realisation S^4 ; in particular Y_1 is not contractible.*

In (i), the cuspidal curve is certified once and reused: $(Z, t) = (s^3, -s^2)$ with inverse $s = -Z/t$ away from the cusp point, so $[\text{cusp}] = \mathbb{L}$ — with the recorded control that the curve is genuinely singular and not an $\mathbb{A}^1$ in disguise.

6.2 The identification

Theorem 6.2 ([1], applied). *Under the linear change of coordinates*

$$(x_1, t_1, x_2, t_2, z) = (t, H, q, -Z, p + \frac{1}{2}), \quad (13)$$

certified an isomorphism of $\mathbb{A}^5$ in both directions, the equation $x_1^3 t_1 + x_2 t_2 = z^2 - \frac{1}{4}$ of [1, Ex. 4.2.5] is W exactly. The polynomial $z^2 - \frac{1}{4}$ has distinct roots and unit discriminant, which is their smoothness hypothesis. Hence

$$Y_1 = Q_{(3,1), z^2 - 1/4},$$

and by [1] its $\mathbb{A}^1$ -homotopy type is $\mathbb{P}^1 \wedge \mathbb{P}^1$, with complex realisation S^4 .

The family $Q_{m,P} = \{\sum_i x_i^{m_i} t_i = P(z)\}$, its smoothness criterion, its $\mathbb{A}^1$ -homotopy type $(\mathbb{P}^{1 \wedge n})^{\vee(\deg P - 1)}$ and the open classification Question 4.2.6 — can non-isomorphic $Q_{m,P}$ be distinguished as m varies, for fixed P ? — are all [1]’s. We reprove none of them and we make no claim on the classification question.

Remark 6.3 (Prior art, stated plainly). Y_1 is prior art as a variety: it is a member of a published family, and its smoothness and homotopy type are published. What appears to be new is its role, namely that this known $Q_{m,P}$ -fourfold arises as the smooth intermediate completion in a commuting affine-modification square attached to the Koras–Russell cylinder. Accordingly the following are *not* claimed and are false: that this is a new fourfold of this form, a previously unknown deformed quadric, a new motivic-sphere model, or a first study of this class.

Combining: the square of §5 realises the chain

$$\mathbb{A}^4 \longleftarrow Y_1 \text{ (a motivic } S^4) \longleftarrow X \times \mathbb{A}^1 \text{ (}\mathbb{A}^1\text{-contractible, by [7, 11])}, \quad (14)$$

with each arrow an affine modification. The rest of the paper is about how the middle term’s sphere is created and then destroyed.

7 The motivic ledger

7.1 Four corners, four arrows

Proposition 7.1 (The corners).

	<i>class</i>	χ_c	Sing
$\mathbb{A}^4$	$\mathbb{L}^4$	1	$\emptyset$
$X_1 \times \mathbb{A}^1$	$\mathbb{L}^4$	1	<i>a line</i>
Y_1	$\mathbb{L}^4 + \mathbb{L}^2$	2	$\emptyset$
$X \times \mathbb{A}^1$	$\mathbb{L}^4$	1	$\emptyset$

Every entry is computed by explicit stratification with two-way parametrisations and an independent cross-check. The units and the class group of Y_1 were not determined and are not needed.

Proposition 7.2 (The arrows). *Every arrow is verified to be an isomorphism off its divisor, so the scissor relation $[V'] - [V] = [E] - [D]$ applies to all four, with D the divisor in the target and E the exceptional locus in the source:*

<i>arrow</i>	<i>along</i>	$[D]$	$[E]$	$[E] - [D]$
$\sigma_1: X_1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^4$	$\{p+1=0\}$	$\mathbb{L}^3$	$\mathbb{L}^3$	0
$\sigma_2: X \times \mathbb{A}^1 \rightarrow X_1 \times \mathbb{A}^1$	$\{t=0\}$	$\mathbb{L}^3$	$\mathbb{L}^3$	0
$\sigma_3: Y_1 \rightarrow \mathbb{A}^4$	$\{t=0\}$	$\mathbb{L}^3$	$\mathbb{L}^3 + \mathbb{L}^2$	$+\mathbb{L}^2$
$\sigma_4: X \times \mathbb{A}^1 \rightarrow Y_1$	$\{p+1=0\}$	$\mathbb{L}^3 + \mathbb{L}^2 - \mathbb{L}$	$\mathbb{L}^3 - \mathbb{L}$	$-\mathbb{L}^2$

So the singular route is K_0 -neutral at each step, while the smooth route creates an $\mathbb{L}^2$ packet and then removes it.

Throughout the rest of the paper, D_i denotes the divisor of σ_i in its target, E_i the exceptional locus of σ_i in its source, and C_i the centre of σ_i . In particular $D_4 = Y_1 \cap \{p+1=0\}$, a threefold, and $C_4 = \text{cusp} \times \mathbb{A}_H^1$.

7.2 Where the packet enters, and where it dies

Proposition 7.3 (Entry). *The surface Q appearing in σ_3 's exceptional locus has $[Q] = \mathbb{L}^2 + \mathbb{L}$, against $\mathbb{L}^2$ for a plane. The map $Q \rightarrow \mathbb{A}_q^1$ has fibre $\mathbb{A}^1$ over $q \neq 0$ but splits into the two disjoint lines $\{p = 0\}$ and $\{p = -1\}$ over $q = 0$. That extra line, times $\mathbb{A}_H^1$, is exactly the $+\mathbb{L}^2$ — the class-level shadow of the $(\deg P - 1) = 1$ wedge summand of $[1]$.*

Proposition 7.4 (Exit). *In shear coordinates, in which P becomes $x_s(1 - x_s H_s - 2w_s Z_s) + Z_s^2 + T_s^3$ and $p + 1$ becomes $1 - x_s H_s - 2w_s Z_s$, the exceptional locus of σ_4 is*

$$E_4 = \{x_s H_s + 2w_s Z_s = 1, \quad Z_s^2 + T_s^3 = 0\},$$

and on it the image coordinate q collapses to $-HZ$ — certified by ideal membership, with a boundary check confirming that this is no identity off E_4 . Consequently E_4 maps onto only the cusp locus $\{Z^2 + t^3 = 0, q = -HZ\}$ inside the threefold D_4 , missing a single point, and is an $\mathbb{A}^1$ -bundle over its image; the cross-check $[\text{image}] \cdot \mathbb{L} = (\mathbb{L}^2 - 1)\mathbb{L}$ holds.

Proposition 7.5 (Where the surgery is centred). *D_4 meets the $t = 0$ fibre of Y_1 in $\{p = -1, t = 0, qZ = 0\} \times \mathbb{A}_H^1$, whose $q = 0$ branch is the $p = -1$ line of the very pair whose doubling produced the packet in Proposition 7.3. The surgery is centred on one of the two components that created the class, and Proposition 5.6's (qZ^2) restriction is the same locus read with its multiplicity.*

7.3 Hodge–Deligne polynomials, and why they add nothing

Proposition 7.6 (A collapse, and it is the answer). *A twenty-one entry ledger — four corners, four divisors, four exceptional loci, four centres, the compressed centre, the crossing and its reduced restriction, the packet and the cusp — shows that every stratum occurring in the square is a product of affine spaces, tori and points, hence Hodge–Tate. Therefore $E(V)$ is just $[V]$ with $\mathbb{L} \mapsto uv$, verified entry by entry, and Hodge–Deligne polynomials carry nothing beyond the class level: the exact symbolic machinery tops out there. Two centres are classed here for the first time, $[C_2] = \mathbb{L}^2 + \mathbb{L}$ and $[C_4] = \mathbb{L}^2$.*

What the computation does sharpen is the type of the packet: $+\mathbb{L}^2 \mapsto (uv)^2$ is a Tate class of weight 4 and Hodge type $(2, 2)$ — exactly the reduced motive of $\mathbb{P}^1 \wedge \mathbb{P}^1$, which is what the homotopy type of Theorem 6.2 predicts. The class ledger and the homotopy statement agree at the only level at which both are visible. We record, as an observation and explicitly not as a law, that $[C_4] = \mathbb{L}^2$ equals the packet class; σ_1 's centre also has class $\mathbb{L}^2$ and removes nothing, so the coincidence is flagged and not interpreted.

7.4 The mechanism at Euler-characteristic level

Theorem 7.7. *Let U be the common open set of Y_1 and $X \times \mathbb{A}^1$, with $[U] = \mathbb{L}^4 - \mathbb{L}^3 + \mathbb{L}$. Then*

$$\chi_c(Y_1) = \chi_c(D_4) + \chi_c(U) = 1 + 1 = 2, \quad \chi_c(X \times \mathbb{A}^1) = \chi_c(E_4) + \chi_c(U) = 0 + 1 = 1,$$

so the entire Euler difference is $\chi_c(D_4) - \chi_c(E_4) = 1$. And $\chi_c(E_4) = 0$ rather than 1 for exactly one reason: E_4 is an $\mathbb{A}^1$ -bundle over $C_4 = \text{cusp} \times \mathbb{A}_H^1$ minus a single point, and is empty over that point,

$$(p, q, t, Z, H) = (-1, 0, 0, 0, 0),$$

which lies on the carrier of the packet. Over every other point of C_4 the fibre is verified to be an $\mathbb{A}^1$.

The load-bearing check is the counterfactual: were the exceptional locus the full $\mathbb{A}^1$ -bundle over C_4 , the target would have $\chi_c = 2$ rather than 1. *The missing point is necessary for contractibility to be possible.*

At this level of resolution one can go no further, and we said so at the time: χ_c and Grothendieck classes see Euler characteristics, not cycles. The reading suggested by the book-keeping — that the S^4 generator is carried by the packet, and that the exceptional locus missing that one point is what destroys it — was recorded as a candidate and asserted of nothing. §8 replaces it with cycle-level statements. It does *not* establish the causal reading: what it proves is that the generator is supported on the modification divisor, that the inverse image of its divisor representative is a punctured plane whose own fundamental class vanishes in the source, and that the defect point is that puncture. Deriving the vanishing *from* the defect would require the computation named in §8.4, which we do not carry out.

8 The generator, and the cycle that dies

In this section H_*^{BM} is integral Borel–Moore homology of the complex points. Every algebraic identity used is machine-certified; the topological steps are numbered lemmas over those identities.

8.1 Groundwork

Lemma 8.1. $\sigma_4: X \times \mathbb{A}^1 \rightarrow Y_1$ *is not proper.*

Proof. A proper affine morphism is finite, and a finite birational morphism onto a normal target is an isomorphism. Y_1 is smooth, hence normal; σ_4 is birational; and σ_4 is not an isomorphism, since the classes differ by Proposition 7.2. $\square$

So there is no Borel–Moore pushforward along σ_4 , and the comparison must go through the common open set. The organising choice of this section, and the reason it stays elementary, is this: *localise at the fibre over the origin, not at the singular modification centre.* The fibre is two disjoint smooth planes; the centre is singular. The singular centre never enters the argument below.

Let

$$\pi: Y_1 \longrightarrow \mathbb{A}_{t,q}^2 \tag{15}$$

be the projection, and write $Y_1^* = \pi^{-1}(\mathbb{A}^2 \setminus \{0\})$.

Lemma 8.2. *The fibre of π over a point with $t \neq 0$ or $q \neq 0$ is a plane, by the explicit graphs $H = (p + p^2 + qZ)/t^3$ and $Z = (t^3H - p - p^2)/q$ respectively. Over the origin the fibre degenerates to $p + p^2 = 0$, giving the doubled fibre with components*

$$F_0 = \{p = 0\}, \quad F_{-1} = \{p = -1\},$$

each isomorphic to $\mathbb{A}_{Z,H}^2$, disjoint, and reduced — the discriminant of $p + p^2$ being 1.

Moreover π is a smooth morphism of relative dimension two. In the fibre coordinates (p, Z, H) the relative differential of W is

$$\left(\frac{\partial W}{\partial p}, \frac{\partial W}{\partial Z}, \frac{\partial W}{\partial H} \right) = (-1 - 2p, -q, t^3),$$

which vanishes exactly on $\{p = -\frac{1}{2}, q = 0, t = 0\}$ with Z, H free; there $W \equiv \frac{1}{4} \neq 0$, so that locus is disjoint from $Y_1 = V(W)$. Hence the relative differential is nowhere zero on Y_1 , π is smooth, and in particular flat.

Lemma 8.3. $V_0 = Y_1 \setminus F_{-1}$ and $V_{-1} = Y_1 \setminus F_0$ form an open cover of Y_1 with $V_0 \cap V_{-1} = Y_1^*$; the two graphs of Lemma 8.2 trivialise Y_1^* over $\{t \neq 0\}$ and over $\{q \neq 0\}$, so Y_1^* is an $\mathbb{A}^2$ -bundle over $\mathbb{A}^2 \setminus \{0\}$ and hence $Y_1^* \simeq S^3$.

The involution $\iota: p \mapsto -1 - p$ is certified an automorphism of Y_1 that swaps the two components. Class-level consistency was checked throughout: $[V_0] = [V_{-1}] = \mathbb{L}^4$, with $\chi_c = 1$, consistent with the $\mathbb{A}^1$ -contractibility of the one-branch deletion in [1, Prop. 4.3.1]; and $[Y_1^*] = \mathbb{L}^4 - \mathbb{L}^2$, with $\chi_c = 0$, consistent with S^3 . The entire $\mathbb{L}^2$ packet sits off $\{t \neq 0\}$: the support datum is the origin fibre, precisely.

8.2 The generator

Theorem 8.4. $H_4^{\text{BM}}(Y_1; \mathbb{Z}) \cong \mathbb{Z}$, with $[F_0] \mapsto 1$ and $[F_{-1}] \mapsto -1$. Each component class is a generator; they satisfy $[F_0] = -[F_{-1}]$; and $[F_{-1}] - [F_0] \mapsto -2$ is twice a generator — of index two, and not a generator.

Proof, as certified. (1) $Y_1 \simeq S^4$ by Theorem 6.2, and $Y_1^* \simeq S^3$ by Lemma 8.3.

- (2) Poincaré duality on the smooth oriented 8-manifold gives $H_4^{\text{BM}}(Y_1) = \mathbb{Z}$, $H_4^{\text{BM}}(F_0 \sqcup F_{-1}) = \mathbb{Z}^2$ and $H_5^{\text{BM}}(Y_1^*) = \mathbb{Z}$.
- (3) π is smooth of relative dimension two with reduced fibre over the origin (Lemma 8.2), so the smooth Gysin map $\pi^!: H_k^{\text{BM}}(\mathbb{A}^2) \rightarrow H_{k+4}^{\text{BM}}(Y_1)$ is defined and carries the class of the origin to $[F_0] + [F_{-1}]$. Hence $[F_0] + [F_{-1}] = \pi^![\text{origin}] = 0$, because $H_0^{\text{BM}}(\mathbb{A}^2) = H^4(\mathbb{C}^2) = 0$. (Smoothness, rather than flatness alone, is what makes the Gysin map available here; see [9, §19.1] for the cycle map and the localisation sequence, and [3, §2.6] for Borel–Moore homology and its duality.)
- (4) The localisation sequence for the closed doubled fibre inside Y_1 with open complement Y_1^* reads $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{Z} \rightarrow 0$.
- (5) The kernel of $\mathbb{Z}^2 \rightarrow H_4^{\text{BM}}(Y_1)$ is primitive of rank one and contains $(1, 1)$ by (3), hence is $\langle (1, 1) \rangle$; so $H_4^{\text{BM}}(Y_1) = \mathbb{Z}^2 / \langle (1, 1) \rangle \cong \mathbb{Z}$ with the stated images.

□

Two independent cross-checks close, and neither is automatic. First, the sequence in (4) forces $H_5^{\text{BM}}(Y_1^*) \cong \mathbb{Z}$, while Poincaré duality gives $H^3(S^3) = \mathbb{Z}$ independently; had the fibre structure or the homotopy type been wrong, this would not have closed. Second, ι acts by -1 on $H_4^{\text{BM}}(Y_1)$ and squares to $+1$. A Mayer–Vietoris reading of the same cover gives the same answer and the same description of the generator, as the connecting class of $\mathbb{A}_{t,q}^2 \setminus \{0\}$.

Remark 8.5 (Robustness, and why the natural guess fails). Since ι swaps the components, the kernel in step (5) must be ι -stable, and the only primitive options are $\langle (1, 1) \rangle$ and $\langle (1, -1) \rangle$. Step (3) selects the first; under the second the difference $[F_{-1}] - [F_0]$ would be zero. So on either reading the difference class fails to generate, and the conclusion does not rest on step (3) alone. The reason the difference is the natural guess, and the reason it is wrong, are the same: a difference of branches is the usual shape, but here the two branches are already negatives of one another — because the fibre class vanishes — so the difference doubles instead of cancelling.

8.3 The cycle that dies

Theorem 8.6. *Let $D_4 \subset Y_1$ be the divisor of σ_4 and U the common open set.*

- (i) $[F_0] = -[F_{-1}]$ generates $H_4^{\text{BM}}(Y_1; \mathbb{Z}) \cong \mathbb{Z}$ (Theorem 8.4).
- (ii) F_0 is disjoint from D_4 , and F_{-1} lies entirely inside it. So the generator is the pushforward of the fundamental class of F_{-1} from inside the surgery locus — even though it also has a representative disjoint from that locus. That is the structural reason the surgery can reach it.
- (iii) Hence the generator restricts to zero on U .
- (iv) The inverse image of that representative is $\sigma_4^{-1}(F_{-1}) \cong \mathbb{G}_m \times \mathbb{A}^1$ — the same plane with the fibre over the defect point $(-1, 0, 0, 0, 0)$ deleted. It is a closed subvariety of $X \times \mathbb{A}^1$, and the pushforward of its fundamental class along that closed immersion is zero, because $H_4^{\text{BM}}(X \times \mathbb{A}^1) = 0$.

Remark 8.7 (What (iv) does *not* say). σ_4 is not proper (Lemma 8.1), so there is no Borel–Moore pushforward σ_{4*} and no sense in which the generator of $H_4^{\text{BM}}(Y_1)$ “pushes forward” to $X \times \mathbb{A}^1$. Statement (iv) is about the class of the locus $\sigma_4^{-1}(F_{-1})$ in its *own* ambient space, pushed forward along a closed immersion — which is proper, so the map exists. Nothing is transported along σ_4 in homology.

Remark 8.8 (The vanishing, minimally cited). $H_4^{\text{BM}}(X \times \mathbb{A}^1; \mathbb{Z}) = 0$ needs nothing motivic. $X \times \mathbb{A}^1$ is a smooth complex fourfold, so Poincaré duality for Borel–Moore homology [3, §2.6] gives $H_4^{\text{BM}}(X \times \mathbb{A}^1) \cong H^4(X \times \mathbb{A}^1; \mathbb{Z})$; and X is topologically contractible, hence so is $X \times \mathbb{A}^1$, so that cohomology group vanishes. That is the whole argument, and it is elementary.

The contractibility of X is classical, and is the only input here. It is recorded as well known in [5, §1], who observe that X is consequently diffeomorphic to $\mathbb{A}^3$, citing Dimca [4]. For context rather than as an input: X is $\mathbb{A}^1$ -contractible by Dubouloz–Fasel [7], extending Hoyois–Krishna–Østvær [11], whose theorem is stated for *finite suspensions* of Koras–Russell threefolds. Neither motivic statement is used anywhere in this paper.

Remark 8.9 (The functorial form, with no properness). There is a dual statement that does travel along σ_4 , because cohomological pullback is functorial for *every* morphism. Let $\text{PD}[F_{-1}] \in H^4(Y_1; \mathbb{Z})$ be the Poincaré dual of $[F_{-1}] \in H_4^{\text{BM}}(Y_1)$, available because Y_1 is a smooth oriented 8-manifold [3, §2.6]; the cycle class itself and the localisation sequence are as in [9, §19.1]. Then

$$\sigma_4^* \text{PD}[F_{-1}] = 0 \quad \text{in } H^4(X \times \mathbb{A}^1; \mathbb{Z}),$$

simply because $H^4(X \times \mathbb{A}^1; \mathbb{Z}) = 0$ by Remark 8.8. No properness hypothesis enters, and the generator of $H^4(Y_1; \mathbb{Z})$ — the dual of the generator of $H_4^{\text{BM}}(Y_1)$ — is what dies under pullback. This is the precise sense in which the modification kills the class, and it is the form we recommend a reader carry away.

The defect is completely explicit: $\sigma_4^{-1}(F_{-1})$ maps onto the *punctured line* $\{Z = 0, H \neq 0\}$ inside F_{-1} with $\mathbb{A}^1$ fibres; every point of F_{-1} with $Z \neq 0$ has *empty* preimage; and the puncture *is* the defect point of Theorem 7.7. The origin fibre’s H_3^{BM} jumps from 0 to $\mathbb{Z}$ — the loop in $\mathbb{G}_m$ that the defect creates.

Set against §7: Theorem 7.7 gave the Euler bookkeeping, that the whole difference is $\chi_c(D_4) - \chi_c(E_4) = 1$ and is caused by one missing point. Theorem 8.6 upgrades that to integral homology and names the cycle carrying it. The missing point is now the puncture of an explicit cycle rather than a numerical shadow.

8.4 The statement, and the gap inside it

The result of this section, stated once and completely, with its limitation inside it rather than after it:

$H_4^{\text{BM}}(Y_1; \mathbb{Z}) \cong \mathbb{Z}$ is generated by the fundamental class of either component of the doubled fibre over the origin of $\mathbb{A}_{t,q}^2$, with $[F_0] = -[F_{-1}]$. The generator is *supported on the modification divisor* — it has the representative F_{-1} contained in D_4 — and restricts to zero on the common open set. The inverse image of that representative is $\sigma_4^{-1}(F_{-1}) \cong \mathbb{G}_m \times \mathbb{A}^1$, the same plane with the fibre over the defect point $(-1, 0, 0, 0, 0)$ deleted, and the fundamental class of that locus, pushed forward along its closed immersion into $X \times \mathbb{A}^1$, is zero, since $H_4^{\text{BM}}(X \times \mathbb{A}^1) = 0$ by duality and topological contractibility. Dually, and with no properness hypothesis, $\sigma_4^* \text{PD}[F_{-1}] = 0$ in $H^4(X \times \mathbb{A}^1; \mathbb{Z})$. The exceptional defect thus accounts exactly for the Euler and Grothendieck-class change, and locates its cycle-level content at one named point. Two things are *not* asserted: no class is pushed forward along σ_4 , which is not proper; and deriving the vanishing from the modification geometry alone, rather than from the contractibility of $X \times \mathbb{A}^1$, would require the Borel–Moore homology of the modified punctured family, which we do not compute.

The reason for that last clause is worth stating precisely, because it is a real limitation and not a formality. The modification changes the fibre of π over *every* point of $\mathbb{A}_{t,q}^2$, not only over the origin: over $t \neq 0$ the plane loses its $p = -1$ line and gains two lines exchanged by the double cover $Z_s^2 = -t^3$; over $t = 0$ with $q \neq 0$ it simply loses that line. All three facts are certified. Hence the two localisation sequences do not differ only at the defect, and step (iv) leans on the cited contractibility. To make the implication run *from* the modification geometry *to* the vanishing one needs H_*^{BM} of the modified punctured family $(\pi \circ \sigma_4)^{-1}(\mathbb{A}^2 \setminus \{0\})$ computed independently: a monodromy computation over $\mathbb{A}^2 \setminus \{0\}$ with the double cover as its only non-elementary ingredient. It is well posed and elementary in kind; we have not carried it out.

9 What this does and does not say

It says that the published action on the Koras–Russell cylinder has a completely mapped quotient geometry; that the cylinder is an explicit canonical affine modification of $\mathbb{A}^4$ whose centre cannot be rectified to a smooth one; that the modification factors in two orders, through a singular threefold cylinder and through a smooth fourfold; that the smooth fourfold is a known model of the motivic 4-sphere and that its sphere is created by one step of the square and destroyed by the other; that at Euler-characteristic level the destruction reduces to a single missing point; and that at integral Borel–Moore level that point is the puncture of an explicit cycle representing the generator, whose Poincaré dual pulls back to zero.

It does not say anything about whether $X \times \mathbb{A}^1 \cong \mathbb{A}^4$ or $X \times \mathbb{A}^2 \cong \mathbb{A}^5$. Both remain exactly as open as they were. What is closed is one route: the flow-compatible route through this action, at $k = 1$ and at $k = 2$, by Theorems 3.14, 3.15 and 3.16, which carry no degree bound. The further elementary moves we looked for were sought *within declared, bounded pools* (Remarks 3.19 and 5.8); those searches found nothing, and that is a statement about the pools searched and not about the existence of such a move. The dissection results are properties of the action and do not separate $X \times \mathbb{A}^1$ from $\mathbb{A}^4$.

Positioned once more, and in the terms of §1.3: this is a concrete singular-defect instance of the established cofiber mechanism for affine modifications.

A Methods

A.1 The harness

This paper was written with substantial assistance from AI systems, in the role and under the disciplines described in this appendix. The author is solely responsible for the mathematical content and for every claim made. For the archival record, the systems used were: **Claude Fable 5** (`claude-fable-5`, Anthropic) as mission control; **Claude Opus 5** (`claude-opus-5`, Anthropic) as the executor of each bounded visit and as the author of the manuscript draft; and **GPT-5.6 Sol** (OpenAI) as an adversarial reviewer, in an advisory capacity only. No AI system is an author of this paper.

The work was produced by an AI research harness operating under a written protocol, and the protocol is part of the result’s warrant. A Claude Fable 5 session acted as mission control: it set the scope of each visit, ruled on findings, and never executed. Separate Claude Opus 5 sessions executed bounded visits against a written plan, with the human author relaying between them. Five such visits produced the content of this paper, one per section-group of §§3–8.

A.2 Certification discipline

Every algebraic identity in the paper is verified by machine, in exact arithmetic over $\mathbb{Q}$, by a runner that emits a ledger of individually named checks. The disciplines the work was held to, all of them adopted after they had cost something:

- **Twin implementations.** Load-bearing predicates are computed by two independent implementations which must agree; the agreement is itself a recorded check.
- **Positive controls.** Every search that can return “nothing found” also runs on an object where something *is* to be found. An empty verdict with no passing control is not recorded as a finding.
- **Boundary counter-checks.** For each certified identity, a nearby false variant is exhibited and shown to fail, so that a check which would pass vacuously is detected.
- **Declared caps.** Bounded searches declare their pool and their degree bounds *before* running, and their outcomes are recorded as measurements of what was searched, never as negative results. Every such statement in this paper is marked.
- **Frozen, replay-verified ledgers.** Ledgers are written read-only and re-generated in a fresh process in an isolated directory before the corresponding commit; a ledger that does not replay byte-identically is a defect. No wall-clock value may appear in a ledger, for that reason.
- **Machine first.** Where a hand expectation and a machine check disagreed, the machine was believed and the hand claim was logged rather than quietly reworded. This happened in four of the five visits, and the corrections are recorded in the write-ups accompanying the repository; two of them are visible in this paper as Remark 4.5 and Proposition 3.9.

A.3 Adversarial review

Four rounds of adversarial review were carried out by GPT-5.6 Sol (OpenAI) in an advisory capacity, each returned to the primary sources for verification before anything was adopted; the verification records, including the claims that were *not* confirmed and were carried as verify-on-use, are in the repository. The reviews contributed the prior-art demarcation that Remark 6.3 states, the terminology restrictions of Remark 2.3, the invariant battery used in Proposition 7.1, and the question that §8 answers. One substantive inference of the review’s — that the difference class $[F_{-1}] - [F_0]$ generates — was overturned by the computation, and Theorem 8.4 records the corrected statement; see Remark 8.5.

In the second review round the reviewer also *re-ran* part of the verification layer in its own sandbox — both standing gates and eight of the certificate runners — and obtained ledgers identical to the ones shipped here. That is a partial external replication of the computational claims, carried out independently of the author, and it is recorded as such.

A.4 Limitations of the method

The symbolic machinery is exact but it tops out at the class level, which is the content of Proposition 7.6: it sees Euler characteristics and Grothendieck classes, not cycles. Everything in §8 above the certified algebraic identities is human-and-model mathematics on top of those identities, in numbered lemmas that state which certified inputs they consume. The gap in §8.4 is stated inside the theorem for the same reason.

B The verification layer

The accompanying repository contains the runners, their ledgers, and a single entry point `verify.sh` which creates a fresh environment, runs the whole suite, and reports pass counts against the expected totals. The layer is organised by visit:

prefix	content	section here
<code>rd_*</code>	the action: content, kernel, plinth, slice, brackets	§3
<code>rg_*</code>	the boundary: diagnostics, tower, $k = 1$, $k = 2$	§§3–4
<code>rm_*</code>	the modification: canonical data, compression, crossing	§§4–5
<code>rs_*</code>	the ledger: invariant battery, arrows, E -polynomials	§§6–7
<code>rf_*</code>	the generator: groundwork, generator, death	§8
<code>pub_*</code>	the revision repairs: factoriality, smoothness of π	§§3, 8

together with the twin checkers and their cross-validation (`checker.py`, `checker_twin.py`, `cross_check.py`) and the Danielewski isomorphism certificate (`danielewski.py`) used as a standing environment gate. Running `verify.sh` re-runs all 22 runners in a scratch directory, replays each freshly produced ledger against the frozen certificate shipped with it, and checks the tallies; the expected totals are 883 checks with 713 PASS and 0 FAIL. `VERIFY.md` in the repository states what each check class means, how to read a ledger, and what a discrepancy indicates.

Data and code availability

All code, ledgers, and the source of this paper are in the repository accompanying it. Corrections and prior-art pointers are welcome; if any statement here has appeared earlier elsewhere, we will cite it prominently.

Acknowledgements

The starting datum of this paper — the conjugacy, the derivation, and the elements p, q, Z — is due to Adrien Dubouloz, Lucy Moser-Jauslin and Pierre-Marie Poloni [5], and I thank them for a paper whose explicit computations made everything here possible. The identification in §6 rests entirely on the family of Aravind Asok, Adrien Dubouloz and Paul Arne Østvær [1]. The final theorem’s ambient vanishing uses only the classical topological contractibility of X together with duality; I record with thanks the $\mathbb{A}^1$ -contractibility results of Adrien Dubouloz and Jean Fasel [7], extending Marc Hoyois, Amalendu Krishna and Paul Arne Østvær [11], which set the context for this circle of questions even though no statement of theirs is used here. I thank GPT-5.6 Sol (OpenAI) for four rounds of adversarial review that materially shaped the paper: the prior-art demarcation, the terminology discipline, the invariant battery, and the question that §8 answers. The work was carried out with Claude Fable 5 and Claude Opus 5 (Anthropic) as described in Appendix A. Errors that remain are mine.

References

- [1] A. Asok, A. Dubouloz and P. A. Østvær, *Geometric models for algebraic suspensions*, Int. Math. Res. Not. IMRN **2023**, no. 20, 17788–17821; arXiv:2112.08241.
- [2] J. Blanc and P.-M. Poloni, *Bivariabes and Vénéreau polynomials*, Ann. Fac. Sci. Toulouse Math. (6) **31** (2022), no. 5, 1391–1418; arXiv:2004.10739.
- [3] N. Chriss and V. Ginzburg, *Representation Theory and Complex Geometry*, Birkhäuser, Boston, 1997; §2.6 (Borel–Moore homology and Poincaré duality).
- [4] A. Dimca, *Hypersurfaces in $\mathbb{C}^n$ diffeomorphic to $\mathbb{R}^{4n-2}$* , Max-Planck-Institut für Mathematik preprint, Bonn, 1990; cited in this connection by [5].
- [5] A. Dubouloz, L. Moser-Jauslin and P.-M. Poloni, *Inequivalent embeddings of the Koras–Russell cubic 3-fold*, Michigan Math. J. **59** (2010), no. 3, 679–694; arXiv:0903.4278.
- [6] A. Dubouloz, *The cylinder over the Koras–Russell cubic threefold has a trivial Makar–Limanov invariant*, Transform. Groups **14** (2009), no. 3, 531–539; arXiv:0807.4085.
- [7] A. Dubouloz and J. Fasel, *Families of $\mathbb{A}^1$ -contractible affine threefolds*, Algebr. Geom. **5** (2018), no. 1, 1–14; arXiv:1512.01933.
- [8] A. Dubouloz, S. Pauli and P. A. Østvær, *$\mathbb{A}^1$ -contractibility of affine modifications*, Internat. J. Math. **30** (2019), no. 14, 1950069, 34 pp.; arXiv:1805.08959.
- [9] W. Fulton, *Intersection Theory*, 2nd ed., Ergeb. Math. Grenzgeb. (3) **2**, Springer, 1998; Ch. 19, §19.1 (the cycle map and Borel–Moore homology).

- [10] N. Gupta, *The Zariski cancellation problem and related problems in affine algebraic geometry*, arXiv:2208.14736.
- [11] M. Hoyois, A. Krishna and P. A. Østvær, $\mathbb{A}^1$ -contractibility of Koras–Russell threefolds, *Algebr. Geom.* **3** (2016), no. 4, 407–423; arXiv:1409.1293.
- [12] S. Kaliman and M. Zaidenberg, *Affine modifications and affine hypersurfaces with a very transitive automorphism group*, *Transform. Groups* **4** (1999), no. 1, 53–95; arXiv:math/9801076.
- [13] Sh. Kaliman, St. Vénéreau and M. Zaidenberg, *Simple birational extensions of the polynomial ring $\mathbb{C}^{[3]}$* , arXiv:math/0104204.
- [14] D. Lewis, *Strongly residual coordinates over $A[x]$* , arXiv:1304.1765.
- [15] L. Makar-Limanov, *On the hypersurface $x + x^2y + z^2 + t^3 = 0$ in $\mathbb{C}^4$ or a $\mathbb{C}^3$ -like threefold which is not $\mathbb{C}^3$* , *Israel J. Math.* **96** (1996), 419–429.
- [16] K. Masuda, *Faithfully flat quotient morphisms by $\mathbb{G}_a$ -actions on factorial affine varieties*, to appear in *J. Algebra*; arXiv:2512.06687.
- [17] K. Shakhmatov, *Cylinders and the zero locus of the plinth ideal*, arXiv:2605.00138.